Cell free miRNAs are pharmacodynamic biomarkers for enhanced Dicer activity by Enoxacin in
human patients with Amyotrophic lateral sclerosis

Iddo Magen[1,2], Hannah Marlene Kaneb[3], Maria Masnata[3], Nisha Pulimood[3], Anna Emde[1,2],
Angela Genge[3], Eran Hornstein[1,2]
1 Department of Molecular Genetics, Weizmann Institute of Science, Rehovot, Israel
2 Department of Molecular Neuroscience, Weizmann Institute of Science, Rehovot, Israel
3 Montreal Neurological Institute-Hospital, McGill University, Montreal, Canada.

Correspondence for molecular research:
Eran Hornstein, Arnold R. Meyer Building, Room 313, Weizmann Institute of Science, Rehovot
7610001, Israel, email: eran.hornstein@weizmann.ac.il
Correspondence for clinical research:
Angela Genge, Montreal Neurological Institute and Hospital, Department of Neurology, 3801
University Street, Suite 207, Montreal, QC, CA H3A 2B4, email: angela.genge@mcgill.ca

Running head: miRNAs as biomarkers for Enoxacin treatment in ALS
Number of words in abstract: 180
Number of words in main text: 5701 (excluding title page, abstract, references, figure
legends and acknowledgments)
Number of figures: 7
Number of tables: 2

ABSTRACT

Background and Objectives: The activity of the RNAse III DICER is downregulated in sporadic and genetic forms of ALS. Accordingly, hundreds of miRNAs are broadly downregulated, and their mRNA targets are de-repressed. Enoxacin is a fluoroquinolone, which increases DICER activity and miRNA biogenesis. Here, we tested for the first time the molecular effect of Enoxacin on miRNA biogenesis in patients with ALS and demonstrated that Enoxacin engagement with DICER can be pharmacodynamically monitored, via miRNA levels in human patients.

Methods: In an investigator-initiated, first-in-patient study, REALS1, we explored miRNAs as pharmacodynamic biomarkers for Dicer activation. Thus, patients with sporadic ALS were dosed with oral Enoxacin twice daily for 30 days in a double blind, randomized clinical trial.

Results: The study demonstrated a global increase in cell-free miRNA levels in both plasma and CSF at all time points following Enoxacin treatment (400 mg or 800 mg/day), compared to baseline. Additionally, no serious adverse events were reported among patients.

Conclusion: Enhancing DICER activity by Enoxacin, increases miRNA biogenesis in human patients with ALS. These results encourage testing Enoxacin efficacy in larger trials.

Keywords: Amyotrophic lateral sclerosis; Enoxacin; DICER; miRNAs; pharmacodynamic marker; target engagement.

INTRODUCTION

Amyotrophic lateral sclerosis (ALS) is a devastating neurodegenerative syndrome of the human motor neuron system[1]. Other than Tofersen therapy for ALS with superoxide dismutase 1 (SOD1) mutations[2], disease-modifying treatments do not exist. Studies over the last 20 years have uncovered ALS-causing mutations in a number of genes[3], many of which encode for RNA-binding proteins, e.g. TAR DNA-binding protein 43 (TDP-43), Heterogeneous ribonucleoprotein particle (HNRNP) A2/B1, HNRNP A1 and FUS . It is therefore hypothesized that dysregulation of RNA activity may be involved in the pathogenesis of ALS[4–8]. Accordingly, dysregulated splicing has been recently suggested to play an important role in pathogenesis because of the loss of normal TDP-43[9–12]. However, other facets of RNA dysregulation are not fully explored.

microRNAs (miRNAs) are endogenous non-protein-coding small RNAs. miRNAs silence messenger RNA (mRNA) expression by controlling translational inhibition and mRNA decay. The involvement of miRNAs in ALS pathogenesis is suggested by a global downregulation of miRNAs in motor neurons of patients with ALS[13–16].

DICER is an RNAse III, which works in concert with TDP-43 to promote miRNA biogenesis[17]. DICER cleaves the precursor miRNAs (pre-miRNAs), giving rise to single-stranded mature miRNAs[18]. The expression of ALS-causing mutant forms of TDP-43, FUS or SOD1, resulted in inhibition of DICER activity and accumulation of pre-miRNA species[13]. Consistently, spinal motor neuron-specific knockout of Dicer in adult mice resulted in progressive loss of motor neurons, denervation, and muscular atrophy[4].

While almost all miRNAs are dependent on DICER for biogenesis and function, miR-451a is Dicer-independent, as it is processed by Argonaute (Ago)[19]. Importantly, Dicer-dependent miRNAs compete with miR-451a in binding to the effector RNA induced silencing complex (RISC). Therefore, an increase in DICER activity results in relative upregulation of mature miRNAs and relative depletion of miR-451a[20].

Enoxacin is a fluoroquinolone, an antibiotic that was identified as an agonist of DICER complex activity, via increasing the binding of pre-miRNAs[21–23]. Enoxacin elevated miRNA levels in rat frontal cortex[24] and rescued expression of miRNAs that were downregulated by ALS-associated mutations in a cell line[13] or in induced pluripotent stem cell (iPSC)-derived motor neurons[25]. This suggests that the general upregulation of miRNA levels can be beneficial by regulating key genes that modulate pathological phenotypes. Furthermore, Enoxacin mitigated some of the symptoms associated with motor deterioration in the SOD1G93A and TDP-

43A315T ALS mouse models[13]. These preclinical data support the potential use of Enoxacin

as a novel disease-modifying drug and prompted us to initiate a study of Enoxacin in patients

with ALS.

Cell-free miRNAs are studied as potential biomarkers in neurodegeneration[25–57]. Accordingly,

we have recently shown that circulating miR-181 can serve as an ALS prognostication

biomarker in three patient cohorts [58,59], that miRNAs can predict the response to Spinraza

therapy in spinal muscular atrophy[60] and can assist the diagnosis of frontotemporal dementia[61].

Therefore, we hypothesized that cell-free miRNAs might also serve pharmacodynamic analysis

and could potentially reveal the engagement of Enoxacin onto its therapeutic target, DICER, in

humans.

Here, we report the findings from a study of orally administered Enoxacin in patients with ALS.

We found that Enoxacin globally increases miRNA levels in plasma and CSF and that the

magnitude of the molecular changes in miRNAs is associated with the concentration of

Enoxacin in plasma. Enoxacin was also safe and tolerable. Thus, our study provides means to

quantify target engagement by DICER agonists and encourages further studies to assess the

efficacy of Enoxacin in patients with ALS.

MATERIALS AND METHODS

Trial design and oversight

REALS1 (ClinicalTrials.gov identifier: NCT04840823) was a randomized, double-blind, parallel

groups, phase 1b/2a study of Enoxacin in patients with ALS. Enoxacin was dosed orally at a

400mg, 800mg or 1200mg per day, which is the range used in humans, and the trial was

limited to 30 days of dosing, as this period is known to be safe in humans[62, 63]. All enrolled

patients were actively dosed and patient's baseline miRNA measurement, prior to initiation of

therapy, was established on day 1 as a reference. No placebo group was included. The study

protocol was approved by the Research Ethics Board at the Clinical Research Unit at The

Neuro (Montreal Neurological Institute-Hospital (MNI, CRU / 2020-6359). The study enrolled

participants at MNI, Montreal, Quebec, Canada from April 2021 through March 2023. All

participants provided written informed consent, and experiments conformed to the principles

set out in the World Medical Association (WMA) Declaration of Helsinki and the Department of

Health and Human Services Belmont Report. Participants were evaluated and monitored in the

clinic on days 1, 7, 14, 21, and 30 of treatment and at a follow-up visit 14 days after the last

dose.

Sample size determination

This study was a small, early-phase study to assess dose range and design in preparation for
a large-scale future efficacy trial. There was no effect size available for calculation at the time
the protocol was written. 36 patients (12 per treatment arm), were considered adequate for the
initial assessment of safety, tolerability, pharmacokinetics (PK), and pharmacodynamics (PD)
of three doses of Enoxacin in adults with ALS. However, the number of recruited patients was
dramatically affected by restrictions due to COVID and the burden the pandemic imposed,
particularly on the participation of Canadian patients with ALS in a full 30-day study in Montreal
Neurological Institute.

Participant inclusion criteria

Participants were diagnosed with either familial or sporadic ALS and were on a stable dose of
Riluzole for at least 30 days prior to screening. Participants with a forced vital capacity (FVC)
lower than 50% of what is expected by age, height, weight, and sex were excluded. reported
hypersensitivity/allergy to fluoroquinolone was an exclusion criterion. Diagnosis of an
additional neurodegenerative disease, pulmonary disorder not attributed to ALS, severe renal
impairment or impaired liver function also resulted in exclusion. Patients currently enrolled in
another clinical trial involving an experimental drug or device were excluded as well.

Randomization and blinding

Patients were randomly allocated to one of the dosing groups and were administered with a
single morning dose and a single evening dose of Enoxacin for 30 days (Figures 1, 2). A single
morning dose (200mg, 400mg or 600mg) was administered on days 1 and 30 to determine
Enoxacin PK – maximal concentration (C_{max}) and time to reach maximal concentration (t_{max})
over 24 hours. The study statistician generated a random allocation sequence, enrolled
participants, and randomly assigned them to interventions as per the sequence. Block
randomization with block size of 3 was used to achieve balance and maintain the blind. The
study statistician and a delegated individual not involved in the day to day running of the study
were unblinded to the randomization code for the purposes of the generation and verification of
this code, however they had no contact with study participants and were unaware of the
identity of the study participants, whose treatment were assigned based on this allocation
sequence.

The sponsor, study participants, principal investigators, coordinators, clinical laboratory staff
and all other study site staff except for the study pharmacist(s), were blinded to participant

treatment dose assignment throughout the study. To preserve blinding, an identical placebo

tablet composed of excipient only was provided to patients assigned to doses that did not

require 3 'active' 200mg Enoxacin tablets to complete one dose. Thus, patients receiving

200mg Enoxacin x 2 / day took one 200mg Enoxacin tablet and two placebo tablets twice

daily, patients receiving 400mg x 2/day took two 200mg Enoxacin tablets and one placebo

tablet twice daily, and patients receiving 600mg x2/day took three Enoxacin tablets twice daily.

Enoxacin and placebo tablets were in-kind gifts from Apotex, Inc., Toronto, Canada.

Study endpoints

We sought unbiased analysis of miRNA levels in the plasma on days 1, 7, 14, 21 and 30 and

in the CSF on days 1 and 28. To allow for pharmacokinetic-pharmacodynamic (PK-PD)

relationship, PK analysis of Enoxacin levels was done in plasma on days 1, 7, 14, 21 and 30.

Safety and tolerability of Enoxacin included the following assessments: adverse events (AEs),

including AEs related to the drug, and serious AEs, i.e. AEs that affect patient's safety and

require medical intervention. Adverse events (AEs) were coded to preferred terms from the

MedDRA library (version 23.0). Additional safety assessments included physical examination;

body weight; 12-lead electrocardiogram (ECG) parameters; vital signs; laboratory safety

assessments; and FVC predicted percentage. The tolerability of Enoxacin was determined

based on rates of AEs and the discontinuation of the study drug due to drug-related AEs. An

illustration of the study design and its endpoints is seen in Figure 2. Secondary outcomes

included Amyotrophic Lateral Sclerosis Functional Rating Scale-Revised (ALSFRS-R) score at

baseline and at the end of the follow-up period on day 30 and PK analysis of Riluzole on days

1 and 30.

Biofluid collection

Phlebotomies were performed before Enoxacin morning dosing on days 1, 7 (+/- two days), 14

(+/- two days), 21 (+/- two days) and 30. After blood withdrawal, cells were pelleted by

centrifugation at 2000 RPM for 10 minutes at 4°C. The plasma was transferred to a separate

tube and stored at -80°C.

Lumbar puncture (LP) to collect Cerebrospinal fluid (CSF) was performed only for participants

who specifically consented to this procedure, before dosing on day 1, and two hours (+/-one

hour) post dosing on day 28. CSF samples were centrifuged at 2000 RPM for 10 minutes at

4°C to pellet cells and stored at -80°C.

Pharmacokinetic (PK) analysis

Trough plasma concentrations of Enoxacin (ng/ml) were measured before administration of the
morning dose on days 1 (before treatment onset), 7, 14, 21, and 30 to determine steady-state
levels. Plasma concentrations were also measured 1, 2, 4, 6, 8, and 24 hours after dosing on
days 1 and 30 to determine C_{max} and t_{max} after a single dose of Enoxacin. Enoxacin
concentrations were measured by liquid chromatography with tandem mass spectrometry (LC-
MS/MS) at SGS Life Science Services (Saint-Benoît, France). Plasma concentrations of
Riluzole (ng/ml) were measured 2, 4, 6 and 8 hours after Enoxacin dosing on days 1 and 30 to
determine potential inhibitory effect of Enoxacin on Riluzole metabolism, as reported
previously[64]. Riluzole was measured by Protein-precipitation extraction / HPLC with MS/MS at
Aliri Bioanalysis (Salt Lake City, Utah).

Biomarker assays
miRNA profiling. Total RNA was extracted from 0.5-1 ml of plasma or CSF using the
miRNeasy micro kit (Qiagen, Hilden, Germany) by an experimenter blinded to the identity of
samples. Baseline plasma and CSF samples, as well as post-treatment CSF samples, were
processed in technical duplicates, and the resulting data were averaged before analysis.
Repeated samples from the same individuals were processed in the same batch to avoid
confounding effects of batch-to-batch variation on differences between repeated samples.
Small RNA-seq libraries were prepared from 5μl total RNA using the QIAseq miRNA UDI
Library Kit and QIAseq miRNA 96 Index Kit IL UDI-A (Qiagen) in a blinded fashion. 3' and 5'
adapter ligation was followed by reverse transcription. Precise linear quantification of miRNA
was achieved by 12-nucleotide long unique molecular identifiers (UMIs) within the reverse
transcription primers[65]. cDNA libraries were amplified by 22 cycles (plasma) or 23 cycles (CSF)
of PCR, with two unique dual indices (UDIs) of 10 nucleotides each, followed by on-bead size
selection and cleaning. Library concentration was determined with a Qubit fluorometer (dsDNA
high sensitivity assay kit; Thermo Fisher Scientific, Waltham, MA) and library size with
Tapestation D1000 (Agilent). Libraries with different indices were multiplexed and sequenced
on Novaseq 6000 S1 flow cell (Illumina), with 72-bp single read and 10-bp read for index 1 and
index 2. Fastq files were de-multiplexed and human miRNAs were mapped using miRBase[66]
reference and the RNA-seq Analysis & Biomarker Discovery Portal (GeneGlobe, Qiagen).

Neurofilament light chain (NfL) analysis. NfL concentrations were measured in duplicates
by a single molecule array (Simoa, Quanterix, Boston, MA, USA), using a bead-conjugated
immunocomplex and NF-light Advantage Kit for HD-1/HD-X adjusted for SR-X (Uman

Diagnostics Umea, Sweden). The immunocomplex was applied to a multi-well array designed
to enable imaging of every single bead. A calibration curve constructed using serially diluted
bovine NfL standard (UmanDiagnostics) enabled precise quantification of the average number
of enzymes per bead per sample of plasma or CSF (diluted 1:4 and 1:100, respectively). The
experimenter was blinded to labels and analysis comprised a single batch of reagents.

Statistical analysis
After alignment of sequencing reads to the human genome and averaging data from technical
duplicates, only miRNAs with UMI counts >50 in at least 50% of plasma samples, or an
average UMI count >20 across all samples in the CSF were included in the analysis. To
reduce subject-to-subject variability, counts were batch-corrected using the ComBat function,
whereby each subject is considered a batch, as in our previous works[59,60]. Batch-corrected
counts were further divided, either by the corresponding levels of miR-451a, a dicer-
independent miRNA[19] to create a "DICER score" that takes into account two opposing effects
of dicer enhancement; or by a size factor generated by the DESeq2 algorithm which takes into
account library size geometric mean[67], under the assumption that miRNA counts followed a
negative binomial distribution. Total miRNA abundance was calculated by summing the
abundances of all miRNAs (whether divided by miR-451a or normalized by DESeq2).

Principal Component Analysis (PCA) was employed to reduce dataset dimensionality and to
identify data patterns that correspond to the largest variances. Data were standardized to have
a mean of zero and a standard deviation of one, ensuring that all variables contributed equally
to the analysis and then eigenvalues and eigenvectors of the covariance matrix were
calculated to determine the principal components. A transformation into a new coordinate
system facilitated the visualization of only the two principal components with the largest
variances being presented.

K-mean clustering was performed using R. Multiple miRNA expression was partitioned into
predefined distinct non-overlapping clusters by minimizing the sum of distances between the
data points and their respective cluster centroids. Data points consisted of each miRNA's
mean abundance and fold change post-treatment vs baseline. Data were first normalized to
prevent scale dependencies that could bias the distance calculations. Based on preliminary
analysis data were divided into k=2 distinct clusters.

Plasma NfL levels following treatment were analyzed with one-way repeated measures
ANOVA. Plasma abundance of multiple miRNA species following treatment was analyzed by

repeated measures two-way ANOVA wherein time point was included as a repeated factor and

miRNA species as an additional factor. Pairwise comparisons were performed between post-

treatment points (days 7, 14, 21 and 30) and baseline.

Comparison of plasma abundance of multiple miRNAs averaged over days 7, 14, 21 and 30 to

baseline abundance, as well as of CSF abundance of multiple miRNAs post-treatment to

baseline, was performed with paired t-tests for the different miRNA species, followed by

correction to multiple hypotheses.

Post-treatment plasma total miRNA abundance was analyzed with a non-parametric Friedman

test, followed by pairwise time point comparisons (post-treatment vs. baseline). The

significance of averaged plasma total miRNA abundance (days 7, 14, 21 and 30 averaged), as

well as CSF total miRNA abundance and NfL (post-treatment vs. baseline), were determined

with paired t-tests. Number of increasing vs decreasing miRNAs following Enoxacin was

compared with the one-sample sign test, under the null hypothesis that half of the miRNAs

increase and half of them decrease. Average fold change across all miRNAs was compared to

a theoretical mean of 1 using the one-sample t-test.

In the case of multiple hypotheses testing (i.e. multiple time points tested vs baseline or

multiple miRNAs tested between time points), p-values were corrected according to Benjamini

and Hochberg[68], or the two-stage linear step-up procedure of Benjamini, Krieger, and

Yekutieli[69]. The false discovery rate (FDR) was set at 0.1. Statistical analyses were performed

with R (version 4.2.0, R Foundation for Statistical Computing, Vienna, Austria) or GraphPad

Prism version 10.2.3 (GraphPad Software, Boston, Massachusetts USA). Graphs were

generated using GraphPad Prism version 10.2.3.

RESULTS

Enoxacin elevates cell-free miRNA levels in the plasma, demonstrating enhancement of
Dicer activity

A total of 13 patients living with ALS were screened for the study, of whom 8 who passed
screening were included and enrolled (Figure 1). Recruitment occurred between April 2021
and March 2023, and the last patient follow up was in April 2023. Participants' baseline
characteristics are presented in Table 1. One patient in the 600mg x2/day group dropped out
early from the study (see below), leaving only n=1 patient in this group who completed the
dosing period. We therefore performed in-depth analysis of data associated with the other two
doses. Due to the small number of observations in the 200mg x2/day and 400mg x2/day
groups (n=3 per dose), samples from the six patients were analyzed together. Because
Enoxacin is an agonist of the Dicer complex[21,23,25] and is expected to upregulate miRNA
biogenesis in tissues[13],   we hypothesized that it would elevate the levels of many miRNAs and
that accordingly next-generation sequencing of cell-free miRNAs could serve as a
pharmacodynamic (PD) marker of target (DICER) engagement. Plasma samples were
collected by phlebotomy at baseline before treatment initiation, and on days 7, 14, 21, and 30
before the morning dose. Small RNA libraries were prepared and sequenced on the Illumina
NovaSeq platform with a 72-bp single read, followed by mapping of gained sequences into the
human genome. As many as 2632 individual miRNA species were aligned to the human
genome (GRCh37/hg19) across all samples. We included in the downstream analysis only
miRNAs with UMI counts >50 in at least 50% of plasma samples (N=92 miRNAs). The
repeated sampling from subjects allowed us to estimate the mean shift in read counts
associated with each subject. We adjusted the original counts using the mean shift to reduce
the confounding effect of subject-to-subject variability. We analyzed the difference in
abundance of miRNAs between post-treatment time points and pre-treatment baseline.
Because miR-451a is a DICER independent miRNA[19], which competes with dozens of DICER-
dependent miRNAs on binding Argonaute proteins in the miRNA effector complex[20], the
enhancement of DICER activity results in upregulation of all miRNAs other than miR-451a that
is downregulated. Accordingly, we devised a DICER score that displays the ratio of all miRNA
counts, divided by the sample levels of miR-451a.

Of the 92 miRNAs included in the analysis, 31/92, 35/92, 43/92, 32/92 increased significantly
in their DICER scores on days 7, 14, 21, and 30 respectively (Figure 3A-D) and 59 miRNAs
increased significantly when DICER score was averaged on measurements performed under
therapy (Figure 3E). Importantly, no miRNAs decreased in a significant manner at either time

point compared to baseline. Furthermore, the average DICER score of 89 out of 92 miRNAs

increased relative to baseline (97% of miRNAs, one-sample sign test: $Z=9$, $p<0.0001$). The

mean effect size was a 2.5-fold increase in the DICER score (one-sample t-test, $p<0.0001$).

These results demonstrate that Enoxacin increases miRNA levels in human patients with ALS.

In order to reduce the dimensionality of the dataset, which is composed of multiple miRNAs,

we ran principal component analysis (PCA) on the DICER scores averaged on measurements

performed under therapy, (days 7, 14, 21 and 30) relative to the baseline DICER scores

(Figure 3F). Baseline samples and post-treatment samples were well-separated. Since the

post-treatment samples are better clustered than baseline samples, Enoxacin reduces

variability in DICER scores, and this might be due to enhancing dicer activity to the maximum.

Notable miRNAs that increased at all time points are the miR-320 family members (miR-320a,

miR-320b, miR-320c, and miR-320d), the muscle-enriched miR-206 and miR-378a and the

blood enriched miR-486-5p. We next grouped the 92 miRNAs into distinct groups, based on

similar mean abundance and fold change. This step simplifies data interpretation and

highlights miRNAs that consistently behave similarly under the effect of Enoxacin. Specifically,

miR-16, miR-423, and miR-486-5p clustered together, in combination with let-7b/a (Figure 4)

and may serve as potential biomarkers set for the molecular response to Enoxacin.

We also compared the total abundance of miRNAs on days 7, 14, 21, and 30 and at all post-

treatment time points averaged, to baseline. Total DICER score was calculated by summing

the DICER scores of all individual miRNAs. There was an overall increase in total DICER

score when each post-treatment time point was compared to the baseline (Friedman test:

$p<0.01$, Figure 5A). The highest increases in total DICER scores were observed on days 14

and 21: 3.4-fold and 3.1-fold, respectively, from 170±45 at baseline to 583±381 on day 14, and

526±226 on day 21 (q value = 0.007 and 0.004, respectively, after multiple hypothesis

correction). In addition, the total DICER score at all post-treatment time points averaged was

2.5-fold higher than baseline (422±17.5 vs 170±45, paired t-test: $p<0.0001$, Figure 5B). When

total miRNA abundance was calculated by summing the miRNA levels normalized by DESeq2,

the average abundance across all post-treatment time points increased relative to baseline

(paired t-test: $p=0.042$, Supp. Figure 1).

Taken together, Enoxacin elevates miRNA levels in plasma at all time points compared to

baseline; miR-451a levels are reduced, and a robust effect is monitored when we consider a

DICER score that divides levels of all miRNAs by miR-451a levels. The data suggest target

engagement of Enoxacin with Dicer in humans and corroborates pre-clinical works measuring

miRNAs in animal tissues.

Enoxacin elevates miRNA levels in the CSF

We next studied miRNA abundance in the CSF in samples collected from patients who

specifically consented to this procedure (n=4). CSF reflects the cerebral microenvironment and

thus can be a better approximation of the Enoxacin effect in the brain. Similar to plasma, miR-

451a significantly decreased by 40%, from 14.8±1.8 UMIs at baseline to 8.85±1.6 UMIs post-

treatment (Figure 6A, paired t-test: p=0.04). We then divided all miRNA levels by miR-451a to

create a DICER score, as was done in plasma. Total DICER score in the CSF significantly

increased by 70%, from 731±37 at baseline to 1259±86 post-treatment (Figure 6B, paired t-

test: p=0.018).

We further focused on 24 miRNAs that increased in plasma at all time points (source data

table 1) and were also detected in the CSF (average UMI count >20 across all samples). Eight

out of these 24 miRNAs were significantly elevated in the CSF: four members of the miR-320

family (miR-320a-d), let-7c, miR-2110, miR-378a and miR-423 (Figure 6C, adj. p value < 0.05).

miR-378a increased by 3.4-fold, compared to baseline.

We also measured NfL. Within the short period of therapy, NfL levels were not expected to

change in the plasma or CSF. Indeed, we did not reveal any significant differences at any time

point relative to baseline (Supp. Figure 2).

PK study of Enoxacin and Riluzole

We quantified the average Enoxacin plasma concentrations (before the morning dose,

Supplementary table 1) and revealed that the average C_{max} was reached within two hours of

administration. Enoxacin trough concentrations were rather constant from day 7 until the end

of drug dosing on day 30. Therefore, Enoxacin levels were at a steady state at the time points

in which pharmacodynamic markers were measured.

The concentrations of Riluzole, a common therapy for patients with ALS, were measured at 2h,

4h, 6h and 8h after Enoxacin administration on days 1 and 30. Higher Riluzole concentrations

on day 30 (Supplementary table 2), suggests an Enoxacin-Riluzole interaction, in agreement

with reported in vitro effects[64]. However, measured Riluzole levels did not exceed the C_{max}

reported to be safe, given a dose of 50mg BID twice daily[70,71]. This suggests Riluzole levels

are within the safe range, in agreement with lack of Riluzole-related adverse events.

PK-PD relationship

To study the association between plasma trough Enoxacin concentrations and miRNAs, we
performed PK-PD analysis. One data point was omitted due to a technical failure in measuring
Enoxacin. 13 miRNAs that were upregulated at least 2.5 fold on day 14, where the maximal
effect of Enoxacin on total miRNA abundance was observed (Figure 5A), were significantly
correlated with Enoxacin concentrations at this time point (Pearson correlations ≥ 0.89,
FDR<0.1, Figure 7). The association between Enoxacin levels and miRNA abundance
supports the interpretation that Enoxacin enhances miRNA biogenesis.

Safety and tolerability of Enoxacin in people living with ALS

All participants experienced adverse events (AEs; Supplementary table 3), but most of the AEs
reported were mild (36/52 events, 69%) or moderate (13/52, 25%). Severe AEs (3/52, 6%)
were not life-threatening, or serious (0/52). The three severe AEs were as follows: deep vein
thrombosis in one patient (dosed at 200mg x 2/day), probably not related to Enoxacin and
resolved after increasing the dose of an anticoagulant, apixaban, which the patient has been
taking regularly for at least six years. Nausea in another patient (dosed at 200mg x 2/day), was
resolved after taking dimenhydrinate (Dramamine). Dizziness in a third patient (dosed at
400mg x 2/day), decreased in severity during the next visit without medical intervention.

One patient in the 600mg x 2/day dosing group, which was excluded from analysis (see
methods) experienced mild or moderate nausea and dizziness that were assessed as related
to Enoxacin. These were defined as non-serious (Supplementary table 3), were not life
threatening, but were symptomatically severe enough to cause early discontinuation of
Enoxacin before completion of the dosing period.

AEs related to Enoxacin did not cause an early drop-out in any other patient and all
participants completed the dosing period and did not miss any dose of Enoxacin, with the
exception of a single patient that was treated with the highest dose of 600mg x 2/day. In
addition, throughout the dosing period none of the patients presented significant abnormalities
in ECG, vital signs (blood pressure and pulse rate), or blood tests. All the AEs seen were
previously observed in the original development of Enoxacin for antimicrobial indications. In
addition, although potential interaction of Enoxacin with Riluzole was suggested in vitro[64], none
of the AEs were deemed related to a possible change in Riluzole levels.

Taken together, enhancing DICER activity by Enoxacin increases miRNA biogenesis in human
patients with ALS, without any safety concerns.

DISCUSSION

In our study, we measured miRNAs as biomarkers and molecular endpoints for the effects of

Enoxacin. Because of the known effect of Enoxacin on DICER and its substrate, miRNAs, we

profiled miRNAs as pharmacodynamic and target engagement biomarkers. Most of the

miRNAs were increased as a result of Enoxacin treatment in both plasma and CSF, supporting

the engagement of Enoxacin with its target. The extent of the effect of Enoxacin on miRNAs.

i.e. the increase in their levels was correlated with the concentration of Enoxacin in the plasma,

which further supports target engagement.

A main question of our study was whether Enoxacin could have an effect in the CSF, a

potential reflection of the central nervous system as the target tissue. Previous works show

that the ratio of CSF to plasma levels is 33.3% in dogs[72], and 20% in rats[73,74], suggesting

substantial penetration of Enoxacin into the CSF from the blood. One study found a ratio of

29% in the CSF and 10% in the brains of rats[75]. In agreement with these studies, we observed

an effect of Enoxacin on miRNA levels in the CSF, and three of the miRNAs that were

increased in the CSF - let-7c, miR-16 and miR-320a - are abundant in both the human brain

and spinal cord (>1000 copies)[76]. Together, the findings suggest that Enoxacin can affect

miRNA biogenesis within the central nervous system when administered orally.

Some of the miRNAs that were increased by Enoxacin treatment were reported to be

neuroprotective, in the context of ALS or brain injuries. For instance, miR-139 was

downregulated in postmortem motor neurons of ALS patients by 8-fold[15], and in ALS iPSC-

derived motor neurons, leading to upregulation of the Wnt/β-catenin pathway which is a driver

of motor neuron degeneration[77]. Thus, restoring miR-139 levels in ALS may be beneficial to

motor neurons by regulating the Wnt/β-catenin pathway. Other miRNA that were increased by

Enoxacin were miR-130a, which displays neuroprotective activity after cerebral ischemia

upstream of the PTEN/PI3K/AKT pathway[78–80], Let-7c, miR-378a, miR-486 and miR-93 that

were reported to regulate microglial activation, suppress neuroinflammation and inhibit

neuronal apoptosis in cerebral ischemia and traumatic brain injury[81–85] and miR-30e that

regulated neuroinflammation in a Parkinson's disease model[86].

Some of the miRNAs that were increased by Enoxacin were reduced in biofluids and tissues

from ALS patients or ALS mouse models, suggesting that Enoxacin recovers the expression of

these miRNAs. For example, miR-378a levels are reduced in CSF from ALS patients[57] and

were increased in the CSF following Enoxacin in our study. miR-320 levels are reduced in the

serum or CSF of patients with ALS[46] or frontotemporal dementia (FTD) [87,88] and in the spinal

cord and brainstem of SOD1 G93A mice[89] and are increased in response to Enoxacin therapy.

While the safety profile of Enoxacin is known and this drug has been in clinical use for many
years, we also examined for the first time the safety and tolerability of Enoxacin in patients with
ALS. At 600mg x2/day one patient dropped out before study completion, due to AEs that were
deemed related to the drug, but these AEs were not life-threatening and no other drug-related
AEs were observed in other patients.

We did not expect any effect on NfL levels in our trial, as even in the VALOR trial for Tofersen
therapy in SOD1-ALS, NfL responded to therapy only after 12 weeks[2]. Indeed, a single month
of treatment was insufficient to demonstrate beneficial effects of Enoxacin on disease
trajectory. The single month study encourages a longer study of the efficacy of Enoxacin
therapy.

Limitations
The small cohort size hindered the determination of whether the effect of Enoxacin on miRNAs
was dose-dependent. The COVID pandemic is the main reason for the relatively low enrolment
in 2021 and 2022 and the expiry of the experimental drug batch prevented the extension of the
trial further.

Unfortunately, no female patients were enrolled in the study; therefore, we do not know
whether there are sex-associated differences in the response to Enoxacin. Finally, the study
did not explore efficacy as an endpoint due to the short therapy duration and the small number
of patients.

In summary, Enoxacin presents a promising therapeutic option for ALS with an established
safety profile, robust mechanistic data, and evidence for target engagement in both human
patient plasma and CSF. The ready-to-use pharmacodynamic biomarker package enhances
its appeal as a testable compelling addition to the ALS treatment landscape. The recent
approval of Tofersen as a therapy for ALS by the FDA, based on its ability to lower blood levels
of NfL[2] highlights the importance of biomarkers as an endpoint for clinical trials.

Declarations

Ethics approval and consent to participate

The study protocol was approved by the Research Ethics Board at the Clinical Research Unit

at The Neuro (Montreal Neurological Institute-Hospital (MNI, CRU / 2020-6359). All

participants provided written informed consent, and experiments conformed to the principles

set out in the World Medical Association (WMA) Declaration of Helsinki and the Department of

Health and Human Services Belmont Report.

Availability of data and materials

Data is available upon reasonable request from the author.

Competing Interests

EH and AE have a patent titled: "Methods of diagnosing and treating motor neuron diseases

and other cellular stress-related diseases" (US patent US10159670B2) which is related to this

work. All other authors declare no competing interests.

Funding

This study was funded by ALS Association clinical trial award (program: 'Repurposing

Enoxacin therapy for patients with ALS' 22-CTA-614), ALS Canada-Brain Canada Discovery

grant (program: 'Advanced Pharmacokinetics and Pharmacodynamics for phase Ib/IIa trial of

repurposed Enoxacin therapy for patients with ALS'), ERA-Net for Research Programs on

Rare Diseases (eRARE FP7) via the Israel Ministry of Health and Muscular Dystrophy

Canada, (CIHR) and Fonds de recherche du Québec - Santé (FRQS). IM was supported by

Teva Pharmaceutical Industries as part of the Israeli National Network of Excellence in

Neuroscience (fellowship no. 117941).

Research at the Hornstein lab is further funded by Andi and Larry Wolfe Centre for

Neuroimmunology and Neuromodulation, the Binational Science Foundation (BSF);

Association Française Contre les Myopathies (AFM) grants 24882, 28680; Muscular Dystrophy

Association (MDA) grant 1280000; Target ALS; Israel Science Foundation (ISF 3497/21,

424/22); ALS Canada; Minna-James-Heineman Stiftung through Minerva, Minerva Foundation,

with funding from the Federal German Ministry for Education and Research; Robert Packard

Center for ALS Research at Johns Hopkins; McGill University; EU - ERA-Net; Radala

Foundation for ALS Research; Additional support generously provided by the Kekst Family

Institute for Medical Genetics. Weizmann SABRA - Yeda-Sela - WRC Program, the Estate of

Emile Mimran, and The Maurice and Vivienne Wohl Biology Endowment. Nella and Leon

Authors' Contributions

486 IM, AE, AG and EH contributed to the conception and design of the study; IM, HMK, NP, MM,
487 AG and EH contributed to the acquisition and analysis of data; IM, AG and EH contributed to
488 drafting the text or preparing the figures.

Acknowledgements

EH is the Mondry Family Professorial Chair and Head of the Nella and Leon Benoziyo Centre
for Neurological Diseases and of the Andi and Larry Wolfe Centre for Neuroimmunology and
Neuromodulation. We thank Orla Hardiman (Trinity College Dublin), Dame Pamela Shaw
(University of Sheffield), and Leonard van den Berg (University Medical Center Utrecht) for
their advice on the study design. We are grateful to Susan Stern, National Executive Director
and CEO of Weizmann Canada, and Prof. Erik Storkebaum (Radboud University, Nijmegen)
for their involvement. Enoxacin was an in-kind gift from the late Barry Sherman, Alexandra
Krawczyk, the Honey & Barry Sherman Legacy Foundation and Apotex Inc., Toronto, Canada.
We thank Drs. Brigitte Happ, Kostas Kaloulis, Sonia Poli and Dawn Toronto for their
consultation on drug development and regulatory affairs. We thank Yahel Cohen and Dr.
Nancy S Yacovzda (WIS) for advice and for comments on the manuscript.

Abbreviations

Adverse event, AE

ALS functional rating score revised, ALSFRS-R

Amyotrophic lateral sclerosis, ALS

Argonaute, Ago

Cerebrospinal fluid, CSF

Electrocardiogram, ECG

False discovery rate, FDR

Frontotemporal dementia, FTD

Forced vital capacity, FVC

Heterogeneous ribonucleoprotein particle, HNRNP

Induced pluripotent stem cell, iPSC

Lumbar puncture, LP

microRNA, miRNA

messenger RNA, mRNA

Neurofilament light chain, NfL

Principal component analysis, PCA

Pharmacodynamic(s), PD

Pharmacokinetic(s), PK

precursor miRNAs, pre-miRNA

RNA induced silencing complex, RISC

TAR DNA binding protein 43, TDP-43

Unique dual index, UDI

Unique molecular identifier, UMI

REFERENCES

1. Tsai, M.-J., Hsu, C.-Y. & Sheu, C.-C. Amyotrophic Lateral Sclerosis. *N. Engl. J. Med.* **377**, 1602 (2017).

2. Miller, T. M. *et al.* Trial of Antisense Oligonucleotide Tofersen for SOD1 ALS. *N. Engl. J. Med.* **387**, 1099–1110 (2022).

3. Al-Chalabi, A., van den Berg, L. H. & Veldink, J. Gene discovery in amyotrophic lateral sclerosis: implications for clinical management. *Nat. Rev. Neurol.* **13**, 96–104 (2017).

4. Haramati, S. *et al.* miRNA malfunction causes spinal motor neuron disease. *Proc. Natl. Acad. Sci. U. S. A.* **107**, 13111–13116 (2010).

5. Lagier-Tourenne, C., Polymenidou, M. & Cleveland, D. W. TDP-43 and FUS/TLS: emerging roles in RNA processing and neurodegeneration. *Hum. Mol. Genet.* **19**, R46–64 (2010).

6. Lagier-Tourenne, C. & Cleveland, D. W. Rethinking ALS: the FUS about TDP-43. *Cell* **136**, 1001–1004 (2009).

7. Lemmens, R., Moore, M. J., Al-Chalabi, A., Brown, R. H., Jr & Robberecht, W. RNA metabolism and the pathogenesis of motor neuron diseases. *Trends Neurosci.* **33**, 249–258 (2010).

8. Strong, M. J. The evidence for altered RNA metabolism in amyotrophic lateral sclerosis (ALS). *J. Neurol. Sci.* **288**, 1–12 (2010).

9. Irwin, K. E. *et al.* A fluid biomarker reveals loss of TDP-43 splicing repression in presymptomatic ALS-FTD. *Nat. Med.* **30**, 382–393 (2024).

10. Keuss, M. J. *et al.* Loss of TDP-43 induces synaptic dysfunction that is rescued by UNC13A splice-switching ASOs. *bioRxivorg* (2024) doi:10.1101/2024.06.20.599684.

11. Costantino, I., Meng, A. & Ravits, J. Pathologic changes in ELAVL3 expression and splicing occur in TDP43 and non-TDP43 proteinopathies (P4-9.007). *Neurology* **102**, (2024).

12. Cao, M. C. *et al.* A panel of TDP-43-regulated splicing events verifies loss of TDP-43 function in amyotrophic lateral sclerosis brain tissue. *Neurobiol. Dis.* **185**, 106245 (2023).

13. Emde, A. *et al.* Dysregulated miRNA biogenesis downstream of cellular stress and ALS-causing mutations: a new mechanism for ALS. *EMBO J.* **34**, 2633–2651 (2015).

14. Campos-Melo, D., Droppelmann, C. A., He, Z., Volkening, K. & Strong, M. J. Altered microRNA expression profile in Amyotrophic Lateral Sclerosis: a role in the regulation of NFL mRNA levels. *Mol. Brain* **6**, 26 (2013).

15. Figueroa-Romero, C. *et al.* Expression of microRNAs in human post-mortem amyotrophic lateral sclerosis spinal cords provides insight into disease mechanisms. *Mol. Cell. Neurosci.* **71**, 34–45 (2016).

16. Reichenstein, I. *et al.* Human genetics and neuropathology suggest a link between miR-218 and amyotrophic lateral sclerosis pathophysiology. *Sci. Transl. Med.* **11**, (2019).

17. Kawahara, Y. & Mieda-Sato, A. TDP-43 promotes microRNA biogenesis as a component of the Drosha and Dicer complexes. *Proc. Natl. Acad. Sci. U. S. A.* **109**, 3347–3352 (2012).

18. Bernstein, E., Caudy, A. A., Hammond, S. M. & Hannon, G. J. Role for a bidentate ribonuclease in the initiation step of RNA interference. *Nature* **409**, 363–366 (2001).

19. Yang, J.-S. *et al.* Conserved vertebrate mir-451 provides a platform for Dicer-independent, Ago2-mediated microRNA biogenesis. *Proc. Natl. Acad. Sci. U. S. A.* **107**, 15163–15168 (2010).

20. Kretov, D. A. *et al.* Ago2-Dependent Processing Allows miR-451 to Evade the Global MicroRNA Turnover Elicited during Erythropoiesis. *Mol. Cell* **78**, 317–328.e6 (2020).

21. Melo, S. *et al.* Small molecule enoxacin is a cancer-specific growth inhibitor that acts by enhancing TAR RNA-binding protein 2-mediated microRNA processing. *Proc. Natl. Acad. Sci. U. S. A.* **108**, 4394–4399 (2011).

22. Schaeffer, A. J. The expanding role of fluoroquinolones. *Am. J. Med.* **113 Suppl 1A**, 45S–54S (2002).

23. Shan, G. *et al.* A small molecule enhances RNA interference and promotes microRNA processing. *Nat. Biotechnol.* **26**, 933–940 (2008).

24. Smalheiser, N. R., Zhang, H. & Dwivedi, Y. Enoxacin Elevates MicroRNA Levels in Rat Frontal Cortex and Prevents Learned Helplessness. *Front. Psychiatry* **5**, 6 (2014).

25. Rizzuti, M. *et al.* MicroRNA expression analysis identifies a subset of downregulated

miRNAs in ALS motor neuron progenitors. *Sci. Rep.* **8**, 10105 (2018).

26. Arakawa, Y. *et al.* Association between oxidative stress and microRNA expression pattern of ALS patients in the high-incidence area of the Kii Peninsula. *Brain Res.* **1746**, 147035 (2020).

27. Banack, S. A., Dunlop, R. A. & Cox, P. A. An miRNA fingerprint using neural-enriched extracellular vesicles from blood plasma: towards a biomarker for amyotrophic lateral sclerosis/motor neuron disease. *Open Biol.* **10**, 200116 (2020).

28. Benigni, M. *et al.* Identification of miRNAs as Potential Biomarkers in Cerebrospinal Fluid from Amyotrophic Lateral Sclerosis Patients. *Neuromolecular Med.* **18**, 551–560 (2016).

29. Cloutier, F., Marrero, A., O'Connell, C. & Morin, P., Jr. MicroRNAs as potential circulating biomarkers for amyotrophic lateral sclerosis. *J. Mol. Neurosci.* **56**, 102–112 (2015).

30. Daneshafrooz, N. *et al.* Identification of let-7f and miR-338 as plasma-based biomarkers for sporadic amyotrophic lateral sclerosis using meta-analysis and empirical validation. *Sci. Rep.* **12**, 1373 (2022).

31. de Andrade, H. M. T. *et al.* MicroRNAs-424 and 206 are potential prognostic markers in spinal onset amyotrophic lateral sclerosis. *J. Neurol. Sci.* **368**, 19–24 (2016).

32. De Felice, B. *et al.* miR-338-3p is over-expressed in blood, CFS, serum and spinal cord from sporadic amyotrophic lateral sclerosis patients. *Neurogenetics* **15**, 243–253 (2014).

33. De Luna, N. *et al.* Downregulation of miR-335-5P in Amyotrophic Lateral Sclerosis Can Contribute to Neuronal Mitochondrial Dysfunction and Apoptosis. *Sci. Rep.* **10**, 4308 (2020).

34. Dobrowolny, G. *et al.* A longitudinal study defined circulating microRNAs as reliable biomarkers for disease prognosis and progression in ALS human patients. *Cell Death Discov* **7**, 4 (2021).

35. Freischmidt, A. *et al.* Serum microRNAs in sporadic amyotrophic lateral sclerosis. *Neurobiol. Aging* **36**, 2660.e15–20 (2015).

36. Giagnorio, E. *et al.* MiR-146a in ALS: Contribution to Early Peripheral Nerve Degeneration and Relevance as Disease Biomarker. *Int. J. Mol. Sci.* **24**, (2023).

37. Joilin, G., Leigh, P. N., Newbury, S. F. & Hafezparast, M. An Overview of MicroRNAs as Biomarkers of ALS. *Front. Neurol.* **10**, 186 (2019).

38. Joilin, G. *et al.* Identification of a potential non-coding RNA biomarker signature for amyotrophic lateral sclerosis. *Brain Commun* **2**, fcaa053 (2020).

39. Katsu, M. *et al.* MicroRNA expression profiles of neuron-derived extracellular vesicles in plasma from patients with amyotrophic lateral sclerosis. *Neurosci. Lett.* **708**, 134176 (2019).

40. Liguori, M. *et al.* Dysregulation of MicroRNAs and Target Genes Networks in Peripheral Blood of Patients With Sporadic Amyotrophic Lateral Sclerosis. *Front. Mol. Neurosci.* **11**, 288 (2018).

41. Lo, T.-W. *et al.* Extracellular Vesicles in Serum and Central Nervous System Tissues Contain microRNA Signatures in Sporadic Amyotrophic Lateral Sclerosis. *Front. Mol. Neurosci.* **14**, 739016 (2021).

42. Malacarne, C. *et al.* Dysregulation of Muscle-Specific MicroRNAs as Common Pathogenic Feature Associated with Muscle Atrophy in ALS, SMA and SBMA: Evidence from Animal Models and Human Patients. *Int. J. Mol. Sci.* **22**, (2021).

43. Matamala, J. M. *et al.* Genome-wide circulating microRNA expression profiling reveals potential biomarkers for amyotrophic lateral sclerosis. *Neurobiol. Aging* **64**, 123–138 (2018).

44. Panio, A., Cava, C., D'Antona, S., Bertoli, G. & Porro, D. Diagnostic Circulating miRNAs in Sporadic Amyotrophic Lateral Sclerosis. *Front. Med.* **9**, 861960 (2022).

45. Pregnolato, F. *et al.* Exosome microRNAs in Amyotrophic Lateral Sclerosis: A Pilot Study. *Biomolecules* **11**, (2021).

46. Raheja, R. *et al.* Correlating serum micrornas and clinical parameters in amyotrophic lateral sclerosis. *Muscle Nerve* **58**, 261–269 (2018).

47. Ravnik-Glavač, M. & Glavač, D. Circulating RNAs as Potential Biomarkers in Amyotrophic Lateral Sclerosis. *Int. J. Mol. Sci.* **21**, (2020).

48. Ricci, C., Marzocchi, C. & Battistini, S. MicroRNAs as Biomarkers in Amyotrophic Lateral Sclerosis. *Cells* **7**, (2018).

49. Saucier, D. *et al.* Identification of a circulating miRNA signature in extracellular vesicles collected from amyotrophic lateral sclerosis patients. *Brain Res.* **1708**, 100–108 (2019).

50. Sheinerman, K. S. *et al.* Circulating brain-enriched microRNAs as novel biomarkers for detection and differentiation of neurodegenerative diseases. *Alzheimers. Res. Ther.* **9**, 89

(2017).

51. Soliman, R. *et al.* Assessment of diagnostic potential of some circulating microRNAs in Amyotrophic Lateral Sclerosis Patients, an Egyptian study. *Clin. Neurol. Neurosurg.* **208**, 106883 (2021).

52. Sproviero, D. *et al.* Different miRNA Profiles in Plasma Derived Small and Large Extracellular Vesicles from Patients with Neurodegenerative Diseases. *Int. J. Mol. Sci.* **22**, (2021).

53. Takahashi, I. *et al.* Identification of plasma microRNAs as a biomarker of sporadic Amyotrophic Lateral Sclerosis. *Mol. Brain* **8**, 67 (2015).

54. Tasca, E., Pegoraro, V., Merico, A. & Angelini, C. Circulating microRNAs as biomarkers of muscle differentiation and atrophy in ALS. *Clin. Neuropathol.* **35**, 22–30 (2016).

55. Toivonen, J. M. *et al.* MicroRNA-206: a potential circulating biomarker candidate for amyotrophic lateral sclerosis. *PLoS One* **9**, e89065 (2014).

56. Waller, R. *et al.* Serum miRNAs miR-206, 143-3p and 374b-5p as potential biomarkers for amyotrophic lateral sclerosis (ALS). *Neurobiol. Aging* **55**, 123–131 (2017).

57. Waller, R. *et al.* Small RNA Sequencing of Sporadic Amyotrophic Lateral Sclerosis Cerebrospinal Fluid Reveals Differentially Expressed miRNAs Related to Neural and Glial Activity. *Front. Neurosci.* **11**, 731 (2017).

58. Benatar, M. *et al.* Prognostic clinical and biological markers for amyotrophic lateral sclerosis disease progression: validation and implications for clinical trial design and analysis. *EBioMedicine* **108**, 105323 (2024).

59. Magen, I. *et al.* Circulating miR-181 is a prognostic biomarker for amyotrophic lateral sclerosis. *Nat. Neurosci.* **24**, 1534–1541 (2021).

60. Magen, I. *et al.* Muscle microRNAs in the cerebrospinal fluid predict clinical response to nusinersen therapy in type II and type III spinal muscular atrophy patients. *Eur. J. Neurol.* **29**, 2420–2430 (2022).

61. Magen, I. *et al.* microRNA-based predictor for diagnosis of frontotemporal dementia. *Neuropathol. Appl. Neurobiol.* **49**, e12916 (2023).

62. Scholar, E. Enoxacin. in *xPharm: The Comprehensive Pharmacology Reference* 1–6

(Elsevier, 2007).

63. Enoxacin Dosage. *Drugs.com* https://www.drugs.com/dosage/enoxacin.html.

64. Sanderink, G. J., Bournique, B., Stevens, J., Petry, M. & Martinet, M. Involvement of human CYP1A isoenzymes in the metabolism and drug interactions of riluzole in vitro. *J. Pharmacol. Exp. Ther.* **282**, 1465–1472 (1997).

65. Coenen-Stass, A. M. L. *et al.* Evaluation of methodologies for microRNA biomarker detection by next generation sequencing. *RNA Biol.* **15**, 1133–1145 (2018).

66. Kozomara, A. & Griffiths-Jones, S. miRBase: annotating high confidence microRNAs using deep sequencing data. *Nucleic Acids Res.* **42**, D68–73 (2014).

67. Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biol.* **15**, 550 (2014).

68. Benjamini, Y., Drai, D., Elmer, G., Kafkafi, N. & Golani, I. Controlling the false discovery rate in behavior genetics research. *Behav. Brain Res.* **125**, 279–284 (2001).

69. Benjamini, Y., Krieger, A. M. & Yekutieli, D. Adaptive linear step-up procedures that control the false discovery rate. *Biometrika* **93**, 491–507 (2006).

70. Le Liboux, A. *et al.* Single- and multiple-dose pharmacokinetics of riluzole in white subjects. *J. Clin. Pharmacol.* **37**, 820–827 (1997).

71. Wymer, J., Apple, S., Harrison, A. & Hill, B. A. Pharmacokinetics, Bioavailability, and Swallowing Safety With Riluzole Oral Film. *Clin Pharmacol Drug Dev* **12**, 57–64 (2023).

72. Tran Van Tho, Armengaud, A. & Davet, B. Diffusion of enoxacin into the cerebrospinal fluid in dogs with healthy meninges and with experimental meningitis. *J. Antimicrob. Chemother.* **14 Suppl C**, 57–62 (1984).

73. Delon, A. *et al.* Pharmacokinetic-pharmacodynamic contributions to the convulsant activity of fluoroquinolones in rats. *Antimicrob. Agents Chemother.* **43**, 1511–1515 (1999).

74. Marchand, S., Pariat, C., Boulanger, A., Bouquet, S. & Couet, W. A pharmacokinetic/pharmacodynamic approach to show that not all fluoroquinolones exhibit similar sensitivity toward the proconvulsant effect of biphenyl acetic acid in rats. *J. Antimicrob. Chemother.* **48**, 813–820 (2001).

75. Kawakami, J., Ohashi, K., Yamamoto, K., Sawada, Y. & Iga, T. Effect of acute renal failure

on neurotoxicity of enoxacin in rats. *Biol. Pharm. Bull.* **20**, 931–934 (1997).

76. Keller, A. *et al.* miRNATissueAtlas2: an update to the human miRNA tissue atlas. *Nucleic Acids Res.* **50**, D211–D221 (2022).

77. Hawkins, S. *et al.* Upregulation of β-catenin due to loss of miR-139 contributes to motor neuron death in amyotrophic lateral sclerosis. *Stem Cell Reports* **17**, 1650–1665 (2022).

78. Wang, Y. *et al.* miR-130a alleviates neuronal apoptosis and changes in expression of Bcl-2/Bax and caspase-3 in cerebral infarction rats through PTEN/PI3K/Akt signaling pathway. *Exp. Ther. Med.* **19**, 2119–2126 (2020).

79. Zhang, C.-Y. *et al.* Effect of miR-130a on neuronal injury in rats with intracranial hemorrhage through PTEN/PI3K/AKT signaling pathway. *Eur. Rev. Med. Pharmacol. Sci.* **23**, 4890–4897 (2019).

80. Zheng, T. *et al.* MiR-130a exerts neuroprotective effects against ischemic stroke through PTEN/PI3K/AKT pathway. *Biomed. Pharmacother.* **117**, 109117 (2019).

81. Ma, Q. *et al.* Blood microRNA-93 as an indicator for diagnosis and prediction of functional recovery of acute stroke patients. *J. Clin. Neurosci.* **62**, 121–127 (2019).

82. Ni, J. *et al.* MicroRNA let-7c-5p protects against cerebral ischemia injury via mechanisms involving the inhibition of microglia activation. *Brain Behav. Immun.* **49**, 75–85 (2015).

83. Tian, F., Yuan, C., Hu, L. & Shan, S. MicroRNA-93 inhibits inflammatory responses and cell apoptosis after cerebral ischemia reperfusion by targeting interleukin-1 receptor-associated kinase 4. *Exp. Ther. Med.* **14**, 2903–2910 (2017).

84. Zhang, N. *et al.* MicroRNA-378 Alleviates Cerebral Ischemic Injury by Negatively Regulating Apoptosis Executioner Caspase-3. *Int. J. Mol. Sci.* **17**, (2016).

85. Zhu, G. *et al.* MSCs-derived exosomes containing miR-486-5p attenuate cerebral ischemia and reperfusion (I/R) injury. *Gene* **906**, 148262 (2024).

86. Li, D. *et al.* MicroRNA-30e regulates neuroinflammation in MPTP model of Parkinson's disease by targeting Nlrp3. *Hum. Cell* **31**, 106–115 (2018).

87. Denk, J. *et al.* Specific serum and CSF microRNA profiles distinguish sporadic behavioural variant of frontotemporal dementia compared with Alzheimer patients and cognitively healthy controls. *PLoS One* **13**, e0197329 (2018).

88. Tan, Y. J. *et al.* Altered Cerebrospinal Fluid Exosomal microRNA Levels in Young-Onset Alzheimer's Disease and Frontotemporal Dementia. *J Alzheimers Dis Rep* **5**, 805–813 (2021).

89. Zhou, F. *et al.* Screening the expression characteristics of several miRNAs in G93A-SOD1 transgenic mouse: altered expression of miRNA-124 is associated with astrocyte differentiation by targeting Sox2 and Sox9. *J. Neurochem.* **145**, 51–67 (2018).

MAIN TABLES AND FIGURES

Characteristic	200mg x 2/day (N=3)	400mg x 2/day (N=3)	600mg x 2/day (N=2)	Total (N=8)
Sex, male	3 (100%)	3 (100%)	2 (100%)	8 (100%)
Age, years	67 (6.6), 60-73	60.3 (6.8), 55-68	69.5 (2.1), 68-71	65.1 (6.6), 55-73
Height, cm	179 (7.5), 171-185	163 (9.8), 152-172	174 (8.7), 168-180	172 (11.7), 152-185
Weight kg	73 (11), 62-84	83.9 (17.2), 66-100	78.8 (3.4), 76.4-81.2	78.6 (12.1), 62-100
Symptom duration, months	19.1 (6.5), 11.6-23.3	34.9 (33.4), 10.3-72.9	11 (0.4), 10.7-11.3	23 (21), 10.3–72.9
Diagnostic delay, months	11.3 (9.1), 8.3 – 19.3	14.5 (18.6), 2.9 - 36	9.4 (1.2), 8.5 – 10.3	12.2 (8.9), 2.9 - 36
El Escorial, definite	1 (33%)	1 (33%)	0	2 (25%)
Site of onset, bulbar	0	1 (33%)	2 (100%)	3 (38%)
Concomitant ALS medication use				
Riluzole only	2 (67%)	2 (67%)	2 (100%)	6 (75%)
Riluzole and Edaravone	1 (33%)	1 (33%)	0	2 (25%)
ALSFRS-R total score	31.3 (6.4), 24-36	32.3 (4.7), 27-36	38.5 (4.9), 35-42	33.5 (5.6), 24-42
ΔFRS, points per months	-1.1 (0.86), -2.1 – (0.54)	-0.95 (0.97), -2 – (-0.16)	-0.86 (0.42), -1.15 – (-0.56)	-0.97 (0.72), -2.1 – (-0.16)
FVC, % predicted	82.6 (27.2), 51.7-103	82.6 (17.6), 64-98.9	58 (1.4), 57-59	76.5 (20.8), 51.7-103
King's stage 2/3/4 (N)	1/1/1	1/1/1	0/1/1	2/3/3

Table 1. Characteristics of patients at baseline. Continuous data are presented with mean (standard deviation) and range, categorical data are presented as frequency (percentage).

ΔFRS = (48-ALSFRS-R total score)/ symptom duration.

Figure 1. CONSORT flow diagram of the REALS1 trial.

Figure 2. Illustration of study design. Dosing was performed over 30 days with 14 days follow up for reporting adverse effects. Blood and CSF were collected pre-dosing (day 1). CSF was collected again on day 30. Blood was withdrawn on days 7, 14, 21 and 30 with -/+ 2 days variation.

Figure 3. Change in specific miRNA levels in the plasma, compared to baseline levels, on days 7 **(A)**, 14 **(B)**, 21 **(C)**, 30 **(D)** and in all four time points averaged, compared to baseline **(E)**. miRNA counts were corrected to patients and divided by miR-451a levels to generate a "DICER score". Log 2 transformed fold change (y-axis) is shown against Log 10 transformed DICER scores (x-axis). Red indicates miRNAs which are significantly changed vs baseline (FDR<0.1 after multiple hypotheses correction). **(F)** principal component analysis of data presented in E by individual patients. Dots are annotated with the patient number. Green/blue dots denote baseline/post-treatment samples. Data in panels A-D were analyzed with two-way repeated measure ANOVA, data in E were analyzed with multiple t-tests. Multiple hypothesis correction was performed by a Two-stage linear step-up procedure of Benjamini, Krieger and Yekutieli.

Figure 4. Centroids of k-mean clustering of changes in specific miRNA levels in the plasma, compared to baseline levels, on days 7 **(A)**, 14 **(B)**, 21 **(C)**, 30 **(D)** and in all four time points averaged, compared to baseline **(E)**. miRNA counts were corrected to patients and divided by miR-451a levels. K-mean clustering was performed for the significantly increased miRNAs, with k=2 clusters. Log 2 transformed fold change (y-axis) is shown against Log 10 transformed DICER score (x-axis). Next to each centroid are annotations, or number of the miRNAs included in the corresponding cluster.

Figure 5. (A) Total DICER score in the plasma at baseline and on days 7, 14, 21 and 30 before morning dose (N=6 patients per time point). *FDR<0.1, **FDR<0.01 for comparisons vs baseline, Friedman test followed by Benjamini-Hochberg correction. **(B)** Total DICER score in the plasma at baseline and post treatment (average of days 7, 14, 21 and 30). ***p<0.0001 vs baseline, paired t-test.

Figure 6. (A) CSF miR-451a decreases after Enoxacin treatment **(B)** Total DICER score in the CSF at baseline and at the completion of 1-month treatment with Enoxacin. *p<0.05, paired t-test. Data are presented as the mean ± SEM of four patients per time point. **(C)** change in specific DICER scores after 1-month treatment compared to baseline levels. miRNA counts in each sample were corrected to patients and divided by the corresponding levels of miR-451a. Log 2 transformed fold change (y-axis) is shown against log 10 transformed of DICER scores (x-axis). Red indicates significantly changed miRNAs (FDR < 0.1, multiple paired t-test followed by the two-stage linear step-up procedure of Benjamini, Krieger and Yekutieli). miR-320 family members are shown with their suffix letter (a, b, c, d) next to the data point.

Figure 7. Pearson correlation between Enoxacin levels and fold-change values of miRNAs measured on day 14. p-values shown are unadjusted. All p-values displayed an FDR<0.1 after Benjamini-Hochberg correction.

SUPPLEMENTARY TABLES AND FIGURES

Enoxacin pharmacokinetics

C_{trough} (ng/ml)				C_{max} (ng/ml)		t_{max} (hr)	
Day 7	Day 14	Day 21	Day 30	Day 1	Day 30	Day 1	Day 30
628	563	548	685	1324	1752	2	2 (1)
(286)	(250)	(288)	(296)	(477)	(475)	(1.1)	

Supplementary table 1. Pharmacokinetic (PK) parameters of Enoxacin in plasma: trough concentrations on days 7, 14, 21 and 30 before morning dose, maximal concentrations and time to reach maximal concentrations on days 1 and 30 after Enoxacin morning dose. Data are presented as mean (sd).

Riluzole concentration (ng/ml)

	Day 1				Day 30		
2h	**4h**	**6h**	**8h**	**2h**	**4h**	**6h**	**8h**
93.53	82.4	78.0	66.67	237.3**	209.3*	177.8*	156.7*
(35.7)	(31.7)	(36.7)	(34.1)	(84.4)	(96.7)	(99.1)	(90.6)

Supplementary table 2. Plasma Riluzole concentrations on days 1 and 30 at the indicated times after Enoxacin morning dose. Data presented as mean (sd). *FDR<0.05, **FDR<0.01 compared to the respective hour on day 1, repeated measure ANOVA followed by Benjamini-Hochberg correction.

Dosing group	200mg x 2/day (N=3)		400mg x 2/day (N=3)		600mg x 2/day (N=2)	
Adverse event	**No of patients (%)**	**No of times reported**	**No of patients (%)**	**No of times reported**	**No of patients (%)**	**No of times reported**
Any adverse event	3 (100%)	13	3 (100%)	27	2 (100%)	12
Serious adverse event	0	0	0	0	0	0
Adverse event leading to discontinuation	0	0	0	0	1 (50%)	2
Severity						
Mild	2 (67%)	10	3 (100%)	16	2 (100%)	10
Moderate	1 (33%)	1	2 (67%)	10	1 (50%)	2
Severe	2 (67%)	2	1 (33%)	1	0	0
System organ class						
Nervous system	1 (33%)	1	2 (67%)	12	1 (50%)	3
Gastrointestinal	2 (67%)	6	2 (67%)	2	2 (100%)	8
Musculoskeletal	1 (33%)	2	2 (67%)	5		
Injury, poisoning and procedural complications	2 (67%)	2	1 (33%)	2		
General system disorders			2 (67%)	3	1 (50%)	1
Respiratory			1 (33%)	1		
Psychiatric			1 (33%)	2		
Vascular	1 (33%)	1				
Appetite	1 (33%)	1				
Related to drug	1 (33%)	5	2 (67%)	7	1 (50%)	7
System organ class Gastrointestinal						
Nausea	1 (33%)	4			1 (50%)*	3
Nervous system disorder						
Dizziness			1 (33%)	4	1 (50%)*	3
Headache			1 (33%)	2		
General system disorders						
Fatigue			1 (33%)	1		
Excess sweating					1 (50%)	1
Appetite						
Decreased appetite	1 (33%)	1				

Supplementary table 3. Overview of adverse events in patients with ALS treated with Enoxacin.

*Increase in the severity of these AEs caused early termination.

Supplementary figure 1. Total miRNA levels in the plasma, normalized by DESeq2, at baseline and post treatment (average of days 7, 14, 21 and 30). *p<0.05 vs baseline, paired t-test.

Supplementary figure 2. NfL levels measured in plasma **(A)** and CSF **(B)**. Enoxacin did not affect NfL levels in plasma (one-way repeated measure ANOVA: p=0.27), or in the CSF (p=0.37, paired sample t-test).