Epidermal Growth Factor in the Brain: A Promising Biomarker for Depression

Short title: Epidermal Growth Factor and Depression

Shu-xian Xu, M.D. & Ph.D.[1]; Honggang Lyu, M.D.[1]; Mian-mian Chen, M.D.[1]; Kun Li, M.D.[1]; Lihua Yao, M.D.[1]; Chao Wang, M.D. & Ph.D.[1]; Xin-hui Xie, M.D. & Ph.D.[1]*; Zhongchun Liu, M.D. & Ph.D.[1,2]*

Affiliations:

1. Department of Psychiatry, Renmin Hospital of Wuhan University, Wuhan, Hubei, PR China
2. Taikang center for life and medical sciences, Wuhan University, Wuhan, PR China.

*Corresponding authors:

Zhongchun Liu: Address: 1) Department of Psychiatry, Renmin Hospital of Wuhan University, No. 99 Jiefang Road, Wuchang District, Wuhan, Hubei, PR China. ZIP: 430060. Telephone: 86-8804191181399. 2) Taikang center for life and medical sciences, Wuhan University, Wuhan, PR China. ZIP: 430071.
E-mail: zcliu6@whu.edu.cn

Xin-hui Xie: Address: 1) Department of Psychiatry, Renmin Hospital of Wuhan University, No. 99 Jiefang Road, Wuchang District, Wuhan, Hubei, PR China. ZIP: 430060. Telephone: 86-8804191181399. Email: xxh.med@gmail.com

Abstract

Background: This study aimed to examine the neurotrophic factors secreted from astrocytes in depression by analyzing astrocyte-derived extracellular vesicles (ADEVs) isolated from plasma, and to explore the causal relationship between the expression of neurotrophic factors in the brain and depression.

Methods: A total of 40 patients with treatment-resistant depression (TRD) and 35 matched healthy controls (HCs) were recruited at baseline, and 34 TRD patients completed the post-electroconvulsive therapy (ECT) visits. The concentrations of five neurotrophic factors in ADEVs were measured. A correlation analysis was performed between neurotrophic factors in ADEVs and our previously published data on the neurogenesis marker doublecortin (DCX) in neuron-derived extracellular vesicles (NDEVs). Subsequently, Mendelian randomization (MR) study and cell experiments were conducted.

Results: Our findings revealed a decrease in the level of epidermal growth factor (EGF) in ADEVs among TRD patients compared to controls, with an increase observed post-ECT. MR suggested that decreased expression levels of the *EGF* gene in the cortex constitute a risk factor for depression. We observed a positive correlation between the levels of EGF in ADEVs and DCX in NDEVs. Subsequently, cell experiments suggested that EGF can activate EGF receptor (EGFR) to trigger the PI3K-Akt pathway, participating in the promotion of DCX.

Conclusions: This study provides the *in vivo* evidences supporting that a reduction in EGF levels in the central nervous system could potentially contribute to depression and serve as a biomarker for it. Additionally, the EGF/EGFR signaling pathway may be involved in regulating early neurogenesis traits in depression.

Keywords: treatment-resistant depression; neurotrophic factors; astrocyte-derived extracellular vesicle; neurogenesis; Mendelian randomization

Introduction

Major depressive disorder (MDD) is among the most prevalent mental disorders, marked by significant and enduring depression, loss of interest, and anhedonia[1, 2]. It is also a leading cause of disability worldwide and contributes substantially to the global burden of disease[3].

Various hypotheses have been proposed to explain the underlying causes of depression[4, 5]. Among them, the neurotrophic hypothesis of depression suggests that MDD-related synaptic and brain changes are primarily due to disrupted neurotrophic support[5]. Neurotrophic factors are of major importance because it modulates the plasticity and neurogenesis, inhibits cell death cascades and increases cell survival proteins that are responsible for proliferation and maintenance of central nervous system (CNS) neurons[4, 6].

Astrocytes, known for their critical role in regulating neurotrophic factors and gliotransmitters that maintain brain homeostasis[7, 8], may potentially contribute to the neuropathology of depression[9, 10]. Astrocytes produce various neurotrophic factors, including brain-derived neurotrophic factor (BDNF), glial cell line-derived neurotrophic factor (GDNF), epidermal growth factor (EGF), fibroblast growth factor (FGF), insulin-like growth factor (IGF-1), vascular endothelial growth factor (VEGF), and nerve growth factor (NGF), all of which are implicated in the etiology and treatment of depression[11, 12]. Among them, EGF is a protein that could be valuable for the early diagnosis and treatment assessment of depression. For instance, MDD patients with the rs11569017-TT genotype showed low plasma levels of EGF when compared to controls[13]. MDD patients exhibit reduced EGF receptor *ErbB3* expression, which increases and correlates with clinical improvement following 12 weeks of antidepressant treatment [14]. This suggests that investigating variations in the *EGF* gene and their receptor expression could serve as a clinical predictor of recovery in depression.

Unfortunately, research on other neurotrophic factors in depression remains limited, with inconsistent findings. Peripheral blood samples, used in many previous studies[15], might not accurately reflect brain neurotrophic factor levels due to their widespread expression in various tissues[16-18]. Investigating these factors directly within the brain is likely crucial for a deeper understanding of depression. The difficulty of directly measuring neurotrophic factor levels in the CNS hinders the acquisition of *in vivo* molecular evidence in patients with MDD. However, recent breakthroughs in isolating brain-derived extracellular vesicles (EVs) offer a revolutionary non-invasive approach for assessing CNS status *in vivo*. These EVs hold immense promise as a "liquid biopsy" to evaluate molecular alterations within the brain[19-21].

In this study, we initially found a decrease in the level of EGF in astrocyte-derived extracellular vesicles (ADEVs) among treatment-resistant depression (TRD) patients compared to controls, with an increase observed post-electroconvulsive therapy (ECT).

Subsequently, a Mendelian randomization (MR) study was conducted to explore the causal effect of *EGF/EGF receptor (EGFR)* gene expression on depression from a genetic perspective. As we also observed a positive correlation between EGF levels in ADEVs and the early neurogenesis marker doublecortin (DCX) in neuron-derived extracellular vesicles (NDEVs). Cell experiments using SH-SY5Y cells were then performed to investigate the relationship between EGF and DCX.

Methods

Participants details

This observational longitudinal cohort was conducted at Renmin Hospital of Wuhan University in accordance with the Declaration of Helsinki[22] as our previous study[23, 24]. The Human Ethics Committee of Renmin Hospital of Wuhan University approved the study protocol. Patients or their legal guardians provided informed consents. This report follows the STrengthening the Reporting of Observational studies in Epidemiology (STROBE) statement[25].

A total of 40 patients with TRD and 35 age-, and sex-matched healthy controls (HCs) were recruited. Subjects were excluded if they had comorbidity with other major psychiatric disorders. The HCs were required to be in good physical health, with no personal or first-degree family history of psychiatric disorders.

Thirty-four patients with TRD completed the post-ECT visits. Six milliliters of blood were drawn at baseline (T1), and 24 hours after the final ECT session (T2) to mitigate acute treatment effects. Cohort details and ECT procedures can be found in supplementary materials and our previous studies[23, 24].

Isolation and confirmation of ADEVs

We employed different isolation methods for ADEVs used in protein measurements and cell treatments. A two-step ADEVs protocol based on our prior work[24] was used for protein analysis. In contrast, ADEVs for cell treatment underwent ultracentrifugation (UC)[26] to isolate the total plasma EVs, followed by the same immunoprecipitation method to isolate ADEVs. This approach avoided potential contamination and cellular effects associated with polymer precipitation methods commonly used in clinical settings. Confirmation of ADEV isolation was performed using transmission electron microscopy (TEM), nanoparticle tracking analysis (NTA), and western blotting. Please refer to the supplementary materials and our previous studies[24] for more details.

Protein measurements in ADEVs

The levels of CD81, an EV marker, was measured using enzyme-linked immunosorbent assay (ELISA) kit (American Research Products Inc., Catalog# CSB-EL004960HU). Neurotrophic

129 factors BDNF, FGF-2, GDNF, EGF and NGF-β in ADEVs were measured using 5-Plex
130 Human ProcartaPlex™ Panel (Thermo Fisher Scientific).

Causal genetic evidence from Mendelian randomization

A MR study was conducted to validate the association between the neurotrophic factors
identified above and depression from a genetic perspective. By utilizing genetic variants as
instrumental variables, MR can provide more robust causal inference between exposures and
outcomes compared to traditional observational studies, thereby mitigating common
confounding factors such as residual confounding, reverse causation bias, and measurement
error. In this study, we leveraged brain-tissue specific expression quantitative trait loci (eQTL)
data and genome-wide association study (GWAS) summary statistics on depression to
perform two-sample MR (**sFigure 1**). Wald ratio and inverse variance weighting (IVW) were
employed as the primary analytical methods in the main analysis. Additionally, sensitivity
analyses were conducted. Details are provided in the supplementary materials.

Culture and treatments of SH-SY5Y cells

The SH-SY5Y cell line was obtained from Wuhan Procell Biotech Co., Ltd., China. These
cells were cultured in basic DMEM/F-12 medium (Gibco, Catalog# C11330500BT)
supplemented with 10% fetal bovine serum (FBS, WISENT, Catalog# 086-150) and 100 U/ml
Penicillin-Streptomycin (Gibco, Catalog# 15140122) under 5% CO_2 at 37 °C. The medium
was changed every 2-3 days, and when the cell density reached 80-90%, they were passaged
at a 1:3 ratio.

Next, the well-growing SH-SY5Y cells were seeded in a 6-well plate at a density of 6
$*10^5$ cells per well with 2.5 ml medium for 24 hours. All cells used in the experiment were in
the logarithmic growth phase. The cells were divided into two groups: the HC group (treated
with ADEVs from 6 HCs) and the TRD group (treated with ADEVs from 6 patients with
TRD). The EGF levels in ADEVs from HCs were higher than those in TRD. ADEVs were
diluted with culture medium to a final concentration of 20 μg/ml. After incubating with the
respective agents for 24 hours, the cells were harvested for subsequent analysis.

Furthermore, we performed another experiment, the well-growing SH-SY5Y cells were
seeded in a 6-well plate at a density of 6 $*10^5$ cells per well with 2.5 ml medium for 24 hours.
All cells used in the experiment were in the logarithmic growth phase. Recombinant human
EGF protein was diluted to a final concentration of 10 ng/ml to treat SH-SY5Y cells. The
control group received an equal volume of growth medium instead of the EGF solution. After
24 hours of treatment, the cells were harvested for analysis.

Western blotting

166 Previous studies found that EGF binds to its receptor EGFR, activating the PI3K-Akt and

MAPK-Erk pathways which are crucial for neurogenesis in depression(27-29). Therefore, we analyzed the proteins in these two pathways.

The following proteins were detected in SH-SY5Y cells: DCX (Abcam, Catalog# ab207175), GAPDH (Huabio, Catalog# HA721131), EGFR (Affinity, Catalog# AF6043), Phospho-EGFR (Tyr1068) (Affinity, Catalog# AF3045), Akt (Cell Signaling, Catalog# 9272S), Phospho-Akt (Ser473) (Cell Signaling, Catalog# 4060S), PI3K p85 (Affinity, Catalog# AF6241), Phospho-PI3K p85 (Affinity, Catalog# AF3241), Erk1/2 (Cell Signaling, Catalog# 9102S), Phospho-Erk1/2 (Thr202/Tyr204) (Cell Signaling, Catalog# 4370T).

Statistical analyses

For the clinical data, in the main analysis of protein levels in ADEVs at baseline, following the MIESV 2018(30), the outcomes were defined as the ratio of protein concentrations to CD 81 concentrations. General linear models (GLMs) were used to examine differences between two groups. In the GLMs, we set the outcomes as dependent variables and group as the independent variable, while controlling for covariates for sex and age. To represent the differences between the two groups, we used model-fitted estimated marginal means (EMMs) and their corresponding 95% confidence intervals (CIs). Additionally, to ensure robustness, a sensitivity analysis was conducted, with protein concentration levels per 1 ml plasma set as dependent variables. Receiver operating characteristic curves (ROCs) and corrected area under the curve (AUC) were employed to further explore the potential of these neurotrophic factors in ADEVs as biomarkers. In the exploratory analysis, Spearman coefficient was used to examine the correlation between EGF in ADEVs and DCX in NDEVs (as reported in our previous report(23)). For the comparisons between pre- and post-ECTs (paired longitudinal data), paired *t*-tests were used, with Cohen's *d* and its associated 95% CIs utilized to quantify effect sizes. Similar to the comparisons at baseline, a sensitivity analysis was conducted with the concentrations of protein levels per 1 ml plasma as dependent variables. False discovery rate (FDR) correction (Benjamini and Hochberg method) was employed to mitigate false positive outcomes. In comparisons of data from cell experiments, the Wilcoxon tests were employed. A two-tailed *p*-values less than 0.05 was considered significant. All statistical analyses were performed using R version 4.1.0 (R Project for Statistical Computing) within Rstudio version 1.4.1106 (Rstudio).

Results

EGF levels in ADEVs represent a promising potential biomarker for depression

Among 40 patients with TRD, six patients either did not undergo ECT treatment or did not complete the post-ECT visit. Detailed baseline characteristics of the participants was provided in **sTable 1**.

At baseline, the level of EGF in the TRD group was significantly lower than in the HC

group. This effect was consistent across different normalization methods. Changes in other neurotrophic factors were more variable and yielded different outcomes depending on the normalization method used (**Table 1** and **Figure 1**). The AUCs for EGF were larger than those for other neurotrophic factors, with values of 0.99 (95% CI: 0.98-1.00) and 0.93 (95% CI: 0.86-0.99), respectively (**Figure 1**). This indicates that EGF in ADEVs has a strong discrimination ability. Additionally, the EGF level in ADEVs showed a positive correlation (ρ = 0.38, $p < 0.001$) with the DCX level in NDEVs (**sFigure 2**).

Following ECT, there was a significant increase in EGF levels, this effect was also consistent across different normalization methods (**Table 2** and **Figure 2**). In summary, among the five neurotrophic factors tested, EGF showed the most robust results both at baseline and in terms of changes before and after ECT.

Causal influence of brain-specific *EGF* gene expression on depression

Given the significant correlation between EGF expression and depression above, we employed MR to investigate the causal influence of brain-tissue specific *EGF* and *EGFR* genes expression on depression. Adhering to MR assumptions, we identified cis-eQTL data for *EGF* and *EGFR* genes in the brain cortex, respectively (**sTable 2**). Our MR analysis suggests that decreased expression of the *EGF* gene in the cortex may be a potential risk factor for depression (Wald ratio OR= 0.95, $p = 0.009$) (**sFigure 3**). However, neither method found a significant causal relationship between *EGFR* and depression.

Subsequently, we attempted to identify potential mechanistic pathways between *EGF/EGFR* genes and depression through MELODI Presto(31). The analysis revealed that potential intermediary factors between these genes and depression were primarily associated with metabolic processes, cellular function, and neurological disorders. Interestingly, the results suggest an overlap between the *EGF*-depression interaction and the serotonin transporter pathway, potentially implicating *EGF* as a target for antidepressant development (**sTable 3**).

EGF activates EGFR to participate in the promotion of DCX expression

ADEVs purified using UC were confirmed via TEM, NTA, and western blotting validating successful ADEVs collection (**sFigure 4**).

As shown in **Figure 3A-F**, after 24 hours of intervention with ADEVs from patients with TRD and HCs, the expression level of the DCX protein in SH-SY5Y cells intervened by ADEVs in the HC group showed a significant increase compared to the TRD group ($p = 0.009$). Moreover, there was a significant increase in the phosphorylation level of the EGFR ($p = 0.041$), along with simultaneous increases in the phosphorylation levels of downstream pathway proteins PI3K ($p = 0.004$) and Akt ($p = 0.015$). However, there was no significant difference in the phosphorylation level of Erk between the two groups ($p = 0.818$).

Following a 24-hour intervention with recombinant human EGF (**Figure 3G-L**), the
expression level of the DCX protein in SH-SY5Y cells significantly increased compared to
the control group ($p = 0.041$), accompanied by a significant elevation in the phosphorylation
level of EGFR ($p = 0.015$). Furthermore, the phosphorylation levels of downstream pathway
proteins PI3K ($p = 0.015$) and Akt ($p = 0.004$) were also significantly increased concurrently.
Nonetheless, there was no significant difference in the phosphorylation level of Erk between
the two groups ($p = 0.240$).
250

Discussion

Our study, using plasma ADEVs, indicates that decreased levels of EGF in the CNS could
serve as a biomarker for depression. Subsequently, MR results suggest that decreased
expression of the *EGF* gene in the cortex may contribute to the development of depression.
We also discovered a positive correlation between EGF levels in ADEVs and DCX,
suggesting that the EGF/EGFR signaling pathway may play a role in regulating early
neurogenesis traits in depression.

This study's strength lies in directly acquiring CNS evidences through ADEVs and MR
based on brain-tissue data, bypassing limitations of peripheral sources. Neurotrophic factors,
like EGF, are often found at higher levels in peripheral tissues compared to the brain(17, 18).
This widespread expression can confound results when solely using peripheral measures, a
common limitation in depression studies. Additionally, the blood-brain barrier further
complicates assessing CNS neurotrophic factors(32). Therefore, our study offered valuable
insights into the brain-specific relationship between neurotrophic factors and depression.

EGF and depression

Previous research on EGF and depression has yielded conflicting results. While one
study found elevated EGF levels in adolescents with depression or bipolar disorder(33),
another study focusing on elderly patients did not detect differences compared to HCs(34),
consistent with another case-control study(35). Additionally, two studies reported decreased
peripheral blood EGF levels in depressed patients(13, 36), consistent with findings of
downregulated *EGF* mRNA levels in peripheral blood(37). Additionally, cognitive alterations
in MDD patients have been associated with the SNP rs2250724 of *EGF* gene(38).

Heterogeneity in EGF findings likely stems from several factors. First, variations in
study populations can influence results. Serum EGF levels demonstrably decline with age(39),
highlighting potential age-related heterogeneity. Second, comorbid conditions like anxiety
might confound EGF expression. A study reported elevated EGF in patients with mild to
moderate depression and anxiety(40), while others found decreases in depression without
anxiety(13, 36). Most importantly, previous research has primarily focused on assessing EGF
levels in peripheral blood, where they can be influenced by various tissues(17). This

highlights the importance of directly assessing EGF expression within the CNS, precisely the approach employed in our study. We utilized ADEVs for the first time to detect EGF within the brains of depression patients, and observed a significant decrease in EGF levels in TRD patients compared to HCs. ROC curve analyses revealed high sensitivity and specificity of EGF in ADEVs in identifying TRD, suggesting it as a promising biomarker for depression.

Our study further revealed that levels of EGF in ADEVs increased following ECTs in patients with depression. Earlier studies have observed acute changes in EGF levels after ECT. For example, one study reported a significant decrease in serum EGF concentrations following ECTs(41), while another focusing on schizophrenia showed no change in serum EGF levels between antipsychotic therapy alone and combined ECT and antipsychotic treatment(42). Notably, these prior studies relied on peripheral EGF measurements, which may not reflect central EGF dynamics. However, our findings, based on ADEVs which directly reflect the CNS, suggest EGF as a potential marker for ECT response, highlighting the importance of CNS-level assessment for understanding and treating depression.

Additionally, our MR analysis indicated that genetically predicted decreases in *EGF* gene expression constitute a potential risk factor for depression, further supporting our clinical observations. As eQTL effects on gene expression can vary depending on the tissue examined, potentially introducing confounding factors and leading to incorrect inferences. To address this limitation, we specifically investigated the causal relationship between *EGF* eQTLs derived from brain-tissue specifically in the cortex and depression. To strengthen our results, we used the MELODI Presto analysis tool to further prioritize the *EGF* gene, highlighting its potential interactions with antidepressive agents. This provides additional evidence supporting the involvement of *EGF* in the mechanisms of antidepressant activity. This approach offers insights into the biological pathways linking *EGF* and depression, indicating that changes in the *EGF* expression could play a crucial role in the pathogenesis of depression.

The observed genetic associations with reduced *EGF* expression reinforce the need for additional preclinical studies to investigate how *EGF*-based interventions might benefit patients with depression, and they also shed light on how *EGF*-related pathways contribute to the onset and treatment of depression.

EGF/EGFR, neurogenesis and depression

Our previous study using NDEVs, firstly provided human *in vivo* evidence of reduced markers of early neurogenesis DCX in patients with TRD, and observed increased traits of immature neurons following ECT(23). One potential mechanism linking neurogenesis impairments with depression is through the loss of neurotrophic factors and the disruption of their associated signaling pathways(5). This study discovered a notable positive correlation between EGF levels in ADEVs and DCX levels in NDEVs. This finding suggests a potential

link between EGF and early neurogenesis. Then, we validated that ADEVs from HCs, with higher EGF levels, increased DCX expression in SH-SY5Y cells compared to ADEVs from patients with TRD. Similarly, EGF protein alone also stimulated DCX expression in SH-SY5Y cells, potentially via the EGFR-PI3K-Akt pathway.

Astrocytes are the predominant cell type within the neurogenic niche, creating a specialized environment for neurogenesis(43). Previous studies have found that astrocytes can secret EGF to increase neurogenesis(8, 44). EGF can promotes the differentiation, maturation, and survival of various neurons, exerting neuromodulatory effects on different types of neurons in the CNS(44). The exogenous EGF has specific effects on precursor cells *in vivo*(45). Administering EGF into the ventricles for 2 weeks expands precursor cell populations in the subventricular zone (SVZ), decreases the total number of newborn neurons reaching the olfactory bulb, and significantly increases the production of olfactory bulb astrocytes. Moreover, EGF increases the number of newborn cells in the striatum either by migrating SVZ cells or by stimulating local precursor cells, while also boosting the number of newborn glial cells and reducing newborn neurons(46). These findings illuminate EGF's role in neurogenesis, and emerging evidence suggests that neurogenesis may play a role in depression(47).

However, there is limited research exploring the relationship between EGF and neurogenesis in depression. One study shown that FGF, EGF, BDNF, ketamine, melatonin and the ketamine/melatonin combination stimulate neurogenesis and cell cluster formation in cultured human olfactory neuronal precursors, thus offering insights into the mechanism underlying their antidepressant effects(48). Once EGF binds to its receptor, EGFR, it forms a homodimer, which undergoes autophosphorylation in the intracellular kinase domain, activating signaling cascades within the cell and even in the nucleus(49). Upon EGFR activation, the main downstream signaling pathways include the PI3K-Akt pathway, the Ras-Raf-MAPK-Erk pathway, and the JAK/STAT signaling pathway(29). Studies have shown that the PI3K-Akt and MAPK-Erk signaling pathways play crucial roles in neurogenesis, with several therapeutic targets for depression acting on these pathways to promote neuronal growth and survival(27, 28). Our findings suggest that EGF might exert its effect on early neurogenesis in depression by specifically activating the EGFR, which in turn triggers the PI3K-Akt pathway. This activation pattern contrasts with the MAPK-Erk pathway, often implicated in other cell growth and differentiation processes, indicating that the selective activation of PI3K-Akt by EGF could be a distinctive mechanism through which early neurogenesis is influenced in the context of depression. ECT may stimulate this signaling pathway by promoting EGF secretion, leading to increased early neurogenesis and potentially achieving antidepressant effects.

EV, a candidate key mediator in mediating EGF-neurogesis signaling

During the process of EGF/EGFR signaling, EVs may play a significant role. EVs
derived from neurons and neuroglial cells exert crucial bidirectional regulatory effects in the
physiological and pathological processes of the CNS(50). These nanoscale messengers not
only ferry information from neurons to glial cells but also mediate the reverse pathway,
influencing neuronal function and establishing a dynamic intercellular network(20). For
instance, neurons can transfer miR-124b to astrocytes via EVs, enhancing the expression of
glutamate transporter-1 in astrocytes, thereby regulating synaptic plasticity(51). Additionally,
ADEVs can enhance neuronal survival under ischemic and hypoxic conditions by delivering
antiviral proteins(52). Regarding our study, results of cell experiments suggest a possible
antidepressant mechanism between astrocytes and neurogenesis, EGF carried by ADEVs
activates the EGFR signaling pathway (including PI3K-Akt) in neurons, thereby promoting
neuronal proliferation and growth.

This study proposes a novel framework for investigating CNS diseases in human
subjects. By leveraging brain-derived extracellular vesicles (BDEVs), we can gain initial
insights into the *in vivo* CNS state, facilitating the study of real-world pathological conditions.
Furthermore, complementing BDEV analysis with *in vitro* cell experiments offers a
comprehensive understanding of CNS biology. Additionally, the potential of EVs as drug
delivery vehicles, with their ability to cross the blood-brain barrier and target specific cell
types, opens exciting avenues for developing targeted therapeutic strategies for CNS diseases.
376

Limitations
This study has several limitations. Firstly, the limited quantity of ADEVs restricted
high-throughput omics analyses, also hindering the detection of additional neurotrophic
factors. Additionally, plasma ADEVs represent the entire brain, precluding identification of
specific neurotrophic factor source regions. Elucidating these regional origins is helpful for
understanding their roles in depression. Secondly, for the MR study, the major GWAS data
populations were predominantly European, necessitating further validation of these MR
findings in other populations. Also, the limited number of instruments for cis-eQTL data from
cortex tissue and other brain tissues to investigate their association with depression,
warranting further examination in future studies. Furthermore, large-scale independent
tissue-specific eQTL datasets and employing additional genetic methods like
transcriptome-wide association studies(53) to estimate gene expression effects on depression
should be considered in future studies. We look forward to future research addressing this
issue more specifically. Thirdly, for cell experiments, resource constraints necessitated using
the SH-SY5Y cell line instead of primary neural stem cells, potentially compromising the
accuracy of neurogenesis data. Future studies should prioritize neural stem cells for a more
faithful representation of neurogenesis processes. Additionally, serum-containing media
hindered precise control over EGF levels, potentially influencing our results. Employing

serum-free media would enhance experimental control. Finally, the complex composition of
ADEVs limited our ability to identify the specific pro-neurogenic molecule(s). Future studies
isolating and analyzing individual ADEV components are necessary to pinpoint the molecules
responsible for the observed increase in neurogenesis.
399
Conclusions
This study provides *in vivo* evidence for a potential role of reduced CNS EGF levels in
TRD patients, suggesting EGF as a promising biomarker. Then using MR, we revealed a
causal influence of low brain-specific *EGF* gene expression on depression. Furthermore, our
findings highlight the involvement of the EGF/EGFR signaling pathway in regulating early
neurogenesis, a process implicated in depression. These insights may offer critical insights for
developing novel antidepressant therapies targeting the EGF/EGFR pathway using EVs.
407
408

Acknowledgements

This work was supported by grants from the National Natural Science Foundation of China (U21A20364). No funding was received from commercial organizations.

Conflict of interest disclosure

The authors declare no competing interests.

Authors' contributions

SXX conducted methodology, investigation, data curation, and contributed to the original draft and conceptualization. HL contributed to data curation and participated in manuscript review and editing. MMC and KL conducted investigation and data curation. LY and CW participated in manuscript review and editing. XHX provided conceptualization, performed formal analysis, visualization and supervision. ZL provided project administration, supervision, and acquired funding.

References

1. Otte C, Gold SM, Penninx BW, Pariante CM, Etkin A, Fava M, et al. (2016): Major depressive disorder. *Nat Rev Dis Primers*. 2:16065.

2. Diseases GBD, Injuries C (2020): Global burden of 369 diseases and injuries in 204 countries and territories, 1990-2019: a systematic analysis for the Global Burden of Disease Study 2019. *Lancet*. 396:1204-1222.

3. Friedrich MJ (2017): Depression Is the Leading Cause of Disability Around the World. *JAMA*. 317:1517.

4. Chopra K, Kumar B, Kuhad A (2011): Pathobiological targets of depression. *Expert Opin Ther Targets*. 15:379-400.

5. Fries GR, Saldana VA, Finnstein J, Rein T (2023): Molecular pathways of major depressive disorder converge on the synapse. *Mol Psychiatry*. 28:284-297.

6. Skaper SD (2018): Neurotrophic Factors: An Overview. *Methods Mol Biol*. 1727:1-17.

7. Cope EC, Gould E (2019): Adult Neurogenesis, Glia, and the Extracellular Matrix. *Cell Stem Cell*. 24:690-705.

8. Araki T, Ikegaya Y, Koyama R (2021): The effects of microglia- and astrocyte-derived factors on neurogenesis in health and disease. *Eur J Neurosci*. 54:5880-5901.

9. Wang Q, Jie W, Liu JH, Yang JM, Gao TM (2017): An astroglial basis of major depressive disorder? An overview. *Glia*. 65:1227-1250.

10. Rajkowska G, Stockmeier CA (2013): Astrocyte pathology in major depressive disorder: insights from human postmortem brain tissue. *Curr Drug Targets*. 14:1225-1236.

11. Galvez-Contreras AY, Campos-Ordonez T, Lopez-Virgen V, Gomez-Plascencia J, Ramos-Zuniga R, Gonzalez-Perez O (2016): Growth factors as clinical biomarkers of prognosis and diagnosis in psychiatric disorders. *Cytokine Growth Factor Rev*. 32:85-96.

12. Martin JL, Magistretti PJ, Allaman I (2013): Regulation of neurotrophic factors and energy metabolism by antidepressants in astrocytes. *Curr Drug Targets*. 14:1308-1321.

13. Tian W, Zhang J, Zhang K, Yang H, Sun Y, Shen Y, et al. (2012): A study of the functional significance of epidermal growth factor in major depressive disorder. *Psychiatr Genet*. 22:161-167.

14. Milanesi E, Minelli A, Cattane N, Cattaneo A, Mora C, Barbon A, et al. (2012): ErbB3 mRNA leukocyte levels as a biomarker for major depressive disorder. *BMC Psychiatry*. 12:145.

15. Wang H, Yang Y, Pei G, Wang Z, Chen N (2023): Neurotrophic basis to the pathogenesis of depression and phytotherapy. *Front Pharmacol*. 14:1182666.

16. Colitti M (2017): Distribution of BDNF and TrkB isoforms in growing antler tissues of red deer. *Ann Anat*. 213:33-46.

17. Fukuyama R, Shimizu N (1991): Expression of epidermal growth factor (EGF) and the EGF receptor in human tissues. *J Exp Zool*. 258:336-343.

18. The Genotype-Tissue Expression (GTEx) Portal https://www.gtexportal.org/home/.

19. Shi M, Sheng L, Stewart T, Zabetian CP, Zhang J (2019): New windows into the brain: Central nervous system-derived extracellular vesicles in blood. *Prog Neurobiol*. 175:96-106.

20. Saint-Pol J, Gosselet F, Duban-Deweer S, Pottiez G, Karamanos Y (2020): Targeting and Crossing the Blood-Brain Barrier with Extracellular Vesicles. *Cells*. 9.

21. Brenna S, Krisp C, Altmeppen HC, Magnus T, Puig B (2021): Brain-Derived Extracellular Vesicles in Health and Disease: A Methodological Perspective. *Int J Mol Sci*. 22.

22. World Medical A (2013): World Medical Association Declaration of Helsinki: ethical principles for medical research involving human subjects. *JAMA*. 310:2191-2194.

23. Xie XH, Xu SX, Yao L, Chen MM, Zhang H, Wang C, et al. (2024): Altered in vivo early neurogenesis traits in patients with depression: Evidence from neuron-derived extracellular vesicles and electroconvulsive therapy. *Brain Stimul*. 17:19-28.

24. Xu SX, Xie XH, Yao L, Wang W, Zhang H, Chen MM, et al. (2023): Human in vivo evidence of reduced astrocyte activation and neuroinflammation in patients with treatment-resistant depression following electroconvulsive therapy. *Psychiatry Clin Neurosci*. 77:653-664.

25. von Elm E, Altman DG, Egger M, Pocock SJ, Gotzsche PC, Vandenbroucke JP, et al. (2007): The Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) statement: guidelines for reporting observational studies. *Lancet*. 370:1453-1457.

26. Greening DW, Xu R, Ji H, Tauro BJ, Simpson RJ (2015): A protocol for exosome isolation and characterization: evaluation of ultracentrifugation, density-gradient separation, and immunoaffinity capture methods. *Methods Mol Biol*. 1295:179-209.

27. Liu Q, Zhang J, Xiao C, Su D, Li L, Yang C, et al. (2022): Akebia saponin D protects hippocampal neurogenesis from microglia-mediated inflammation and ameliorates depressive-like behaviors and cognitive impairment in mice through the PI3K-Akt pathway. *Front Pharmacol*. 13:927419.

28. Xia CX, Gao AX, Dong TT, Tsim KW (2023): Flavonoids from Seabuckthorn (Hippophae rhamnoides L.) mimic neurotrophic functions in inducing neurite outgrowth in cultured neurons: Signaling via PI3K/Akt and ERK pathways. *Phytomedicine*. 115:154832.

29. Wee P, Wang Z (2017): Epidermal Growth Factor Receptor Cell Proliferation Signaling Pathways. *Cancers (Basel)*. 9.

30. Thery C, Witwer KW, Aikawa E, Alcaraz MJ, Anderson JD, Andriantsitohaina R, et al. (2018): Minimal information for studies of extracellular vesicles 2018 (MISEV2018): a position statement of the International Society for Extracellular Vesicles and update of the MISEV2014 guidelines. *J Extracell Vesicles*. 7:1535750.

31. Elsworth B, Gaunt TR (2021): MELODI Presto: a fast and agile tool to explore semantic triples derived from biomedical literature. *Bioinformatics*. 37:583-585.

32. Thorne RG, Frey WH, 2nd (2001): Delivery of neurotrophic factors to the central

nervous system: pharmacokinetic considerations. *Clin Pharmacokinet*. 40:907-946.

33. Skibinska M, Kapelski P, Dmitrzak-Weglarz M, Lepczynska N, Pawlak J, Twarowska-Hauser J, et al. (2021): Elevated Epidermal Growth Factor (EGF) as Candidate Biomarker of Mood Disorders-Longitudinal Study in Adolescent and Young Adult Patients. *J Clin Med*. 10.

34. Wu HE, Teixeira AL, Barroso L, Silva APM, de Souza Nicolau M, Ferreira JDR, et al. (2019): Epidermal growth factor and fibroblast growth factor-2 circulating levels in elderly with major depressive disorder. *Psychiatry Res*. 272:141-143.

35. Domenici E, Wille DR, Tozzi F, Prokopenko I, Miller S, McKeown A, et al. (2010): Plasma protein biomarkers for depression and schizophrenia by multi analyte profiling of case-control collections. *PLoS One*. 5:e9166.

36. Sohan M, Qusar M, Shahriar M, Islam SMA, Bhuiyan MA, Islam MR (2023): Association of reduced serum EGF and leptin levels with the pathophysiology of major depressive disorder: A case-control study. *PLoS One*. 18:e0288159.

37. Woo HI, Lim SW, Myung W, Kim DK, Lee SY (2018): Differentially expressed genes related to major depressive disorder and antidepressant response: genome-wide gene expression analysis. *Exp Mol Med*. 50:1-11.

38. Tian WM, Zhang KR, Zhang J, Shen Y, Xu Q (2010): Association between the epidermal growth factor gene and intelligence in major depression patients. *Chin Med Sci J*. 25:105-108.

39. Kjaer IM, Olsen DA, Alnor A, Brandslund I, Bechmann T, Madsen JS (2019): EGFR and EGFR ligands in serum in healthy women; reference intervals and age dependency. *Clin Chem Lab Med*. 57:1948-1955.

40. Memon AA, Sundquist K, Ahmad A, Wang X, Hedelius A, Sundquist J (2017): Role of IL-8, CRP and epidermal growth factor in depression and anxiety patients treated with mindfulness-based therapy or cognitive behavioral therapy in primary health care. *Psychiatry Res*. 254:311-316.

41. Stelzhammer V, Guest PC, Rothermundt M, Sondermann C, Michael N, Schwarz E, et al. (2013): Electroconvulsive therapy exerts mainly acute molecular changes in serum of major depressive disorder patients. *Eur Neuropsychopharmacol*. 23:1199-1207.

42. Zhang X, Xiao W, Chen K, Zhao Y, Ye F, Tang X, et al. (2020): Serum Epidermal Growth Factor is Low in Schizophrenia and Not Affected by Antipsychotics Alone or Combined With Electroconvulsive Therapy. *Front Psychiatry*. 11:104.

43. Verkhratsky A, Matteoli M, Parpura V, Mothet JP, Zorec R (2016): Astrocytes as secretory cells of the central nervous system: idiosyncrasies of vesicular secretion. *EMBO J*. 35:239-257.

44. Wong RW, Guillaud L (2004): The role of epidermal growth factor and its receptors in mammalian CNS. *Cytokine Growth Factor Rev*. 15:147-156.

45. Kuhn HG, Winkler J, Kempermann G, Thal LJ, Gage FH (1997): Epidermal growth

factor and fibroblast growth factor-2 have different effects on neural progenitors in the adult rat brain. *J Neurosci*. 17:5820-5829.

46. Kuhn HG, Dickinson-Anson H, Gage FH (1996): Neurogenesis in the dentate gyrus of the adult rat: age-related decrease of neuronal progenitor proliferation. *J Neurosci*. 16:2027-2033.

47. Berger T, Lee H, Young AH, Aarsland D, Thuret S (2020): Adult Hippocampal Neurogenesis in Major Depressive Disorder and Alzheimer's Disease. *Trends Mol Med*. 26:803-818.

48. Estrada-Reyes R, Quero-Chavez DB, Alarcon-Elizalde S, Cercos MG, Trueta C, Constantino-Jonapa LA, et al. (2022): Antidepressant Low Doses of Ketamine and Melatonin in Combination Produce Additive Neurogenesis in Human Olfactory Neuronal Precursors. *Molecules*. 27.

49. Stoll SW, Stuart PE, Swindell WR, Tsoi LC, Li B, Gandarillas A, et al. (2016): The EGF receptor ligand amphiregulin controls cell division via FoxM1. *Oncogene*. 35:2075-2086.

50. Wu J, Li S, Zhang Y (2023): Research progress in role of exosomes exosomes in mental disorders. *Zhong Nan Da Xue Xue Bao Yi Xue Ban*. 48:771-781.

51. Morel L, Regan M, Higashimori H, Ng SK, Esau C, Vidensky S, et al. (2013): Neuronal exosomal miRNA-dependent translational regulation of astroglial glutamate transporter GLT1. *J Biol Chem*. 288:7105-7116.

52. Guitart K, Loers G, Buck F, Bork U, Schachner M, Kleene R (2016): Improvement of neuronal cell survival by astrocyte-derived exosomes under hypoxic and ischemic conditions depends on prion protein. *Glia*. 64:896-910.

53. Gusev A, Ko A, Shi H, Bhatia G, Chung W, Penninx BW, et al. (2016): Integrative approaches for large-scale transcriptome-wide association studies. *Nat Genet*. 48:245-252.

567   **Figure 1.** Comparisons of target protein concentrations in astrocyte-derived extracellular

568   vesicles (ADEVs) between treatment-resistant depression (TRD) and healthy control (HC)

569   group at baseline. Note: n.s.: not significant; $*p_{\text{false discovery rate (FDR)}} < 0.05$; $**p_{\text{FDR}} < 0.01$;

570    $***p_{\text{FDR}} < 0.001$.

572
Figure 2. Changes of target protein concentrations in ADEVs between pre- and post-ECT in
patients with TRD. Note: n.s.: not significant; **p*FDR < 0.05.

Figure 3. Western blotting results of cell expriments. A-F: After 24 hours of intervention with ADEVs from patients with TRD and HCs in SH-SY5Y cells, the expression levels of doublecortin (DCX) protein in the HC group significantly increased compared to the TRD group. The phosphorylation levels of EGFR, PI3K, and Akt also significantly increased, while there was no significant difference in the phosphorylation levels of Erk between the two groups. G-L: After 24 hours of intervention with ADEVs from patients with TRD and HCs in SH-SY5Y cells, the expression levels of DCX protein in the HC group significantly increased compared to the TRD group. The phosphorylation levels of EGFR, PI3K, and Akt also significantly increased, while there was no significant difference in the phosphorylation levels of Erk between the two groups. Note: n.s.: not significant; *$p_{FDR} < 0.05$; **$p_{FDR} < 0.01$.

588 Table 1. Baseline comparisons of neurotrophic factors in ADEVs between TRD and HC groups

variable	HC group (EMM (95% CI))	TRD group (EMM (95% CI))	Difference (95% CI)	p	p_{FDR}
Main analysis: Normalized to CD81 ratio					
BDNF	0.0025 (0.0022 to 0.0029)	0.0012 (0.0009 to 0.0015)	-0.0013 (-0.0018 to -0.0009)	<0.001	<0.001
EGF	0.0011 (0.001 to 0.0012)	0.0003 (0.0002 to 0.0004)	-0.0008 (-0.0009 to -0.0006)	<0.001	<0.001
FGF-2	0.0039 (0.0032 to 0.0045)	0.0031 (0.0025 to 0.0037	-0.0008 (-0.0016 to 0.0001)	0.097	0.097
GDNF	0.0167 (0.0145 to 0.0189)	0.0121 (0.01 to 0.0141)	-0.0047 (-0.0076 to -0.0018)	0.002	0.004
NGF-β	0.0043 (0.0036 to 0.0050)	0.0035 (0.0028 to 0.0042)	-0.0008 (-0.0017 to 0.0001)	0.094	0.097
Sensitive analysis: Normalized to 1 ml plasma					
BDNF	9.84 (8.16 to 11.52)	8.65 (7.1 to 10.2)	-1.19 (-3.41 to 1.03)	0.298	0.298
EGF	3.98 (3.62 to 4.34)	2.01 (1.67 to 2.34)	-1.97 (-2.45 to -1.5)	<0.001	<0.001
FGF-2	13.94 (10.56 to 17.32)	21.39 (18.27 to 24.51)	7.45 (2.98 to 11.92)	0.002	0.002
GDNF	60.55 (53.47 to 67.64)	82.77 (76.24 to 89.31)	22.22 (12.85 to 31.59)	<0.001	<0.001
NGF-β	15.42 (11.71 to 19.13)	24.56 (21.14 to 27.98)	9.14 (4.23 to 14.05)	<0.001	0.001

592 Table 2. Changes of neurotrophic factors in ADEVs of TRD after ECTs.

Variable	Change from baseline (95% CI))	Cohen's d (95% CI))	p	p_{FDR}
Main analysis: Normalized to CD81 ratio				
BDNF	0.00001 (-0.00022 to 0.00024)	0.01 (-0.33 to 0.36)	0.934	0.934
EGF	0.00027 (0.00019 to 0.00036)	1.14 (0.69 to 1.57)	<0.001	<0.001
FGF-2	-0.00047 (-0.00118 to 0.00024)	-0.23 (-0.58 to 0.11)	0.189	0.321
GDNF	0.00066 (-0.00101 to 0.00233)	0.14 (-0.2 to 0.48)	0.424	0.53
NGF-β	-0.00048 (-0.00122 to 0.00026)	-0.23 (-0.58 to 0.12)	0.193	0.321
Sensitive analysis: Normalized to 1 ml plasma				
BDNF	-1.31 (-3.03 to 0.41)	-0.27 (-0.61 to 0.08)	0.131	0.131
EGF	1.2 (0.74 to 1.67)	0.91 (0.5 to 1.31)	<0.001	<0.001
FGF-2	-5.7 (-9.63 to -1.76)	-0.51 (-0.87 to -0.15)	0.006	0.007
GDNF	-9.15 (-15.48 to -2.82)	-0.51 (-0.87 to -0.15)	0.006	0.007
NGF-β	-6.74 (-11.19 to -2.29)	-0.54 (-0.9 to -0.17)	0.004	0.007

594 **Supplementary materials**

600 **sFigure 1.** Mendelian randomization framework.

610   **sFigure 2.** The spearman coefficient between EGF in ADEVs and the DCX in NDEVs.

Exposure	Methods	snp		OR (95% CI)	P-value
Cortex EGF gene expression					
	Wald ratio	rs2190907	■	0.95 (0.91 to 0.99)	0.009
	MR RAPS	rs2190907	■	0.95 (0.90 to 0.99)	0.021
Cortex EGFR gene expression					
	Wald ratio	rs73135226	■	1.00 (0.97 to 1.03)	0.985
	MR RAPS	rs73135226	■	1.00 (0.97 to 1.03)	0.986

```
          0.8      1     1.2
        ←─────────────────────→
        Protect Factor  Risk Factor
```

612
613

sFigure 3. Results of Mendelian randomization study.

sFigure 4. Confirmation of ADEVs utilizing UC. A) The results of NTA show the diameter

distribution of the EVs, which also corroborate with the prior knowledge of exosomes-like

small EVs (peak diameter: 82.3 nm, concentration: $2.4*10^9$ particles/ml); B) The transmission

electron microscopy image of ADEVs isolated from a health control, the image clearly shows

the exosomes-like shape (orange arrows); C) Western blotting results of total EV (TEV),

supernatant of TEV, and ADEV from one healthy control. Three exosomes markers, CD9,

Alix, TSG 101, and an astrocyte marker glial fibrillary acidic protein (GFAP) was identified.

sTable 1. Baseline characteristics of included participants for clinical trial.

sTable 2. Genetic instruments for *EGF/EGFR* expression in brain tissues.

sTable 3. MELODI Presto results for *EGF/EGFR* genes and depression.

Supplementary document: Detailed Methods.