Development of Alzheimer's Disease Risk Score for Future Primary Care:

A White-Box Approach

Yumiko Wiranto, M.A.[a*+], Devin R Setiawan, B.S.[b+], Amber Watts, Ph.D.[a,c], Arian Ashourvan, Ph.D.[a], and for the Alzheimer's Disease Neuroimaging Initiative[1]

[a]Department of Psychology, University of Kansas, Lawrence, Kansas, USA

[b]Department of Electrical Engineering and Computer Science, University of Kansas, Lawrence, Kansas, USA

[c]University of Kansas, Alzheimer's Disease Research Center, Fairway, Kansas, USA

[1]Data used in preparation of this article were obtained from the Alzheimer's Disease Neuroimaging Initiative (ADNI) database (adni.loni.usc.edu). As such, the investigators within the ADNI contributed to the design and implementation of ADNI and/or provided data but did not participate in analysis or writing of this report.

[+] These authors contributed equally to this work.

* Corresponding author

Yumiko Wiranto

Department of Psychology, University of Kansas

Email: yumiko.wiranto@ku.edu

Word count: 2592 words

Key Points

Question: Can accessible information, such as demographics, cognitive tests, and functioning questionnaire, yield in reliable results for predicting Alzheimer's disease development using interpretable machine learning?

Findings: The results of 665 participants from the Alzheimer's Disease Neuroimaging Initiative demonstrated robust performance of determining Alzheimer's disease development using four separate measures of (F)unctioning, (LA)nguage, (M)emory, and (E)xecutive functioning or the FLAME scorecard. It remains reliable when externally validated with a separate dataset of 3657 participants from the National Alzheimer's Coordinating Center.

Meaning: The FLAME scorecard shows potential to be implemented in integrated primary care settings to promote early detection and intervention of cognitive decline due to Alzheimer's disease.

Abstract

Importance: Interpretable scoring system can contribute to bridge the gap between the timeliness and complexity of diagnosing Alzheimer's disease (AD) and promote early intervention at non-specialist settings.

Objective: To develop a risk score to predict the likelihood of AD with interpretable machine learning using variables that are obtainable at integrated primary care settings.

Design: A secondary data analysis including cohort studies from the Alzheimer's Disease Neuroimaging Initiative (ADNI) and the National Alzheimer's Coordinating Center (NACC) extracted in August 2023 and March 2024.

Setting: The ADNI and NACC are multi-site cohort studies in North America.

Participants: Participants with normal cognition or mild cognitive impairment at baseline visit were identified. Participants with the same diagnosis overtime were assigned to the stable group, and those converted to AD were placed in the progressive group.

Main Outcome(s) and Measure(s): Cognitive tests and daily functioning measured with Functional Assessment Questionnaire (FAQ) at baseline visit.

Results: A total of 676 participants from ADNI and 4592 participants from NACC were identified. After removing incomplete data, 665 ADNI (mean age [SD]: 73.44 [6.90]; 293 [44.1%] female; 374 stable and 291 progressive) and 3657 NACC participants (mean age [SD]: 70.96 [10.03]; 2405 [65.8%] female; 2445 stable and 1212 progressive) remained. Combinations of 4 measures were selected to generate 10 scorecards using FasterRisk algorithm, showing strong performance (area under the curve [AUC] = 0.868-0.892) in ADNI and remaining robust when validated in NACC (AUC = 0.795). The features were Category Animal $\leq$ 20 (2 points), Trail Making Test B $\leq$ 143 (-3 points), Logical Memory Delayed $\leq$ 3 (4 points), Logical Memory Delayed $\leq$ 8 (3 points), and FAQ $\leq$ 2 (-5 points). The probable AD risk corresponded to total

points: 7.4% (–8), 25.3% (–4), 50% (–1), 74.7% (2), and > 90% ($\geq$ 6). We refer to this model as the (F)unctioning, (LA)nguage, (M)emory, and (E)xecutive functioning or FLAME scorecard.

Conclusions and Relevance: Our findings highlight the potential to predict AD development using obtainable information, allowing for applicability at integrated primary care. While our scope centers on AD, this foundation paves the way for other dementia types.

Keywords: Alzheimer's Disease; Interpretable Machine Learning; Integrated Primary Care; Cognition

Introduction

As the prevalence of Alzheimer's disease (AD) continues to rise, timely early detection becomes increasingly urgent. One potential barrier to timely diagnosis is the initial point of contact for many patients: their primary care physicians (PCPs). When individuals first notice memory-related issues, the first healthcare professional they typically go to is their PCPs. However, many PCPs may not feel confident in delivering a conclusive diagnosis, resulting in delays in obtaining appropriate care.[1,2] Additionally, even if PCPs refer their patients for further evaluations, the waitlist for specialists is long, and patients may miss a critical window for early interventions or mitigations of risk factors for AD. Therefore, the solution lies in bridging this diagnostic gap at the primary care level by developing an easily administered and interpretable method to screen for AD risk and involving behavioral health consultants (BHCs), typically clinical psychologists, to carry out the screening and collaborate with PCPs to mitigate the risk factors.

The advancement of machine learning models offers a vast avenue for aiding the diagnostic process due to their speed and data-driven decisions that often excel in comparison to humans.[3] Recent efforts to develop machine learning models to assist clinicians in identifying early-stage AD have demonstrated robust accuracy.[4,5] However, the use of these models has raised important issues pertaining to the needs for inaccessible data and interpretability due to their "black box" nature in generating the output, potentially leading to a lack of trust in the outputs from clinicians.[6,7] Interpretable machine learning models (IML; i.e., white-box approach), on the other hand, do not suffer from the same issues as it provides the "why" of outputs, offering insights into how specific features contribute to predictions and allowing for transparent and understandable decision-making processes, which has been shown to promote trust between clinicians and the machine learning outputs.[8]

In this study, we developed risk scores that were presented in a scorecard model to assess the risk of developing AD using the FasterRisk algorithm. Risk scores are predictive models that have been used in various fields, including medicine, to aid decision-making processes through basic mathematical calculation.[9–11] The FasterRisk algorithm is a recent advancement that significantly improves the creation of high-quality risk scores that exceeds the performance of traditional methods, such as rounding logistic regression coefficients or non-data-driven approaches.[12] We selected the following variables to develop the scorecards due to their practicability and comprehensive representation of factors influencing AD: demographic information, cognitive tests from various domains, and daily functioning. We designed the scorecards to help inform clinicians of the probable risk of developing AD based on a patient's presentation, aiding in decisions about when to refer patients to specialists and involving BHCs to initiate interventions while the patients are on the waitlist for specialists. We predicted that our framework could generate a scoring system with robust predictive power using accessible variables.

Materials and methods

Participants

We included data from 676 baseline visits from several Alzheimer's Disease Neuroimaging Initiative database (ADNI 1, 2, GO, and 3; adni.loni.usc.edu) as of August 2023 and 4592 baseline visits from the National Alzheimer's Coordinating Center (NACC) Uniform Data Set (UDS) extracted in March 2024. ADNI was launched in 2003 as a public-private partnership, led by Principal Investigator Michael W. Weiner, MD. The primary goal of ADNI has been to test whether serial magnetic resonance imaging (MRI), positron emission

tomography (PET), other biological markers, and clinical and neuropsychological assessment can be combined to measure the progression of mild cognitive impairment (MCI) and early AD.

ADNI's broader criteria include age 55-90, a minimum of 6 years of education, consistent medication for the past 4 weeks, Hachinski scale < 4 (to rule out vascular dementia), and Geriatric Depression Scale < 6; more information can be found www.adni-info.org. We conducted performance validation of the model using an independent cohort, NACC UDS, which is a multi-site study that has been collecting data from Alzheimer's Disease Research Centers (ADRCs) since 2005. In general, NACC UDS inclusion criteria for enrollment include an absence of neurological disorders, severe mental illness, learning disability, and substance abuse. The data consists of participants' demographics, cognitive status, cognitive performance, and other clinical history and information; more information can be found www.naccdata.org.

We included participants in our analysis who were classified by ADNI and NACC as having normal cognition (NC) or amnestic MCI. Participants were divided into two groups: stable and progressive. The stable group consisted of individuals who remained at the same diagnosis level over time. The progressive group included those who developed AD at later visits. Specifically, participants who progressed from aMCI to AD or NC to AD were placed in the progressive group. To be included in the stable group, the sample had to contain data indicating at least 3 years of aMCI and 8 years of NC. Individuals who reverted to aMCI from NC were not included in the analysis. We excluded participants with non-amnestic MCI or those who developed other types of dementia. Participants with invalid or missing values were identified and removed from both ADNI (n = 11; 1.63%) and NACC (n = 1219; 25%).

Neuropsychological Tests and Functioning

We selected a range of neuropsychological tests that tapped into a variety of cognitive domains, such as attention, executive function, memory (short-term and long-term), and language/verbal fluency. The selected tests were the Logical Memory Immediate (LIMM), Logical Memory Delayed (LDEL), Category Animal (CATANIMSC), Trail Making Test A (TMT A), and Trail Making Test B (TMT B). These tests were selected because they were administered across all ADNI and NACC cohorts. Additionally, we included the instrumental activities of daily living measured with the Functional Activities Questionnaire (FAQ). Some adjustments were made for the NACC dataset because some tests were replaced or not administered. For example, the later NACC cohorts (UDS v.3) completed Craft Story 21 instead of Logical Memory (UDS v.2). We converted the test scores to match Logical Memory based on the conversion table provided by NACC crosswalk study.[13]

Data Preprocessing

To prepare the data for analysis, we converted categorical variables into numerical representations through Scikit-learn Labelencoder. The next preprocessing step was applying binarization using the FasterRisk built-in binarization module to convert the features from continuous into binary features (Figure 1). This ensures the proper input data format for the algorithm.

Risk Scorecard Generation and Optimization

We employed the FasterRisk algorithm to develop an interpretable, point-based risk scorecard, a model type widely used for transparent clinical decision-making.[12] Specific risk factors are assigned simple, whole-number points (e.g., +5 points for a given condition). A clinician sums up these points to calculate a patient's total risk score, which corresponds to a risk percentage, making each decision transparent and easy to explain. The algorithm then constructs

risk scores through an iterative process. It starts by identifying the single most predictive risk factor to form a base model. Then, it incrementally adds new factors one by one, keeping only those that improve the model's performance. The algorithm also explores other combinations by swapping existing features with other promising candidates. This approach was selected because it significantly outperforms RiskSlim, a previous state-of-the-art model for finding risk scores.[14] The scorecard's complexity is controlled by a 'sparsity' parameter, k, which dictates the exact number of predictive features included. To find the optimal balance between simplicity and accuracy, we systematically tested models with 1 to 10 features using 5-fold cross-validation (Figure 1).

Final Model Evaluation

The final scorecard was trained on 80% of ADNI data and its performance was internally validated on the held-out 20% test set (Figure 1). To ensure the stability of our results, we generated and analyzed the top 10 scorecards produced by the algorithm. By doing this, we were able to examine the set of most consistently predictive features. The model's performance was evaluated using accuracy, where a 50% risk threshold determined classification, and also the Area Under the Receiver Operating Curve (AUC), which quantifies the model's overall ability to discriminate between classes. We validated the best scorecard out of the 10 generated scorecards on NACC dataset. All computations were performed on Python 3.12.3 and data preprocessing was done using Numpy 1.23.5.

<p align="center">**Results**</p>

Participant Characteristics

We included data from 665 ADNI participants (164 NC and 501 aMCI) and 3657 NACC participants (2846 NC and 810 aMCI) from 41 ADRCs at baseline visits. Over time, 43.8% ($n =$

291) of ADNI and 33.1% (*n* = 1212) of NACC participants were diagnosed with AD. As shown

in Table 1, ADNI participants were 44.1% female, with a mean age of 73.44 years and an

average of 16.15 years of education, and NACC participants were 65.80% female, with a mean

age of 73.44 years and an average of 16.15 years of education. There were statistically

significant group differences between stable and progressive in ADNI (age at *p* < 0.01 and

education at *p* < 0.05) and NACC (age and education at *p* < 0.001). The average transitions for

the aMCI-AD and the NC-AD groups are 1.09 years and 7.10 years for ADNI and 3.02 years and

7.20 years for NACC.

Group Differences in Functioning and Cognitive Performance at Baseline

We compared group differences in cognitive performance and daily functioning using

analysis of covariance (ANCOVA) between stable and progressive groups. The FAQ was

adjusted for age, and cognitive measures were adjusted for age and education. All measures

showed statistical differences between stable and progressive groups in both ADNI and NACC

(*p* < 0.001). The progressive groups showed lower performance and scores in verbal

fluency/language, TMT A, TMT B, immediate and delayed memory, and daily functioning.

Higher values on TMT A and TMT B indicate worse performance.

Alzheimer Prediction Risk Score

Based on the FasterRisk algorithm, a sparsity level of 5 was selected for the most optimal

combination for the generation of the final scorecards to predict AD development. Ten

scorecards were generated with a test AUC ranging from 0.868 to 0.892 and accuracy between

81.20% and 84.21% (Supplementary Figure 1). The scorecard with the highest test AUC (0.892)

shown in Table 2 represents Category Animal ≤ 20 (2 points), Trail Making Test B ≤ 143 (-3

points), Logical Memory Delayed ≤ 3 (4 points), Logical Memory Delayed ≤ 8 (3 points), and

FAQ $\leq$ 2 (-5 points). Positive points contribute to increased total score of the scorecard, while negative points decrease it. The total score corresponds to the risk percentage shown in Table 2. The probable AD development risk was 7.4% for a total score of -8, 19.1% for a score of -5, 32.7% for a score of -3, 50% for a score of -1, 74.7% for a score of 2, 85.8% for a score of 4, and greater than 90% for a score of 6, 7, or 9 (Table 2). We refer to this scorecard model as the FLAME scorecard, a mnemonic representing functioning, language, memory, and executive function. We validated this scorecard with NACC dataset, yielding a test AUC of 0.795, suggesting good generalizability of the model.

Discussion

Our study presents a novel approach to predicting the risk of developing AD that offers potential to be applied in integrated primary care settings by employing a set of obtainable variables, including demographics, daily functioning, and cognitive performance. By utilizing the FasterRisk algorithm, we generated ten scorecards, each achieved high predictive accuracy (81.20%–84.21%) and strong discriminative performance (AUC = 0.868–0.892), reflecting a robust balance between sensitivity and specificity in identifying individuals at risk of developing AD. We further validated one of the ten scorecards on a separate cohort, indicating good generalizability. Additionally, the FLAME scorecard compared favorably to established machine learning models for predicting AD development,[15,16] and compared similarly or slightly below previously studied interpretable machine learning models incorporating biological (AUC = 0.86) and neuroimaging data (accuracy = 87.5%).[17,18]

A key advantage of the FLAME scorecard lies in its potential clinical practicality at integrated primary care settings. We selected variables that can be implemented and fit in one appointment with a behavior health consultant (BHC), which typically lasts approximately 30

minutes. The BHC would also have sufficient time to gather brief information related to patients'

concerns and general mental wellbeing. Following this assessment, the BHCs can input the

results into the scorecard and collaborate with the PCPs to determine appropriate next steps,

which may vary depending on the findings and presenting concerns. For instance, the PCPs may

refer patients to neurologists or other specialists, start a pharmacological treatment to slow down

memory decline (e.g., acetylcholinesterase inhibitors), reduce the use of anticholinergic

medications, order p-tau 217 blood test, or recommend patients to make follow-up appointments

with the BHCs to address behavioral factors that could mitigate memory decline (e.g., sleep

hygiene, engaging in physical activity, medication adherence, mood symptoms). BHCs may also

address safety concerns (e.g., driving, living arrangement) and make appropriate

recommendations. These services could serve as valuable interim interventions while patients are

placed on the waitlist for specialists.

Another advantage of the FLAME scorecard lies in its interpretability, which provides

clinicians with a clear explanation of the specific features influencing its predictions. This

transparency promotes human-computer interaction, in this case, trust between clinicians and the

machine learning outputs.[8] Furthermore, these scorecards offer flexibility in their implementation,

which allows clinicians to incorporate their expertise into the scorecards when predicting the risk

of AD development. For example, on the FLAME scorecard, having a FAQ score of 2 or below

would decrease the total by 5 points. If a patient indicates that the reason for requiring assistance

on some activities is due to physical limitations rather than decreased mental capacity, then the

clinician can consider this context when evaluating the predicted risk of AD. The balance

between performance, interpretability, and flexibility positions our scorecard as a promising tool

for practical clinical application, where understanding the rationale behind predictions is paramount for effective and informed decision support.

There are some limitations in our study. The scorecards generated in this study are only applicable to typical AD with an amnestic profile. Recent reviews highlighted that a subset of AD populations presents an atypical profile, such as deficits in executive functioning, language, visuospatial, and other non-memory domains, and this subset is typically younger than those who develop typical AD[19,20]. This scorecard might also be limited to discern typical AD presentation from limbic predominant age-related TDP-43 encephalopathy (LATE) due to similarity in their cognitive profiles.[21,22] Future studies including participants with atypical profiles or LATE are necessary to better inform PCPs of which follow-up tests to order to help confirm their diagnosis, which will cut down some costs compared to sending the patients to all tests/procedures. Additionally, the demographic composition of the ADNI and NACC samples, predominantly White and highly educated individuals, highlights the need for further validation in more diverse populations to ensure the generalizability of our findings. Lastly, it was beyond the scope of our study to determine a definitive cutoff value for clinical decision-making.

Our study lays the groundwork for a more accessible and population-wide approach to screening for Alzheimer's disease. Moving forward, we aim to collaborate with primary care physicians and behavioral health consultants to collect both qualitative and quantitative data on the feasibility and potential impact of implementing these scorecards in routine clinical practice. This collaboration will provide valuable insights into the practical challenges and opportunities for integrating our tool into the healthcare system.

Conclusions

Our study generated a robust scoring system for predicting the likelihood of developing Alzheimer's disease using accessible and cost-efficient variables through interpretable machine learning. This framework's interpretability may aid clinicians in integrated primary care settings in providing early detection to their patients, including those residing in resource-constrained areas.

Acknowledgments

Data collection and sharing for this project was funded by the Alzheimer's Disease Neuroimaging Initiative (ADNI) (National Institutes of Health Grant U01 AG024904) and DOD ADNI (Department of Defense award number W81XWH-12-2-0012). ADNI is funded by the National Institute on Aging, the National Institute of Biomedical Imaging and Bioengineering, and through generous contributions from the following: AbbVie, Alzheimer's Association; Alzheimer's Drug Discovery Foundation; Araclon Biotech; BioClinica, Inc.; Biogen; Bristol-Myers Squibb Company; CereSpir, Inc.; Cogstate; Eisai Inc.; Elan Pharmaceuticals, Inc.; Eli Lilly and Company; EuroImmun; F. Hoffmann-La Roche Ltd and its affiliated company Genentech, Inc.; Fujirebio; GE Healthcare; IXICO Ltd.; Janssen Alzheimer Immunotherapy Research & Development, LLC.; Johnson & Johnson Pharmaceutical Research & Development LLC.; Lumosity; Lundbeck; Merck & Co., Inc.; Meso Scale Diagnostics, LLC.;☐ NeuroRx Research; Neurotrack Technologies; Novartis Pharmaceuticals Corporation; Pfizer Inc.; Piramal Imaging; Servier; Takeda Pharmaceutical Company; and Transition Therapeutics. The Canadian Institutes of Health Research is providing funds to support ADNI clinical sites in Canada. Private sector contributions are facilitated by the Foundation for the National Institutes of Health (www.fnih.org). The grantee organization is the Northern California Institute for Research and Education, and the study is coordinated by the Alzheimer's Therapeutic Research Institute at the University of Southern California. ADNI data are disseminated by the Laboratory for Neuro Imaging at the University of Southern California.☐

The NACC database is funded by NIA/NIH Grant U24 AG072122. NACC data are contributed by the NIA-funded ADRCs: P30 AG062429 (PI James Brewer, MD, PhD), P30 AG066468 (PI Oscar Lopez, MD), P30 AG062421 (PI Bradley Hyman, MD, PhD), P30

AG066509 (PI Thomas Grabowski, MD), P30 AG066514 (PI Mary Sano, PhD), P30 AG066530 (PI Helena Chui, MD), P30 AG066507 (PI Marilyn Albert, PhD), P30 AG066444 (PI David Holtzman, MD), P30 AG066518 (PI Lisa Silbert, MD, MCR), P30 AG066512 (PI Thomas Wisniewski, MD), P30 AG066462 (PI Scott Small, MD), P30 AG072979 (PI David Wolk, MD), P30 AG072972 (PI Charles DeCarli, MD), P30 AG072976 (PI Andrew Saykin, PsyD), P30 AG072975 (PI Julie A. Schneider, MD, MS), P30 AG072978 (PI Ann McKee, MD), P30 AG072977 (PI Robert Vassar, PhD), P30 AG066519 (PI Frank LaFerla, PhD), P30 AG062677 (PI Ronald Petersen, MD, PhD), P30 AG079280 (PI Jessica Langbaum, PhD), P30 AG062422 (PI Gil Rabinovici, MD), P30 AG066511 (PI Allan Levey, MD, PhD), P30 AG072946 (PI Linda Van Eldik, PhD), P30 AG062715 (PI Sanjay Asthana, MD, FRCP), P30 AG072973 (PI Russell Swerdlow, MD), P30 AG066506 (PI Glenn Smith, PhD, ABPP), P30 AG066508 (PI Stephen Strittmatter, MD, PhD), P30 AG066515 (PI Victor Henderson, MD, MS), P30 AG072947 (PI Suzanne Craft, PhD), P30 AG072931 (PI Henry Paulson, MD, PhD), P30 AG066546 (PI Sudha Seshadri, MD), P30 AG086401 (PI Erik Roberson, MD, PhD), P30 AG086404 (PI Gary Rosenberg, MD), P20 AG068082 (PI Angela Jefferson, PhD), P30 AG072958 (PI Heather Whitson, MD), P30 AG072959 (PI James Leverenz, MD).

AW was supported by an NIH grant from NIGMS and OD 1P20GM152280.

Conflict of Interest

The authors have no conflict of interest to report.

References

1. Bradford A, Kunik ME, Schulz P, Williams SP, Singh H. Missed and Delayed Diagnosis of Dementia in Primary Care: Prevalence and Contributing Factors. *Alzheimer Disease & Associated Disorders*. 2009;23(4):306. doi:10.1097/WAD.0b013e3181a6bebc

2. Koch T, Iliffe S, the EVIDEM-ED project. Rapid appraisal of barriers to the diagnosis and management of patients with dementia in primary care: a systematic review. *BMC Family Practice*. 2010;11(1):52. doi:10.1186/1471-2296-11-52

3. Kumar Y, Koul A, Singla R, Ijaz MF. Artificial intelligence in disease diagnosis: a systematic literature review, synthesizing framework and future research agenda. *J Ambient Intell Human Comput*. 2023;14(7):8459-8486. doi:10.1007/s12652-021-03612-z

4. Helaly HA, Badawy M, Haikal AY. Deep Learning Approach for Early Detection of Alzheimer's Disease. *Cogn Comput*. 2022;14(5):1711-1727. doi:10.1007/s12559-021-09946-2

5. Kavitha C, Mani V, Srividhya SR, Khalaf OI, Tavera Romero CA. Early-Stage Alzheimer's Disease Prediction Using Machine Learning Models. *Front Public Health*. 2022;10. doi:10.3389/fpubh.2022.853294

6. LeCun Y, Bengio Y, Hinton G. Deep learning. *Nature*. 2015;521(7553):436-444. doi:10.1038/nature14539

7. Friedman JH. Greedy function approximation: A gradient boosting machine. *The Annals of Statistics*. 2001;29(5):1189-1232. doi:10.1214/aos/1013203451

8. Nasarian E, Alizadehsani R, Acharya UR, Tsui KL. Designing interpretable ML system to enhance trust in healthcare: A systematic review to proposed responsible clinician-AI-collaboration framework. *Information Fusion*. 2024;108:102412. doi:10.1016/j.inffus.2024.102412

9. Moreno RP, Metnitz PGH, Almeida E, et al. SAPS 3--From evaluation of the patient to evaluation of the intensive care unit. Part 2: Development of a prognostic model for hospital mortality at ICU admission. *Intensive Care Med*. 2005;31(10):1345-1355. doi:10.1007/s00134-005-2763-5

10. Six AJ, Backus BE, Kelder JC. Chest pain in the emergency room: value of the HEART score. *Neth Heart J*. 2008;16(6):191-196.

11. Than M, Flaws D, Sanders S, et al. Development and validation of the Emergency Department Assessment of Chest pain Score and 2 h accelerated diagnostic protocol. *Emerg Med Australas*. 2014;26(1):34-44. doi:10.1111/1742-6723.12164

12. Liu J, Zhong C, Li B, Seltzer M, Rudin C. FasterRisk: Fast and Accurate Interpretable Risk Scores. *arXiv*. Preprint posted online October 11, 2022. Accessed June 14, 2024. http://arxiv.org/abs/2210.05846

13. Monsell SE, Dodge HH, Zhou XH, et al. Results from the NACC Uniform Data Set neuropsychological battery Crosswalk Study Running head: Neuropsychological Battery Crosswalk Study Results. *Alzheimer Dis Assoc Disord*. 2016;30(2):134-139. doi:10.1097/WAD.0000000000000111

14. Ustun B, Rudin C. Learning Optimized Risk Scores. *Journal of Machine Learning Research*. 2019;20(150):1-75.

15. Fraser KC, Meltzer JA, Rudzicz F. Linguistic Features Identify Alzheimer's Disease in Narrative Speech. *Journal of Alzheimer's Disease*. 2016;49(2):407-422. doi:10.3233/JAD-150520

16. Ye J, Farnum M, Yang E, et al. Sparse learning and stability selection for predicting MCI to AD conversion using baseline ADNI data. *BMC Neurology*. 2012;12(1):46. doi:10.1186/1471-2377-12-46

17. Vlontzou ME, Athanasiou M, Dalakleidi KV, Skampardoni I, Davatzikos C, Nikita K. A comprehensive interpretable machine learning framework for mild cognitive impairment and Alzheimer's disease diagnosis. *Sci Rep*. 2025;15(1):8410. doi:10.1038/s41598-025-92577-6

18. Das D, Ito J, Kadowaki T, Tsuda K. An interpretable machine learning model for diagnosis of Alzheimer's disease. *PeerJ*. 2019;7:e6543. doi:10.7717/peerj.6543

19. Graff-Radford J, Yong KXX, Apostolova LG, et al. New insights into atypical Alzheimer's disease in the era of biomarkers. *The Lancet Neurology*. 2021;20(3):222-234. doi:10.1016/S1474-4422(20)30440-3

20. Shea YF, Pan Y, Mak HKF, et al. A systematic review of atypical Alzheimer's disease including behavioural and psychological symptoms. *Psychogeriatrics*. 2021;21(3):396-406. doi:10.1111/psyg.12665

21. Kapasi A, Yu L, Boyle PA, Barnes LL, Bennett DA, Schneider JA. Limbic-predominant age-related TDP-43 encephalopathy, ADNC pathology, and cognitive decline in aging. *Neurology*. 2020;95(14):e1951-e1962. doi:10.1212/WNL.0000000000010454

22. Serio MA, Dethloff DR, Curry GC, Beesley ML, Shekoohi S, Kaye AD. Emerging Concepts in Diagnosis, Pathologic Features, and Treatment of Limbic-Predominant Amnestic Neurodegenerative Syndrome (LANS): A Narrative Review. *Adv Ther*. Published online August 29, 2025. doi:10.1007/s12325-025-03337-x

Table 1.

Table of ADNI and NACC demographic, cognition, and functioning by groups

Variables (ADNI)	Stable (*n* = 374)	Progressive (*n* = 291)	All (*n* = 665)
Age (years)	72.8 (6.74) **	74.27 (7.03) **	73.44 (6.90)
Education (years)	16.34 (2.72) *	15.89 (2.82) *	16.15 (2.77)
Female (%)	169 (45.2%)	124 (42.6%)	293 (44.1%)
Race (White%)	350 (93.6%)	274 (94.2%)	624 (93.83%)
FAQ	0.98 (2.56) ***	4.90 (5.00) ***	2.70 (4.29)
Category Animal	19.91 (5.43) ***	16.11 (4.94) ***	18.25 (5.55)
TMT A	35.29 (14.42) ***	45.9 (24.2) ***	39.93 (20.01)
TMT B	87.91 (46.57) ***	134.63 (73.83) ***	108.36 (64.32)
LM immediate	11.66 (3.84) ***	7.63 (3.88) ***	9.90 (4.35)
LM delayed	9.83 (4.21) ***	4.59 (4.13) ***	7.54 (4.92)
Variables (NACC)	Stable (*n* = 2445)	Progressive (*n* = 1212)	All (*n* = 3657)
Age (years)	68.16 (9.55) ***	76.61 (8.49) ***	70.96 (10.03)
Education (years)	16.61 (6.15) ***	15.85 (5.6) ***	16.36 (5.98)
Female (%)	1646 (67.3%)	759 (62.6%)	2405 (65.8%)
Race (White%)	2074 (84.8%)	1075 (88.7%)	3149 (86.11%)
FAQ	0.26 (1.33) ***	2.59 (4.66) ***	1.03 (3.1)
Category Animal	21.87 (5.66) ***	17.70 (5.66) ***	20.49 (5.99)
TMT B	80.35 (79.88) ***	140.73 (168.42) ***	100.36 (120.29)
LM delayed	12.91 (4.19) ***	7.68 (5.26) ***	11.18 (5.19)

ADNI = Alzheimer's Disease Neuroimaging Initiative; NACC = National Alzheimer's Coordinating Center; FAQ = Functional Activities Questionnaire; TMT A = Trail Making Test A; TMT B = Trail Making Test B; LM = Logical Memory

Differences between stable and progressive, * $p < 0.05$, **$p<0.01$, ***$p<0.001$□□

Cognitive measures were adjusted for age and education.

FAQ was adjusted for age.

Table 2.

Scorecard with the highest AUC and Risk Score to assess AD development probability

Variables	Points	
1. Category Animal $\leq$ 20	+ 2 points	…
2. Trail Making Test B $\leq$ 143	-3 points	+ …
3. Logical Memory Delayed $\leq$ 3	+ 4 points	+ …
4. Logical Memory Delayed $\leq$ 8	+ 3 points	+ …
5. FAQ $\leq$ 2	-5 points	+
	SCORE	=

LDEL = Logical Memory delayed recall; Test; FAQ=Functional Activities Questionnaire

Score	-8	-6	-4	-3	-1	1	2	4	6	9
Risk (%)	7.4	14.2	25.3	32.7	50.0	67.3	74.7	85.8	92.6	97.3

Figure 1.

Pipeline of conducting FasterRisk algorithm to generate the FLAME scorecard and its clinical application.