A new preprocedural predictive risk model for post-endoscopic retrograde cholangiopancreatography pancreatitis: The SuPER model

Mitsuru Sugimoto[1]*, Tadayuki Takagi[1], Tomohiro Suzuki[2], Hiroshi Shimizu[2], Goro Shibukawa[3], Yuki Nakajima[3], Yutaro Takeda[4], Yuki Noguchi MD[4], Reiko Kobayashi MD[4], Hidemichi Imamura[4], Hiroyuki Asama[5], Naoki Konno[5], Yuichi Waragai[6], Hidenobu Akatsuka[7], Rei Suzuki[1], Takuto Hikichi[8], and Hiromasa Ohira[1]

[1]Department of Gastroenterology, Fukushima Medical University, School of Medicine, Fukushima, Japan

[2]Department of Gastroenterology, Fukushima Rosai Hospital, Iwaki, Japan

[3]Department of Gastroenterology, Aizu Medical Center, Fukushima Medical University, Aizu, Japan

[4]Department of Gastroenterology, Ohta Nishinouchi Hospital, Koriyama, Japan

[5]Department of Gastroenterology, Fukushima Redcross Hospital, Fukushima, Japan

[6]Department of Gastroenterology, Soma General Hospital, Soma, Japan

[7]Department of Gastroenterology, Saiseikai Fukushima General Hospital, Fukushima, Japan

[8]Department of Endoscopy, Fukushima Medical University Hospital, Fukushima, Japan

* **For correspondence**

kita335@fmu.ac.jp

Abstract

Background: Post–endoscopic retrograde cholangiopancreatography (ERCP) pancreatitis (PEP) is a severe and deadly adverse event following ERCP. The ideal method for predicting PEP risk before ERCP has yet to be identified. We aimed to establish a simple PEP risk score model (SuPER model: Support for PEP Reduction) that can be applied before ERCP.

Methods: This multicenter study enrolled 2074 patients who underwent ERCP. Among them, 1037 patients each were randomly assigned to the development and validation cohorts. In the development cohort, the risk score model for predicting PEP was established via logistic regression analysis. In the validation cohort, the performance of the model was assessed.

Results: In the development cohort, five PEP risk factors that could be identified before ERCP were extracted and assigned weights according to their respective regression coefficients: -2 points for pancreatic calcification, 1 point for female sex, and 2 points for intraductal papillary mucinous neoplasm, a native papilla of Vater, or the pancreatic duct procedures (treated as "planned pancreatic duct procedures" for calculating the score before ERCP). The PEP occurrence rate was 0% among low-risk patients (≤ 0 points), 5.5% among moderate-risk patients (1 to 3 points), and 20.2% among high-risk patients (4 to 7 points). In the validation cohort, the C statistic of the risk score model

was 0.71 (95% CI 0.64–0.78), which was considered acceptable. The PEP risk

classification (low, moderate, and high) was a significant predictive factor for PEP that

was independent of intraprocedural PEP risk factors (precut sphincterotomy and

inadvertent pancreatic duct cannulation) (OR 4.2, 95% CI 2.8–6.3; $P < 0.01$).

Conclusions: The PEP risk score allows an estimation of the risk of PEP prior to ERCP,

regardless of whether the patient has undergone pancreatic duct procedures. This simple

risk model, consisting of only five items, may aid in predicting and explaining the risk

of PEP before ERCP and in preventing PEP by allowing selection of the appropriate

expert endoscopist and useful PEP prophylaxes.

Introduction

Endoscopic retrograde cholangiopancreatography (ERCP) is widely performed as an important diagnostic and therapeutic procedure for pancreaticobiliary diseases. Among endoscopic procedures, ERCP-related procedures are relatively risky. The high-risk adverse events of ERCP include duodenal perforation and bleeding after endoscopic sphincterotomy (EST) and post-ERCP pancreatitis (PEP). The rate of PEP occurrence is 3.1–13.0% (*Andriulli et al. , 2007, Freeman et al. , 1996, Glomsaker et al. , 2013, Katsinelos et al. , 2014, Kochar et al. , 2015, Loperfido et al. , 1998*). PEP can even become life-threatening. The fatality rate of PEP is 0.1–0.7% (*Andriulli et al. , 2007, Kochar et al. , 2015*). Therefore, the decision to perform ERCP should be made carefully, considering each patient's risk factors for PEP.

To predict an individual patient's PEP risk, six scoring systems have been devised (*Archibugi et al. , 2023, Chiba et al. , 2021, DiMagno et al. , 2013, Friedland et al. , 2002, Fujita et al. , 2021, Zheng et al. , 2020*). The first risk scoring system for PEP occurrence was established in 2002. In that study, pain during the procedure, pancreatic duct cannulation, a history of PEP, and the number of cannulation attempts were identified as risk factors for PEP. After the first scoring system was reported, each new scoring system used risk factors that were extracted via multivariate analyses. These included various patient characteristics before ERCP and postprocedural risk factors.

Postprocedural risk factors, such as precut sphincterotomy, procedure time, and difficult

cannulation, have been proposed, but it is difficult to predict these risk factors and

determine the PEP risk before ERCP. Thus, a new prediction scoring system for PEP

before ERCP is desirable. If the risk of PEP can be predicted before ERCP, then the

expert endoscopist can perform ERCP from the start, and high-PEP-risk procedures (for

example, precut sphincterotomy, multiple cannulation attempts, and inadvertent

pancreatic duct cannulation) can be avoided (*Testoni et al. , 2010, Wang et al. , 2009*). If

biliary cannulation without the use of at least one high-PEP-risk procedure is difficult,

other treatments (for example, percutaneous transhepatic biliary drainage (PTBD) or

endoscopic ultrasound (EUS)-guided biliary drainage (EUS-BD)) could be considered.

Therefore, we aimed to establish a PEP prediction model using only risk

factors that can be gathered before ERCP. Our model was developed and validated with

multicenter data from Japan.

Methods

We performed a multicenter retrospective study at six institutions in Japan. This study

was approved by the institutional review board of Fukushima Medical University and

that of each partner medical institution. All patients agreed to undergo ERCP after

providing written consent.

Patients

Among 2,176 patients who underwent ERCP between November 2020 and October

2022, 2,074 were included in this study. The other 102 patients were excluded for the

following reasons: past history of choledochojejunostomy, acute pancreatitis,

choledochoduodenal fistula, difficulty finding the Vater papilla, past history of

pancreatojejunostomy, or past history of pancreatogastrostomy (**Figure 1**). The

predictive PEP score is not necessary for the excluded patients. The reasons were as

follows. Biliary duct cannulation was not attempted in patients for whom it was difficult

to identify the Vater papilla. The biliary tract was separated from the pancreas in

patients with a past history of choledochojejunostomy, pancreatojejunostomy, or

pancreatogastrostomy. PEP risk was thought to be clearly low in these patients and

patients who underwent bile duct cannulation via the choledochoduodenal fistula. PEP

diagnosis is difficult in patients with acute pancreatitis, whose diagnosis is currently in

progress.

Study design

We randomly sampled 50% of the patients as the development cohort and 50% as the validation cohort (**Figure 1**). In the development cohort, we established a risk scoring system for predicting PEP before ERCP, which was named the support for PEP reduction model (SuPER model). The validation cohort was used to confirm the effectiveness of the scoring system. PEP diagnosis and severity were assessed according to Cotton's criteria (*Cotton et al. , 1991*). Patients who experienced abdominal pain and had hyperamylasemia (more than three times the normal upper limit) at least 24 hours after ERCP were diagnosed with PEP. Mild PEP was defined as pancreatitis that required prolongation of the planned hospitalization by 2–3 days. Moderate PEP was defined as pancreatitis that required 4–10 days of hospitalization. Severe PEP was defined as pancreatitis that required more than 10 days of hospitalization or intervention or hemorrhagic pancreatitis, phlegmon, or pseudocysts.

To establish the risk score, the risk factors for PEP were investigated via data from the development cohort. To determine the PEP risk score, factors that might be associated with PEP occurrence were investigated. To predict the PEP risk score before ERCP, factors related to patient characteristics and previously scheduled procedures, as reported in the Japanese guidelines for acute pancreatitis and PEP, were selected (*Itoi et al. , 2024, Takada et al. , 2022*). The patients' risk factors included age < 50 years,

female sex, a past history of pancreatitis, a past history of PEP, a past history of

gastrectomy, pancreatic cancer, intraductal papillary mucinous neoplasm (IPMN), a

native papilla of Vater, absence of chronic pancreatitis (CP), normal serum bilirubin ($\leq$

1.2 mg/dl), and periampullary diverticulum (*Ding et al. , 2015, Freeman et al. , 2001,*

Freeman et al. , 1996, Fujita et al. , 2022, Fujita et al. , 2021, Masci et al. , 2003, Wang

et al. , 2009, Williams et al. , 2007, Zheng et al. , 2020). Pancreatic divisum was

excluded from the patient risk factor list because pancreatic divisum was observed in

only two patients. Pancreatic calcification and a diameter of the main pancreatic duct >

3 mm were considered to indicate CP (*Beyer et al. , 2023, Sarner and Cotton, 1984*).

These imaging findings were confirmed by CT, MRI, and EUS before ERCP. The CT

and MRI findings were reviewed by radiologists. IPMNs were diagnosed according to

the results of CT, MRI, and EUS. As pre-ERCP prophylaxes for PEP, protease inhibitors

(gabexate mesilate or nafamostat mesilate), intravenous hydration, and NSAID

suppositories have been used (*Fujita et al. , 2022*). As planned procedure-related risk

factors, EST, endoscopic papillary balloon dilation (EPBD), endoscopic papillary large

balloon dilation (EPLBD) using a $\geq$ 12 mm balloon catheter (*Itoi et al. , 2018*), biliary

stone removal, ampullectomy, biliary stent material (plastic stent, self-expandable

metallic stent (SEMS), or covered SEMS (CSEMS)), inside stent placement, and

procedures on the pancreatic duct were evaluated (*Freeman et al. , 2001, Freeman et al. , 1996, Harewood et al. , 2005, Kato et al. , 2022, Masci et al. , 2003, Masci et al. , 2001, Testoni et al. , 2010, Williams et al. , 2007*). A biliary stent above the Vater papilla was also assessed as a prophylactic measure against PEP (*Ishiwatari et al. , 2013*).

To demonstrate the independence of the established risk classification, the relationships between it and intraprocedural PEP risk factors (including precut sphincterotomy and inadvertent pancreatic duct cannulation) (*Testoni et al. , 2010, Wang et al. , 2009*) were investigated. Because of the retrospective nature of the data, the exact number of cannulations and the number of cannulation attempts were not available. Therefore, multiple cannulation attempts and a prolonged cannulation time could not be investigated as intraprocedural PEP risk factors.

Sample size

The primary aim of this study was to establish a PEP prediction model that could be used to calculate a risk score before ERCP. To construct a prediction model via logistic regression analysis, 10 events per explanatory variable were needed (*Wynants et al. , 2015*). Seven variables were evaluated in the development cohort, so 70 PEP patients were included. Five variables were evaluated in the validation cohort, so 50 PEP

patients were necessary. According to a previous systematic review, the rate of PEP

occurrence was 9.7% (*Kochar et al. , 2015*). Therefore, at least 722 and 521 patients

were included in the development and validation cohorts, respectively.

Statistical analysis

In the development cohort, univariate and multivariate logistic regression analyses were

performed to identify the risk factors for PEP. The factors that had a *p* value < 0.10 in

the univariate analysis were included in the multivariate analysis. To construct the

scoring system for PEP risk, the factors with $p < 0.10$ in the multivariate analysis were

ultimately included in the risk score model. The factors selected in the multivariate

analysis were assigned points according to the regression coefficient (each variable's

risk points = the ratio of the variable's regression coefficient/minimum regression

coefficient). The sum of the assigned points was calculated for each patient, and the

patients were classified into three groups (low risk, moderate risk, and high risk)

according to the expected rate of PEP occurrence (*Friedland et al. , 2002*). The risk

classification system (SuPER model) was also applied to the validation cohort.

With respect to both the development and validation cohorts, the effectiveness

of the risk score model was evaluated as follows. The correlations between the risk

score, risk classification and PEP occurrence were evaluated via the Cochran–Armitage

trend test. The predictive accuracy of the risk score was assessed via the C statistic. The

goodness of fit of the model was evaluated via the Hosmer□Lemeshow test. The

independence of the established risk classification from the unexpected intraprocedural

PEP risk factors was assessed via multivariate logistic regression analyses.

Patients with missing data for variables selected in the risk score model were

removed from the final cohort.

Statistical analyses were performed via EZR version 1.62 (Saitama Medical

Centre, Jichi Medical University, Saitama, Japan) and SPSS version 26.0 (IBM Corp.,

Armonk, NY, USA). A p value < 0.05 indicated statistical significance.

Results

Patient characteristics and ERCP outcomes in each cohort

The patient characteristics and ERCP outcomes of each cohort are shown in **Table 1**. A

total of 1037 patients were assigned to each of the development and validation cohorts,

including 70 (6.8%) and 64 (6.2%) patients diagnosed with PEP, respectively. Age,

history of pancreatitis, and procedures involving the pancreatic duct were significantly

different between the development cohort and the validation cohort. The pre-ERCP

prophylactic measures used at each hospital differed, and not all patients received

prophylaxis.

Construction of the PEP risk scoring system

According to the univariate analyses, age < 50 years, female sex, IPMN, a native papilla

of Vater, pancreatic calcification, EST, and procedures on the pancreatic duct had *p*

values < 0.10 (**Table 2**). According to the multivariate analysis, female sex, IPMN, a

native papilla of Vater, pancreatic calcification, and procedures on the pancreatic duct

had *p* values < 0.10. These factors were assigned risk points according to their

respective regression coefficients.

The risk score of each patient was calculated as the total number of risk points

and ranged from -2 to 7 points (**Table 3**). The risk score was found to be correlated with

PEP occurrence ($p < 0.01$, Cochran–Armitage trend test). The patients were classified as

low (≤ 0 points), moderate (1–3 points), or high risk (4–7 points) for PEP according to

the risk score. The PEP rates were 0% (0/327) among the low-risk patients, 5.5%

(27/492) among the moderate-risk patients, and 20.2% (39/193) among the high-risk

patients. The risk classification was correlated with PEP occurrence ($p < 0.01$,

Cochran–Armitage trend test).

The C statistic of the risk score model was sufficiently high at 0.77 (95%

confidence interval (CI) 0.72–0.82) (**Table 4**). The goodness of fit of the risk score

model was also confirmed by the Hosmer–Lemeshow test ($p = 0.59$).

Validation of the PEP risk scoring system

The risk score was associated with PEP occurrence in the validation cohort ($p < 0.01$,

Cochran–Armitage trend test) (**Table 3**). We found that 2.4% (8/331) of the patients at

low risk, 5.3% (27/513) of those at moderate risk, and 18.0% (29/161) of those at high

risk experienced PEP. The risk classification was also correlated with PEP occurrence in

the validation cohort ($p < 0.01$, Cochran–Armitage trend test).

The C statistic of the risk score was 0.71, which was also high in the validation

cohort (**Table 4**). The PEP risk score model showed good fitness according to the

Hosmer–Lemeshow test ($p = 0.40$). According to the above results, the preprocedural

PEP risk can be calculated, as shown in **Figure 2**. The pancreatic duct procedure was

treated as a "planned pancreatic duct procedure". Otherwise, we can calculate both the

PEP risk score, which involves the pancreatic duct procedure, and the score, which does

not involve the pancreatic duct procedure.

Risk classification and unexpected PEP risk factors

The relationships between the established risk classification and intraprocedural PEP

risk factors are shown in **Appendix 1—Table 1**. For all patients in the development

cohort and the validation cohort, the risk classification was significantly associated with

the occurrence of PEP. On the other hand, precut sphincterotomy and inadvertent

pancreatic duct cannulation were not significantly associated with the occurrence of

PEP.

Discussion

In this multicenter study, we aimed to establish a preprocedural PEP prediction model.

As a result, we created a risk scoring system (the SuPER model) using five items that

could be measured before performing ERCP. The term "pancreatic duct procedures"

was "planned pancreatic duct procedures" in the model. When unintended pancreatic

duct procedures are considered, we can calculate both the PEP risk score with

pancreatic duct procedures and that without pancreatic duct procedures. With this score,

PEP occurrence can be accurately predicted to some degree. Although the development

cohort and validation cohort had significantly different patient characteristics and ERCP

outcomes, the risk score was good in both cohorts. In addition, the established PEP risk classification was associated with PEP occurrence independent of unpredictable intraprocedural PEP risk procedures.

This risk scoring and classification of PEP has several advantages. First, the score is calculated using only five items, all of which can be easily assessed via medical interviews and imaging (for example, CT). One scoring system included sphincter of Oddi dysfunction (SOD) as a test item (*DiMagno et al. , 2013*). The diagnosis of SOD requires sphincter of Oddi manometry and fulfillment of the criteria for biliary pain, but sphincter of Oddi manometry is not widely used (*Cotton et al. , 2016*). The diagnostic criterion for biliary pain included 8 items, and that for SOD included 15 items. Among the items of the SuPER risk scoring system, pancreatic calcification was assigned -2 points. Its low weighting could be explained by the following. The international conceptual model of CP can be divided into four stages: acute pancreatitis–recurrent acute pancreatitis, early CP, established CP, and end-stage CP (*Whitcomb et al. , 2016*). Established CP patients have already passed the acute pancreatitis–recurrent acute pancreatitis course, and pancreatic calcification has been reported in established CP patients. Acinar dysfunction has also been observed in these patients (*Whitcomb et al. , 2016*). Therefore, patients with pancreatic calcification may have a lower incidence of

PEP.

Second, the SuPER risk score can be determined before the ERCP procedure,

as the established risk classification was found to be the sole significant factor

predicting the occurrence of PEP independent of intraprocedural PEP risk factors. As

described in the Background section, precut sphincterotomy, multiple cannulation

attempts, and a cannulation time greater than 10 minutes were identified as high-risk

factors that cannot be accounted for prior to ERCP (*Testoni et al. , 2010, Wang et al. ,*

2009). Although the established PEP risk classification was independent of the included

intraprocedural risk factors (precut sphincterotomy and inadvertent pancreatic duct

cannulation), detailed data on the number of cannulation attempts and the cannulation

time were not available. Therefore, to avoid intraoperative procedures associated with a

high risk of PEP occurrence, an expert endoscopist can initially perform ERCP for

high-PEP-risk patients. In addition, PEP prophylaxis can be administered beforehand

for high-PEP-risk patients. As effective prophylaxes for PEP, rectal NSAIDS and

pancreatic stent placement have been reported (*Akshintala et al. , 2021, Choi et al. ,*

2023, Elmunzer et al. , 2008, Koshitani et al. , 2022, Murray et al. , 2003, Sperna

Weiland et al. , 2022, Sugimoto et al. , 2019, Yang et al. , 2020). In this report, rectal

NSAID use was not identified as a significant factor preventing PEP. One reason for this

is that in past reports describing the use of rectal NSAIDs to prevent PEP, patients at

high risk for PEP were often treated. In contrast, this study included all patients who

underwent ERCP. Another reason might be the difference in dose. One hundred

milligrams of rectal diclofenac have been used in most past reports, whereas the

efficacy of low-dose rectal diclofenac (25–50 mg) for preventing PEP is under

discussion (*Maeda et al. , 2021, Okuno et al. , 2018, Otsuka et al. , 2012, Sakai et al. ,*

2023, Takaori et al. , 2021, Tomoda et al. , 2021). In this study, 12.5–50 mg of

diclofenac was used. In Japan, the approved diclofenac dose covered by insurance is 50

mg or less, with the dose typically being lower for elderly patients. Therefore,

diclofenac doses ranging from 12.5–50 mg were prescribed by the doctors depending on

the age and size of the patients. Pancreatic stent placement itself is one of the

procedures performed on the pancreatic duct and is a higher-risk procedure for PEP than

endoscopic biliary procedures without an approach to the pancreatic duct (**Appendix**

2—Table 2). Moreover, pancreatic stent placement has become a prophylactic treatment

for PEP in patients who have undergone pancreatography or wire placement to the

pancreatic duct (*Mazaki et al. , 2014, Sugimoto et al. , 2019*). As described above,

pancreatic stent placement was performed along with high-risk-PEP procedures (i.e.,

guidewire placement to the pancreatic duct or pancreatography); therefore, pancreatic

stent placement was grouped with the other endoscopic retrograde pancreatography

procedures as "procedures on the pancreatic duct". In a recent randomized trial

involving 1950 patients, the combination of rectal NSAIDs and prophylactic pancreatic

stents was more effective for preventing PEP than NSAIDs alone (*Elmunzer et al. ,*

2024). For high-PEP-risk patients with high scores, the combination of prophylactic

methods might be desirable.

This study has several limitations. First, the study was retrospective, and there

were missing data. However, the results reported are trustworthy. The percentage of

patients who did not meet the inclusion criteria was not greater than 5%, and the

percentage of missing data was not greater than 1%. As described in the Materials and

Methods section, patients with missing data for the variables selected in the risk score

model were removed from the final cohort. The reliability of the SuPER risk score

model was also statistically confirmed. Second, some factors cannot be assessed before

ERCP. Additional procedures can be conducted during ERCP, and unplanned

pancreatography is often performed in patients who are scheduled for endoscopic

cholangiography or biliary treatment. However, the established PEP risk classification

was independent of the included intraprocedural risk factors. A planned procedure for

accessing the pancreatic duct is listed in the SuPER risk model. Therefore, we can

predict the SuPER risk score and classification of patients regardless of whether they have undergone pancreatic duct procedures. Third, this study was performed in a single country. In the future, prospective validation studies over wider geographic regions are needed.

In conclusion, a simple and useful PEP scoring system (SuPER model) with only five clinical items was developed in this multicenter study. This scoring system may aid in predicting and explaining PEP risk and in selecting appropriate prophylaxes for PEP and endoscopic pancreatobiliary procedures for each patient.

Acknowledgments

We thank all the staff at the Department of Gastroenterology of Fukushima Medical University, the Department of Endoscopy of Fukushima Medical University Hospital, the Department of Gastroenterology of Fukushima Rosai Hospital, the Department of Gastroenterology of Aizu Medical Center, Fukushima Medical University, the Department of Gastroenterology of Ohtanishinouchi Hospital, Koriyama, the Department of Gastroenterology of Fukushima Redcross Hospital, the Department of Gastroenterology of Soma General Hospital, the Department of Gastroenterology of Saiseikai Fukushima General Hospital, and the Gastroenterology Ward of Fukushima

Medical University Hospital. We also thank American Journal Experts for providing

English language editing services.

Additional information

Competing interests

The authors declare that they have no competing interests to report.

Funding

None.

Author contributions

M.S. wrote the paper and designed and performed the research. T.T. wrote the paper and

designed and oversaw the research. T.S., H.S., G.S., Y.S., Y.N., Y.T., Y.N., R.K., H.I.,

H.A., N.K., Y.W., and H.A. performed the research. R.S. provided clinical advice. T.H.

supervised the report. H.O. supervised the reporting and writing of the paper. All the

authors read and approved the final manuscript.

Ethics

The study protocol was reviewed and approved by the Institutional Review Board of Fukushima Medical University (number 2453). The analysis used anonymous clinical data obtained after all the participants agreed to treatment with written consent; thus, patients were not required to provide informed consent for the study. The details of the study can be found on the homepage of Fukushima Medical University.

Data availability

The datasets generated and/or analyzed during the current study are available from the corresponding author upon reasonable request.

References

Akshintala VS, Sperna Weiland CJ, Bhullar FA, Kamal A, Kanthasamy K, Kuo A,
Tomasetti C, Gurakar M, Drenth JPH, Yadav D, Elmunzer BJ, Reddy DN,
Goenka MK, Kochhar R, Kalloo AN, Khashab MA, van Geenen EJM, Singh
VK. 2021. Non-steroidal anti-inflammatory drugs, intravenous fluids, pancreatic
stents, or their combinations for the prevention of post-endoscopic retrograde
cholangiopancreatography pancreatitis: a systematic review and network
meta-analysis. *Lancet Gastroenterol Hepatol* **6**:733-742. DOI:
10.1016/s2468-1253(21)00170-9

Andriulli A, Loperfido S, Napolitano G, Niro G, Valvano MR, Spirito F, Pilotto A,
Forlano R. 2007. Incidence rates of post-ERCP complications: a systematic
survey of prospective studies. *Am J Gastroenterol* **102**:1781-1788. DOI:
10.1111/j.1572-0241.2007.01279.x

Archibugi L, Ciarfaglia G, Cárdenas-Jaén K, Poropat G, Korpela T, Maisonneuve P,
Aparicio JR, Casellas JA, Arcidiacono PG, Mariani A, Stimac D, Hauser G, Udd
M, Kylänpää L, Rainio M, Di Giulio E, Vanella G, Lohr JM, Valente R, Arnelo
U, Fagerstrom N, De Pretis N, Gabbrielli A, Brozzi L, Capurso G, de-Madaria E.
2023. Machine learning for the prediction of post-ERCP pancreatitis risk: A

proof-of-concept study. *Digestive & Liver Disease* **55**:387-393. DOI: 10.1016/j.dld.2022.10.005

Beyer G, Kasprowicz F, Hannemann A, Aghdassi A, Thamm P, Volzke H, Lerch MM, Kühn JP, Mayerle J. 2023. Definition of age-dependent reference values for the diameter of the common bile duct and pancreatic duct on MRCP: a population-based, cross-sectional cohort study. *Gut* **72**:1738-1744. DOI: 10.1136/gutjnl-2021-326106, PMID: PMC10423481

Chiba M, Kato M, Kinoshita Y, Shimamoto N, Tomita Y, Abe T, Kanazawa K, Tsukinaga S, Nakano M, Torisu Y, Toyoizumi H, Sumiyama K. 2021. The milestone for preventing post-ERCP pancreatitis using novel simplified predictive scoring system: a propensity score analysis. *Surg Endosc* **35**:6696-6707. DOI: 10.1007/s00464-020-08173-4

Choi JH, Lee SH, Kim JS, Park N, Lee MH, Lee MW, Cho IR, Paik WH, Ryu JK, Kim YT. 2023. Combinatorial Effect of Prophylactic Interventions for Post-ERCP Pancreatitis among Patients with Risk Factors: A Network Meta-Analysis. *Gut Liver* **17**:814-824. DOI: 10.5009/gnl220268, PMID: PMC10502499

Cotton PB, Elta GH, Carter CR, Pasricha PJ, Corazziari ES. 2016. Rome IV. Gallbladder and Sphincter of Oddi Disorders. *Gastroenterology*. DOI:

10.1053/j.gastro.2016.02.033

Cotton PB, Lehman G, Vennes J, Geenen JE, Russell RC, Meyers WC, Liguory C,

Nickl N. 1991. Endoscopic sphincterotomy complications and their

management: an attempt at consensus. *Gastrointest Endosc* **37**:383-393. DOI:

10.1016/s0016-5107(91)70740-2

DiMagno MJ, Spaete JP, Ballard DD, Wamsteker EJ, Saini SD. 2013. Risk models for

post-endoscopic retrograde cholangiopancreatography pancreatitis (PEP):

smoking and chronic liver disease are predictors of protection against PEP.

Pancreas **42**:996-1003. DOI: 10.1097/MPA.0b013e31827e95e9, PMID:

PMC3701741

Ding X, Zhang F, Wang Y. 2015. Risk factors for post-ERCP pancreatitis: A systematic

review and meta-analysis. *Surgeon* **13**:218-229. DOI:

10.1016/j.surge.2014.11.005

Elmunzer BJ, Foster LD, Serrano J, Coté GA, Edmundowicz SA, Wani S, Shah R, Bang

JY, Varadarajulu S, Singh VK, Khashab M, Kwon RS, Scheiman JM,

Willingham FF, Keilin SA, Papachristou GI, Chak A, Slivka A, Mullady D,

Kushnir V, Buxbaum J, Keswani R, Gardner TB, Forbes N, Rastogi A, Ross A,

Law J, Yachimski P, Chen YI, Barkun A, Smith ZL, Petersen B, Wang AY,

Saltzman JR, Spitzer RL, Ordiah C, Spino C, Durkalski-Mauldin V. 2024. Indomethacin with or without prophylactic pancreatic stent placement to prevent pancreatitis after ERCP: a randomised non-inferiority trial. *Lancet* **403**:450-458. DOI: 10.1016/s0140-6736(23)02356-5, PMID: PMC10872215

Elmunzer BJ, Waljee AK, Elta GH, Taylor JR, Fehmi SM, Higgins PD. 2008. A meta-analysis of rectal NSAIDs in the prevention of post-ERCP pancreatitis. *Gut* **57**:1262-1267. DOI: 10.1136/gut.2007.140756

Freeman ML, DiSario JA, Nelson DB, Fennerty MB, Lee JG, Bjorkman DJ, Overby CS, Aas J, Ryan ME, Bochna GS, Shaw MJ, Snady HW, Erickson RV, Moore JP, Roel JP. 2001. Risk factors for post-ERCP pancreatitis: a prospective, multicenter study. *Gastrointest Endosc* **54**:425-434. DOI: 10.1067/mge.2001.117550

Freeman ML, Nelson DB, Sherman S, Haber GB, Herman ME, Dorsher PJ, Moore JP, Fennerty MB, Ryan ME, Shaw MJ, Lande JD, Pheley AM. 1996. Complications of endoscopic biliary sphincterotomy. *N Engl J Med* **335**:909-918. DOI: 10.1056/nejm199609263351301

Friedland S, Soetikno RM, Vandervoort J, Montes H, Tham T, Carr-Locke DL. 2002. Bedside scoring system to predict the risk of developing pancreatitis following

ERCP. *Endoscopy* **34**:483-488. DOI: 10.1055/s-2002-32004

Fujita K, Yazumi S, Matsumoto H, Asada M, Nebiki H, Matsumoto K, Maruo T,
Takenaka M, Tomoda T, Onoyama T, Kurita A, Ueki T, Katayama T, Kawamura
T, Kawamoto H. 2022. Multicenter prospective cohort study of adverse events
associated with biliary endoscopic retrograde cholangiopancreatography:
Incidence of adverse events and preventive measures for post-endoscopic
retrograde cholangiopancreatography pancreatitis. *Dig Endosc* **34**:1198-1204.
DOI: 10.1111/den.14225, PMID: PMC9540598

Fujita K, Yazumi S, Uza N, Kurita A, Asada M, Kodama Y, Goto M, Katayama T,
Anami T, Watanabe A, Sugahara A, Mukai H, Kawamura T. 2021. New practical
scoring system to predict post-endoscopic retrograde cholangiopancreatography
pancreatitis: Development and validation. *JGH Open* **5**:1078-1084. DOI:
10.1002/jgh3.12634, PMID: PMC8454475

Glomsaker T, Hoff G, Kvaloy JT, Soreide K, Aabakken L, Soreide JA. 2013. Patterns
and predictive factors of complications after endoscopic retrograde
cholangiopancreatography. *Br J Surg* **100**:373-380. DOI: 10.1002/bjs.8992

Harewood GC, Pochron NL, Gostout CJ. 2005. Prospective, randomized, controlled
trial of prophylactic pancreatic stent placement for endoscopic snare excision of

the duodenal ampulla. *Gastrointest Endosc* **62**:367-370. DOI:

10.1016/j.gie.2005.04.020

Ishiwatari H, Hayashi T, Ono M, Sato T, Kato J. 2013. Newly designed plastic stent for

endoscopic placement above the sphincter of Oddi in patients with malignant

hilar biliary obstruction. *Dig Endosc* **25 Suppl 2**:94-99. DOI: 10.1111/den.12080

Itoi T, Ikeura T, Irisawa A, Iwasaki E, Katanuma A, Kitamura K, Takenaka M,

Takeyama Y, Hirota M, Masamune A, Mayumi T, Mukai S, Morizane T, Yasuda

I, Ryozawa S, Yoshida M, Ito T, Yasuda K, Ito T, Iwano K, Entani T, Kamata K,

Kayashima A, Koiwai A, Tanisaka Y, Toyonaga H, Nakamaru K, Nagashima K,

Hayashi N, Horibe M, Honma T, Mizuide M, Minaga K, Yamamiya A. 2024.

Clinical practice guideline for post-ERCP pancreatitis 2023 (Japanese). *Suizo*

39:79-158

Itoi T, Ryozawa S, Katanuma A, Okabe Y, Kato H, Horaguchi J, Tsuchiya T, Gotoda T,

Fujita N, Yasuda K, Igarashi Y, Fujimoto K. 2018. Japan Gastroenterological

Endoscopy Society guidelines for endoscopic papillary large balloon dilation.

Dig Endosc **30**:293-309. DOI: 10.1111/den.13029

Kato S, Kuwatani M, Onodera M, Kudo T, Sano I, Katanuma A, Uebayashi M, Eto K,

Fukasawa M, Hashigo S, Iwashita T, Yoshida M, Taya Y, Kawakami H, Kato H,

Nakai Y, Kobashigawa K, Kawahata S, Shinoura S, Ito K, Kubo K, Yamato H, Hara K, Maetani I, Mukai T, Shibukawa G, Itoi T. 2022. Risk of Pancreatitis Following Biliary Stenting With/Without Endoscopic Sphincterotomy: A Randomized Controlled Trial. *Clinical Gastroenterology & Hepatology* **20**:1394-1403 e1391. DOI: 10.1016/j.cgh.2021.08.016

Katsinelos P, Lazaraki G, Chatzimavroudis G, Gkagkalis S, Vasiliadis I, Papaeuthimiou A, Terzoudis S, Pilpilidis I, Zavos C, Kountouras J. 2014. Risk factors for therapeutic ERCP-related complications: an analysis of 2,715 cases performed by a single endoscopist. *Ann Gastroenterol* **27**:65-72, PMID: PMC3959534

Kochar B, Akshintala VS, Afghani E, Elmunzer BJ, Kim KJ, Lennon AM, Khashab MA, Kalloo AN, Singh VK. 2015. Incidence, severity, and mortality of post-ERCP pancreatitis: a systematic review by using randomized, controlled trials. *Gastrointest Endosc* **81**:143-149 e149. DOI: 10.1016/j.gie.2014.06.045

Koshitani T, Konaka Y, Ohishi T, Yasuda T, Morinushi T, Mita M, Masuda M, Nakano K, Fuki S, Nakagawa S. 2022. Prophylaxis of Post-Endoscopic Retrograde Cholangiopancreatography Pancreatitis Using Temporary Pancreatic Stents Versus Rectal Nonsteroidal Anti-inflammatory Drugs: A Randomized Controlled Trial. *Pancreas* **51**:663-670. DOI: 10.1097/mpa.0000000000002090

Loperfido S, Angelini G, Benedetti G, Chilovi F, Costan F, De Berardinis F, De Bernardin M, Ederle A, Fina P, Fratton A. 1998. Major early complications from diagnostic and therapeutic ERCP: a prospective multicenter study. *Gastrointest Endosc* **48**:1-10

Maeda N, Higashimori A, Nakatani M, Mizuno Y, Nakamura Y, Ikeda D, Maruyama H, Morimoto K, Fukuda T, Watanabe T, Fujiwara Y. 2021. A 25 mg rectal dose of diclofenac for prevention of post-ERCP pancreatitis in elderly patients. *Scand J Gastroenterol* **56**:1109-1116. DOI: 10.1080/00365521.2021.1946134

Masci E, Mariani A, Curioni S, Testoni PA. 2003. Risk factors for pancreatitis following endoscopic retrograde cholangiopancreatography: a meta-analysis. *Endoscopy* **35**:830-834. DOI: 10.1055/s-2003-42614

Masci E, Toti G, Mariani A, Curioni S, Lomazzi A, Dinelli M, Minoli G, Crosta C, Comin U, Fertitta A, Prada A, Passoni GR, Testoni PA. 2001. Complications of diagnostic and therapeutic ERCP: a prospective multicenter study. *Am J Gastroenterol* **96**:417-423. DOI: 10.1111/j.1572-0241.2001.03594.x

Mazaki T, Mado K, Masuda H, Shiono M. 2014. Prophylactic pancreatic stent placement and post-ERCP pancreatitis: an updated meta-analysis. *J Gastroenterol* **49**:343-355. DOI: 10.1007/s00535-013-0806-1

Murray B, Carter R, Imrie C, Evans S, O'Suilleabhain C. 2003. Diclofenac reduces the

incidence of acute pancreatitis after endoscopic retrograde

cholangiopancreatography. *Gastroenterology* **124**:1786-1791. DOI:

10.1016/s0016-5085(03)00384-6

Okuno M, Shiroko J, Taguchi D, Yamaguchi K, Takada J, Imai S, Sato H, Thanabashi S.

2018. The Effectiveness of the Rectal Administration of Low-dose Diclofenac

for the Prevention of Post-endoscopic Retrograde Cholangiopancreatography

Pancreatitis. *Intern Med* **57**:2289-2294. DOI: 10.2169/internalmedicine.0554-17,

PMID: PMC6148161

Otsuka T, Kawazoe S, Nakashita S, Kamachi S, Oeda S, Sumida C, Akiyama T, Ario K,

Fujimoto M, Tabuchi M, Noda T. 2012. Low-dose rectal diclofenac for

prevention of post-endoscopic retrograde cholangiopancreatography

pancreatitis: a randomized controlled trial. *J Gastroenterol* **47**:912-917. DOI:

10.1007/s00535-012-0554-7

Sakai H, Iwai N, Sakagami J, Okuda T, Ohara T, Hattori C, Taniguchi M, Oka K, Hara T,

Tsuji T, Komaki T, Kagawa K, Dohi O, Yasuda H, Konishi H, Itoh Y. 2023.

Rectal administration of low-dose diclofenac does not reduce post-endoscopic

retrograde cholangiopancreatography pancreatitis: a propensity score matching

analysis. *Surg Endosc* **37**:2698-2705. DOI: 10.1007/s00464-022-09718-5

Sarner M, Cotton PB. 1984. Classification of pancreatitis. *Gut* **25**:756-759. DOI:

10.1136/gut.25.7.756, PMID: PMC1432589

Sperna Weiland CJ, Smeets X, Verdonk RC, Poen AC, Bhalla A, Venneman NG, Kievit

W, Timmerhuis HC, Umans DS, van Hooft JE, Besselink MG, van Santvoort

HC, Fockens P, Bruno MJ, Drenth JPH, van Geenen EJM. 2022. Optimal timing

of rectal diclofenac in preventing post-endoscopic retrograde

cholangiopancreatography pancreatitis. *Endosc Int Open* **10**:E246-E253. DOI:

10.1055/a-1675-2108, PMID: PMC8920594

Medical and served as a consultant

for Medtronic, Cook Medical and Boston Scientific, outside the submitted work.

Dr. Besselink has received research funding form Intuitive, Ethicon

Endo-Surgery, and Medtronic, outside the submitted work. Dr. Bruno has

received research funding from Boston Scientific, Cook Medical, Pentax

Medical, InterScope, ChiRho- Clin, and 3M and served as a consultant for

Boston Scientific, Cook Medical, and Pentax Medical, outside the submitted

work. Dr. Fockens has received consultancy fees from Cook Medical and

Olympus, outside the submitted work. Dr. Drenth has received research funding

from Gilead to support Hepatitis C elimination in the Netherlands, outside the

submitted work. Dr. van Geenen has received research funding from Mylan,

Boston Scientific, and Olympus and served as a consultant for

MTW-Endoskopie, outside the submitted work. Christina J. Sperna Weiland,

Xavier J.N.M. Smeets, Robert C. Verdonk, Alexander C. Poen, Abha Bhalla,

Niels G. Venneman Wietske Kievit, Hester C. Timmerhuis, Devica S. Umans,

and Hjalmar C. van Santvoort do not have potential conflicts of interest or

disclosures to report.

Sugimoto M, Takagi T, Suzuki R, Konno N, Asama H, Sato Y, Irie H, Watanabe K,

Nakamura J, Kikuchi H, Takasumi M, Hashimito M, Hikichi T, Ohira H. 2019.

Pancreatic stents to prevent post-endoscopic retrograde

cholangiopancreatography pancreatitis: A meta-analysis. *World J Metaanal*

7:249-258

Takada T, Isaji S, Mayumi T, Yoshida M, Takeyama Y, Itoi T, Sano K, Iizawa Y,

Masamune A, Hirota M, Okamoto K, Inoue D, Kitamura N, Mori Y, Mukai S,

Kiriyama S, Shirai K, Tsuchiya A, Higuchi R, Hirashita T. 2022. JPN clinical

practice guidelines 2021 with easy-to-understand explanations for the

management of acute pancreatitis. *J Hepatobiliary Pancreat Sci* **29**:1057-1083.

DOI: 10.1002/jhbp.1146

Takaori A, Ikeura T, Hori Y, Ito T, Nakamaru K, Masuda M, Mitsuyama T, Miyoshi H, Shimatani M, Takaoka M, Okazaki K, Naganuma M. 2021. Rectally Administered Low-Dose Diclofenac Has No Effect on Preventing Post-Endoscopic Retrograde Cholangiopancreatography Pancreatitis: A Propensity Score Analysis. *Pancreas* **50**:1024-1029. DOI: 10.1097/mpa.0000000000001877

Testoni PA, Mariani A, Giussani A, Vailati C, Masci E, Macarri G, Ghezzo L, Familiari L, Giardullo N, Mutignani M, Lombardi G, Talamini G, Spadaccini A, Briglia R, Piazzi L. 2010. Risk factors for post-ERCP pancreatitis in high- and low-volume centers and among expert and non-expert operators: a prospective multicenter study. *Am J Gastroenterol* **105**:1753-1761. DOI: 10.1038/ajg.2010.136

Tomoda T, Kato H, Miyamoto K, Matsumi A, Ueta E, Fujii Y, Saragai Y, Yamazaki T, Uchida D, Matsumoto K, Horiguchi S, Tsutsumi K, Okada H. 2021. Efficacy of low dose rectal diclofenac for preventing post-endoscopic retrograde cholangiopancreatography pancreatitis: Propensity score-matched analysis. *Dig Endosc* **33**:656-662. DOI: 10.1111/den.13828

Wang P, Li ZS, Liu F, Ren X, Lu NH, Fan ZN, Huang Q, Zhang X, He LP, Sun WS, Zhao Q, Shi RH, Tian ZB, Li YQ, Li W, Zhi FC. 2009. Risk factors for

ERCP-related complications: a prospective multicenter study. *Am J Gastroenterol* **104**:31-40. DOI: 10.1038/ajg.2008.5

Whitcomb DC, Frulloni L, Garg P, Greer JB, Schneider A, Yadav D, Shimosegawa T. 2016. Chronic pancreatitis: An international draft consensus proposal for a new mechanistic definition. *Pancreatology* **16**:218-224. DOI: 10.1016/j.pan.2016.02.001, PMID: PMC6042966

Williams EJ, Taylor S, Fairclough P, Hamlyn A, Logan RF, Martin D, Riley SA, Veitch P, Wilkinson ML, Williamson PR, Lombard M. 2007. Risk factors for complication following ERCP; results of a large-scale, prospective multicenter study. *Endoscopy* **39**:793-801. DOI: 10.1055/s-2007-966723

Wynants L, Bouwmeester W, Moons KG, Moerbeek M, Timmerman D, Van Huffel S, Van Calster B, Vergouwe Y. 2015. A simulation study of sample size demonstrated the importance of the number of events per variable to develop prediction models in clustered data. *J Clin Epidemiol* **68**:1406-1414. DOI: 10.1016/j.jclinepi.2015.02.002

Yang J, Wang W, Liu C, Zhao Y, Ren M, He S. 2020. Rectal Nonsteroidal Anti-Inflammatory Drugs for Endoscopic Retrograde Cholangiopancreatography Postoperative Pancreatitis Prevention: A Network Meta-Analysis. *J Clin*

Gastroenterol **54**:305-313. DOI: 10.1097/mcg.0000000000001322

Zheng R, Chen M, Wang X, Li B, He T, Wang L, Xu G, Yao Y, Cao J, Shen Y, Wang Y,

Zhu H, Zhang B, Wu H, Zou X, He G. 2020. Development and validation of a

risk prediction model and scoring system for post-endoscopic retrograde

cholangiopancreatography pancreatitis. *Ann Transl Med* **8**:1299. DOI:

10.21037/atm-20-5769, PMID: PMC7661903

Figure legends

Figure 1. Flowchart of the inclusion criteria. ERCP, endoscopic retrograde

cholangiopancreatography.

Figure 2. Example of the preprocedural PEP risk checklist. ERCP, endoscopic

retrograde cholangiopancreatography. IPMN, intraductal papillary mucinous neoplasm;

PEP, post-ERCP pancreatitis.

Tables

Table 1. Comparison of patient characteristics and ERCP outcomes between the development and validation cohorts.

	Development cohort (n=1037)	Validation cohort (n=1037)	*P* value
Patient factors			
Age, y, mean ± SD	73.8 ± 12.7	75.1 ± 12.5	0.02
Sex, n, male/female	642/395	629/408	0.59
History of pancreatitis, n (%)	73 (7.0)	45 (4.4)	0.01
History of PEP, n (%)	26 (2.5)	24 (2.3)	0.89
History of gastrectomy, n (%)	82 (7.9)	88 (8.5)	0.69
Billroth-I reconstruction, n	24	25	
Billroth-II reconstruction, n	23	25	
Roux-en-Y reconstruction, n	33	36	
Double tract, n	1	1	
Gastric tube reconstruction, n	1	1	
Pancreatic cancer, n (%)	145 (14.0)	174 (16.8)	0.09
IPMN, n (%)	17 (1.6)	8 (0.8)	0.11
Native papilla of Vater, n (%)	535 (51.6)	494 (47.7)	0.08
Total bilirubin, mg/dl, mean ± SD	3.5 ± 5.3	3.6 ± 5.0	0.45
Diameter of the MPD, mm, mean ± SD [a]	2.84 ± 2.63	3.1 ± 2.9	0.10
Pancreatic calcification, n (%) [b]	107 (10.6)	87 (8.7)	0.15
Periampullary diverticulum, n (%)	207 (20.0)	224 (21.6)	0.39
Pre-ERCP prophylaxis			
Protease inhibitors, n (%)	709 (68.4)	703 (67.8)	0.81
Intravenous hydration, n (%)	22 (2.1)	14 (1.4)	0.24
NSAID suppository, n (%)	53 (5.1)	45 (4.3)	0.47
Factors related to the planned procedure			
EST, n (%)	449 (43.3)	434 (41.9)	0.53
EPBD, n (%)	31 (3.0)	40 (3.9)	0.33
EPLBD, n (%)	56 (5.4)	55 (5.3)	1.0
Biliary stone removal, n (%)	327 (31.5)	342 (33.0)	0.51

Ampullectomy, n (%)	5 (0.5)	5 (0.5)	1.0
Biliary stent, n (%)	594 (57.3)	611(58/9)	0.48
Plastic stent, n (%)	445 (42.9)	436 (42.0)	0.72
SEMS, n (%)	119 (11.5)	122 (11.8)	0.89
CSEMS, n (%)	36 (3.5)	44 (4.2)	0.43
Biliary stent above the papilla, n (%)	45 (4.3)	47 (4.5)	0.92
Procedures on the pancreatic duct, n (%)	285 (27.5)	237 (22.9)	0.017
PEP occurrence, n (%)	70 (6.8)	64 (6.2)	0.66
Mild, n	60	53	
Moderate, n	8	7	
Severe, n	2	4	

ERCP, endoscopic retrograde cholangiopancreatography; SD, standard deviation; PEP, post-ERCP pancreatitis; IPMN, intraductal papillary mucinous neoplasm; MPD, main pancreatic duct; EST, endoscopic sphincterotomy; EPBD, endoscopic papillary balloon dilation; SEMS, self-expandable metallic stent; CSEMS, covered SEMS.

[a] Data were available for 1671 patients.

[b] Data were available for 2017 patients.

Table 2. Logistic regression analysis of predictive factors for PEP in the development cohort.

	Univariate analysis			Multivariate analysis				
	OR	95% CI	*P* value	OR	95% CI	*P* value	Regression coefficient	Points
Age < 50 years	2.42	0.99-6.0	0.053	1.76	0.67-4.63	0.25	0.56	-
Female	1.91	1.17-3.10	<0.01	1.72	1.03-2.89	0.039	0.55	1
History of pancreatitis	1.26	0.53-3.0	0.61					
History of PEP	1.84	0.54-6.28	0.33					
History of gastrectomy	0.89	0.35-2.27	0.81					
Pancreatic cancer	1.03	0.51-2.06	0.94					
IPMN	8.15	2.92-22.7	<0.01	3.04	0.97-9.52	0.056	1.11	2
Native papilla of Vater	4.49	2.42-8.30	<0.01	2.72	1.30-5.71	<0.01	1.0	2
Total bilirubin ≤ 1.2 mg/dl	1.13	0.69-1.84	0.62					
Diameter of the MPD > 3 mm [a]	1.31	0.76-2.25	0.33					
Pancreatic calcification [b]	0.36	0.11-1.17	0.089	0.32	0.10-1.1	0.072	-1.13	-2
Periampullary diverticulum	0.65	0.33-1.30	0.22					
Protease inhibitors	0.72	0.44-1.19	0.20					
Intravenous hydration	1.39	0.32-6.08	0.66					
NSAID suppository before ERCP	1.47	0.57-3.83	0.43					
EST	1.71	1.05-2.79	0.03	0.83	0.45-1.52	0.54	-0.19	-
EPBD	< 0.01	0-Infinity	0.98					
EPLBD	0.24	0.03-1.76	0.16					
Biliary stone removal	0.68	0.39-1.19	0.18					

	OR	95% CI	p-value	OR	95% CI	p-value		Points
Ampullectomy	3.49	0.39-31.6	0.27					
Biliary stent	0.93	0.57-1.52	0.78					
Plastic stent	0.72	0.44-1.20	0.21					
SEMS	1.66	0.87-3.20	0.13					
CSEMS	0.81	0.19-3.43	0.77					
Biliary stent above the papilla	0.30	0.04-2.24	0.24					
Procedures on the pancreatic duct	4.77	2.89-7.89	<0.01	3.49	1.99-6.12	<0.01	1.25	2

PEP, post–endoscopic retrograde cholangiopancreatography pancreatitis; OR, odds ratio; CI, confidence interval; IPMN, intraductal papillary mucinous neoplasm; MPD, main pancreatic duct; EST, endoscopic sphincterotomy; EPBD, endoscopic papillary balloon dilation; EPLBD, endoscopic papillary large balloon dilation; SEMS, self-expandable metallic stent; CSEMS, covered SEMS.

[a] Data were available for 985 patients in the development cohort.

[b] Data were available for 1012 patients in the development cohort.

Table 3. Patient distribution in terms of risk score and classification.

Risk score	Development cohort (n = 1012) [a]			Validation cohort (n = 1005) [b]		
	PEP occurrence, N	PEP rate (95% CI) (%)	*P* value [c]	PEP occurrence, N	PEP rate (95% CI) (%)	*P* value [c]
-2	0/29	0 (0-11.9)	< 0.01	0/33	0 (0-10.6)	< 0.01
-1	0/9	0 (0-33.6)		0/5	0 (0-52.2)	
0	0/289	0 (0-1.3)		8/293	2.7 (1.2-5.3)	
1	6/140	4.3 (1.6-9.1)		5/160	3.1 (1.0-7.1)	
2	8/202	4.0 (1.7-7.7)		14/195	7.2 (4.0-11.8)	
3	13/150	8.7 (4.7-14.4)		8/158	5.1 (2.2-9.7)	
4	18/97	18.6 (11.4-27.7)		14/84	16.7 (9.4-26.4)	
5	17/83	20.5 (12.4-30.8)		14/71	19.7 (11.2-30.9)	
6	3/9	33.3 (7.5-70.1)		0/3	0 (0-70.8)	
7	1/4	25.0 (0.6-80.6)		1/3	33.3 (0.8-90.6)	

Risk classification	Risk score	PEP occurrence, N	PEP rate (95% CI) (%)	*P* value [c]	PEP occurrence, N	PEP rate (95% CI) (%)	*P* value [c]
Low	≤ 0	0/327	0 (0-1.1)	< 0.01	8/331	2.4 (1.0-4.7)	< 0.01
Moderate	1-3	27/492	5.5 (3.6-7.9)		27/513	5.3 (3.5-7.6)	
High	4-7	39/193	20.2 (14.8-26.6)		29/161	18.0 (12.4-24.8)	

PEP, post–endoscopic retrograde cholangiopancreatography pancreatitis; CI, confidence interval.

[a] There were missing data for 25 patients.

[b] Data for 32 patients were missing.

[c] The correlations between the risk score or classification and PEP occurrence were evaluated via the Cochran–Armitage test.

Table 4. Goodness of fit of the risk score model.

	Development cohort	Validation cohort
C-statistic (95% CI)	0.77 (0.72–0.82)	0.71 (0.64–0.78)
Hosmer☐Lemeshow test, *p* value	0.59	0.40

CI, confidence interval.