Benchmarking the Confidence of Large Language Models in Clinical Questions

Mahmud Omar[1], Reem Agbareia[2], Benjamin S Glicksberg[1], Girish N Nadkarni[1*], Eyal Klang[1*].

1- Division of Data-Driven and Digital Medicine (D3M), Department of Medicine, Icahn School of Medicine at Mount Sinai, NY.

2- Tel Aviv Sourasky Medical Center, Ichilov, Tel-Aviv.

* Equal Contribution

Acknowledgment – We thank Dr. Uriel Katz and the co-authors of the paper "GPT versus Resident Physicians — A Benchmark Based on Official Board Scores, DOI: 10.1056/AIdbp2300192" for sharing the MCQ dataset.

Financial disclosure – This research received no specific grant from any funding agency in the public, commercial, or not-for-profit sectors.

Competing interest – None declared.

Ethical approval was not required for this research, as it utilized already published synthetic data.

Corresponding author:

Mahmud Omar M.D.

E-mail: Mahmudomar70@gmail.com

Abstract

Background and Aim: The capabilities of large language models (LLMs) to self-assess their own confidence in answering questions in the biomedical realm remain underexplored. This study evaluates the confidence levels of 12 LLMs across five medical specialties to assess their ability to accurately judge their responses.

Methods: We used 1,965 multiple-choice questions assessing clinical knowledge from internal medicine, obstetrics and gynecology, psychiatry, pediatrics, and general surgery areas. Models were prompted to provide answers and to also provide their confidence for the correct answer (0-100). The confidence rates and the correlation between accuracy and confidence were analyzed.

Results: There was an inverse correlation (r=-0.40, p=0.001) between confidence and accuracy, where worse performing models showed paradoxically higher confidence. For instance, a top performing model, GPT4o had a mean accuracy of 74% with a mean confidence of 63%, compared to a least performant model, Qwen-2-7B, which showed mean accuracy 46% but mean confidence 76%. The mean difference in confidence between correct and incorrect responses was low for all models, ranging from 0.6% to 5.4%, with GPT4o having the highest differentiation of 5.4%.

Conclusion: Better performing LLMs show more aligned overall confidence levels. However, even the most accurate models still show minimal variation in confidence between right and wrong answers. This underscores an important limitation in current LLMs' self-assessment mechanisms, highlighting the need for further research before integration into clinical settings.

Keywords: Large Language Models (LLMs), Safe AI, AI Reliability, Clinical knowledge

Introduction

With their capacity to understand and generate human-like text, Large Language Models (LLMs) are poised to support healthcare professionals in complex clinical decisions (1,2). A wide array of LLMs is now accessible, including open source models, offering solutions that cater to both the public and medical professionals (1).

The efficacy of these models has been demonstrated in a variety of tasks, albeit with some limitations (3,4). For instance, LLMs like Generative Pre-trained Transformers (GPT) have shown promise in providing diagnostic assistance and answering medical queries (3,5–7). Katz et al. demonstrated that GPT-4 not only improved clinically compared to its predecessor GPT-3.5, but also matched physician performance in certain areas (8). However, there is evidence of "hallucinations" and inaccuracies in model outputs, which could lead to harm in clinical decision-making (9,10). Specifically, LLMs occasionally generated completely fabricated evidence (e.g., information and references) and presented it as factual (9,10).

One way of building confidence in applying models in health care is explainable AI (11,12). However, easily explainable outputs are difficult due to the complexity of how LLMs process and output data (11,13,14). Recent studies show that LLMs tend to be overly confident (15), and it is hard to ascertain where that confidence stems from in a black box system.

The goal of the current study was to benchmark widely used LLMs in terms of accuracy and associated confidence in answering clinical questions. Our aim was to determine if these models can accurately judge when to be confident in their responses, and in doing so, allow for better explainability in their application.

Materials and Methods

Study Design and Data Source

This study utilizes a public compiled dataset from a previous study by Katz et al., which includes 655 questions across five medical specialties: internal medicine, obstetrics and gynecology (OBGYN), psychiatry, pediatrics, and general surgery (8). Crafted from internationally recognized textbooks and guidelines, this dataset serves as a standardized framework for assessment (16–20). To enhance benchmarking reliability, each original question was rephrased twice using GPT-4 Application Programming Interface (API) in Python, yielding 1965 questions. The prompts ensured preservation of the original medical content and correct answers (21). A random 20% sample from each field underwent manual validation by two expert physicians, confirming accuracy and consistency with the original questions.

Model Setup and Configuration

The LLMs employed in this study were prompted to return the correct answer along with a confidence score for each choice ("A", "B", "C", "D"), expressed as a percentage. The open access models were executed using API codes in a dedicated server with 4xH100 80GB GPUs, with the corresponding codebase accessible on GitHub. We utilized Python (3.10) for data analyses. The commercial models were used using the companies API interfaces. We used several Python libraries to facilitate data processing, model interaction, and analysis: NumPy (1.26.4), Pandas (2.1.4, Scikit-Learn (1.3.0), Hugging Face's Transformers (4.37.2) and torch (2.2.2+cu121), and the Json module (2.0.9). We used the default hyperparameters for each model (22). For the open access models, we used the "instruct" versions, which perform better on zero-shot questioning.

Benchmarked LLMs

We evaluated 12 LLMs representing a range of model sizes and architectures (**Figure 1**). This selection includes both widely used and recently developed models. The benchmarked models are shown in **Table 1**.

Table 1: A list of the benchmarked LLMs.

Model	Description
GPT-4 0613 and GPT-4o	Latest iterations of OpenAI's GPT series.
GPT-3.5	Predecessor to GPT-4, still widely used in various applications.
Claude Sonnet 3.5 and Claude Opus 3	Recent models from Anthropic.
Gemini 1.5 Pro	Google's advanced model.
Llama-3-70B and Llama-3-8B	Different sizes of Meta's open-source Llama-3 series.
Llama3-OpenBio-70B	A domain-specific variant of the Llama model.
Mixtral-8x7B	An open-source mixture of experts (MOE) model developed by Mistral AI.
Qwen2-72B and Qwen2-7B	Different sizes of Alibaba's Qwen model series.

Figure 1: A flowchart representing the evaluation methodology.

Statistical analysis

Pearson correlation coefficient was used to correlate means of confidence and accuracy across models and medical fields. Chi-square tests assessed overall performance differences within each field, using proportions of correct responses. Post-hoc pairwise comparisons with Bonferroni correction identified specific inter-model differences. Confidence levels were compared between correct and incorrect responses for each model using two-sample t-tests. Mean confidence scores were

calculated for higher-tier and lower-tier models, as well as across all models. Performance consistency was evaluated by comparing confidence gaps between correct and incorrect responses. All statistical tests used a significance level of α = 0.05. Analyses were performed using R version 4.1.2 (R Core Team, 2021).

Results

Confidence analysis

Table 2 summarizes accuracies and confidence levels across the models and **Supplementary Table 1** presents the comprehensive data across all inspected fields and all models. An inverse correlation is demonstrated (r = -0.40, p < 0.01), where better performing models generally show lower confidence.

Table 2: accuracies and confidence levels across the models.

MODEL	ACCURACY	TOTAL CONFIDENCE	CORRECT CONFIDENCE	INCORRECT CONFIDENCE
CLAUDE 3.5 SONNET	74	69.7	70.5	67.4
GPT4O	73.8	63	64.4	59
CLAUDE 3 OPUS	71.7	68.5	68.9	67.3
GPT4	66	84.1	84.5	83.3
LLAMA-3-70B	63.4	57.3	59.5	53.6
LLAMA OPENBIO	59.2	77.9	77.7	78.1
GEMINI	59.1	86.5	87.2	85.5
QWEN2-72B	57.8	57.7	58.6	56.5
MIXTRAL 8X7B	50.6	84.3	85.5	83
GPT-3.5	49	82.3	81.6	82.9
LLAMA-3-8B	48.4	80	79.7	80.3
QWEN2-7B	46	75.5	74.4	76.4

The mean confidence across all 12 models was 76.1% when correct and 74.4% when incorrect. The top six preforming models showed a mean confidence of 72.5% when correct and 69.4% when incorrect, while lower-tier models displayed 79.6% confidence when correct and 79.5% when incorrect. (**Table 3**).

Table 3: Confidence means of LLMs between correct and incorrect answers.

Model	Confidence When Incorrect (mean, SD)	Confidence When Correct (mean, SD)	p-value
GPT-4o	58.99 ± 14.31	64.38 ± 16.11	<0.01
Llama-3-70B	53.59 ± 22.38	59.50 ± 23.54	<0.01
Claude 3.5 Sonnet	67.37 ± 9.08	70.52 ± 11.07	<0.01
Gemini	85.55 ± 16.23	87.17 ± 16.58	0.35
Claude 3 Opus	67.32 ± 13.06	68.90 ± 15.65	0.61
GPT-4	83.34 ± 23.30	84.52 ± 22.43	0.07
Qwen2-72B	56.49 ± 18.55	58.59 ± 20.03	<0.01
Qwen2-7B	76.37 ± 17.11	74.45 ± 20.30	0.01
Mixtral 8x7B	82.99 ± 16.52	85.49 ± 14.62	0.04
Llama-3-8B	80.25 ± 17.40	79.67 ± 21.59	0.31
Llama OpenBio	78.14 ± 27.59	77.73 ± 28.78	0.83
GPT-3.5	82.85 ± 27.17	81.63 ± 28.66	0.81

Four models (GPT-4, Llama-3-70B, Claude 3.5 Sonnet, and Qwen2-72B) demonstrated significantly higher confidence when correct (p < 0.01). Gemini exhibited the highest overall confidence levels (85.6% when incorrect, 87.2% when correct). Qwen2-7B was unique in displaying higher confidence when incorrect (76.4% vs 74.5% when correct, p= 0.01).

GPT-3.5 and Llama-OpenBio-70B revealed minimal differences in confidence between correct and incorrect answers (p = 0.8). The largest confidence gap was

observed in GPT-4 (5.4 percentage points), while Llama-3-8B had the smallest gap

(0.6 percentage points) (**Figure 2**).

*(Left) - Displays the average confidence and 95% confidence intervals for each model, categorized by correct answers (green) and incorrect answers (red).

(Right) - Shows the differences in average confidence for each model, where green indicates higher confidence in correct answers, and red indicates higher confidence in incorrect answers.

Figure 2: LLMs confidence results between correct and incorrect answers.

Models' performances across fields

Significant differences were seen in model performance across all five medical

specialties ($p < 0.01$). GPT-4o and Claude 3.5 Sonnet consistently outperformed other

models. In Internal Medicine, GPT-4o (70.9%) and Claude 3.5 Sonnet (73.5%)

showed no significant difference ($p = 1.0$) but outperformed lower-tier models like

qwen-7b (43.7%, p < 0.001). For OBGYN, Claude 3.5 Sonnet (71.0%) significantly outperformed most models, including GPT-4 (54.0%, p < 0.001). In Pediatrics, the top five models (GPT-4o, Llama3-70b, Claude 3.5 Sonnet, Claude Opus, GPT-4) showed no significant differences among themselves (all p > 0.05) but outperformed lower-tier models. Psychiatry results mirrored this pattern, with GPT-4O (84.4%) and Claude 3.5 Sonnet (82.4%) leading. In Surgery, GPT-4O (70.9%) and Claude 3.5 Sonnet (70.5%) again showed no significant difference (p = 1.0) but outperformed lower-performing models like qwen-7b (45.6%, p < 0.01) (**Tables 4-5**).

Discussion

In our evaluation, an inverse correlation was demonstrated between accuracy and confidence for LLMs. Notably, some lower-complexity models displayed higher confidence in incorrect answers. However, across all models, there was limited differentiation in confidence levels between right and wrong answers. For instance, GPT-4o, a top performing model, also displayed the largest confidence gap of 5.4 percentage points between correct and incorrect answers. However, this margin is too narrow for practical clinical decision-making. These results highlight potential risks in clinical applications, where model confidence, regardless of answer correctness, could lead to misinformed decisions.

Katz et al. reported GPT-4 outperforming physicians in psychiatry and performing comparably in general surgery and internal medicine (8). Our study corroborates GPT-4's strong performance, particularly in psychiatry where it achieved 84.4% accuracy. However, our findings suggest more cautious interpretation is needed, given the high confidence levels observed for incorrect answers. Xiong et al.'s work on

LLM confidence elicitation aligns with our observations of overconfidence (15). They noted improved calibration and failure prediction as model capability increased, which parallels our finding of better confidence calibration in more complex models. The implications for clinical practice warrant careful consideration. While the performance leap of newer models is promising, their inability to accurately self-assess confidence across wrong answers poses risks. Two possible strategies to address these challenges can be human-in-the-loop protocols and implementing ensemble methods (23).

Human-AI collaboration may offer a balanced approach to leveraging AI strengths while maintaining necessary human oversight in healthcare (24). Sezgin et al. suggested human-in-the-loop approach, ensuring that AI systems are supervised by human expertise. However, effective implementation faces challenges. Careful design of user interfaces is important to prevent automation bias (24,25). There are also concerns about potential erosion of clinical skills with over reliance on AI (26).

Ensemble methods, which aggregate multiple models, present another possible strategy (27). Mahajan et al. conducted review of ensemble learning techniques in disease prediction (28). They found that stacking, an ensemble method that combines multiple classifiers, showed the most accurate performance in 19 out of 23 cases. The voting approach was identified as the second-best ensemble method. However, ensemble methods are computationally intensive and may introduce latency in real-time clinical applications (29).

Both strategies would require extensive clinical trials for validation and the development of model-specific calibration curves for each medical specialty.

Our study has several limitations. The dataset was limited to 1965 multiple choice questions across five medical specialties, which may not fully represent the breadth of clinical scenarios. A combination of automatic rephrasing and manual validation could introduce bias (21). Additionally, the study used default model hyperparameters, potentially limiting performance optimization. The generalizability of results to real-world clinical settings remains uncertain. The study did not explore techniques such as fine-tuning and retrieval-augmented generation (RAG), which might improve model accuracy and confidence calibration (30). Finally, the study did not consider the computational cost and time efficiency of deploying these models in practical healthcare scenarios.

In conclusion, better performing LLMs show more aligned overall confidence levels. However, even the most accurate models still show minimal variation in confidence between right and wrong answers. This underscores an important limitation in current LLMs' self-assessment mechanisms, highlighting the need for further research before integration into clinical settings.

References

1.   Thirunavukarasu AJ, Ting DSJ, Elangovan K, Gutierrez L, Tan TF, Ting DSW. Large language models in medicine. Nat Med. 2023 Aug;29(8):1930–40.
2.   Clusmann J, Kolbinger FR, Muti HS, Carrero ZI, Eckardt JN, Laleh NG, et al. The future landscape of large language models in medicine. Commun Med. 2023 Oct 10;3(1):141.
3.   Kanjee Z, Crowe B, Rodman A. Accuracy of a Generative Artificial Intelligence Model in a Complex Diagnostic Challenge. JAMA. 2023 Jul 3;330(1):78–80.
4.   Singhal K, Azizi S, Tu T, Mahdavi SS, Wei J, Chung HW, et al. Large language models encode clinical knowledge. Nature. 2023 Aug 3;620(7972):172–80.
5.   Johnson D, Goodman R, Patrinely J, Stone C, Zimmerman E, Donald R, et al. Assessing the Accuracy and Reliability of AI-Generated Medical Responses: An Evaluation of the Chat-GPT Model. Res Sq. 2023 Feb 28;rs.3.rs-2566942.
6.   Giannos P. Evaluating the limits of AI in medical specialisation: ChatGPT's performance on the UK Neurology Specialty Certificate Examination. BMJ Neurol Open. 2023 Jun 15;5(1):e000451.
7.   Hoch CC, Wollenberg B, Lüers JC, Knoedler S, Knoedler L, Frank K, et al. ChatGPT's quiz skills in different otolaryngology subspecialties: an analysis of 2576 single-choice and multiple-choice board certification preparation questions. Eur Arch Oto-Rhino-Laryngol Off J Eur Fed Oto-Rhino-Laryngol Soc EUFOS Affil Ger Soc Oto-Rhino-Laryngol - Head Neck Surg. 2023 Sep;280(9):4271–8.
8.   Katz U, Cohen E, Shachar E, Somer J, Fink A, Morse E, et al. GPT versus Resident Physicians — A Benchmark Based on Official Board Scores. NEJM AI. 2024 Apr 25;1(5):AIdbp2300192.
9.   Omar M, Nassar S, Hijaze K, Glicksberg BS, Nadkarni GN, Klang E. Generating Credible Referenced Medical Research: A Comparative Study of Openai's Gpt-4 and Google's Gemini [Internet]. Rochester, NY; 2024 [cited 2024 Apr 22]. Available from: https://papers.ssrn.com/abstract=4780940
10.   Azamfirei R, Kudchadkar SR, Fackler J. Large language models and the perils of their hallucinations. Crit Care. 2023 Mar 21;27:120.
11.   Yang G, Ye Q, Xia J. Unbox the black-box for the medical explainable AI via multi-modal and multi-centre data fusion: A mini-review, two showcases and beyond. Int J Inf Fusion. 2022 Jan;77:29–52.
12.   Soroush A, Glicksberg BS, Zimlichman E, Barash Y, Freeman R, Charney AW, et al. Large Language Models Are Poor Medical Coders — Benchmarking of Medical Code Querying. NEJM AI. 2024 Apr 25;1(5):AIdbp2300040.
13.   Schwartz IS, Link KE, Daneshjou R, Cortés-Penfield N. Black Box Warning: Large Language Models and the Future of Infectious Diseases Consultation. Clin Infect Dis. 2023 Nov 16;
14.   Poon AIF, Sung JJY. Opening the black box of AI-Medicine. J Gastroenterol Hepatol. 2021 Mar;36(3):581–4.
15.   Xiong M, Hu Z, Lu X, Li Y, Fu J, He J, et al. Can LLMs Express Their Uncertainty? An Empirical Evaluation of Confidence Elicitation in LLMs [Internet]. arXiv; 2024 [cited 2024 Aug 8]. Available from: http://arxiv.org/abs/2306.13063
16.   Townsend CM, Beauchamp RD, Evers BM, Mattox KL. Sabiston Textbook of Surgery: The Biological Basis of Modern Surgical Practice. Elsevier Health Sciences; 2016. 2191 p.
17.   Loscalzo J. Harrison's principles of internal medicine. No Title [Internet].

[cited 2024 Jul 2]; Available from: https://cir.nii.ac.jp/crid/1130573781693502243

18.   Kliegman RM, Behrman RE, Jenson HB, Stanton BMD. Nelson Textbook of Pediatrics E-Book. Elsevier Health Sciences; 2007. 3200 p.

19.   Association AP. Diagnostic and Statistical Manual of Mental Disorders. Text Revis [Internet]. 2000 [cited 2024 Jul 2]; Available from: https://cir.nii.ac.jp/crid/1573950399819987840

20.   Gabbe SG, Niebyl JR, Simpson JL, Landon MB, Galan HL, Jauniaux ERM, et al. Obstetrics: Normal and Problem Pregnancies E-Book. Elsevier Health Sciences; 2016. 1426 p.

21.   Soni S, Roberts K. Paraphrasing to improve the performance of Electronic Health Records Question Answering. AMIA Summits Transl Sci Proc. 2020 May 30;2020:626–35.

22.   OpenAI, Achiam J, Adler S, Agarwal S, Ahmad L, Akkaya I, et al. GPT-4 Technical Report [Internet]. arXiv; 2024 [cited 2024 Aug 10]. Available from: http://arxiv.org/abs/2303.08774

23.   Longhurst CA, Singh K, Chopra A, Atreja A, Brownstein JS. A Call for Artificial Intelligence Implementation Science Centers to Evaluate Clinical Effectiveness. NEJM AI. 2024 Jul 25;1(8):AIp2400223.

24.   Sezgin E. Artificial intelligence in healthcare: Complementing, not replacing, doctors and healthcare providers. Digit Health. 2023 Jul 2;9:20552076231186520.

25.   Straw I. The automation of bias in medical Artificial Intelligence (AI): Decoding the past to create a better future. Artif Intell Med. 2020 Nov;110:101965.

26.   Čartolovni A, Malešević A, Poslon L. Critical analysis of the AI impact on the patient–physician relationship: A multi-stakeholder qualitative study. Digit Health. 2023 Dec 19;9:20552076231220833.

27.   Yang H, Li M, Zhou H, Xiao Y, Fang Q, Zhang R. One LLM is not Enough: Harnessing the Power of Ensemble Learning for Medical Question Answering. medRxiv. 2023 Dec 24;2023.12.21.23300380.

28.   Mahajan P, Uddin S, Hajati F, Moni MA. Ensemble Learning for Disease Prediction: A Review. Healthcare. 2023 Jun 20;11(12):1808.

29.   Edeh MO, Dalal S, Dhaou IB, Agubosim CC, Umoke CC, Richard-Nnabu NE, et al. Artificial Intelligence-Based Ensemble Learning Model for Prediction of Hepatitis C Disease. Front Public Health. 2022 Apr 27;10:892371.

30.   Glicksberg BS, Timsina P, Patel D, Sawant A, Vaid A, Raut G, et al. Evaluating the accuracy of a state-of-the-art large language model for prediction of admissions from the emergency room. J Am Med Inform Assoc JAMIA. 2024 May 21;ocae103.

Tables

Table 1: A list of the benchmarked LLMs.

Model	Description
GPT-4 0613 and GPT-4o	Latest iterations of OpenAI's GPT series.
GPT-3.5	Predecessor to GPT-4, still widely used in various applications.
Claude Sonnet 3.5 and Claude Opus 3	Recent models from Anthropic.
Gemini 1.5 Pro	Google's advanced model.
Llama-3-70B and Llama-3-8B	Different sizes of Meta's open-source Llama-3 series.
Llama3-OpenBio-70B	A domain-specific variant of the Llama model.
Mixtral-8x7B	An open-source model developed by Mistral AI.
Qwen2-72B and Qwen2-7B	Different sizes of Alibaba's Qwen model series.

Table 2: accuracies and confidence levels across the models.

MODEL	ACCURACY	TOTAL CONFIDENCE	CORRECT CONFIDENCE	INCORRECT CONFIDENCE
CLAUDE 3.5 SONNET	74	69.7	70.5	67.4
GPT4O	73.8	63	64.4	59
CLAUDE 3 OPUS	71.7	68.5	68.9	67.3
GPT4	66	84.1	84.5	83.3
LLAMA-3-70B	63.4	57.3	59.5	53.6
LLAMA OPENBIO	59.2	77.9	77.7	78.1
GEMINI	59.1	86.5	87.2	85.5
QWEN2-72B	57.8	57.7	58.6	56.5
MIXTRAL 8X7B	50.6	84.3	85.5	83

GPT-3.5	49	82.3	81.6	82.9
LLAMA-3-8B	48.4	80	79.7	80.3
QWEN2-7B	46	75.5	74.4	76.4

Table 3: Confidence means of LLMs between correct and incorrect answers.

Model	Confidence When Incorrect (mean, SD)	Confidence When Correct (mean, SD)	p-value
GPT-4o	58.99 ± 14.31	64.38 ± 16.11	<0.01
Llama-3-70B	53.59 ± 22.38	59.50 ± 23.54	<0.01
Claude 3.5 Sonnet	67.37 ± 9.08	70.52 ± 11.07	<0.01
Gemini	85.55 ± 16.23	87.17 ± 16.58	0.35
Claude 3 Opus	67.32 ± 13.06	68.90 ± 15.65	0.61
GPT-4	83.34 ± 23.30	84.52 ± 22.43	0.07
Qwen2-72B	56.49 ± 18.55	58.59 ± 20.03	<0.01
Qwen2-7B	76.37 ± 17.11	74.45 ± 20.30	0.01
Mixtral 8x7B	82.99 ± 16.52	85.49 ± 14.62	0.04
Llama-3-8B	80.25 ± 17.40	79.67 ± 21.59	0.31
Llama OpenBio	78.14 ± 27.59	77.73 ± 28.78	0.83
GPT-3.5	82.85 ± 27.17	81.63 ± 28.66	0.81

Table 4: Top-performing models benchmarking results across different specialties.

Specialty	Model	GPT-4	Llama3-70b	Claude 3.5 Sonnet	Gemini 1.5 Pro	Claude 3 Opus	GPT-4
Surgery	Overall	70.92	61.47	70.45	60.76	67.14	69.50
	Original	70.92	55.32	66.67	58.87	65.96	68.79

		Qwen-2-72b	Qwen-7b	Mistral	Llama 8b	Llama Bio	GPT-3.5
	Rephrase 1	68.79	65.25	68.79	59.57	67.38	68.79
	Rephrase 2	73.05	63.83	75.89	63.83	68.09	70.92
Internal Medicine	Overall	75.13	58.47	73.54	58.99	70.45	64.02
	Original	73.81	57.14	71.43	56.35	66.67	61.11
	Rephrase 1	74.60	59.52	73.81	63.49	68.79	66.67
	Rephrase 2	76.98	58.73	75.40	57.14	75.89	64.29
Obstetrics and Gynecology	Overall	65.71	55.64	70.98	58.75	65.47	53.96
	Original	69.06	53.96	72.66	57.55	68.35	56.12
	Rephrase 1	62.59	56.12	71.22	59.71	68.35	53.96
	Rephrase 2	65.47	56.83	69.06	58.99	59.71	51.80
Pediatrics	Overall	71.72	68.35	71.38	57.58	68.01	67.00
	Original	68.69	64.65	70.71	54.55	67.68	64.65
	Rephrase 1	72.73	70.71	70.71	58.59	71.72	66.67
	Rephrase 2	73.74	69.70	72.73	59.60	64.65	69.70
Psychiatry	Overall	84.44	73.56	82.44	NA	79.78	77.33
	Original	77.33	67.33	83.33	NA	77.33	78.00
	Rephrase 1	88.00	76.00	81.33	NA	81.33	77.33
	Rephrase 2	88.00	77.33	82.67	NA	80.67	76.67

Table 5: Lower-performing models benchmarking results across different specialties.

Specialty	Model	Qwen-2-72b	Qwen-7b	Mistral	Llama 8b	Llama Bio	GPT-3.5
Surgery	Overall	57.45	45.63	46.81	47.52	56.03	53.43
	Original	53.90	46.10	45.39	46.81	58.87	52.48
	Rephrase 1	61.70	47.52	46.10	48.23	54.61	53.90
	Rephrase 2	56.74	43.26	48.94	47.52	54.61	53.90
Internal Medicine	Overall	47.88	43.65	44.97	40.48	53.70	45.77

	Original	46.03	45.24	46.83	40.48	55.56	46.83
	Rephrase 1	49.21	42.86	43.65	40.48	52.38	43.65
	Rephrase 2	48.41	42.86	44.44	40.48	53.17	46.83
Obstetrics and Gynecology	Overall	55.16	43.65	44.36	43.17	53.72	42.93
	Original	56.83	43.17	43.88	44.60	56.83	46.04
	Rephrase 1	55.40	43.88	46.04	39.57	50.36	41.01
	Rephrase 2	53.24	43.88	43.17	45.32	53.96	41.73
Pediatrics	Overall	59.60	39.73	52.19	48.48	61.28	49.16
	Original	52.53	36.36	47.47	44.44	60.61	49.49
	Rephrase 1	61.62	45.45	53.54	51.52	62.63	46.46
	Rephrase 2	64.65	37.37	55.56	49.49	60.61	51.52
Psychiatry	Overall	67.33	54.22	63.33	60.89	70.44	57.56
	Original	67.33	54.00	62.00	60.00	67.33	62.67
	Rephrase 1	65.33	53.33	62.67	60.00	72.00	51.33
	Rephrase 2	69.33	55.33	65.33	62.67	72.00	58.67

Supplementary Table 1

Model	Field	Accuracy	Total Confidence	Correct Confidence	Incorrect Confidence
	Surgery	70.9	64.5	66.4	59.9
	OBGYN	65.7	62.3	64.1	59
GPT4o	Internal Medicine	75.1	62.8	64.1	58.8
	Pediatrics	71.7	64.6	66.8	59.1
	Psychiatry	84.4	61.3	61.9	57.6
	Surgery	61.5	64.1	67.9	58
Llama-3-70B	OBGYN	55.6	64.6	66	62.8
	Internal Medicine	58.5	30.6	31.3	29.6

	Pediatrics	68.4	65.4	67.2	61.5
	Psychiatry	73.3	61.5	62.5	58.7
	Surgery	70.4	69.4	70.4	67.2
	OBGYN	71	68.4	69.4	65.9
Sonnet-3.5	Internal Medicine	73.5	69	69.2	68.5
	Pediatrics	71.4	70	71.3	66.8
	Psychiatry	82.4	71.5	72	69.2
	Surgery	60.8	87.1	88.6	84.6
	OBGYN	58.8	87.3	87.9	86.5
Gemini	Internal Medicine	59	83.4	83.6	83.1
	Pediatrics	57.6	88.6	88.7	88.4
	Psychiatry				
	Surgery	67.1	70.6	71.9	68
	OBGYN	71	68.4	69.4	65.9
Opus	Internal Medicine	71.7	68	68.2	67.6
	Pediatrics	68	70.3	71.5	67.6
	Psychiatry	79.1	65.6	65.1	67.5
	Surgery	66.7	86.2	86.3	86
	OBGYN	54	86.5	84.8	88.5
GPT-4	Internal Medicine	64	83.4	85.7	79.4
	Pediatrics	67	85.8	87.9	81.6
	Psychiatry	77.3	79.4	80.2	76.8
	Surgery	57.4	59.5	60.5	58.2
	OBGYN	55.2	59	57.5	60.8
Qwen2-72B	Internal Medicine	48.1	57.3	58.6	56.1
	Pediatrics	59.6	56.2	58.5	52.7
	Psychiatry	67.3	56.1	57.9	52.5
	Surgery	45.6	76	73.8	77.8

	OBGYN	43.6	78.1	76.2	79.6
Qwen2-7B	Internal Medicine	43.9	77.5	79.4	76
	Pediatrics	39.7	77.1	75.7	78
	Psychiatry	54.2	69.9	69.7	70.1
	Surgery	46.8	84.9	85.5	84.3
	OBGYN	44.4	83.9	84	83.8
mixtral_8x7B	Internal Medicine	45.2	84.8	85.8	84
	Pediatrics	52.2	84.1	85.7	82.2
	Psychiatry	63.3	83.7	86.1	79.5
	Surgery	47.5	82.5	82.5	82.5
	OBGYN	43.2	80.2	80.1	80.4
Llama-3-8B	Internal Medicine	40.5	81.8	82.4	81.4
	Pediatrics	48.5	84.6	86.3	83.1
	Psychiatry	60.9	72.7	72.3	73.4
	Surgery	56	78.1	79.7	76.2
	OBGYN	53.7	80.8	79.9	81.9
Llama_openbio	Internal Medicine	54	77.9	78.2	77.6
	Pediatrics	61.3	80.6	79.9	81.7
	Psychiatry	70.4	73.2	73.2	73
	Surgery	48.7	86.3	86.2	86.4
	OBGYN	42.9	83.4	83.5	83.2
GPT-3.5	Internal Medicine	45.8	84.7	85	84.5
	Pediatrics	49.2	89.2	88	90.5
	Psychiatry	57.6	70.7	70.9	70.6