Geospatial Modeling Study Assessing Population Level Accessibility to Medical College Hospitals in India

Hospitals in India

Short title: Medical College Hospitals Accessibility India

Harsh Thakkar[1], Chaitanya Reddy[1] +, Varun Raj Passi[1] +, Aamir Miyajiwala[1], Sahil Kale[1],

Ankit Raj[1], Siddhesh Zadey[1,2]

      1 Association for Socially Applicable Research (ASAR), Pune, Maharashtra, India

      2 Dr. D. Y. Patil Dental College and Hospital, Dr. D. Y. Patil Vidyapeeth, Pune,

Maharashtra, India

  + Contributed equally

Corresponding author:

Siddhesh Zadey BSMS, MScGH;

ASAR Office Address: D2 Sai Heritage, New DP Road, Aundh, Pune, Maharashtra, India

411007

Email: sidzadey@asarforindia.org

Article Type: Original Article

Word Count: 4852

Keywords: India; Health Services Accessibility; Teaching Hospital; Geographic Information

System

Declarations

Ethics and Guidelines

Consent to participate: Not Applicable

Consent for publication: Not Applicable

Ethical Approval: Not Applicable

Availability of data and materials: The datasets used and/or analysed during the current

study are available from the corresponding author on reasonable request.

Competing interests: Siddhesh Zadey represents the Association for Socially Applicable

Research (ASAR) on the drafting committee of the Maharashtra State Mental Health Policy

and the Global Alliance for Surgical, Obstetric, Trauma, and Anesthesia Care (G4 Alliance).

He serves as the chair of the South Asia Working Group of the G4 Alliance. He is on the

board of ASAR and the advisory board of Nivarana. He has previously received honoraria

from Think Global Health and The Hindu. Other authors declare no competing interests.

Funding: None

Authors' contributions:

Study concept and design: Siddhesh Zadey

Acquisition, analysis, or interpretation of data: Chaitanya Reddy, Harsh Thakkar, Aamir

Miyajiwala, Sahil Kale, Ankit Raj, Siddhesh Zadey

Drafting of the manuscript: Chaitanya Reddy, Varun Raj Passi

Critical revision of the manuscript for important intellectual content: All authors

Statistical analysis: Harsh Thakkar, Aamir Miyajiwala, Sahil Kale, Ankit Raj, Siddhesh

Zadey

Administrative, technical, or material support: Siddhesh Zadey

Study supervision: Siddhesh Zadey

Acknowledgments: Not Applicable

Clinical Trial Number: Not Applicable

ABSTRACT

Background: While studies have investigated the availability of Medical College Hospitals

(MCHs) in India, data on geographic accessibility are limited. Our study examines current

geographic accessibility to these MCHs across 36 states and union territories and 735

districts.

Methods: We validated MCH data from the National Health Profile 2022, using travel-time

rasters from the Malaria Atlas Project (2019) and population data from WorldPop (2020). We

examined the density of MCHs per million population and the median travel time to the

nearest MCH. We assessed the Access Population Coverage (APC), defined as the proportion

of the population within 30, 60, 90, and 120 minutes by motorized transport, and within 30

and 60 minutes by walking, to the nearest MCH.

Results: In 2022, India had an average density of 0.47 MCHs per million. The median travel

time to the nearest MCH was 67.94 minutes by motorized transport and 589.82 minutes by

walking. 71.76% of the population could access the nearest MCH within 60 minutes by

motorized transport. The APC was 64.20% within 60 minutes by motorized transport in rural

areas vs. 92.34% in urban areas. The APC within 60 minutes by motorized transport for

public MCHs was 63.62%, while that for private was 45.95%.

Conclusions: Median travel time and APC help assess geographic accessibility. Our study

found wide disparities in MCH access across Indian states and rural vs. urban areas. These

analyses can guide the optimal placement of new MCHs.

1. Background

Medical college hospitals (MCH) play an essential role in healthcare systems worldwide. They provide hands-on clinical training to medical students, act as centers of research setting standards of care, (1) and offer specialized clinical care to patients. In lower-middle-income countries (LMICs) MCHs often act as the last points of referral, providing the highest level of specialized care available. (2)

India was estimated to have 648 MCHs in 2022. (3) MCHs are essential in providing subsidized or free tertiary care services, as a large proportion of them (estimated to be 53.39% in 2022) are publicly owned. Privately owned MCHs often provide subsidized care in exchange for government concessions. They play a crucial role in emergencies like trauma, maternal and newborn care, and non-communicable diseases, as public tertiary centers have limited capacity. MCHs also improve local primary health care in their surrounding region by providing outreach services. (4) Therefore, MCHs act as essential stakeholders in the healthcare system of their geographical area. Provinces with higher MCH density in India perform better on major health indicators, such as the maternal mortality ratio (MMR), infant mortality rate (IMR), and the percentage of institutional deliveries. (5)

Given the importance of MCHs in India's public health system, it is important to study their growth and distribution in the country. There has been a rapid growth in the total number of MCHs in India in the last few decades. An important feature of this growth is the increasing involvement of the private sector in medical education. (4) However, the distribution and growth of MCHs are not even. MCHs are heavily clustered in provincial capitals and major cities that already have established and well-functioning MCHs. Out of 207 newly established MCHs between 2009 and 2019, 139 (67.15%) were located within 50km of an old

MCH. This unequal distribution of MCHs is starker for private MCHs, with 60.6% of all

private MCHs in the country located in a handful of provinces in the southern part of India,

i.e., Maharashtra, Andhra Pradesh, Telangana, Karnataka, Kerala, Tamil Nadu, and

Puducherry. (5)

Availability refers to the extent to which resources are available to meet patients' needs. The

number of health facilities per defined population size is referred to as the health facility

density and is a measure of the availability of care. Geographic accessibility refers to the

proximity to care, with an element of time, and delineates how easily patients can reach the

provider's location. (6) While studies have mapped the distribution and density of MCHs in

India, (4,5) geographic accessibility in terms of time and the population covered remains to

be studied. Timely access to tertiary healthcare services is necessary for optimal outcomes,

particularly in time-sensitive emergencies such as stroke, (7) acute coronary syndromes, (8)

surgical, and maternal and child health emergencies. Poor access to specialized care also

increases costs due to travel and longer lost wages. (9–11) Consequently, geographic

accessibility to MCHs constitutes an important component of moral capital in the country.

(12)

We evaluated geographic accessibility to MCHs using density per million people, travel time

to the nearest MCH, and access population coverage (APC). For travel time and APC, two

modes of transport were considered: motorized and walking. These modes correspond to the

two travel-time friction-surface rasters available through the Malaria Atlas Project (MAP),

(13) which is one of the most widely validated and frequently used global datasets for

geospatial accessibility modelling. Motorized and walking surfaces capture two extremes of

mobility in the Indian context: (i) the dominant mode with optimal speeds for medical travel

(motorized), and (ii) a lowest mobility scenario (walking) to reflect populations without

transport access, especially in rural and remote regions. These rasters incorporate terrain,

slope, land cover, and road networks, which makes them suitable for nationwide modelling.

Estimates were provided for 735 districts across 36 Indian states and union territories. We

also analyzed disparities between rural and urban areas and public versus private hospitals.

2. Methods

2.1 Data Sources

To comprehensively map MCH densities and their travel times, the study required the collection, extraction, and collation of data from the National Health Profile (NHP) Report 2022 published by the Central Bureau of Health Intelligence. (3) The report lists active MCHs, including addresses, ownership, beds, and training intake. Government-owned MCHs and those jointly managed by the government and non-profits were classified as public; all others were classified as private. Population data came from the 2020 WorldPop, providing high-resolution (1 km²) UN-adjusted counts for India. (14) Motorized and walking friction rasters (1 km²) were sourced from the 2019 Malaria Atlas Project (MAP). MAP, an international scientific initiative, focuses on mapping global malaria responses. (13) Friction rasters are informational maps that show how much it costs to go through them, usually expressed in time units. (15) These friction rasters account for several physical and environmental variables that may impede or prolong travel. The administrative borders of India were drawn from the publicly available shapefile. (4,16) A global raster with ordinal catchment area (CA) categories based on population densities and proximity to high-density urban areas provided standardized rural-urban regions. (17) We classified CA categories >7 as rural.

2.2 Outcomes

We examined three primary outcomes: MCH density, measured as the number of MCHs per million people to assess geographic distribution; travel time to the nearest MCH, reporting median and interquartile range (IQR) values; and Access Population Coverage (APC), defined as the percentage of the population with timely access to the nearest MCH. Timely access thresholds were set at 30, 60, 90, and 120 minutes for motorized transport and 30 and

60 minutes for walking, combining population and timeliness aspects to evaluate

accessibility. (18–21)

2.3 Primary Analysis

Before geocoding, MCH addresses were cleaned for machine readability by standardizing

formats and correcting errors. Geocoding used the Google Maps API to select the best match

for each address. Coordinates were checked for duplicates and implausible locations outside

India.

Population data were obtained from WorldPop 2020 and rural-urban CA rasters. The global

CA raster (1 km² resolution) was clipped to India's boundary and split into rural (CA >7) and

urban (CA ≤7) layers. WorldPop 2020 data were overlaid on these layers and resampled to

align. Rural populations were weighted by 1, urban by 'NA', and the totals were summed.

National, state, and district populations were aggregated using administrative boundaries.

Travel times were calculated using the Dijkstra algorithm on MAP friction surfaces,

estimating the shortest time from each 1 km² pixel to the nearest MCH. Walking and

motorized transport rasters provided travel times in minutes. Median and IQR values were

computed at various levels. APCs used thresholds of 120 minutes (motorized) and 30 minutes

(walking) in the primary analysis. Accessibility rasters were dichotomized, matched to

population rasters, and used to calculate timely access proportions.

Figure 1 describes the analysis pipeline, using rural motorized APC estimation as an

example. We have previously discussed the details in our research papers and used them to

investigate access to surgical facilities in India. (20–25)

Figure 1: Raster-based estimation pipeline used for creating criteria-specific regional population counts. Here, we focus on rural motorized APC as an example.

2.4 Sensitivity Analyses

To assess the robustness of our primary estimates, we conducted a sensitivity analysis using multiple APC scenarios. First, APC was calculated assuming motorized transportation for time thresholds of 30, 60, and 90 minutes. Recognizing that a substantial share of the population may lack access to motorized transport, we included APC when walking was the mode of travel, applying 60 minute thresholds. This approach allowed us to examine how assumptions about travel mode and time thresholds influence APC estimates and to evaluate the stability of our findings across alternative accessibility scenarios.

2.5 Correlation Analysis with District-Level Universal Health Coverage (UHC) Index

We conducted a district-level correlation analysis to examine the relationship between

geographic accessibility to MCHs, measured by APC values, and the district-level Universal

Health Coverage (UHC) index (24), which served as the reference indicator. We chose the

UHC index since it is the relevant indicator of overall health system performance. To ensure

comparability, districts for which UHC index values were not reported were excluded,

resulting in a final sample (n) of 687 districts. This analysis was undertaken to assess whether

variations in geographic access to MCH are aligned with broader dimensions of health

service coverage and system strength captured by the UHC index. APC estimates were

matched to corresponding district-level UHC index values using name-based district

identifiers. This ecological exploratory association analysis used the nonparametric Spearman

rank correlation coefficient (denoted by 'R'), given the nonnormal data distribution. A p-

value of <0.05 was considered statistically significant.

3. Results

3.1 MCH Density in India

According to NHP 2022, India had 648 MCHs. Tamil Nadu had the highest number of MCHs (71), followed by Karnataka (67) and Uttar Pradesh (67). Nagaland, Ladakh and Lakshadweep had no MCHs. The national density was 0.47 MCHs per million people **(eFigure 1A)**.

Twenty-three states/union territories exceeded the national MCH density average. Puducherry had the highest density (6.47), while Bihar had the lowest (0.16). Among 735 districts, 359 lacked MCHs. Puducherry also led in district-level density (7.35 per million). Only 104 districts (14.14%) had a density above one, while 264 (35.91%) surpassed the national average. **(eFigure 1B)**.

In rural areas, 210 (32.41%) MCHs were located. Seven states (19.44%) had no rural MCHs per million people, while eight (22.22%) surpassed the national average. Puducherry had the highest rural MCH density at 9.91. In urban areas, three states (8.33%) had no MCHs per million people. However, 29 states (80.55%) had densities above the national average. Sikkim had the highest urban MCH density at 20.71.

Of the 648 MCHs, 346 (53.40%) were public, and 302 (46.60%) were private. Sikkim had no public MCHs, but 11 states (30.55%) had public MCH densities above the national average. The Andaman and Nicobar Islands had the highest public MCH density at 2.76.

Seven states (19.44%) had no private MCHs, while five states (13.88%) exceeded the national density. Puducherry had the highest private MCH density at 5.03.

3.2 Travel time to the nearest MCH

3.2.1 Motorized Transport

The national median (Interquartile Range or IQR) travel time to the nearest MCH by

motorized transport was 67.94 (41.32, 110.42) minutes **(eFigure 2A)**. Seventeen states/union

territories reported shorter travel times than the national value. Access was poor in Ladakh,

where the median travel time exceeded 591.74 minutes. Five states had median travel times

under 30 minutes, and 17 under 60 minutes. At the district level, 60.81% districts had median

travel times below the national median value. Mumbai in Maharashtra reported the shortest

median travel time of 1.89 (1.16, 2.67) minutes, whereas Anjaw in Arunachal Pradesh had

the longest travel time of 1576.80 (1136.23, 2054.06) minutes **(eFigure 2B)**. Rural areas had

a median travel time of 69.04 (42.54, 111.38) minutes compared with 25.92 (8.79, 49.47)

minutes in urban areas. Seventeen rural and 33 urban states/union territories reported median

travel times shorter than the national median value **(eFigure 2C)**.

The median travel time to public MCHs was 77.46 (47.86, 120.38) minutes **(eFigure 3A)**,

compared with 106.69 (61.67, 181.94) minutes for private MCHs **(eFigure 3B)**. Chandigarh

reported the shortest median travel time to public MCH at 6.33 minutes (4.16, 7.93),

compared with Ladakh, which reported the longest median travel time at 591.74 minutes. For

private MCHs, travel time was shortest in Puducherry at 8.25 (4.70, 12.23) minutes and

longest in Arunachal Pradesh at 856.29 minutes. At the district level, median travel times for

public MCHs ranged from 1.89 minutes in Mumbai to 1576.80 minutes in Anjaw **(eFigure**

3C). Similarly, for private MCHs, median travel times ranged from 5.69 minutes in

Hyderabad to 1895.79 minutes in Anjaw, highlighting severe disparities **(eFigure 3D)**.

3.2.2 Walking

The national median travel time (IQR) to the nearest MCH by walking was 589.82 (359.49,

913.74) minutes. Eighteen states/union territories had shorter travel times, and 17 had longer

travel times than the national median value. Lakshadweep had no median travel time value

because there were no MCHs. Chandigarh reported the shortest travel time at 58.60 (39.14,

75.13) minutes, whereas Ladakh reported the longest at 4120.86 (3353.96, 4910.96) minutes

(eFigure 4A). At the district level, 60.95% reported a median travel time below the national

median value. Mumbai had the shortest travel time of 22.13 (11.06, 30.55) minutes, while

Leh recorded the longest at 4466.33 (3786.09, 5176.35) minutes **(eFigure 4B)**. The rural

population experienced substantially longer travel time by walking with a median travel time

of 597.18 (368.41, 919.30) minutes, as compared to the urban value of 269.57 (93.13,

509.30) minutes. Among rural areas, the population residing in Chandigarh had the shortest

median travel time of 59.65 (48.84, 92.89) minutes, while among urban areas, the population

residing in Sikkim had the shortest median travel time of 48.34 (25.78, 215.33) minutes

(eFigure 4C).

For public MCHs, the national median travel time, by walking, was 687.91 (426.49,

1044.83) minutes **(eFigure 5A)**, increasing to 992.37 (567.88, 1672.34) minutes for private

MCHs **(eFigure 5B)**. Chandigarh reported the shortest public MCH time at 58.60 (39.14,

75.13) minutes, while Ladakh had the longest at 4120.86 (3353.96, 4910.96) minutes. For

private MCHs, Puducherry had the shortest median travel time at 90.82 (53.40, 132.16)

minutes, and Arunachal Pradesh had the longest at 5853.52 (4964.69, 6905.61) minutes.

District-level median travel time ranged from 22.13 (15.27, 30.55) minutes in Mumbai for

public MCHs **(eFigure 5C)** to 7322.25 (6815.12, 7973.86) minutes in Upper Dibang Valley

for private MCHs **(eFigure 5D)**.

3.3 Access population coverage (APC)

3.3.1 Motorized transport

By motorized transport, timely access to an MCH was defined as reaching a hospital within

120 minutes. The national APC was 95.94%. Lakshadweep had no MCH access within 120

minutes. In the Andaman and Nicobar Islands, under 1% of the population could reach an

MCH within 120 minutes. In Chandigarh and Delhi, 100% of the population was within 120

minutes of an MCH **(Figure 2A)**. Fourteen (1.90%) districts had no access to an MCH. Less

than 1% of the population in three districts and 100% of the population in 112 (15.23%)

districts had timely access. A total of 307 (41.76%) districts had APCs above 99% **(Figure**

2B).

Nationally, 94.55% of the rural population and 99.86% of the urban population had access to

an MCH (difference is -5.31 percentage points). Rural areas in Lakshadweep, Andaman and

Nicobar Islands, and urban areas in Ladakh had zero APCs. Chandigarh had the highest rural

APC (100%), while Ladakh had the lowest (3.60%). Urban areas in 17 states/union territories

had 100% APC, while Andaman and Nicobar Islands had the lowest (3.05%). Arunachal

Pradesh had the largest rural-urban APC gap (-55.4%), and Chandigarh had the smallest

(0.1%) **(Figure 2C)**.

Figure 2: APC values to access the nearest MCH within 120 minutes by motorized transport for A. States and union territories with the national value of 95.94% (black dotted line). B. Districts and C. Rural and urban areas. The difference is shown as the rural APC minus the urban APC, in percentage points.

Nationwide, 94.46% of the population had access to public MCHs **(Figure 3A)**, while 77.29% could reach private MCHs within 120 minutes **(Figure 3B)**. Lakshadweep had no public MCH access and four states/union territories had no private MCH access. Chandigarh and Delhi had 100% APC for both types of MCHs. At the district level, 15 (2.04%) districts lacked public MCH access, and 134 (18.23%) districts had no private MCHs. Ninety-seven (13.19%) districts with public MCHs **(Figure 3C)** and 81 (11.02%) districts with private MCHs achieved 100% APC **(Figure 3D)**. Compared to the national APC, 455 (61.90%) districts had higher public MCH access, and 412 (56.05%) districts had higher private MCH access.

Figure 3: APC values to access the nearest MCH within 120 minutes by motorized

transport for A. States and union territories with public MCHs. B. States and union

territories with private MCHs. C. Districts with public MCHs and D. Districts with

private MCHs.

3.3.1.1 Sensitivity analyses

Sensitivity analyses for 30, 60, and 90 minute motorized thresholds showed persistent

disparities **(Figure 4)**. Within 30 minutes, national APC was 39.85%, with urban (75.92%)

far exceeding rural (26.54%), a 49.38-point gap. Eleven states/union territories exceeded 50%

APC, while Ladakh and Lakshadweep had none. Public and private MCH had an APC of

32.95% and 25.32%, respectively. At 60 minutes, APC rose to 71.76%, with urban (92.34%)

exceeding rural (62.20%) by 30.14 points. Sixteen states/union territories surpassed 75%

APC, though Ladakh and Andaman and Nicobar Islands lagged. By 90 minutes, APC reached

89.40%, with urban at 98.51% and rural at 86.10%. Despite improved access, rural and

remote areas still faced major challenges.

Figure 4: Sensitivity analyses for motorized transport APC values at different time thresholds (30, 60, 90 and 120 minutes) across various areas (rural and urban) and sectors (public and private).

3.3.1.2 Association with UHC Index

In the correlation analysis with UHC index, the association was strongest for access within 30 minutes with a Spearman Rank correlation (R) of 0.308 ([95% CI: 0.239, 0.374], n=687, p <0.001) **(eFigure 6A).** It remained positive at 60 minutes (R = 0.275 [95% CI: 0.205, 0.343], n=687, p <0.001) **(eFigure 6B)** 90 minutes (R = 0.218 [95% CI: 0.146, 0.288], n=687, p <0.001) **(eFigure 6C)** and 120 minutes (R = 0.125 [95% CI: 0.050, 0.198], n=687, p=0.001) **(eFigure 6D)**.

3.3.2 Walking

By walking, timely access to an MCH meant reaching it within 30 minutes. Nationally, only 4.22% of the population had timely access. Lakshadweep, Ladakh and Nagaland had no

access, while Puducherry (29.24%) and Delhi (16.80%) had the highest APC values. Only

three states/union territories exceeded 10% APC **(Figure 5A)**. At the district level, 350

(47.61%) districts had zero APC. Mumbai had the highest APC (71.86%), while Kullu had an

APC below 1% **(Figure 5B)**. Nationally, rural access (less than 1%) lagged far behind urban

access (14.44%). Puducherry led in rural APC (13.43%), and Sikkim led in urban APC

(40.69%). Sikkim also had the largest rural-urban gap (-38.5%), while Mizoram had the

smallest (0.1%) **(Figure 5C)**.

Figure 5: APC values to access the nearest MCH within 30 minutes by walking for A.

States and union territories with the national value of 4.22% (black dotted line). B.

Districts and C. Rural and urban areas. The difference is shown as the rural APC

minus the urban APC, in percentage points.

For public MCHs, 2.96% of the population had timely access **(Figure 6A)**, compared to

1.53% for private MCHs **(Figure 6B)**. Four states/union territories had no access to public

MCHs, whereas ten had no access to private MCHs. Puducherry led in both public (19.68%)

and private (9.56%) APC. At the district level, 425 (57.82%) districts had no access to public

MCHs, and 550 (74.82%) districts had no access to private MCHs. Mumbai had the highest

public APC (71.83%) **(Figure 6C)**, and Chengalpattu led in private APC (20.96%) **(Figure**

6D). Compared to the national APC, 139 (18.91%) districts had higher APC for public

MCHs, and 93 (12.65%) districts had higher APC for private MCHs.

Figure 6: APC values to access the nearest MCH within 30 minutes by walking for A.

States and union territories with public MCHs. B. States and union territories with

private MCHs. C. Districts with public MCHs and D. Districts with private MCHs.

3.3.2.1 Sensitivity Analyses

Sensitivity analysis for the 60-minute walking threshold **(Figure 7)** showed 10.96% of the

population had timely access. Chandigarh (68.46%) and Puducherry (65.75%) had the highest

APC, while rural access (1.98%) lagged far behind urban access (35.27%). Only 1.90% of

districts exceeded 50% APC, with Shahdara in Delhi leading at 99.80%. Access to public

(7.95%) and private (4.86%) MCHs was limited, with 56.19% of districts lacking public

MCHs and 73.19% lacking private MCHs. Shahdara in Delhi (99.80%) and Bangalore in

Karnataka (51.81%) led in public and private access, respectively, with urban areas

consistently outperforming rural areas.

366
Figure 7: Sensitivity analyses for walking APC values at different time thresholds (30

and 60 minutes) across various areas (rural and urban) and sectors (public and

private).

3.3.2.2 Association with UHC Index

From correlational analysis, districts where a higher proportion of the population could reach

MCHs by walking within shorter time limits tended to have better UHC index scores, with an

R = 0.297, 95% CI: 0.228, 0.363, n=687, p <0.001 for 30 minutes **(eFigure 7A)** and R =

0.316, 95% CI: 0.248, 0.381, n=687, p <0.001 for 60 minutes **(eFigure 7B)**.

4. **Discussion**

Using the NHP 2022 report, we assessed MCH densities per million, median travel times (motorized and walking) and APC up to the district level in India. Nationally, India had 0.47 MCHs per million. Median travel times were 67.94 minutes (motorized) and 589.82 minutes (walking). By motorized transport, APC was 39.85% (30 minutes), 71.76% (60 minutes), 89.40% (90 minutes), and 95.94% (120 minutes). By walking, APC was 4.22% (30 minutes) and 10.96% (60 minutes). We observed a consistent positive association with APC by motorized transport across all travel time thresholds.

The observed positive direction and statistical significance of the correlations provide preliminary evidence for the expected relationship between APC and the UHC index. Districts with higher APC values reported higher UHC index values. The association between APC and UHC index was comparable between motorized transport and walking. The strongest association was seen for the time threshold of 30 minutes by walking . This finding is important because walking access reflects local service availability and is less influenced by transport infrastructure. The analysis did not account for confounding factors that could influence both ease of access and health system performance. The strength of the association also differed across various districts within the same state.

MCH density rose from 0.41 to 0.47 per million since 2019. Puducherry had the highest density, while Nagaland and Ladakh had zero MCHs, unchanged from 2019. (5) Our study aligns with international findings. Bangladesh has 0.44 teaching hospitals per million, similar to India's 0.47. District disparities are significant, with travel times to the nearest tertiary hospital ranging from 30 minutes to 276 minutes. (26,27) The significant disparities in availability and accessibility between rural and urban areas are also consistent. (27) For

example, Dhaka, Bangladesh's capital, hosts 16 (45.71%) of the country's 35 tertiary health

centers. Similarly, 92.5% of Sub-Saharan Africa's population lives within two hours of a

hospital offering essential surgical services, comparable to India's 95.94% accessibility by

motorized transport. (28) Therefore, the patterns and values observed are similar to those of

South Asian countries and other LMICs.

Density shows facility distribution but not accessibility. Our study is the first to calculate

median travel times and APC for MCHs in India. Travel time depends on roads, terrain, and

traffic, not just distance. Geographic features such as rivers and mountains also affect

movement. Recent global modelling work shows that terrain ruggedness, slope and elevation

can substantially increase travel time even where distances are short. (29) Hydrological

barriers such as rivers, wetlands, and coastal inlets also create major discontinuities in travel

networks, often requiring long detours or slower modes of transport. (17) In low- and middle-

income countries where road networks are heterogeneous, these geographic features produce

large variations in travel time to tertiary facilities. (27,28) Therefore, accessibility is better

captured through travel-time modelling than density measures, because friction surfaces

integrate these real-world constraints. APC, which incorporates the population distribution

and the time required to reach the nearest MCH, provides a more meaningful indicator of

accessibility and should guide the strategic placement of new MCHs.

Several states/union territories need policy attention. Ladakh has the longest travel times. In

Andaman and Nicobar, less than 1% could reach an MCH within 120 minutes. Northeastern

states also have poor access, with APCs far below the 71.6% national average. Severe rural-

urban disparities exist. Rural areas had a median travel time of 69.04 minutes, compared to

25.92 minutes in urban areas. Only 62.20% of rural residents could reach an MCH within 60

minutes, versus 92.34% in urban areas. Given that 68.8% of India's population is rural, (30)

this gap is alarming. Limited access to specialized care worsens outcomes, especially in

emergencies, and increases the reliance on unqualified providers. (31)

Rural-urban access gaps may result from poorer rural road infrastructure. In 2018-19, urban

road density was 5,296.3 per 1,000 km², compared to 1,458.1 in rural areas. Growth of

Private MCHs may have also contributed to this gap, as private MCHs, reliant on student fees

and patient revenue, are concentrated in wealthier districts. (32,4) The per-capita net value

added was estimated at INR 40925 and 98435 in rural and urban areas, respectively, in 2011.

(33) Urban populations' higher paying capacity attracts private MCHs. Better urban

infrastructure also draws more private investment. (4) Additionally, health professionals in

India prefer urban areas for better amenities and career growth. This helps city-based MCHs

meet NMC faculty requirements more easily. (5)

Public MCHs are more accessible (63.62%) than private MCHs (45.95%), partly because

they have more facilities (346 vs. 302). Private MCHs tend to be in wealthier districts, where

unregulated placement may not align with healthcare needs, further affecting accessibility.

(4)

The strategic placement of new MCHs can address disparities in geographic accessibility to

MCHs. The use of location-allocation models (LAMs) can help policymakers locate MCHs at

optimal locations, thereby improving accessibility. (34) Plans such as the central

government's proposal to convert district hospitals into MCHs (34), particularly in

underserved districts, can also reduce disparities. Improving roads in rural areas through

schemes such as the Pradhan Mantri Gram Sadak Yojana (PMGSY) can also enhance

accessibility. (35) It has also been noted that alumni of MCHs located in rural areas are more

inclined towards rural practice. (5) Following the World Health Organization's

recommendation of establishing MCHs away from major cities can increase the retention of

doctors in rural areas, improving accessibility. (35)

Limitations and Strengths

The study has limitations but key strengths. It relied on assumptions from source data and

excluded vehicle ownership, facility quality, and non-land transport. Demographic-specific

access was not analyzed. We could not analyse for specific transport modes such as private

motorized transport, intermediate public transport or two-wheelers because validated national

raster data were not available. As a result, our findings capture overall travel time patterns

rather than differences to MCH access between individual transport modes. Despite this, the

study introduced a novel method to assess MCH accessibility in India, analyzing density,

travel times, and APCs across 36 states/union territories and 735 districts. It compared rural

and urban access to public and private MCHs, with validated geolocation strengthening the

findings. Additionally, sensitivity analysis ensured the robustness of our findings.

Future Directions

Future research should examine accessibility at finer geographic and demographic resolutions

better to understand inequities in travel time and service reach. District- or state-specific

analyses using higher-resolution road, traffic, and transport-mode data could help

disaggregate motorized travel into public and private options and more accurately reflect real

mobility patterns. Linking accessibility metrics to outcomes, such as maternal mortality,

neonatal outcomes, emergency referrals, and surgical delays, would also strengthen

validation efforts. Additionally, integrating facility readiness, service availability, and quality

of care into accessibility models would provide a more holistic understanding of MCH access

in India and guide more targeted policy interventions.

5. Conclusion

Median travel time to MCHs and Access Population Coverage (APC) are key tools for evaluating geographic accessibility and guiding policy. Our study revealed poor MCH access in Ladakh, the Andaman and Nicobar Islands, and the northeastern states of Arunachal, Nagaland, Mizoram, and Meghalaya. Rural areas showed significantly worse access than urban areas. Future research should examine accessibility for different demographics and consider the availability and quality of facilities in each MCH.

6. References

1. Cardinal L, Kaell A. The role of medical education in the development of the scientific practice of medicine. J Community Hosp Intern Med Perspect. 2017 Jan;7(1):58–60.

2. Clarke-Deelder E, Afriyie DO, Nseluke M, Masiye F, Fink G. Health care seeking in modern urban LMIC settings: evidence from Lusaka, Zambia. BMC Public Health. 2022 Jun 16;22(1):1205.

3. Central Bureau of Health Intelligence (CBHI), Directorate General of Health Services, Ministry of Health and Family Welfare, Government of India. National Health Profile 2022 17th Issue. Central Bureau of Health Intelligence (CBHI), Directorate General of Health Services, Ministry of Health and Family Welfare, Government of India; 2022.

4. Sabde Y, Diwan V, De Costa A, Mahadik VK. Mapping the rapid expansion of India's medical education sector: planning for the future. BMC Med Educ. 2014 Dec 17;14:266.

5. Sabde Y, Diwan V, Mahadik VK, Parashar V, Negandhi H, Trushna T, et al. Medical schools in India: pattern of establishment and impact on public health - a Geographic Information System (GIS) based exploratory study. BMC Public Health. 2020 May 24;20(1):755.

6. McLaughlin CG, Wyszewianski L. Access to care: remembering old lessons. Health Serv Res. 2002 Dec;37(6):1441–3.

7. Powers WJ, Rabinstein AA, Ackerson T, Adeoye OM, Bambakidis NC, Becker K, et al. Guidelines for the early management of patients with acute ischemic stroke: 2019 update to the 2018 guidelines for the early management of acute ischemic stroke: A guideline for healthcare professionals from the American Heart Association/American Stroke Association. Stroke. 2019 Dec;50(12):e344–418.

8. European Society of Cardiology. 2023 ESC Guidelines for the management of acute coronary syndromes [Internet]. 2023 ESC Guidelines for the management of acute coronary syndromes. 2023 [cited 2024 Jul 14]. Available from: https://www.escardio.org/Guidelines/Clinical-Practice-Guidelines/Acute-Coronary-Syndromes-ACS-Guidelines

9. Mahajan A, Tirakotai W, Masayaanon P. The hidden costs of universal health coverage: solutions from the fight against catastrophic healthcare expenditure in Thailand. BMJ Glob Health. 2023 Jun;8(6).

10. Rocque GB, Williams CP, Miller HD, Azuero A, Wheeler SB, Pisu M, et al. Impact of travel time on health care costs and resource use by phase of care for older patients with cancer. J Clin Oncol. 2019 Aug 1;37(22):1935–45.

11. Kornelsen J, Khowaja AR, Av-Gay G, Sullivan E, Parajulee A, Dunnebacke M, et al. The rural tax: comprehensive out-of-pocket costs associated with patient travel in British Columbia. BMC Health Serv Res. 2021 Aug 21;21(1):854.

522   12.   Bartosch P, Jaye C, Crampton P. Moral economy and moral capital: A new approach
523   to understanding health systems. Soc Sci Med. 2024 Jul;352:117016.

524   13.   The Malaria Atlas Project. Explorer - Malaria Atlas Project [Internet]. [cited 2021 Apr
525   11]. Available from: https://malariaatlas.org/explorer/#/

526   14.   School of Geography and Environmental Science, University of Southampton;
527   Department of Geography and Geosciences, University of Louisville; Departement de
528   Geographie, Universite de Namur, Center for International Earth Science Information
529   Network (CIESIN), Columbia University (2018). WorldPop [Internet]. 2020 [cited
530   2024 Jul 5]. Available from: https://dx.doi.org/10.5258/SOTON/WP00671

531   15.   European Commission. Global Accessibility Map [Internet]. Global Accessibility
532   Map. [cited 2024 Jul 14]. Available from:
533   https://forobs.jrc.ec.europa.eu/gam/description

534   16.   Meyers J. India Official Boundaries 2019 [Internet]. 2020 [cited 2024 Jul 12].
535   Available from: https://github.com/justinelliotmeyers/India_Official_Boundaries_2019

536   17.   Cattaneo A, Nelson A, McMenomy T. Global mapping of urban-rural catchment areas
537   reveals unequal access to services. Proc Natl Acad Sci USA. 2021 Jan 12;118(2).

538   18.   Poddar S, Gorkar A, Passi VR, Thakkar H, Andhale A, Puntambekar V, et al.
539   Geographical Access to COVID-19 Testing Centers in India. COVID-19 Testing
540   Centers Accessibility India. 2025;

541   19.   Sharma P, Thakkar H, Patil A, Chauhan P, Chembon P, Aj S, et al. Access to palliative
542   care in India: situational analysis and modeling of access from public healthcare
543   centers. Res Sq. 2025 Apr 29;

544   20.   Patil S, Miyajiwala A, Arora A, Raj A, Thakkar H, Takalika M, et al. Geospatial
545   accessibility to primary and community health centres in india. medRxiv. 2025 May
546   13;

547   21.   Urs G, Miyajiwala A, Raj A, Zadey S. Timely Geographic Access to Blood Banking
548   Facilities: A Pan-India Modeling Study. medRxiv. 2025 May 19;

549   22.   Poddar S, Gorkar A, Passi VR, Thakkar H, Andhale A, Puntambekar V, et al.
550   Geographical access to COVID-19 testing centers in India. BMC Health Serv Res.
551   2025 Nov 28;25(1):1551.

552   23.   Thakkar PSH, Patil A, Chauhan P, Chembon P, Joseph SA, Rana S, et al. Access to
553   palliative care in India: situational analysis and modeling of access from public
554   healthcare centers. Ecancermedicalscience. 2025 Nov 19;19.

555   24.   Zadey S. Measuring Access to Surgical Care in Rural India: Synthesis of Data and
556   Novel Index [Internet] [Master thesis]. Duke Global Health Institute (DGHI); 2021
557   [cited 2021 Aug 28]. Available from: https://hdl.handle.net/10161/23143

558   25.   Jadhav T, Vissoci JRN, Zadey S. Measuring timely geographical access to surgical
559   care in India: a geospatial modelling study. Lancet Glob Health. 2022 Mar;10 Suppl
560   1:S29.

26. Loveday J, Sachdev SP, Cherian MN, Katayama F, Akhtaruzzaman AKM, Thomas J,
et al. Survey of emergency and essential surgical, obstetric and anaesthetic services
available in bangladeshi government health facilities. World J Surg. 2017
Jul;41(7):1743–51.

27. Rahman M, Islam Z, Alamgir HM. How Geographically Accessible is Tertiary Care
in Low- and Middle-income Countries: The Bangladesh Case. J Health Manag. 2024
Apr 19;

28. Juran S, Broer PN, Klug SJ, Snow RC, Okiro EA, Ouma PO, et al. Geospatial
mapping of access to timely essential surgery in sub-Saharan Africa. BMJ Glob
Health. 2018 Aug 16;3(4):e000875.

29. Weiss DJ, Nelson A, Gibson HS, Temperley W, Peedell S, Lieber A, et al. A global
map of travel time to cities to assess inequalities in accessibility in 2015. Nature. 2018
Jan 18;553(7688):333–6.

30. Chandramouli C. Rural Urban Distribution of Population, Census of India 2011
[Internet]. REGISTRAR GENERAL & CENSUS COMMISSIONER, INDIA -
MINISTRY OF HOME AFFAIRS, NEW DELHI: 15th July 2011; 2011 [cited 2021
Feb 10]. Available from: https://censusindia.gov.in/2011-prov-
results/paper2/data_files/india/Rural_urban_2011.pdf

31. Anand G, Chhajed D, Shah S, Atkins S, Diwan V. Do qualifications matter? A
qualitative study of how villagers decide their health care providers in a developing
economy. PLoS ONE. 2019 Aug 1;14(8):e0220316.

32. Ministry of Road Transport & Highways, Government of India. Basic Road Statistics
of India 2018-19. Ministry of Road Transport & Highways, Government of India;
2019.

33. Ministry of Statistics & Programme Implementation. Per Capita Income in Rural and
Urban Areas [Internet]. 2023 [cited 2024 Jul 12]. Available from:
https://pib.gov.in/PressReleaseIframePage.aspx?PRID=1913325

34. Rahman S, Smith DK. Use of location-allocation models in health service
development planning in developing nations. European Journal of Operational
Research. 2000 Jun;123(3):437–52.

35. WHO. Increasing access to health workers in remote and rural areas through improved
retention: Global Policy Recommendations [Internet]. Increasing access to health
workers in remote and rural areas through improved retention. 2010 [cited 2024 Jul
12]. Available from: https://www.who.int/publications/i/item/increasing-access-to-
health-workers-in-remote-and-rural-areas-through-improved-
retention#:~:text=The%20World%20Health%20Organization%20(WHO,better%20wo
rking%20and%20living%20conditions.