An atlas of genetic effects on the monocyte methylome across European and African populations

Running title: Multi-ancestry monocyte methylome study

Wanheng Zhang[1,2,#], Chuan Qiu[3,#], Xiao Zhang[3], Zichen Zhang[1], Kuan-Jui Su[3], Zhe Luo[3], Minghui Liu[4], Bingxin Zhao[5], Lang Wu[6], Qing Tian[3], Hui Shen[3,*], Chong Wu[1,7,*] and Hong-Wen Deng[3,*]

[1]Department of Biostatistics, The University of Texas MD Anderson Cancer Center, Houston, TX 77030, USA.

[2]Department of Biostatistics and Data Science, School of Public Health, The University of Texas Health Science Center at Houston, Houston, TX 77030, USA

[3]Division of Biomedical Informatics and Genomics, Tulane Center for Biomedical Informatics and Genomics, Deming Department of Medicine, Tulane University, New Orleans, LA 70112, USA

[4]Department of Molecular and Cellular Oncology, The University of Texas MD Anderson Cancer Center, Houston, TX 77030, USA

[5]Department of Statistics and Data Science, University of Pennsylvania, Philadelphia, PA 19104, USA

[6]Cancer Epidemiology Division, Population Sciences in the Pacific Program, University of Hawaii Cancer Center, University of Hawaii at Manoa, Honolulu, HI, 96813, USA

[7]Institute for Data Science in Oncology, The UT MD Anderson Cancer Center, Houston, TX, 77030, USA.

[#]*These authors contribute equally*

Corresponding authors:

Hui Shen (hshen3@tulane.edu), Chong Wu (cwu18@mdanderson.org), and Hong-Wen Deng (hdeng2@tulane.edu)

Hui Shen, Ph.D.
Professor
Associate Director, Center for Biomedical Informatics & Genomics
Associate Director, Tulane Integrated Institute of Data and Health Sciences (TIIDHS)
School of Medicine, Tulane University
1440 Canal Street, Suite 1621, New Orleans, LA 70112, USA
Phone: 504-988-6987
Email: hshen3@tulane.edu

Chong Wu, Ph.D.
Assistant Professor
Department of Biostatistics, University of Texas MD Anderson Cancer Center
7007 Bertner Avenue, Unit 1689, Houston, TX 77030
Phone: 713-409-5160
Email: cwu18@mdanderson.org

Hong-Wen Deng, Ph.D.
Professor
Aron Family Endowed Chair
Director, Center for Biomedical Informatics & Genomics
Director, Tulane Integrated Institute of Data and Health Sciences (TIIDHS)
School of Medicine, Tulane University
1440 Canal Street, Suite 1621, New Orleans, LA 70112, USA
Phone: 504-988-1310
Email: hdeng2@tulane.edu

Abstract

Elucidating the genetic architecture of DNA methylation (DNAm) is crucial for decoding complex disease etiology. However, current epigenomic studies are often constrained by limited methylation coverage and tissue heterogeneity. Here, we present the first comprehensive, multi-ancestry methylome atlas of purified human monocytes, which play a central role in immune response and inflammation, generated through integrated whole-genome bisulfite sequencing and whole-genome sequencing from 160 African Americans (AA) and 298 European Americans (EA). By analyzing over 25 million CpG sites, we identified 3,634,383 and 3,261,407 CpG sites having *cis*-methylation quantitative trait loci (meQTLs) in AA and EA populations, respectively, revealing both shared (1,046,098 sites) and population-specific regulatory patterns. Furthermore, we developed population-specific DNAm imputation models, enabling methylome-wide association studies (MWAS) across 2,677,714 and 1,976,046 CpG sites in AA and EA, respectively. These models were validated through multi-ancestry analysis of 41 complex traits from the Million Veteran Program. All meQTLs, MWAS models, and related resources are publicly available at www.gcbhub.org and https://osf.io/gct57/.

Keywords: meQTL; heritability; WGBS; MWAS; MVP.

Introduction

Unraveling the functional consequences of genetic variation in complex human diseases remains a central challenge in human genetics. While genome-wide association studies (GWAS) have identified numerous single nucleotide polymorphism (SNP)-trait associations across a wide range of complex diseases and traits,[1] the molecular mechanisms through which these variations contribute to disease risk remain largely elusive.[2,3] This knowledge gap hinders the identification of therapeutic targets[4] and underscores the importance of exploring multi-omics biomarkers[5]. Among these, DNA methylation (DNAm), a relatively stable epigenetic modification, has emerged as a critical molecular biomarker in this endeavor. DNAm not only provides insights into the current state of human health[6] but also facilitates the estimation of epigenetic age[7–9] and partially captures the impact of environment and lifestyle exposures[10] on disease pathogenesis.

Recognizing its importance, extensive efforts have been devoted to generating comprehensive, population-scale datasets of the human genome and methylome.[11–19] However, these pursuits have been constrained by several key limitations. Firstly, most epigenomic studies rely on Illumina BeadChip platforms [20], which target only 450,000 or 935,000 predefined CpG sites—capturing only ~1.5-3% of the ~30 million CpG sites in the human genome [21–24]. This limited coverage restricts our ability to fully explore the methylome's extensive landscape. Secondly, the majority of DNAm studies are based on whole blood [25–28] or bulk tissue samples[29–31], incorporating the estimated cell type proportions as covariates in subsequent analysis[32,33]. This approach inherently limits the examination of cell type-specific methylation patterns, which are crucial for elucidating disease mechanisms.[34,35] Monocytes, in particular, play central roles in various physiological and pathological processes, including bone remodeling[36,37], neurodegenerative disorders such as Alzheimer's disease[38,39], inflammatory conditions like rheumatoid arthritis[40], cardiovascular diseases[41], and cancer[42], making them a highly relevant cell type for investigating the interplay between genetic variation, DNAm, and complex diseases. Thirdly, both DNAm and genetic variations can be ancestry-specific[43–45], highlighting the critical need to study DNAm across diverse populations to comprehensively characterize these relationships.

To overcome these limitations, we performed whole genome bisulfite sequencing (WGBS) on purified monocyte DNA and whole genome sequencing (WGS) on blood-derived DNA from 160 African Americans (AA) and 298 European Americans (EA) participants in the Louisiana Osteoporosis Study (LOS) [46]. Our comprehensive, multi-ancestry approach encompassed several key analyses. For each ancestry group, we conducted methylation quantitative trait loci (meQTLs) analyses to identify genetic variants associated with DNAm in both *cis*- and *trans*-regions, and compared findings across populations to uncover shared and ancestry-specific regulatory architecture. Further, we estimated *cis*-heritability (*cis*-h^2) of DNAm to quantify the proportion of methylation variance attributable to *cis*-acting genetic variation. To investigate the connection between DNAm and complex traits, we conducted methylome-wide association studies (MWAS) using a two-step instrumental variable regression framework[47,48]. First, we developed ancestry-specific DNAm imputation models using penalized regression. Second, we tested the association between the predicted DNAm and 41 phenotypes using GWAS summary

statistics from the Million Veteran Program (MVP) dataset [49] for both EA and AA groups. We further analyzed Type 2 diabetes (T2D) results and validated them through colocalization and Mendelian randomization (MR) analyses. This multi-faceted approach provides novel insights into the genetic regulation of monocyte-specific DNAm and its relevance to complex traits across diverse ancestries. An overview of the study design is shown in Fig.1, and the developed models and data resources are publicly available at www.gcbhub.org and https://osf.io/gct57/.

Methods

Study subjects

We recruited a total of 495 male subjects (185 AA and 310 EA), aged 37.2 ± 8.8 years (mean ± SD), from our ongoing Tulane LOS cohort. We excluded individuals with preexisting conditions affecting bone development and immune function, including: (1) cerebral vascular disease, (2) diabetes mellitus, except for easily controlled, noninsulin dependent cases, (3) chronic renal or liver failure, (4) chronic lung disease, (5) chronic obstructive pulmonary disease, (6) any metabolic or inherited bone diseases (e.g., hyper/hypoparathyroidism, Paget's disease, osteomalacia, osteogenesis imperfecta, and hypochondrogenesis), (7) collagen disorder (e.g., rheumatoid arthritis, except for minor cases that involve only hand joint and wrist), (8) chronic gastrointestinal disease, (9) alcohol abuse, (10) treatment with corticosteroid or anticonvulsant therapy for more than 6 months duration, (11) antibiotic usage, (12) gastroenteritis, (13) major surgeries, (14) intercontinental travel in the past 3 months, (15) autoimmune or autoimmune-related diseases (e.g., multiple sclerosis), (16) immune-deficiency conditions (e.g., HIV infection), (17) haematopoietic and lymphoreticular malignancies (e.g., leukaemias), (18) active periods of asthma, or (19) influenza, infected within one week of recruitment. All qualified individuals signed an informed consent document prior to data and biosample collection. The study was approved by the Tulane University Institutional Review Board (IRB #: 10-184088).

Whole Genome Sequencing (WGS)

Genomic DNA was extracted from 10ml of fresh whole blood sample using the Gentra Puregene Blood Kit (Qiagen, USA). DNA concentration and quality were assessed using Nanodrop 2000 spectrophotometer, and samples were stored at -80 °C until further use. For each sample, 300 ng of genomic DNA was used to prepare sequencing libraries. Library preparation involved the generation of DNA Nanoballs (DNBs) through ligation-mediated polymerase chain reaction (LM-PCR), followed by single-strand separation, cyclization, and rolling circle amplification. WGS was performed on the DNBSEQ-500[TM] platform (BGI Americas, USA), with 350 bp paired-end reads in length. Sequencing reads were aligned to the GRCh38/hg38 human reference genome using Burrows-Wheeler Aligner (BWA, v0.7.12) [50]. Genomic variants calling followed the Best Practices of the Genome Analysis Toolkit (GATK, v4.0.3) using HaplotypeCaller [51,52]. The variant quality score recalibration (VQSR) was applied to obtain high-confident variants[51,52].

Isolation of monocytes and nucleic acids

In this study, we focused specifically on peripheral blood monocytes (PBMs), which can act as osteoclast precursors and play important roles in regulating bone metabolism.[35] Briefly,

peripheral blood mononuclear cells (PBMCs) were firstly separated from ~60 ml freshly collected peripheral blood, by a density gradient centrifugation method using Histopaque-1077 (Sigma-Aldrich, USA). The PBMCs were washed with 2 mM EDTA in PBS, and resuspended in 0.5% BSA and 2 mM EDTA in PBS. PBMs were then isolated from the PBMCs with a Monocyte Isolation Kit II (Miltenyi Biotec, Germany) according to the manufacturer's protocol. The kit uses negative selection to deplete unwanted cells (such as T and B cells) from PBMCs while preserving monocytes without surface-bound antibodies or beads. The isolated PBMs were examined under a microscope to assess purity and determine cell count. The genomic DNA was extracted from the fresh PBMs using the AllPrep DNA/RNA/miRNA Universal Kit (Qiagen, USA), following the manufacturer's protocol. Samples were stored at -80 °C until further use.

Whole Genome Bisulfite Sequencing (WGBS)

DNAm profiles were determined by WGBS according to previously published protocols.[53] Briefly, 100 ng genomic DNA isolated from PBMs was fragmented to ~250 bp by sonication using a Bioruptor (Diagenode, Belgium), followed by the blunt-ending, dA addition to 3'-end, and adaptor ligation. The ligated libraries were bisulfite converted using the EZ DNA Methylation-Gold kit (Zymo Research, USA) and sequenced on the Illumina HighSeq 4000 platform (BGI Americas, USA). Raw reads were filtered to remove adaptors, contaminants, and low-quality sequences. Reads with >10% unknown bases or >10% of bases with quality scores <20 were further excluded to obtain high-quality data. The cleaned reads were aligned to the GRCh37/hg19 human reference genome using BSMAP[54], then converted to GRCh38/hg38 using the UCSC LiftOver tool (https://genome.ucsc.edu/cgi-bin/hgLiftOver). The methylation level of each CpG site was determined by the ratio of the number of methylated reads to the total number of reads covering a particular cytosine site.

Data preparation

Genotype data

We followed the GoDMC[11] pipeline for the genotype data processing. Quality control (QC) was conducted on genotype data for chromosomes 1-22. SNPs that failed the Hardy-Weinburg equilibrium ($P < 1 \times 10^{-6}$) and had a minor allele frequency (MAF) < 0.01 were excluded. Duplicated SNPs and those with mismatched alleles were also removed, and indel alleles were recoded as "I" (insertion) and "D" (deletion). Next, we calculated the first 10 genetic principal components (PCs) using SNPs from HapMap3 reference panel, excluding those in regions of long-range linkage disequilibrium (LD) and retaining only SNPs with MAF > 0.2. Ancestry outliers, defined as individuals deviating more than 7 standard deviations from the mean along any PC, were excluded. Genetic PCs were then recalculated for downstream analyses.

DNAm data normalization

To transform our DNAm data into a distribution that more closely approximates a Gaussian distribution, we applied a rank-based inverse normal transformation (INT)[55]. This method comprised two steps. First, the observations are transformed onto the probability scale using the empirical cumulative distribution function (ECDF). Second, the observations are transformed onto the real line, as Z-scores, using the probit function.

Co-variates

To ensure robust analysis and account for potential confounding factors, we implemented a comprehensive covariate adjustment strategy. First, we adjusted for key demographic and lifestyle factors, including age, body mass index (BMI), smoking status, and alcohol consumption. Second, to control for population structure and technical variation, we incorporated ten genetic PCs and ten non-genetic PCs. Genetic PCs were derived to capture the major axes of genetic variation. Non-genetic PCs were obtained by performing principal component analysis (PCA) on DNAm data, using 20,000 most variable CpG sites to extract the top ten components [11]. Incorporating non-genetic PCs derived from DNAm data has been shown to significantly improve the power to detect true associations [56], while posing minimal risk of introducing collider bias [11].

meQTL analyses

We performed a comprehensive analysis of both *cis-* and *trans-*meQTL mappings using the MatrixEQTL[57] package in R, analyzing over 25 million normalized DNAm sites in AA (n = 160) and EA (n = 298) ancestries, separately. *Cis-*regions were defined as ±1 Mb from each CpG sites, with *cis-*acting meQTL (*cis-*meQTL) referring to SNPs located within this window and *trans-*acting meQTLs (*trans-*meQTLs) referring to SNPs located outside of this window. Prior to association testing, DNAm data were residualized by regressing out the covariates described in the previous section. Using MatrixEQTL[57], we identified *cis-*meQTL by testing the association between DNAm levels and genotype via linear regression, adjusting for potential confounders including age, BMI, smoking, alcohol consumption. Furthermore, we controlled for ten genetic and ten nongenetic principal components. Then, for each CpG sites j, the residual value, y_{ji}, was regressed against each SNP k:

$$y_{ji} = \alpha_{jk} + \beta_{jk}x_{ki} + \epsilon_{jki},$$

where genotype values x_{ki} were standardized to have a mean of zero and a standard deviation of one, α_{jk} was the intercept term, and β_{jk} was the effect estimate of each SNP k on each residualized CpG site j. We applied the Benjamini–Hochberg procedure to control the false discovery rate (FDR) at 0.01 for both *cis-* and *trans-*meQTLs.

Replication studies

To validate our findings and assess the consistency of identified meQTLs across different populations and platforms, we conducted comprehensive replication analyses for both the AA and EA cohorts.

For the AA population, we compared our *cis-*meQTL results with those from the Genetic Epidemiology Network of Arteriopathy (GENOA)[18] study. The GENOA study performed *cis-*meQTL mapping analysis on 961 African Americans, measuring DNAm using the Illumina HumanMethylation450 BeadChip platform. For replication analysis, we identified the subset of our *cis-*meQTLs that corresponded to CpG sites also measured in the GENOA study. We then compared the effect sizes and directions between the two studies for these overlapping sites. A successful replication was defined as a *cis-*meQTL association in the GENOA study with FDR < 0.05 and consistent direction of effect. We calculated Pearson's correlation coefficient to quantify the concordance of effect sizes between studies.

For the EA population, we evaluated replication using data from the Genetics of DNA Methylation Consortium (GoDMC)[11]. The GoDMC study measured DNAm in whole blood using the Illumina HumanMethylation450 BeadChip across 27,750 European participants. This large-scale meta-analysis identified 420,509 methylation sites associated with genetic variants. Similar to our AA replication approach, we identified the subset of our EA *cis*-meQTLs that corresponded to CpG sites measured in the GoDMC study. We then compared effect sizes and directions between studies for these overlapping associations. A successful replication was defined as a *cis*-meQTL association in the GoDMC study with FDR < 0.05 and consistent direction of effect. We calculated Pearson's correlation coefficient to assess concordance of effect size estimates between studies.

For *trans*-meQTL replication, we were limited to analyzing only the EA population since detailed trans-meQTL results were publicly available only from the GoDMC study, which exclusively contains European ancestry samples. Using the GoDMC data, we identified overlapping *trans*-meQTL associations with our EA cohort. We compared effect sizes using Pearson's correlation and assessed the proportion of associations that maintained consistent direction of effect.

Fine-mapping

To identify potentially causal variants influencing DNAm, we performed fine-mapping on *cis*-SNPs associated with each CpG site. This analysis specifically focused on CpG sites linked to at least one *cis*-meQTL in AA and EA populations, respectively. We used the *SuSiE* algorithm [58] from the *susieR* package in R to infer single effect components or credible sets for each CpG site and its associated variants. Each credible set was defined to have a 95% probability of encompassing at least one variant with a nonzero causal effect. We limited the number of credible sets per CpG site to a maximum of ten (L = 10), reflecting the assumption that up to ten causal variants may potentially regulate a single CpG site. To validate and compare fine-mapping results, we aggregated all credible sets identified by *SuSiE* for each CpG site by taking their union—thus capturing all variants with potential causal effects across the multiple credible sets (up to L=10). We then compared the sizes of these unified credible set between AA and EA populations to assess differences in fine-mapping resolution.

Cis-h^2

We employed Genome-wide Complex Trait Analysis (GCTA, version 15) [59] to estimate the h^2 – the proportion of phenotypic variance attributable to genetic factors – for each CpG of ~25 million CpG sites analyzed. Again, *cis*-regions were defined as within ±1 Mb of the CpG site, and heritability was estimated using only SNPs within these regions. GCTA constructs a genetic relationship matrix (GRM) from SNP data, which encapsulates the degree of genetic similarity between pairs of individuals. We then utilized this GRM within a restricted maximum likelihood (REML) framework to estimate h^2. This approach allows for the separation of the phenotypic variance into components attributed to genetic variance (captured by the SNPs in the *cis*-regions) and residual variance. By focusing exclusively on *cis*-regional SNPs, our analysis specifically evaluated the local genetic contribution to DNAm variance at each CpG site, providing insights into the genetic architecture of epigenetic modifications. To assess the

robustness and generalizability of our results, we compared the *cis*-h^2 estimates of CpG sites from our WGBS data to those derived from two commonly used methylation platforms: the Infinium MethylationEPIC BeadChip v2.0 that covers ~935,000 CpG sites (referred to as 900K) [60] and the Infinium HumanMethylation450 Beadchip (referred to as 450K) [61].

DNAm imputation model

In this study, we focused on CpG sites with estimated *cis*-h^2 greater than 0.01. To enhance computational efficiency during model training, we adopted a 500-kp window around each target CpG site. For genotype data, following established practices [62,63], we excluded SNPs with a MAF < 1%, ambiguous strand orientation, insertions or deletions (indels), and SNPs absent in the LD reference panel derived from the 1000 Genomes Project [64]. Our DNAm imputation model was constructed using a penalized regression approach, integrating methylation and genotype data within the defined *cis*-regions. Let Y denote the n-dimensional vector of methylation data for a given CpG site, and let X be the $n \times k$ matrix of genotype dosages for k *cis*-SNPs associated with this CpG site:

$$Y = Xw + \varepsilon,$$

where w is a $k \times 1$ vector of SNP effect size to be estimated, and ε is the random error term with a mean of zero. To estimate w, we minimized the following penalized objective function $f(w)$:

$$f(w) = \frac{(Y - Xw)'(Y - Xw)}{N} + J_\lambda(w),$$

where $J_\lambda(w)$ represents a penalty term that regularizes the coefficients to prevent overfitting. Here, we used Elastic-net[65] penalty, which combined both L1 and L2 regularization terms:

$$J_\lambda(w) = \lambda \left(\alpha \sum |w_j| + \frac{1 - \alpha}{2} \sum w_j^2 \right),$$

where λ is a tuning parameter that controls the overall strength of the penalty, α is the elastic-net mixing parameter, which is 0.5 in this study, that determines the trade-off between L1 penalty (lasso) and L2 penalty (ridge). λ is chosen via cross-validation.

The performance of the imputation models was evaluated by predictive R^2, defined as the squared Pearson correlation coefficient between genetically predicted and directly measured DNAm data in nested cross-validation[66]. We only considered models with $R^2 > 0.01$ in the subsequent analysis[47].

MWAS association testing in MVP database

In this section, we tested the association between genetically predicted DNAm levels and trait of interest. For phenotypes with only summary-level GWAS data available, we adopted the methodology described in previous studies[67,68] to mitigate potential biases arising from discrepancies in LD matrices between reference panels and GWAS data. In our MWAS test, we estimated the effect size $\hat{\gamma}$ and its variance for the association between predicted DNAm and the phenotypes by:

$$\hat{\gamma} = \frac{\hat{w}' Z / \sqrt{n_s}}{\sigma_r} \text{ and } \widehat{Var}(\hat{\gamma}) = \left(\frac{1}{n_s} + \frac{1}{n_r} \right) \hat{\gamma}^2 + \frac{\zeta^2}{n_s \sigma_r},$$

where Z is the vectors of z-scores from GWAS, $\hat{w}$ represents the estimated weights from the DNAm imputation model, n_s is the sample size of the GWAS data, n_r is the sample size of the

population reference panel, $\sigma_r = \frac{(G_r\hat{w})'(G_r\hat{w})}{n_r}$, $\zeta^2 = 1 - \frac{2\hat{w}'Z\hat{\gamma}}{\sqrt{n_s}} + \sigma_r\hat{\gamma}^2$, G_r is the standardized genotype matrix of the population reference panel.

We applied this MWAS framework to 41 phenotypes available in the MVP[49] database, for which GWAS summary statistics were available for both African and European ancestries. Details of these GWAS summary statistics are provided in Supplementary Table 1. Prior to association testing, we implemented QC on the GWAS summary statistics, including removal of duplicate entries, correction of strand orientation, and exclusion of ambiguous alleles, ensuring the integrity and reliability of our association analysis. To ensure robust methylation prediction, we excluded CpG sites with the number of non-zero weights from DNAm imputation models smaller than 10.

Post-MWAS analyses

We compared our MWAS results with MR analysis in line with the methodology outlined by Zhao, et al.[69]. We selected a set of *cis*-meQTLs as the MR instrumental variables. We first restricted our analysis to the common set of variants that were shared by *cis*-meQTLs and the GWAS summary statistics. To avoid the potential issue of collinearity, we applied LD clumping and excluded *cis*-meQTLs with high correlation to index genetic variants ($r^2 > 0.001$). Then, we removed variants that were not significant *cis*-meQTLs ($P > 10^{-8}$). To further ensure the validity of the instruments, we applied the Steiger filtering method [70] to exclude instrumental variables with potential reverse causality. Depending on the number of valid instrumental variables, we then applied either the Wald ratio method (for a single instrument) or inverse variance weighting[71] (for multiple instruments) to assess the causal relationship between DNAm and the phenotype.

We also conducted Bayesian colocalization analyses[72] on the significant CpG sites identified in our MWAS and MR analysis to estimate the posterior probability that the DNAm site and phenotype shared the same causal variant, using summary-level *cis*-pQTLs and GWAS data. Specifically, we used the coloc R package (v5.2.2), with its default setups in the coloc function, to estimate the posterior probability of both protein and phenotype being influenced by the same causal variant (i.e., the PPH4). We chose PPH4 > 0.8 as the threshold. DNAm satisfying this threshold would suggest a shared causal variant for the *cis*-meQTLs and GWAS associations.

Results

Data collection and processing

We recruited 495 male subjects aged 20-51 years (185 AA and 310 EA) from the ongoing Tulane LOS cohort[46] with strict inclusion/exclusion criteria; see Method section for details. We restricted our analyses to subjects with purified PBMs. Sequencing reads were processed to remove adaptor sequences, contamination, and low-quality bases, then aligned to the human reference genome GRCh38/hg38. In total, methylation levels for over 25 million CpG sites across the genome were quantified (see Methods for details). DNAm β-values underwent rank-based INT[55] to normalize the data distribution. We also implemented comprehensive QC for the

genetic data. Single nucleotide polymorphisms (SNPs) were excluded based on several criteria: deviation from Hardy-Weinberg equilibrium ($P < 1 \times 10^{-6}$), MAF less than 0.01, and presence of mismatched alleles. Additionally, to mitigate confounding effects from population structure, we removed individuals with a degree of relatedness, defined as an identity by state (IBS) greater than 0.125. This resulted in a final cohort of 160 AA and 298 EA. The characteristics of the final cohort are summarized in Supplementary Table 2.

Identification of *cis*-meQTL reveals methylation landscapes across populations through WGBS

Leveraging the comprehensive coverage of WGBS, we conducted an extensive analysis of *cis*-meQTLs across 25,721,231 CpG sites in both AA (n=160) and EA (n=298) populations. Using MatrixEQTL[57], we identified *cis*-acting methylation quantitative trait loci (*cis*-meQTLs) by testing the association between DNAm and genotype via linear regression, adjusting for potential confounders including age, BMI, smoking, and alcohol consumption. Furthermore, we controlled for ten genetic and ten nongenetic principal components (see Methods).

We first computed meQTLs between each of these CpG sites and common SNPs within cis-regions, which identified 107,154,011 ($P < 3.7 \times 10^{-6}$) and 216,862,547 ($P < 1.1 \times 10^{-5}$) *cis*-meQTL associations in the AA and EA populations, respectively, when controlling FDR < 0.01. In total, 3,634,383 CpG sites in the AA and 3,261,407 in the EA population exhibited at least one significant *cis*-meQTL, corresponding to 14.1% and 12.7% of all tested sites, respectively. Among these, 1,046,098 CpG sites (28.8% of AA and 32.1% of EA) were shared between populations, indicating both common and population-specific genetic regulation of DNAm.

Our analysis uncovered several key patterns in the genetic architecture of methylation regulation. First, we observed a negative correlation between MAF and effect size in both AA ($r = -0.41, \ P < 2.2 \times 10^{-16}$) and EA ($r = -0.39, \ P < 2.2 \times 10^{-16}$) population (Fig. 2a), with larger median effect sizes in AA (0.90) than in EA (0.65, $P < 1 \times 10^{-16}$ by Wilcoxon rank-sum test). These findings extend the principles observed in GWAS studies[73,74] to methylation regulation, while highlighting important population differences. Second, we found that genetic variants closer to CpG sites typically showed stronger associations, with effect sizes decreasing as distance increased (Fig. 2b). This spatial relationship was consistent across both populations (Supplementary Fig. 1). Third, investigation of genomic context using Annotatr[75] revealed that majority of CpG sites with *cis*-meQTLs were located in open sea regions (AA: 92.5%, EA: 94.1%), with smaller proportions in CpG islands (AA: 2.2%, EA: 2.1%) and shores/shelves (AA: 5.3%, EA: 5.6%). By comparison, among all tested CpG sites, 82.9% were located in open sea regions, 9.4% in shores/shelves, and 7.7% in islands. The enrichment of meQTLs in open sea regions aligns with previous findings[18], suggesting that open sea CpGs are more susceptible to genetic regulation (Fig. 2c).

Furthermore, we found substantial population-specific genetic regulation of methylation. Notably, 31.2% of *cis*-meQTLs in the AA population were rare or absent in Europeans, only 9.5% of EA *cis*-meQTLs were rare or absent in Africans. We defined rare as two or fewer individuals carrying the same variant in the Phase-3 1000 Genome Project[64] (1000G) of the corresponding

ancestry. This disparity was even more pronounced for sentinel *cis*-meQTLs, defined as the most significant SNP per CpG site, with 50.0% of AA variants being rare or absent in Europeans compared to 19.6% of EA variants in Africans (Fig. 2d). Despite these differences, for shared variants, effect sizes showed high concordance between populations (correlation $= 0.96, P < 2.2 \times 10^{-16}$, Supplementary Fig. 2).

To narrow down 95% credible sets of causal variants underlying significant *cis*-meQTL, we conducted population-specific fine-mapping for CpG sites with at least one *cis*-meQTL in AA and EA populations using SuSiE[58]. We found that the number of variants within the 95% credible sets was notably smaller for the AA population, with a median of 14 and an interquartile range (IQR) of 8,163, as opposed to the EA population, which exhibited a median of 28 and an IQR of 5,296 (Fig. 2e). This observation likely reflects lower LD in African ancestry populations, though the smaller AA sample size may contribute to this difference.

Our identified cis-meQTLs demonstrated robust replication when compared with previous studies (Fig. 2f). In the AA cohort, we compared our results with the GENOA[76] AA cohort (N = 961). Due to the substantial difference in methylation site coverage between our WGBS approach (over 25 million CpG sites) and the 450K array platform used in GENOA, replication data were available for only 207,533 of our identified cis-meQTLs. We observed a strong correlation of effect sizes (Pearson's r = 0.70, $P < 2.2 \times 10^{-16}$), with 86.3% of associations showing consistent direction of effect. At FDR < 0.05 (P cutoff = 3.7×10^{-6}), 78.1% (162,040) of our discovery meQTLs in AA successfully replicated in the GENOA AA cohort.

For the EA cohort, we compared our findings with the GoDMC[77] study, which measured DNAm in whole blood using the Illumina HumanMethylation 450K BeadChip. Replication data were available for 550,309 of our identified cis-meQTLs in EA. Analysis revealed a strong correlation of effect sizes (Pearson's r = 0.74, $P < 2.2 \times 10^{-16}$), with 90.7% of associations maintaining consistent direction of effect. At FDR < 0.05, ($P < 1.1 \times 10^{-5}$), 95.4% (524,995) of our *cis*-meQTLs in EA replicated in the GoDMC EA cohort. These high replication rates validate the robustness of our WGBS methodology and confirm the consistency of our findings across diverse methylation platforms and populations.

Cis-h^2 of CpG sites

To quantify genetic contributions to DNAm variation, we estimated *cis*-heritability (*cis*-h^2, the proportion of methylation variation explained by *cis*-acting SNPs within 1 Mbp) using REML in GCTA[59]. Across all 25,721,231 CpG sites profiled by WGBS, we observed that 50.5% and 57.1% of CpG sites exhibited *cis*-h^2 < 0.01, 21.1% and 33.8% CpG sites with a *cis*-h^2 between 0.01 and 0.1, and 28.3% and 9.2% CpG sites with a *cis*-h^2 > 0.1 in the AA and EA populations, respectively (Fig. 3a). The average *cis*-h^2 was significantly higher in the AA population (mean=0.09) compared to EA population (mean=0.04; $P < 1 \times 10^{-16}$ by Wilcoxon rank-sum test). We provide the estimated *cis*-h^2 as valuable resources for the community to support subsequent analysis and CpG sites selection.

As expected, integration of heritability estimates with *cis*-meQTL results revealed strong concordance. For highly heritable sites (*cis*-h^2 > 0.5), the majority showed significant *cis*-meQTL associations (81.6% in AA, 99.8% in EA; Fig. 3b). CpG sites with identified *cis*-meQTLs exhibited substantially higher mean heritability (AA: 0.31, EA: 0.18) compared to those without (AA: 0.06, EA: 0.02; Fig. 3c). While AA showed higher overall heritability, we identified fewer *cis*-meQTL associations compared to EA. To assess whether this was driven by sample size differences, we performed a controlled analysis by randomly sampling 160 individuals from the EA population to match the AA sample size. This analysis identified 91,486,150 *cis*-meQTLs in EA, fewer than the 107,154,011 associations detected in AA, indicating that the original disparity resulted from sample size difference. This finding highlights the importance of increasing sample sizes in diverse populations to reduce disparity.

To evaluate methylation quantification approaches, we compared heritability patterns between WGBS and two common array platforms: the MethylationEPIC Infinium v2.0[60], which contains over 935,000 CpG sites (referred to as the 900K set), and the HumanMethylation450 Infinium assay[61], encompassing around 450,000 CpG sites (referred to as the 450K set). The annotated sites for both platforms are available at https://zwdzwd.github.io/InfiniumAnnotation. The distribution of *cis*-h^2 values for CpG sites in the 450K and 900K sets closely mirrored that of the overall WGBS dataset, albeit with slightly lower values (mean *cis*-h^2 of 900K/450K: AA = 0.08/0.08, EA = 0.03/0.03; Fig. 3d). This finding demonstrates that while array-based platforms provide efficient coverage of representative CpG sites, WGBS enables comprehensive genome-wide methylation profiling, allowing identification of additional heritable CpG sites and potentially novel regulatory regions not captured by targeted platforms.

Multi-ancestry MWAS of complex traits in MVP

To illustrate the utility of our data source, we conducted MWAS. Specifically, we first built population-specific methylation prediction models for CpG sites with *cis*-h^2 > 0.01 using Elastic Net regression[65] with nested cross-validation[78] (see Methods). For computational efficiency, we selected SNPs within ±500 kb of each target CpG site as predictors. We first constructed DNAm prediction models (with R^2 > 0.01) for 2,677,714 CpG sites in AA population and 1,976,046 CpG sites in EA populations, with 1,075,702 CpG sites shared between both groups. The mean predictive performance (measured by R^2) was 0.19 for AA and 0.17 for EA, outperforming models built on array-based CpG sites (450K/900K mean R^2: AA=0.11/0.10; EA = 0.10/0.09). These findings align with the observed heritability patterns and demonstrating the enhanced predictive power enabled by WGBS-based comprehensive methylome coverage.

Using these prediction models, we performed MWAS for 41 complex traits using GWAS summary statistics from the MVP dataset[49]. After Bonferroni correction across all models and phenotypes ($P < 4.8 \times 10^{-10}$ for AA and $P < 6.5 \times 10^{-10}$ for EA), we identified 34,174 significant methylation-phenotype associations (Supplementary Tables 3.1 and 3.2), including 31,229 associations in EA and 2,945 in AA (Fig. 4a). The higher number of associations in EA likely reflects the larger sample size in EA GWAS data (mean = 261,202) compared to AA (mean = 54,271).

Notably, our WGBS-based approach uncovered associations not detectable with array-based platforms. For AA, 2,885 (98.0%) and 2,872 (97.5%) significant associations were not captured by the 450K and 900K methylation platforms, respectively. Similarly, for EA, 30,758 (98.5%) and 30,303 (97.0%) associations were beyond the coverage of these platforms. These results highlight the advantage of WGBS for capturing the full landscape of methylation-trait associations across ancestries.

Among the identified associations, 556 unique methylation-phenotype association pairs were significant in both ancestries; the majority (534 out of 556) of these associations displayed consistent directions of effects (Fig. 4b). This consistency underscores a possible shared epigenetic mechanism across ancestries. These consistent associations encompassed 432 distinct DNAm sites and spanned 12 phenotypes categorized into six broad groups: lipids, cardiovascular diseases (CVD), anthropometric measurements, metabolic conditions, renal traits, and addiction. Genomic annotation revealed that most shared associations (529 of 594) involved CpG sites in open sea regions, with fewer sites in shores (31), shelves (19), and islands (15). This pattern highlights the regulatory potential of CpG sites in traditionally understudied genomic regions and reinforces the importance of comprehensive methylome coverage for discovering functional variations.

To aggregate CpG-level associations into gene-level insights, we mapped CpG sites from all associations to the genes and used Aggregated Cauchy Association Test (ACAT)[79] to detect significant gene-phenotype associations.[80] After Bonferroni correction ($P < 2.7 \times 10^{-6}$ for AA and $P < 2.8 \times 10^{-6}$ for EA), we identified 1,106 significant associations for AA population spanning 17 traits across 7 categories (Supplementary Table 4.1). We also identified 8,668 significant associations for EA population encompassing 30 traits and 7 categories (Supplementary Table 4.2). Among these, 746 associations are significant in both ancestries, covering 14 traits across 6 categories (Supplementary Table 5). Notably, the vast majority of gene-phenotype associations remained undetected by conventional methylation arrays: the 450K panel failed to capture 97.2% (1,075/1,106) of significant associations in the AA population and 97.5% (8,451/8,668) in the EA population. Similarly, the 900K panel missed 96.5% (1,067/1,106) and 96.8% (8,391/8,668) of significant associations in AA and EA populations, respectively. These findings highlight the substantial gain in discovery power offered by WGBS over array-based approaches.

We have made all meQTL results, MWAS models, association results, and optimized codes for replicating all results publicly available to facilitate future research endeavors (www.gcbhub.org and https://osf.io/gct57/).

Novel T2D-associated methylation signatures and ancestry-specific regulatory mechanisms
Using our meQTL data and MWAS models, we conducted a comprehensive analysis of T2D. Our MWAS analysis identified 21 CpG sites significantly associated with T2D in the AA population and 4,552 CpG sites in the EA population after Bonferroni correction. Notably, 13

CpG sites showed consistent associations across both populations, suggesting shared epigenetic mechanisms underlying T2D pathophysiology.

To validate the causal nature of these associations, we performed MR analysis, which supported causal relationships for 71.4% (15/21) of significant CpG sites in AA and 73.6% (3,351/4,552) in EA after Bonferroni correction ($P < 1.2 \times 10^{-9}$ for AA and $P < 2.3 \times 10^{-9}$ for EA). Of the 13 cross-ancestry consistent associations, 84.6% (11/13) showed evidence of causality through MR. Complementary Bayesian colocalization analysis revealed that 33.3% (7/21) of AA and 29.7% (1,353/4,552) of EA T2D-associated CpG sites demonstrated strong evidence of shared genetic architecture with T2D GWAS signals (PPH4 > 0.8), indicating these epigenetic associations likely share causal variants with genetic associations.

When aggregating CpG-level associations to genes, we identified 13 genes significantly associated with T2D in both AA and EA populations (Supplementary Table 4). The most significant associations included several established diabetes-related genes with distinct molecular functions: *TCF7L2*, whose association with T2D has been consistently replicated across populations with diverse genetic origins[81] and experimentally validated[82]; *ANK1*, a known T2D susceptibility locus[83] implicated in insulin resistance, a key feature in the development of T2D[84]; and *FTO*, which influences T2D risk primarily through BMI and obesity regulation[85]. We also identified novel associations with potential mechanistic roles: *NKX6-3*, which, along with other specific pancreatic islet β-cell transcription factors, is sensitive to oxidative stress, a condition associated with β-cell dysfunction in both Type 1 and Type 2 diabetes.[86] Previous GWAS studies suggest that *TH* may contribute to T2D susceptibility and insulin resistance. [87,88]

Notably, our analysis revealed seven T2D-associated genes unique to the AA population (Supplementary Table 4.1). Among these, *BCKDHA*, encoding the branched-chain alpha-keto acid (BCAA) dehydrogenase (BCKD) subunit, is particularly interesting. This finding connects with previous observations of elevated BCAA levels in both human and animal models of obesity[89,90], and the documented suppression of BCKD complex activity in liver and adipose tissue during obesity[91]. Given BCKD's expression across multiple tissues including monocytes, this ancestry-specific association suggests potential immune system involvement in T2D pathogenesis that may be particularly relevant in AA populations.

Comparison with array-based approaches demonstrated the superior detection power of WGBS at both CpG site and gene levels. At the CpG site level, conventional methylation arrays captured only 4.8% (1/21) and 2.4% (109/4552) of the significant T2D-associated sites identified by WGBS in AA and EA populations, respectively. At the gene level, the limitation of array-based approaches was particularly evident in T2D analysis, where arrays identified only three genes (*TCF7L2*, *FTO*, and *KCNC2*) in both ancestries, missing numerous biologically relevant associations detected through WGBS, as discussed above.

Identification of *trans*-meQTL for whole genome DNAm sites across two populations

Given our modest sample sizes (AA: n=160; EA: n=298), we conducted an exploratory analysis of *trans*-meQTL associations to gain preliminary insights into long-range methylation regulation. Controlling FDR at 0.01, our analysis revealed 12,859,517 *trans*-meQTL associations (p < 4.2×10^{-11}) involving 1,105,063 CpG sites (4.3% of all tested sites) in the AA population, and 8,850,920 *trans*-meQTL associations (p < 3.6×10^{-10}) involving 640,410 CpG sites (2.5% of all tested sites) in the EA population. The average density of trans-meQTLs per CpG site slightly differed between populations, with 0.50 and 0.34 *trans*-meQTL mappings per CpG site in AA and EA populations, respectively. Notably, we observed minimal overlap between populations, with only 4,502 shared CpG sites showing *trans*-meQTL associations. The majority of associations were population-specific, with 98.9% unique to AA and 97.8% unique to EA populations, although these patterns should be interpreted cautiously given the sample size limitations.

Initial characterization of these putative *trans*-meQTL associations revealed several patterns that warrant further investigation in larger cohorts. We observed a negative correlation between MAF and effect size for variants with MAF > 0.05 (AA: correlation = -0.23, $P = 7.1 \times 10^{-9}$, EA: correlation = -0.48, $P < 2.2 \times 10^{-16}$). Interestingly, variants with low MAFs (~0.01) exhibited unexpectedly small effect sizes (Fig. 5a). The relationship between effect size and genomic distance showed stronger distance-dependent attenuation for *trans*-meQTLs compared to *cis*-meQTLs (Fig. 5b), while effect sizes of shared *trans*-meQTLs demonstrated high concordance between populations (correlation = 0.97, $P < 2 \times 10^{-16}$, Fig. 5c).

For EA, we replicated *trans*-meQTL results from the GoDMC (EA) study (Fig. 5d). Of the trans-meQTLs identified in our EA data, 1,033 were also provided by the GoDMC. Effect sizes were strong correlated (Pearson's r = 0.51, $P < 2.2 \times 10^{-16}$), with 96.5% of associations showing a consistent direction of effect. At FDR < 0.05 ($P < 3.5 \times 10^{-10}$), 849 of the 1,033 *trans*-meQTLs (82.2%) replicated in the GoDMC EA cohort.

Discussion

We present the first comprehensive analysis of genetic influences on monocyte DNAm across AA and EA populations using WGBS. This study contributes several key advances to the field of epigenetics. First, by analyzing over 25 million CpG sites, we identified approximately ten times more *cis*-meQTL associations than previous studies[11], providing unprecedented resolution into the regulation of DNAm. Second, our population-specific DNAm imputation models, validated using the MVP dataset across 41 complex traits, demonstrate the potential of integrative MWAS to complement GWAS in identifying trait-associated CpG sites and genes that may offer insights into disease biology and potential therapeutic avenues. Third, our use of purified monocytes enables for precise characterization of cell-type-specific methylation patterns, which is critical for understanding immune-related diseases such as T2D.

Our cross-ancestry analysis offers valuable insights into population-specific methylation regulation. Notably, the AA population showed higher *cis*-h^2 of DNAm levels compared to EA. Furthermore, approximately 30% of *cis*-meQTLs identified in the AA population were rare or

absent in Europeans, whereas only 8% of EA-specific *cis*-meQTLs were rare in Africans. These findings underscore the critical importance of including diverse populations in epigenetic studies and suggest population-specific regulatory mechanisms that may contribute to differences in disease susceptibility.

The comprehensive coverage (over 25 million CpG sites) of WGBS enabled us to uncover numerous methylation markers beyond the scope of conventional array-based approaches, yielding novel biological insights. Integration of these CpG sites to genes revealed both known and novel gene-trait associations. For example, besides replicating established T2D-associated genes such as *TCF7L2* and *FTO*, we also identified novel, ancestry-specific associations — most notably *BCKDHA* in the AA population which may help explain ancestry-specific differences in disease susceptibility.

Our findings align with recent studies emphasizing that the DNAm landscape is strongly influenced by cell-type-specific regulatory programs[92]. While cell-type identity establishes the baseline methylation framework, we demonstrate that genetic variation plays a crucial role in fine-tuning these patterns, even within a homogeneous cell population like monocytes. This suggests a hierarchical model of methylation regulation where cell-type programs establish the foundational pattern, which is then modulated by genetic factors in a population-specific manner. Our analysis provides direct evidence for this model, revealing how genetic variation can influence methylation patterns within a defined cellular context.

Several limitations of our study merit discussion and offer directions for future work. First, the relatively modest sample sizes, particularly in the AA cohort, may have led to underestimation of *cis*-meQTL associations and limited our ability to detect *trans*-effects, which typically have smaller effect sizes and require larger samples to identify[93]. Nonetheless, this remains one of the largest WGBS studies to date, particularly in its inclusion of both EA and AA populations. Second, the male-predominant composition of our cohort (all males), as well as the MVP (~90% male) dataset may obscure sex-specific methylation differences. Although DNAm patterns are largely consistent across sexes[94], subtle sex-specific effects may still exist.[95] Third, while our focus on purified monocytes provides valuable insights into immune cell regulation, future studies should expand to other cell types and integrate transcriptomic data to build a more holistic view of epigenetic regulation. Our work lays the foundation for broader exploration of tissue- and context-specific methylation landscapes and their roles in complex traits and diseases.

ACKNOWLEDGEMENTS
This work is partially supported by grants from the NIH (U19AG055373, R01AG061917, R01AR069055, P20GM109036, R01CA263494). The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

Figures

Fig.1 Overview of the data collection, preparation, and analysis pipeline.

Fig.2 *cis*-meQTL and fine-mapping analysis across AA and EA populations. **(a)** Relationship between *cis*-meQTL effect sizes and minor allele frequencies (MAF(1-MAF)), with regression trends shown in orange. **(b)** Effect size distribution as a function of distance between SNP and DNAm site in AA (left) and EA (right) populations, measured in base pairs. **(c)** Genomic distribution of CpG sites with *cis*-meQTLs. The bar chart categorizes CpG sites based on the number of *cis*-meQTLs identified, with annotations for four genomic regions: CpG islands, which are regions with a high CpG density; CpG shores, located within 2 kb of islands; CpG shelves, extending an additional 2 kb from shores; and Open Sea, which represents areas more distal to the islands, shores, and shelves. Each region is color-coded. **(d)** Percentage of

nonexistent or rare *cis*-meQTLs across ancestries. It displays the proportion of identified *cis*-meQTLs that are absent or rare within the contrasting population from the 1000 Genomes Project. The four bars represent *cis*-meQTL associations for AA, EA, sentinel *cis*-meQTLs for AA and EA, respectively. Sentinel *cis*-meQTL is defined as the most significant SNPs for each CpG site. **(e)** Fine-mapping results showing the distribution of credible set sizes from SuSiE analysis for CpG sites with significant *cis*-meQTLs in both populations. Box plots show median, quartiles, and 1.5 IQR whiskers; violin width indicates data density. **(f)** Replication effect sizes between WGBS study (EA, n = 185) and GoDMC (EA, n = 27,750). Effect sizes between WGBS study (AA, n = 160) and GENOA (AA, n = 961).

Fig.3 *cis*-h^2. **(a)** Distribution of *cis*-heritability estimates across CpG sites with *cis*-h^2 > 0.01 in AA and EA populations. **(b)** Proportions of CpG sites achieving statistical significance (   ) across varying *cis*-h^2 thresholds, demonstrating the relationship between heritability magnitude and detection power. **(c)** Relationship between CpG site *cis*-h^2 and the presence of significant *cis*-meQTL associations, illustrating the concordance between heritability estimates and genetic regulation. **(d)** Comparative analysis of *cis*-h^2 distribution among CpG sites with and without significant *cis*-meQTL in AA and EA ancestries. **(e)** Comparison of *cis*-h^2 distributions between

WGBS data and conventional array-based platforms (900K and 450K), highlighting the comprehensive coverage of WGBS. Bos plots show median, quartiles, and 1.5×IQR whiskers.

Fig.4 MWAS results for 41 phenotypes in the MVP dataset. **(a)** Distribution of significant MWAS associations after Bonferroni correction ($P < 4.9 \times 10^{-10}$ for AA and $P < 6.6 \times 10^{-10}$ for EA) across different trait categories. **(b)** Cross-ancestry comparison of effect sizes (Z-scores) for CpG-phenotype associations significant in both populations. Colored points represent consistent directional effects across ancestries, while grey points indicate discordant directions. Different colors denote distinct phenotype categories.

Fig.5 Characterization of *trans*-meQTL effects across African American (AA) and European American (EA) populations. (a) Relationship between *trans*-meQTL effect sizes and MAF(1-MAF), demonstrating the impact of allele frequency on long-range methylation. Orange regression lines indicate overall trends. **(b)** Effect sizes distribution as a function of genomic distance between SNP and DNAm site in AA (left) and EA (right). **(c)** Correlation of *trans*-meQTL effect sizes between AA and EA populations for shared associations, demonstrating the consistency of trans effect across ancestries. **(d)** Replication effect sizes of WGBS study (EA, n = 185) and GoDMC study (EA, n = 27,750).

References

1. Buniello, A. *et al.* The NHGRI-EBI GWAS Catalog of published genome-wide association studies, targeted arrays and summary statistics 2019. *Nucleic Acids Res.* **47**, D1005–D1012 (2019).

2. The GTEx Consortium *et al.* The GTEx Consortium atlas of genetic regulatory effects across human tissues. *Science* **369**, 1318–1330 (2020).

3. Sun, B. B. *et al.* Plasma proteomic associations with genetics and health in the UK Biobank. *Nature* **622**, 329–338 (2023).

4. Liggett, S. B. *et al.* A polymorphism within a conserved β_1-adrenergic receptor motif alters cardiac function and β-blocker response in human heart failure. *Proc. Natl. Acad. Sci.* **103**, 11288–11293 (2006).

5. Olivier, M., Asmis, R., Hawkins, G. A., Howard, T. D. & Cox, L. A. The need for multi-omics biomarker signatures in precision medicine. *Int. J. Mol. Sci.* **20**, 4781 (2019).

6. Jones, P. A. Functions of DNA methylation: islands, start sites, gene bodies and beyond. *Nat. Rev. Genet.* **13**, 484–492 (2012).

7. Unnikrishnan, A. *et al.* The role of DNA methylation in epigenetics of aging. *Pharmacol. Ther.* **195**, 172–185 (2019).

8. Lu, A. T. *et al.* DNA methylation GrimAge strongly predicts lifespan and healthspan. *Aging* **11**, 303 (2019).

9. Marioni, R. E. *et al.* DNA methylation age of blood predicts all-cause mortality in later life. *Genome Biol.* **16**, 25 (2015).

10. Schübeler, D. Function and information content of DNA methylation. *Nature* **517**, 321–326 (2015).

11. Min, J. L. *et al.* Genomic and phenotypic insights from an atlas of genetic effects on DNA methylation. *Nat. Genet.* **53**, 1311–1321 (2021).

12. Feingold, E. A. *et al.* The ENCODE (ENCyclopedia of DNA elements) project. *Science* **306**, 636–640 (2004).

13. Bernstein, B. E. *et al.* The NIH roadmap epigenomics mapping consortium. *Nat. Biotechnol.* **28**, 1045–1048 (2010).

14. Stunnenberg, H. G. *et al.* The International Human Epigenome Consortium: a blueprint for scientific collaboration and discovery. *Cell* **167**, 1145–1149 (2016).

15. Moss, J. *et al.* Comprehensive human cell-type methylation atlas reveals origins of circulating cell-free DNA in health and disease. *Nat. Commun.* **9**, 5068 (2018).

16. Investigators, A. The atherosclerosis risk in communit (ARIC) study: design and objectives. *Am. J. Epidemiol.* **129**, 687–702 (1989).

17. Dawber, T. R., Meadors, G. F. & Moore, F. E. Epidemiological Approaches to Heart Disease: The Framingham Study. *Am. J. Public Health Nations Health* **41**, 279–286 (1951).

18. Shang, L. *et al.* meQTL mapping in the GENOA study reveals genetic determinants of DNA methylation in African Americans. *Nat. Commun.* **14**, 2711 (2023).

19. Darrell 1, C. G. A. R. N. T. source sites: D. U. M. S. M. R. 1 F. A. 2 B., J. 3 4 5, E. U. V. M. E. G. 3 4 5 B. D. J. 5 6 M. M. G. 3 O. J., Norman 8, H. F. H. M. T. 7 L., Oliver 11, M. A. C. C. A. K. 9 A. Y. W. 10 B. & Michael 13, U. of C. S. F. V. S. 12 B. M. 13 P. Comprehensive genomic characterization defines human glioblastoma genes and core pathways. *Nature* **455**, 1061–1068 (2008).

20. Pidsley, R. *et al.* Critical evaluation of the Illumina MethylationEPIC BeadChip microarray for whole-genome DNA methylation profiling. *Genome Biol.* **17**, 208 (2016).

21. de Moura, M. C. *et al.* Epigenome-wide association study of COVID-19 severity with respiratory failure. *EBioMedicine* **66**, (2021).

22. Chu, A. Y. *et al.* Epigenome-wide association studies identify DNA methylation associated with kidney function. *Nat. Commun.* **8**, 1286 (2017).

23. Zillich, L. *et al.* Epigenome-wide association study of alcohol use disorder in five brain regions. *Neuropsychopharmacology* **47**, 832–839 (2022).

24. Barbu, M. C. *et al.* Methylome-wide association study of antidepressant use in Generation Scotland and the Netherlands Twin Register implicates the innate immune system. *Mol. Psychiatry* **27**, 1647–1657 (2022).

25. Huan, T. *et al.* Genome-wide identification of DNA methylation QTLs in whole blood highlights pathways for cardiovascular disease. *Nat. Commun.* **10**, 4267 (2019).

26. Bonder, M. J. *et al.* Disease variants alter transcription factor levels and methylation of their binding sites. *Nat. Genet.* **49**, 131–138 (2017).

27. Lin, H. *et al.* Methylome-wide association study of atrial fibrillation in Framingham Heart Study. *Sci. Rep.* **7**, 40377 (2017).

28. Swedish Schizophrenia Consortium *et al.* High density methylation QTL analysis in human blood via next-generation sequencing of the methylated genomic DNA fraction. *Genome Biol.* **16**, 291 (2015).

29. Shi, J. *et al.* Characterizing the genetic basis of methylome diversity in histologically normal human lung tissue. *Nat. Commun.* **5**, 3365 (2014).

30. Dai, J. Y. *et al.* DNA methylation and cis-regulation of gene expression by prostate cancer risk SNPs. *PLoS Genet.* **16**, e1008667 (2020).

31. Schulz, H. *et al.* Genome-wide mapping of genetic determinants influencing DNA methylation and gene expression in human hippocampus. *Nat. Commun.* **8**, 1511 (2017).

32. Houseman, E. A. *et al.* DNA methylation arrays as surrogate measures of cell mixture distribution. *BMC Bioinformatics* **13**, 86 (2012).

33. Fortin, J.-P., Triche Jr, T. J. & Hansen, K. D. Preprocessing, normalization and integration of the Illumina HumanMethylationEPIC array with minfi. *Bioinformatics* **33**, 558–560 (2017).

34. Jakubzick, C. V., Randolph, G. J. & Henson, P. M. Monocyte differentiation and antigen-presenting functions. *Nat. Rev. Immunol.* **17**, 349–362 (2017).

35. Gordon, S. & Taylor, P. R. Monocyte and macrophage heterogeneity. *Nat. Rev. Immunol.* **5**, 953–964 (2005).

36. Tao, L. *et al.* Single-cell RNA sequencing reveals that an imbalance in monocyte subsets rather than changes in gene expression patterns is a feature of postmenopausal osteoporosis. *J. Bone Miner. Res.* zjae065 (2024).

37. Zhou, Y., Deng, H.-W. & Shen, H. Circulating monocytes: an appropriate model for bone-related study. *Osteoporos. Int.* **26**, 2561–2572 (2015).

38. Marventano, I. *et al.* A complex proinflammatory role for peripheral monocytes in Alzheimer's disease. *J. Alzheimers Dis.* **38**, 403–413 (2014).

39. Yan, P. *et al.* Peripheral monocyte–derived cells counter amyloid plaque pathogenesis in a mouse model of Alzheimer's disease. *J. Clin. Invest.* **132**, (2022).

40. Kawanaka, N. *et al.* CD14+,CD16+ blood monocytes and joint inflammation in rheumatoid arthritis. *Arthritis Rheum.* **46**, 2578–2586 (2002).

41. Pamukcu, B., Lip, G. Y. H., Devitt, A., Griffiths, H. & Shantsila, E. The role of monocytes in atherosclerotic coronary artery disease. *Ann. Med.* **42**, 394–403 (2010).

42. Chittezhath, M. *et al.* Molecular profiling reveals a tumor-promoting phenotype of monocytes and macrophages in human cancer progression. *Immunity* **41**, 815–829 (2014).

43. Fraser, H. B., Lam, L. L., Neumann, S. M. & Kobor, M. S. Population-specificity of human DNA methylation. *Genome Biol.* **13**, R8 (2012).

44. Kato, N. *et al.* Trans-ancestry genome-wide association study identifies 12 genetic loci influencing blood pressure and implicates a role for DNA methylation. *Nat. Genet.* **47**, 1282–1293 (2015).

45. Mozhui, K., Smith, A. K. & Tylavsky, F. A. Ancestry dependent DNA methylation and influence of maternal nutrition. *PloS One* **10**, e0118466 (2015).

46. Greenbaum, J. *et al.* A multiethnic whole genome sequencing study to identify novel loci for bone mineral density. *Hum. Mol. Genet.* **31**, 1067–1081 (2022).

47. Wu, L. *et al.* An integrative multi-omics analysis to identify candidate DNA methylation biomarkers related to prostate cancer risk. *Nat. Commun.* **11**, 3905 (2020).

48. Baselmans, B. M. *et al.* Multivariate genome-wide analyses of the well-being spectrum. *Nat. Genet.* **51**, 445–451 (2019).

49. Gaziano, J. M. *et al.* Million Veteran Program: A mega-biobank to study genetic influences on health and disease. *J. Clin. Epidemiol.* **70**, 214–223 (2016).

50. Li, H. & Durbin, R. Fast and accurate short read alignment with Burrows–Wheeler transform. *bioinformatics* **25**, 1754–1760 (2009).

51. DePristo, M. A. *et al.* A framework for variation discovery and genotyping using next-generation DNA sequencing data. *Nat. Genet.* **43**, 491–498 (2011).

52. McKenna, A. *et al.* The Genome Analysis Toolkit: a MapReduce framework for analyzing next-generation DNA sequencing data. *Genome Res.* **20**, 1297–1303 (2010).

53. Yin, Y. *et al.* Impact of cytosine methylation on DNA binding specificities of human transcription factors. *Science* **356**, eaaj2239 (2017).

54. Xi, Y. & Li, W. B. whole genome bisulfite sequence MAPping program. BMC 1027. *Bioinformatics* **10**, 1028 (2009).

55. Pain, O., Dudbridge, F. & Ronald, A. Are your covariates under control? How normalization can re-introduce covariate effects. *bioRxiv* 137232 (2017).

56. Dahl, A., Guillemot, V., Mefford, J., Aschard, H. & Zaitlen, N. Adjusting for principal components of molecular phenotypes induces replicating false positives. *Genetics* **211**, 1179–1189 (2019).

57. Shabalin, A. A. Matrix eQTL: ultra fast eQTL analysis via large matrix operations. *Bioinformatics* **28**, 1353–1358 (2012).

58. Wang, G., Sarkar, A., Carbonetto, P. & Stephens, M. A Simple New Approach to Variable Selection in Regression, with Application to Genetic Fine Mapping. *J. R. Stat. Soc. Ser. B Stat. Methodol.* **82**, 1273–1300 (2020).

59. Yang, J., Lee, S. H., Goddard, M. E. & Visscher, P. M. GCTA: a tool for genome-wide complex trait analysis. *Am. J. Hum. Genet.* **88**, 76–82 (2011).

60. Noguera-Castells, A., García-Prieto, C. A., Álvarez-Errico, D. & Esteller, M. Validation of the new EPIC DNA methylation microarray (900K EPIC v2) for high-throughput profiling of the human DNA methylome. *Epigenetics* **18**, 2185742 (2023).

61. Bibikova, M. *et al.* High density DNA methylation array with single CpG site resolution. *New Genomic Technol. Appl.* **98**, 288–295 (2011).

62. Wu, L. *et al.* A transcriptome-wide association study of 229,000 women identifies new candidate susceptibility genes for breast cancer. *Nat. Genet.* **50**, 968–978 (2018).

63. Aberg, K. A. *et al.* Convergence of evidence from a methylome-wide CpG-SNP association study and GWAS of major depressive disorder. *Transl. Psychiatry* **8**, 162 (2018).

64. Consortium, 1000 Genomes Project. *A Global Reference for Human Genetic Variation. Nature* vol. 526 68 (Nature Publishing Group, 2015).

65. Zou, H. & Hastie, T. Regularization and variable selection via the elastic net. *J. R. Stat. Soc. Ser. B Stat. Methodol.* **67**, 301–320 (2005).

66. Krstajic, D., Buturovic, L. J., Leahy, D. E. & Thomas, S. Cross-validation pitfalls when selecting and assessing regression and classification models. *J. Cheminformatics* **6**, 10 (2014).

67. Xue, H., Shen, X. & Pan, W. Causal Inference in Transcriptome-Wide Association Studies with Invalid Instruments and GWAS Summary Data. *J. Am. Stat. Assoc.* **118**, 1525–1537 (2023).

68. Wu, C., Zhang, Z., Yang, X. & Zhao, B. Large-scale imputation models for multi-ancestry proteome-wide association analysis. *bioRxiv* 2023–10 (2023).

69. Zhao, H. *et al.* Proteome-wide Mendelian randomization in global biobank meta-analysis reveals multi-ancestry drug targets for common diseases. *Cell Genomics* **2**, (2022).

70. Hemani, G., Tilling, K. & Davey Smith, G. Orienting the causal relationship between imprecisely measured traits using GWAS summary data. *PLoS Genet.* **13**, e1007081 (2017).

71. Lawlor, D. A., Harbord, R. M., Sterne, J. A. C., Timpson, N. & Davey Smith, G. Mendelian randomization: Using genes as instruments for making causal inferences in epidemiology. *Stat. Med.* **27**, 1133–1163 (2008).

72. Giambartolomei, C. *et al.* Bayesian test for colocalisation between pairs of genetic association studies using summary statistics. *PLoS Genet.* **10**, e1004383 (2014).

73. Bomba, L., Walter, K. & Soranzo, N. The impact of rare and low-frequency genetic variants in common disease. *Genome Biol.* **18**, 77 (2017).

74. Zhang, J. *et al.* Plasma proteome analyses in individuals of European and African ancestry identify cis-pQTLs and models for proteome-wide association studies. *Nat. Genet.* **54**, 593–602 (2022).

75. Cavalcante, R. G. & Sartor, M. A. Annotatr: genomic regions in context. *Bioinformatics* **33**, 2381–2383 (2017).

76. Shang, L. *et al.* meQTL mapping in the GENOA study reveals genetic determinants of DNA methylation in African Americans. *Nat. Commun.* **14**, 2711 (2023).

77. Min, J. L. *et al.* Genomic and phenotypic insights from an atlas of genetic effects on DNA methylation. *Nat. Genet.* **53**, 1311–1321 (2021).

78. Gamazon, E. R. *et al.* A gene-based association method for mapping traits using reference transcriptome data. *Nat. Genet.* **47**, 1091–1098 (2015).

79. Liu, Y. *et al.* ACAT: a fast and powerful p value combination method for rare-variant analysis in sequencing studies. *Am. J. Hum. Genet.* **104**, 410–421 (2019).

80. Wu, C., Bradley, J., Li, Y., Wu, L. & Deng, H.-W. A gene-level methylome-wide association analysis identifies novel Alzheimer's disease genes. *Bioinformatics* **37**, 1933–1940 (2021).

81. del Bosque-Plata, L., Martínez-Martínez, E., Espinoza-Camacho, M. Á. & Gragnoli, C. The role of TCF7L2 in type 2 diabetes. *Diabetes* **70**, 1220–1228 (2021).

82. Savic, D. *et al.* Alterations in TCF7L2 expression define its role as a key regulator of glucose metabolism. *Genome Res.* **21**, 1417–1425 (2011).

83. Yan, R. *et al.* A novel type 2 diabetes risk allele increases the promoter activity of the muscle-specific small ankyrin 1 gene. *Sci. Rep.* **6**, 25105 (2016).

84. Harder, M. N. *et al.* Type 2 diabetes risk alleles near BCAR1 and in ANK1 associate with decreased β-cell function whereas risk alleles near ANKRD55 and GRB14 associate with decreased insulin sensitivity in the Danish Inter99 cohort. *J. Clin. Endocrinol. Metab.* **98**, E801–E806 (2013).

85. Freathy, R. M. *et al.* Common variation in the FTO gene alters diabetes-related metabolic traits to the extent expected given its effect on BMI. *Diabetes* **57**, 1419–1426 (2008).

86. Guo, S. *et al.* Inactivation of specific β cell transcription factors in type 2 diabetes. *J. Clin. Invest.* **123**, 3305–3316 (2013).

87. Saxena, R. *et al.* Large-scale gene-centric meta-analysis across 39 studies identifies type 2 diabetes loci. *Am. J. Hum. Genet.* **90**, 410–425 (2012).

88. Hanson, R. L. *et al.* 1642-P: Colocalization Analyses of Genetic Associations of Type 2 Diabetes with DNA Methylation on Chromosome 11p in American Indians. *Diabetes* **69**, (2020).

89. Caballero, B., Finer, N. & Wurtman, R. J. Plasma amino acids and insulin levels in obesity: response to carbohydrate intake and tryptophan supplements. *Metabolism* **37**, 672–676 (1988).

90. Wijekoon, E. P., Skinner, C., Brosnan, M. E. & Brosnan, J. T. Amino acid metabolism in the Zucker diabetic fatty rat: effects of insulin resistance and of type 2 diabetes. *Can. J. Physiol. Pharmacol.* **82**, 506–514 (2004).

91. She, P. *et al.* Obesity-related elevations in plasma leucine are associated with alterations in enzymes involved in branched-chain amino acid metabolism. *Am. J. Physiol.-Endocrinol. Metab.* **293**, E1552–E1563 (2007).

92. Loyfer, N. *et al.* A DNA methylation atlas of normal human cell types. *Nature* **613**, 355–364 (2023).

93. Tsai, P.-C. & Bell, J. T. Power and sample size estimation for epigenome-wide association scans to detect differential DNA methylation. *Int. J. Epidemiol.* **44**, 1429–1441 (2015).

94. Singmann, P. *et al.* Characterization of whole-genome autosomal differences of DNA methylation between men and women. *Epigenetics Chromatin* **8**, 43 (2015).

95. Grant, O. A., Wang, Y., Kumari, M., Zabet, N. R. & Schalkwyk, L. Characterising sex differences of autosomal DNA methylation in whole blood using the Illumina EPIC array. *Clin. Epigenetics* **14**, 62 (2022).