How does policy modelling work in practice? A global analysis on the use of epidemiological modelling in health crises

Authors: Liza Hadley*[ab]; Caylyn Rich[c]; Alex Tasker[d]; Olivier Restif[a]; Sebastian Funk[b].

* Corresponding author (LH667@CAM.AC.UK)

[a] Disease Dynamics Unit, University of Cambridge, Cambridge, UK

[b] Centre for the Mathematical Modelling of Infectious Diseases, London School of Hygiene and Tropical Medicine, London, UK

[c] Independent researcher

[d] University of Bristol, Bristol, UK

Abstract:

This study examines the use and translation of epidemiological modelling by policy and decision makers in response to the COVID-19 outbreak. Prior to COVID-19, there was little readiness for global health systems, and many science-policy networks were assembled ad-hoc. Moreover, in the field of epidemiological modelling, one with significant sudden influence, there is still no international guidance or standard of practice on how modelled evidence should guide policy during major health crises. Here we use a multi-country case study on the use of epidemiological modelling in emergency COVID-19 response, to examine the effective integration of crisis science and policy in different countries. We investigated COVID-19 modelling-policy systems and practices in 13 countries, spanning all six UN geographic regions. Data collection took the form of expert interviews with a range of national policy/ decision makers, scientific advisors, and modellers. We examined the current use of epidemiological modelling, introduced a classification framework for outbreak modelling and policy on which best practice can be structured, and provided preliminary recommendations for future practice. Full analysis and interpretation of the breadth of interview responses is presented, providing evidence for the current and future use of modelling in

disease outbreaks. We found that interviewees in countries with a similar size and type of modelling

infrastructure, and similar level of government interaction with modelling reported similar experiences and

recommendations on using modelling in outbreak response. From this, we introduced a helpful grouping of

country experience upon which a tailored future best practice could be structured. We concluded the

article by outlining context-specific activities that modellers and policy actors could consider implementing

in their own countries. This article serves as a first evidence base for the current use of modelling in a

recent major health crisis and provides a robust framework for developing epidemiological modelling-to-

policy best practice.

44 **Funding:** Wellcome Trust.

45 **Key words:** modelling, policy, communication, collaboration, outbreak, decision-making.

1 Introduction

Epidemiological modelling was brought to the forefront of public health decision-making during the COVID-19 pandemic, in many countries for the first time. Epidemiological modelling is a branch of epidemiology that uses mathematics to predict the spread of disease through a population[1]. It is especially useful for estimating the impact of possible prevention and control measures. Many countries relied heavily on this type of scientific advice for resource allocation, planning, and forecasting during the pandemic, especially in the face of unprecedented public health interventions and incomplete knowledge[1–6]. It is highly likely that epidemiological modelling will continue to be needed in future disease control, due to the complex and uncertain nature of both pathogen spread and population behaviour[7–13]. The uptake of modelling, and more broadly science advice as a whole, in policy can be streamlined with improvements to the organisation, translation, and timeliness of scientific evidence. Previous works have identified many open questions[14,15], and it has become clear that best practices are needed for navigating the new prominence of modelling advice in public health decision making. Consequently, our study addresses the *organisation* and *translation* of epidemiological modelling in policy spheres. We used COVID-19 epidemiology as a case study for health crisis systems, and encourage broader public health researchers to build off this work for navigating best practice discussions with their own governments.

As countries have invariably faced differing challenges in COVID-19, a cross-country comparison on the use and organisation of epidemiological modelling in policy is vital. It is essential to document these practices such that we can learn from shared experiences. Of most interest are the perceptions of those who were *directly* involved in COVID-19 modelling response - national epidemiological modellers and the policy/decision makers (government employees) and scientific advisors that they interacted with. (See the principal eligibility criteria in Supplementary Material S1 for an explanation of these roles). Particular academic groups have reflected and reported on their own work (for example Howerton et al. in the US, Sherratt et al. in the UK[16,17]) but there is strong need for a breadth study drawing together global practices.

Dedicated research studies on the use and organisation of modelling in outbreak response have previously

been sparse. Emergent COVID-19 reflective studies include those run by the interagency COVID-19 Multi-

model Comparison Collaboration (CMCC), the KEMRI-Wellcome Trust, and Imperial College London among

others[3,18–21]. In 2020, the CMCC produced early guidance on how models can be used in COVID-19, and set

out three distinct classes of policy questions suitable for modelling (projecting epidemiological impact in

the absence of intervention; scenario analyses and impact of interventions; modelling related to costs)[19].

The group also discussed presentation of results and gave brief examples of the modelling-to-policy

infrastructure and collaborations in a few select countries. Further work is now needed to examine

modelling-to-policy infrastructures and communication pathways in detail, to provide an evidence base of

shared experiences on the integration of scientific advice in health crises.

Our own study builds on this research gap. We examined the use of epidemiological modelling in COVID-19

response in 13 different countries and jurisdictions. Our study aims to:

(a) Compare national practices on the use and organisation of epidemiological modelling in COVID-

19 decision-making (3.1-3.5);

(b) Identify contextual drivers linking the shared experiences of interviewees and develop a

classification framework to structure future guidance (3.2);

(c) Propose preliminary recommendations on the use of epidemiological modelling in health policy,

contextual to the different country settings identified in (b) (3.6).

An adjacent paper reports findings on the topic of communication and visualisation[22].

2 Methods

This Methods section follows the consolidated criteria for reporting qualitative studies (COREQ)[23]. Details

on study design (2.1), country and participant selection (2.2), interview format and location (2.3), data

collection and analysis (2.4), ethics approval (2.5), and inclusivity (2.6) are provided in the following

subsections. Additional information about the study is included in the Supplementary Material and in

Hadley et al., 2025 (thesis forthcoming)[24].

2.1 Study design

This is a qualitative descriptive study. Our study aimed to compare national practices on the use of

epidemiological modelling in COVID-19 decision-making, identify key contexts linking the shared

experiences of interviewees to develop a classification framework, and propose preliminary

recommendations for future practice.

Data collection took the form of in-depth semi-structured interviews with national COVID-19 policy and

decision-makers, science advisors, and epidemiological modellers in 13 different countries and

jurisdictions, spanning all six UN geographic regions. Data analysis used thematic analysis and ideal-type

analysis. Note that our paper is descriptive, aiming to describe the integration of epidemiological modelling

in current health policy systems and advance a typology framework and recommendations for future use.

It is not an explanatory study providing for example reasoning for Covid-19 outcomes in different

countries.

2.2 Country and participant selection

This study defines epidemiological modelling activity in a broad sense, to include statistical modelling and

analytics too. In total, 12 countries and 1 jurisdiction (Hong Kong, China) took part in the study. For brevity,

we use the term 'country' to mean 'country or jurisdiction' throughout this article, with the knowledge that

Hong Kong, China is a jurisdiction or 'special administrative region' and not itself an independent country.

Country identification was performed using a targeted selection process; the lead researchers first carried

out an informal scoping review of the use of epidemiological modelling in national Covid-19 responses to

assemble a longlist of candidate countries for further exploration. Countries with a population of less than

5 million were excluded. Fifteen target countries were chosen to support a target sample size of 20-30

interviewees following current practices in similar qualitative interview research studies[25]. Data saturation

was not a primary goal as this study was intended to be exploratory. Candidate countries were identified

using intersectional selection criteria: covering each UN geographic region, low-and-middle-income

countries and high-income countries, low- and high- COVID-19 prevalence, and diversity of prior

experience of modelling for national decision-making, modelling structures or consortiums (including those

with no formal capacity), pre-COVID-19 pandemic preparedness ratings, and population size. Researchers

then used purposeful sampling to contact key modelling-to-policy actors in each country by email for

interview[26]. If no relevant participants could be sought, a second country was chosen until the study met

the desired study size of 10-15 countries with 2+ participants per country, including at least one country

from each UN geographic region. In total, 27 participants from 13 countries took part. 31 participants

replied to the study recruitment invite but three were unavailable at the time of interview and one could

not be taken forward as a second interviewee from the same country could not be found. Recruitment ran

from 21st April 2023 to 10th May 2024.

Participant names are kept anonymous but participants reported the following as their countries (or

jurisdictions) of practice: Australia, Canada, Colombia, France, Japan, Hong Kong Special Administrative

Region of the People's Republic of China (Hong Kong), Kenya, New Zealand, Peru, Republic of Korea (South

Korea), South Africa, Uganda, and the United Kingdom. At least two respondents were recruited from each

of the above countries. Table M1 describes the countries in our study and Table M2 describes the make-up

of participants. Of the 27 interviewees, 12 self-identified as scientific advisors or policy/ decision makers (3,

9 respectively), and 15 as modellers. Where possible, efforts were made to make sure both modellers and

science advisors or policy/ decision makers were interviewed in each country. By country, the split of

interviewees achieved was fairly even: the majority of countries in our study (8/13) had both a modeller

and a science advisor or policy/ decision maker as interviewees; but it is noted that in three countries, only

modellers were interviewed and in the remaining two countries, only scientific advisors or policy/ decision

makers were interviewed. Interviewees self-identifying as scientific advisors and policy/ decision makers

included national Government Chief Scientific Advisors and heads of the relevant national Health

department or agency, as well as intermediaries or convenors assigned specifically to coordinate or relay

modelling activities.

Country of practice	Country details			
	UN geographic region[a]	Income status[b]	GHS pandemic preparedness rating (2019)[c]	Population size (millions)[d]
Australia	Oceania	HIC	4th	25.8
Canada	Northern America	HIC	5th	38.0
Colombia	LAC	UMIC	65th	51.2
France	Europe	HIC	11th	64.5
Hong Kong, China	Asia	HIC	51st*	7.5
Japan	Asia	HIC	21st	124.9
Kenya	Africa	LMIC	55th	52.5
New Zealand	Oceania	HIC	35th	5.1
Peru	LAC	UMIC	49th	33.5
South Korea	Asia	HIC	9th	51.8
South Africa	Africa	UMIC	34th	59.1
Uganda	Africa	LIC	63rd	45.1
United Kingdom	Europe	HIC	2nd	67.2

Table M1: The 13 countries and jurisdictions involved in our study and relevant demographic statistics. Each interviewee reported their country(ies) of practice as one of the above. LAC = Latin America and the Caribbean; LIC = Low-income country; LMIC = Lower-middle income country; UMIC = Upper-middle income country; HIC = High-income country. Country data is taken from the following sources: (a), (b) & (d) UN geographic region, country income, and population (as of 1st January 2021) is as described in the United Nations World Population Prospects 2022[27,28]; (c) Rankings taken from the 2019 Global Health Security Index, indicating expected overall preparedness for an infectious disease outbreak[29]. Note that Hong Kong's ranking of 51st represents the overall ranking for China - as noted in the main text, Hong Kong is a Special Administrative Region of China.

In Table M1, we see that the 13 countries in our study had a diversity of income status (low, lower-middle, upper-middle, and high), pre-pandemic preparedness ratings (ranging from 2nd to 65th), and population size (ranging from 5.1 million to 124.9 million). However it is noted that additional data on the peculiarities of each country such as public-private partnerships and multisectoral vs multidisciplinary activity was not

191 readily available. Readers interested in public-private partnerships are directed to related study Dabak et

192 al. (in preparation)[30].

Affiliation	Role	No. of interviewees
Academic	Modeller	15
	Independent scientist and advisor	3
Government	Policymaker	5
	Senior government (For example head of a Ministry of Health or Public Health Agency, senior Chief Scientific Advisors, senior members of Cabinet, and related roles)	4

194 *Table M2: Make-up of study participants.*

2.3 Interview format and location

Interviews were in-depth and semi-structured, lasting approximately 1 hour each. Author LH (female PhD

student with training and prior experience in qualitative interviewing) carried out all interviews. The format

was structured discussion on four topics in turn, facilitated by the interviewer. Additionally, the first

interviewee in each country was invited to co-create a 'sketch diagram' with the interviewer mapping out

the key actors in modelling and policy in their country from their perspective. This rough diagram served as

a reference point for discussion, and the main pathways were verbally verified through discussion with

subsequent interviewees. The remainder of the interview followed the standard semi-structured format.

The interview schedule was heavily informed by previous work on 'Challenges for Future Pandemics' in

2020/2021[15]. Hadley et al., 2021 introduce a succinct framework that defines the areas for progress on the

interface of outbreak modelling and policy (reproduced here as Figure M1). This idea-generating piece

directly informed our study's core topics and specific interview questions. Hadley et al., 2021 identified

seven broad challenges of pandemic modelling for policy; we narrowed this down to focus on four related

interview topics, namely 'Structures and pathways to policy' (relating to challenges 1 & 3 in Figure M1),

'Collaboration and knowledge transfer' (2&4), 'Communication and visualisation' (5), and 'Evaluation and

reflection' discussing recommendations for future practice (7). Our study did not discuss technical model

challenges, as this has been frequently researched and reported on in the wider literature (as one example,

see Sherratt et al. discussing European nations[31]). The full interview schedule is included in the

Supplementary Material and questions were piloted on three colleagues before study commencement.

Figure M1: Infographic depicting seven broad challenges of pandemic modelling for policy, around the central equation of

epidemic spread. Reproduced from Hadley et al., 2021[15].

Four interviews were conducted in-person in private workplace meeting rooms; the remaining 23 were

conducted online via video-conferencing software. Majority of interviews were conducted one-to-one in

English, with one using an independent translator to aid communication.

2.4 Data collection and analysis

All interviews were audio recorded, professionally transcribed, qualitatively coded, and then reviewed for

accuracy by the study researchers. Analysis of data followed a two-pronged approach: using thematic

analysis for descriptive findings and ideal-type analysis for a more in-depth understanding of the data,

enabling contextual recommendations to be made on the future use of modelling[32,33]. Thematic analysis is

the main method for identifying, analysing, and reporting patterns within qualitative interview data, and is

best used when the research aim is to explore people's beliefs, knowledge, or experiences from a set of

qualitative data such as interview transcripts[32,34]; hence it was deemed an appropriate method of choice

for our study. Our thematic analysis followed a deductive approach, utilising topics posited in earlier study

Hadley et al., 2021 (described in Section 2.3 above). Two of the researchers (LH, CR) led the initial coding

with manual line-by-line review. Manual coding was chosen over the use of software to enable a deeper

understanding of the material. The first six transcripts were double-coded and then compared to reach a

consensus on the coding framework. Remaining transcripts were then coded and independently reviewed

to enable triangulation and continued code consistency. The researchers also produced analytic memos for

each interview. Initial codes were then merged to account for duplication, from which meaningful basic

themes and organising themes were derived, reviewed, and refined[35]. The final step of thematic analysis

was interpretation and reporting of the major findings.

Ideal-type analysis was carried out by researchers LH and AT, following a modified protocol adapted from

Stapley et al.[33]. Ideal-type analysis is a method for systematically developing a typology (a useful

categorisation) of the subject of interest[33,36,37]. This analytical approach was selected to organise interview

responses and to understand aspects of an interviewee's experiences that could contribute to shared

beliefs and shape recommendations for future practices in outbreak modelling and policy. Guided by the

question "What are the key contexts to consider when comparing shared experiences and

recommendations on the integration of outbreak modelling in policy?", we sought to understand

contextual drivers that would enable the development of tailored guidance for best practice to move

beyond the current approach of blanket recommendations which are rarely contextualised. While

typologies are commonly employed in qualitative data analysis, there is no agreed best practice for

constructing them[38]. Given the subject and design of our study we chose to follow the method of Stapley

et al.[24] to identify key characteristics influencing the subject of interest (beliefs and recommendations on

future practices), collecting examples of optimal cases, and defining contexts within which shared

experiences were observed (forming a description of each 'ideal type'). Ideal types were tested, refined,

and checked for credibility, before using a structured analytical approach to organise into a classification

framework highlighting important similarities and differences between groups. The results of the

classification framework are presented in section 3.2 (Figure R2) and are used to inform the preliminary

contextual recommendations given in Table R3. Further detail on the derivation of this is provided in the

Results and in Supplementary Material S5.

Once data analysis was complete, a draft version of this article was also circulated to all interviewees to

enable verification and opportunity for revisions of the key findings.

Results sections 3.1 and 3.2 present key data from each country and introduce a classification framework

for important contexts to consider in outbreak modelling and policy. Section 3.3 examines any

collaboration of modelling with other scientific disciplines, section 3.4 explores translators and knowledge

brokers, and section 3.5 discusses mistakes. Finally, section 3.6 draws on the complete study analysis,

supplementary results, and pre-existing literature to introduce guidelines for future practice on using

epidemiological modelling in health crises. Results on communication and visual preferences are presented

in our adjacent study[22].

2.5 Ethics statement

The University of Cambridge Psychology Research Ethics Committee (application number PRE.2023.034)

has reviewed this research. The majority of study participants did not know the interviewer prior to study

commencement but all participants were informed of the purpose of the study, reasons for interest, and

proposed outputs at the recruitment stage. All participants provided informed written consent.

2.6 Inclusivity in global research

Additional information regarding the ethical, cultural, and scientific considerations specific to inclusivity in

global research is included in the Supporting Information (Inclusivity in Global Research Checklist).

3 Results

Factors driving shared experiences

Thematic and ideal-type analysis identified two driving parameters linking the shared experiences of

interviewees: size/type of modelling infrastructure, and level of government interaction with modelling.

For example, clusters of interviewees with similar reported experiences, beliefs, and recommendations on

the integration of outbreak modelling in policy had a similar type and size of modelling infrastructure and

similar level of government interaction with modelling. This was evidenced by interviewees' reported

strengths and challenges of modelling-policy systems which commonly framed as relating to either a large

multi-team modelling capacity, limited (single-team) modelling capacity, or disjointed modelling capacity;

and contextual to modellers being either embedded within government or independent. These structural

factors were a strong driver for similarities between groups of interviewees than for example views on

communication and visualisation where recommendations were more universal across interviewees. These

structural factors were also a stronger driver than countries with the same income status, and countries in

the same region. Findings on type of modelling infrastructure and level of government interaction are

presented in section 3.1, and analysis of contextual drivers linking shared experiences in outbreak

modelling and policy is presented in section 3.2.

3.1 Modelling infrastructure and pathways to policy

We observed four distinct types of national **modelling infrastructure**, presented in Figure R1. These are:

one modelling team; multiple small teams functioning as one; a consortium consisting of multiple teams

and multiple models led by a modelling committee; and finally, teams working in isolation, feeding

independently into government. Types 2 and 3 were separated to distinguish a single-model infrastructure

that would likely have no capacity to work on the same questions with a multi-model infrastructure that

does have capacity for self-verification, reproduction, or consensus (in any form). Type 4 considers settings

where modelling activity is disjoint; modellers likely do not confer with each other, and results are instead

'combined' in-house by a government non-modelling committee. Examples are given in Figure R1. Our

classification aims to capture the size of each country's primary modelling capacity, ability or inability to

reproduce model findings, and whether modelling outputs are *synthesised and interpreted by modellers* (1-

3) or *non-modellers* (4) such as other infectious diseases or policy experts.

Figure R1: The four types of national modelling infrastructure observed in our study. One modelling team (1); multiple small

teams functioning as one (no capacity to reproduce) (2); multiple teams and multiple models (capacity to reproduce) led by a

committee of modellers (3); modellers and/or modelling teams do not collaborate, work in isolation, and feed results directly into

government to be combined by a committee of non-modellers (4). Pictured: Individuals with hats represent modelling principal

investigators/ senior modellers. Computers represent models. Boxes represent a modelling committee. For example, in South

Africa (type 2), there was insufficient capacity to examine the same modelling questions with multiple models so teams divided

the research questions among themselves and functioned as one team. In South Korea (type 4), modelling effort was disjointed -

modelling teams did not collaborate or discuss findings and instead sent reports directly to the relevant policy actor.

Additionally, one can classify each country by the **level of government interaction in modelling**. This is

shown in Table R1 (rightmost column), along with summary findings on the structures and pathways to

policy in each country. The leftmost columns concern *modelling* infrastructure while the rightmost columns

consider the *policy infrastructure for modelling* and provide a relative ranking for the level of government

interaction and independence in each setting, based on in-depth discussions with interviewees. One sees

for example that the level of interaction in modelling from government bodies varied greatly (Medium: 3

countries; High: 5; Very high: 2; Embedded: 2). Illustrative definitions of these rankings are provided in

Supplementary Material S3. Note that countries where modelling had little to no use by national

government in COVID-19 were excluded from our study.

Country	Type of modelling infrastructure (type #)	Name of MC or team if given	Main affiliation of modellers	Main government linkage point(s)	Level of (structural) government interaction
Africa					
Kenya	**Multiple teams, multiple models (3)**	Led by Centre for Epidemiological Modelling and Analysis (CEMA), University of Nairobi	Universities or research institutes	Health, also Education, Faith *Multiple linkages reported: MoH mainly, also Ministry of Education, National Council of Churches and religious groups, National Taskforce, and Cabinet at times*	**High.** Kenyan COVID-19 modelling review team sits within the National COVID-19 Task Force.
South Africa	**Small teams working as one (2)**	South African COVID-19 Modelling Consortium (SACMC)	Universities or research institutes	Health, Economics *Ministry of Health advisory groups, MoH direct at times, Treasury*	**High.** SACMC convened by the National Institute of Communicable Diseases at the request of MoH.
Uganda	**Teams working in isolation (4)**	NA	Universities or research institutes	Health *Epidemiological Advisory Group, PRESIDE and National Planning Authority*	**Medium.** Two modellers on relevant SACs.
Asia					
Hong Kong, China	**One team (1)**	Led by LKS Faculty of Medicine, University of Hong Kong researchers	Universities or research institutes	Health *COVID-19 Expert Advisory Panel, and Cabinet at times*	**High.** Highly-integrated key individuals as brokers.
Japan	**One team (1)**	Led by Kyoto University researchers	Universities or research institutes	Health *Ministry of Health*	**Very high.** Modellers in-housed to MoH during first wave and modeller on relevant SAC.
South Korea	**Teams working in isolation (4)**	NA	Universities or research institutes	Health *Korean Disease Control and Prevention Agency (CDC)*	**Medium.** Brokerage carried out by influential ID specialist due to limited modelling expertise.
Europe					
France	**One team** initially (1); then **multiple teams, multiple models (3)**	Led by Institut Pasteur researchers; later the Coordinated Action of Modelling for Disease (ACM)	Universities or research institutes	Health *Conseil Scientifique*	**Medium.** One modeller on national SAC throughout.

United Kingdom	Multiple teams, multiple models (3)	Scientific Pandemic Influenza Group on Modelling, Operational (SPI-M-O)	Universities or research institutes primarily, with government contributors	Health primarily *SAGE (Government Office for Science); Cabinet Office at times*	**High.** Several modellers on national SAC throughout, secondments to Cabinet Office, and regular attendees from Cabinet Office to MC meetings.
Latin America and the Caribbean					
Colombia	Multiple teams, multiple models (3)	Led by National Institute of Health (INS)	Government, with university/ research institute teams contributing on occasion	Health, Education *Ministry of Health, Ministry of Education*	**Embedded** (primary modelling team).
Peru	One team (1)	CDC-Peru modelling unit	Government	Health *Ministry of Health*	**Embedded.**
Northern America					
Canada	Multiple teams, multiple models (3)	Led by Public Health Agency of Canada (PHAC)	Government, with university/ research institute teams contributing regularly	Health *Minister of Health, Cabinet*	**Embedded** (primary modelling team).
Oceania					
Australia	One team (scenarios) (1); multiple teams, multiple models (forecasting) (3)	Doherty Modelling consortium and National COVID-19 Situational Assessment Consortium	Universities or research institutes primarily	Health, Economics *Multiple linkages including Australian Health Protection Principal Committee (AHPPC), Treasury*	**High.** Long-term pre-existing relationship and modellers on multiple SACs.
New Zealand	Multiple teams, multiple models (3)	COVID-19 Modelling Aotearoa (CMA)	Universities or research institutes	Health *Ministry of Health, Department of Prime Minister and Cabinet*	**Very high.** Extremely close science-policy system but still with modellers' independence.

Table R1: Modelling infrastructure and details on the structures and pathways to policy in each country in COVID-19, as described by interviewees. MC = Modelling Committee. SAC = A multidisciplinary Scientific Advisory Committee or expert panel. MoH = Ministry of Health / Department of Health. ID = Infectious Diseases. Illustrative definitions used to identify level of government interaction with modelling are shown in Supplementary Material S3.

3.2 Classification framework for outbreak modelling and policy

Identifying countries with similar experience on the use of modelling will enable the development of more

effective recommendations. As noted in the 'Factors driving shared experiences' subsection, ideal-type

analysis (Supplementary Material S5), suggests that the size/ type of modelling capacity and level of

government interaction with modelling were contextual drivers linking the shared experiences, beliefs, and

recommendations of subsets of interviewees. These are important contexts to consider when grouping

country settings by their current use of outbreak modelling in policy, to produce meaningful further

guidance. Section 3.1 identifies four types of national modelling infrastructure (Figure R1) and four levels

of government interaction with modelling (Table R1), creating sixteen possible categories of shared

experience. Upon systematically comparing and contrasting interview transcripts and summaries, this was

reduced to five key categories of outbreak modelling and policy (Figure R2). See Methods section 2.4 and

Supplementary Material S5 for a description of this analysis.

The following five categories of modelling-policy systems were identified as having similar experiences and

recommendations for future practice: (A) Countries with small modelling capacity and likely high

government linkage; (B) Countries with large modelling capacity and high or very-high government linkage;

(C) Countries where modelling teams worked independently, combined by non-modellers in government;

(D) Countries with a large modelling capacity with a primary government modelling team; (E) Countries

with one government modelling team (Figure R2). This is called a typology in qualitative research[33].

Figure R2: Classification framework for outbreak modelling and policy. Five categories of modelling-policy systems were

identified and organised by similar experiences and recommendations on the use of outbreak modelling in policy. Type of

modelling infrastructure and level of government interaction with modelling were identified as important contextual drivers

when grouping country settings. The five categories of modelling-policy systems identified are: (A) Countries with small modelling

capacity and likely high government linkage, such as Hong Kong; (B) Countries with large modelling capacity and high or very

high government linkage, such as Kenya; (C) Countries where modelling teams worked in isolation, combined by non-modellers in

government, such as South Korea; (D) Countries with a large modelling capacity with a primary government modelling team,

such as Canada; (E) Countries with one government modelling team, such as Peru. Countries were placed in the same category if

interviewees demonstrated similar beliefs and experiences on the use of outbreak modelling in policy. Note: For other pathogens

and other health crises, each category may contain different countries.

Producing a robust categorisation of shared experiences, beliefs, and recommendations in outbreak

modelling and policy as shown in Figure R2 facilitates context-specific recommendations for future

practices. To this end, subsequent results are presented by these categories where it is meaningful to do

so.

3.3 Collaboration with other disciplines

In the majority of cases, modelling activity was focussed on primary health outcomes (i.e. COVID-19

transmission and disease) but there were a few interdisciplinary success stories where modellers

collaborated with other disciplines. Costing, or the idea of evaluating trade-offs (either health or economic

based) was reported by modellers in New Zealand, Australia, Canada, and Peru, but was explicitly indicated

as not existent in the UK, Colombia, and France. In New Zealand, while economic modelling remained the

responsibility of the Treasury, the country's modelling consortium (COVID-19 Modelling Aotearoa, CMA)

collaborated to provide analyses of isolation policy trade-offs and linked informal health modelling within

the CMA to economic modelling within the Treasury.

On the behavioural sciences front, in Australia modellers previously collaborated with social scientists on

the use of citizen juries to inform ethical vaccine allocation, but during the COVID-19 response, such

networks suffered a lack of behavioural social sciences inclusion. A respondent from South Korea also

emphasised the importance of an interdisciplinary approach to modelling: "This is a policy where extensive

educational, social, and economic losses are expected, so it is necessary to listen to the opinions of experts

in each field." Different forms of science can of course be combined 'higher up' in senior government, but

there was a shared sentiment of desire for more cross-pollination in countries that lacked it. For example,

post pandemic, modellers from Kenya reported that they are now actively working with behavioural

scientists in a local School of Anthropology on the current Rift Valley Fever outbreak.

Note: Limitations to the scope of this analysis resulted in a lack of evaluation for the *quality* of

collaboration across disciplines, but rather sought to determine whether collaboration existed.

3.4 Translators and knowledge brokers

We define 'translators' as those individuals who took on the role of packaging information, converting

complex epidemiological modelling findings into meaningful language for policy. In countries where

modellers collaborated and had a level of independence from government (categories A and B in Figure

R2), this 'translator' role was often taken on by the most senior modeller(s) (Australia, France, Kenya,

South Africa) or epidemiologist (Hong Kong). Exceptions include New Zealand's small close-knit science-

policy ecosystem where modellers interacted directly with senior government and first translation was

performed largely by the relevant Chief Scientific Advisor, and the UK whose modelling committee (SPI-M-

O) was jointly convened by a policy co-chair and academic co-chair. In the latter, the policy co-chair and

SPI-M-O secretariat were all civil servants. Uniquely these UK individuals had significant past modelling

expertise and could effectively synthesise key modelling findings that the policy co-chair, crucially *inside*

government, could take into the parts of government that the academics reportedly had limited access to.

Secretariat members ensured that an accurate interpretation of the modelling was relayed to ministers by

meticulously agreeing appropriate slide headings with Cabinet Office officials to ensure the knowledge was

conveyed appropriately; the benchmark became whether policy makers would be able to understand the

implications of the content if they only read the headings.

Countries with more government embeddedness (categories D and E in Figure R2) saw a mix of translators:

the Chief Public Health Officer in Canada, the lead academic modeller in Japan, and the relevant Peru CDC

department director in Peru. Finally, in South Korea (category C - disjointed modelling) where there were

reported to be no dedicated infectious disease modelling teams at the beginning of the COVID-19

pandemic (primarily mathematics or public health research groups with only some modelling expertise),

translation was taken on by an experienced and informed non-modelling infectious diseases principal

investigator. The origin of each of the above interactions varied; as one interviewee commented "it is hard

to create strict rules around who should communicate: it ends up being the persons involved where their

strengths lie and what structures they have around them".

3.5 Mistakes

Many respondents were asked what they perceived to be the 'biggest mistake' when using modelling in

the response. Responses are shown in Table R2 for interest. These comments were shared with the

interviewer in the spirit of identifying shared challenges and should be interpreted not as shortcomings of

a particular individual but as areas where we could collectively improve.

What was your biggest mistake (in relation to the use of modelling)?
Not being transparent enough with lack of knowledge; allowing policy makers to espouse more certainty in the models than existed
Inability to address vaccine hesitancy - lacked incorporating behavioural or economic scientists into modelling which contributed to elevated vaccine hesitancy
Underestimating the number of deaths
Relying on a single model/ not managing to develop a multiple model consortium
Lack of understanding of types of modelling and modelling's limitations (not investing in the pedagogy of modelling)
Not taking an interdisciplinary approach, only collaborating with the Ministry of Health rather than other ministries
Utilising worst-case scenarios to justify actions, especially when presented in a public forum
Erring too far on the pessimistic side of modelling/ the negative outcome or worst-case scenario
Accidentally getting too involved with politically motivated media figures
Lack of resources (financial) to pool expertise
Risk in using models as the primary reason for introducing a lockdown
Not believing vaccines would be available so quickly
Not grasping the impact of interventions
Lack of resourcing creating biases in datasets - marginalised communities couldn't enjoy same level of for example surveillance consequently forcing 'national' rather than 'local' models onto indigenous communities
Some modellers/ modelling groups struggled to recognise the time and place appropriate for complex vs simple models
Narrow scope of metrics, no modelling of socioeconomic impacts
Downplaying models' predictions
"I don't think we made a mistake but with hindsight we've always thought, thank God, we kind of did it right and we didn't get it wrong"
Not being prepared capacity-wise
Poorly communicating the risks

Table R2: Responses to the question "What was your biggest mistake (in relation to the use of modelling)?". All responses have been paraphrased to maintain anonymity.

3.6 Recommendations for future practice

Towards the end of each interview, interviewees discussed their key takeaways and recommendations for

the future of outbreak modelling and policy (presented in full in Supplementary Material S4). This dataset

highlighted four essentials for successful outbreak modelling and policy: data, systems, communication,

and relationships (Figure R3).

Figure R3: Essentials for successful outbreak modelling and policy - data, systems, communication, and relationships. This figure
follows the style of earlier work Hadley et al, 2021.

Expanding on recommendations from interviewees, the results of our study analysis, and additional

literature in this area, we, the authors, introduce the following guidelines for future practice on using

epidemiological modelling in disease outbreaks:

	Guidelines for future practice on using epidemiological modelling in disease outbreaks	
1	**Identify your system**	There is no optimal modelling-policy system. Moving between categories of modelling-policy systems may require significant political and cultural change. Modellers should instead identify the system in which they operate (Figure R2) and utilise the context-specific case studies and guidance presented in this article.
2	**Relationships**	Facilitate long-standing relationships. Consider developing relationships with government groups rather than individuals, to preserve future linkages under government turnover. For systems where modelling sits primarily 'outside' government, it is critical to have an interpreter/translator or advocate (who may or not be a modeller) within policymaking spheres to provide a space for modelling.
3	**Standing capacity for modelling**	More standing capacity for modelling is needed in some countries. For systems with limited modelling capacity, provide secure jobs for modellers. Disjointed modelling effort is encouraged to unify informally. Where there is a lack of capacity to train people in-country, an international modelling network or society could aid development. Large modelling capacity and multi-model consortiums will likely need to be maintained with dedicated funding; where that exists we encourage active sharing of ideas to countries with fewer experts.
4	**Data availability**	Strong data access and availability is essential for modelling. This is a universal recommendation for all settings.
5	**Legitimacy**	For settings where modelling sits 'outside' of government, establish relationships and agreements/ memorandums of understanding ahead of time. Unifying modelling effort through a modelling committee can aid identifiability in policy spheres. Retrospective verification of modelling results is also important for developing trust and legitimacy.
6	**Communicating modelling**	Caution against overbelieved or misinterpreted modelling. For detailed guidance on communicating and interpreting modelling in policymaking, see adjacent study[22].
7	**Interdisciplinarity**	Science advice in health crises should be holistic, looking beyond direct health outcomes. However, research on how modelling can best interact with economics and behavioural sciences is limited. We offer examples (Section 3.3) but further research is needed before a formal guideline can be made here.

Table R3: Preliminary guidelines for future practice on using epidemiological modelling in disease outbreaks. From the results of

our study analysis, recommendations from interviewees, and additional literature in this area, we suggest the following

guidelines for future practice on using epidemiological modelling in disease outbreaks.

<u>4 Discussion</u>

At the time of shock, when the pandemic first hit, nations rapidly organised scientific and modelling

capacity, and crucially this self-organisation happened in *different* ways (Table R1), resulting in *different*

successes and challenges as outlined in detail above. Countries with a similar size and type of modelling

infrastructure, and similar level of government interaction with modelling were found to have reported

similar successes and challenges on using modelling in their COVID-19 response (section 3 'Factors driving

shared experiences'). More universally, there was a desire for epidemiological modelling to have an eye to

wider societal impacts, and in many settings, work formally with other experts. We highlight two success

stories of epidemiological modelling successfully working with economics and social science either in the

COVID-19 pandemic (New Zealand) or as an active post-COVID-19 advancement (Kenya) (section 3.3).

On the communication front, we introduced the idea of *re-translation*, acknowledging the need to ensure

that not only is modelling translated and packaged effectively to policy/ decision makers, but that in

subsequent iterations 'up the chain' the message is maintained. Further results on communication and

visualisation from this study are presented in the adjacent paper Hadley et al.[22]. There is a need now for

further emphasis in Science Communication work on enabling *subsequent* translation of modelling, at least

in policy settings.

The classification framework for outbreak modelling and policy introduced in Figure R2 highlights key

contextual considerations (specifically the size and type of modelling capacity, and independence from

government or embeddedness) across which groups of interviewees reported similar experiences, beliefs,

and recommendations. It is important to note here that interviewees in the same region or in countries

with the same income status did not necessarily report similar experiences in COVID-19: For example,

within the region of Africa, Kenya benefitted from a large multi-team modelling consortium while South

Africa and Uganda had more limited modelling capacity (Table R1). In the region of Asia, modellers in Japan

reportedly had a very high level of integration with the government through the Ministry of Health while in

South Korea, efforts were disjointed and due to insufficient principal investigators in epidemiological

modelling, communication was taken on by an infectious diseases specialist (section 3.4). One can see that

simply grouping countries by income status or region would not accurately capture the nuance of reported

experience in COVID-19 epidemiological modelling and policy. Instead, having a single-team, multi-team, or

disjointed modelling capacity, and being independent from or embedded within government processes did

well-represent the different experiences and recommendations of the study group. The classification

framework introduced in Figure R2 hence provides a useful basis for wider examination of the use and

translation of epidemiological modelling in outbreaks. There is an opportunity for future work here to test

our classification framework with countries beyond the 13 considered, with a larger sample size, and in

other (for example routine public health) disease settings.

Finally, we observed overarching themes of data, communication, systems, and relationships as primary

enablers to the effective use of epidemiological modelling in disease outbreaks. We synthesised specific

and contextual recommendations for future practice on the use of modelling in health crises in the 13

countries examined. Table R3 outlines guidance for the future use of epidemiological modelling in disease

outbreaks. These stem from crucial suggestions from key global actors, many with extensive experience in

multiple health events and from different perspectives.

Looking at related literature in this area, Owek et al. identified pre-existing relationships between

researchers and policy/ decision-makers as an enabling factor for successful knowledge translation, in

addition to co-creation of advice and embeddedness[3]. Our findings echo this, with 'Systems' and

'Relationships' both being pivotal for future outbreak modelling and response. Our findings on systems

also reflect that of formal inquiries taken on by several countries - for example in the UK's COVID-19

Inquiry, modellers (as well as all other scientists, policy/ decision makers, and associated actors) provided

witness statements detailing their involvement in the country's pandemic response from which it is clear

that the systems and relationships built prior to and during COVID-19 were fundamental to modelling's

utility in COVID-19[39].

Additionally there was acknowledgement of the role of translators by the CMCC report discussed earlier in

this article, noting that where needed "decision makers may be supported by advisors or knowledge

brokers with a high literacy to interpret models"[19]. In contrast, Owek et al. found that face-to-face

debriefings between scientists and policy/ decision makers were preferred in Low and Middle Income

Countries over the use of a knowledge broker or intermediary[3]. We observed more of a mix of translators

in our study - of the five countries classified as LMICs in Table M1 (Colombia, Kenya, Peru, Uganda, and

South Africa), translation was taken on by the most senior modeller in three settings but by an

intermediary in the other two. It is worth acknowledging whether academics should be the ones

responsible for translating results into policy language or if this activity could be more effectively in-

housed. There is merit for the latter in the countries we have investigated.

In terms of limitations, we have aimed to capture a range of experiences in our study but there was a

limited number of participants (27) and in most cases, only two participants per country. Consequently, we

did not attribute recommendations to specific countries (with the exception of giving examples) and

instead presented the major analyses by typology (e.g., suggestions for countries with small modelling

capacity, suggestions for countries where modelling sits 'outside' of government, etc.). Note also that

interview data represents the *perceptions* of interviewees, hence this study only evaluates the perceived

use of modelling, albeit from highly-informed individuals. Nonetheless, this may not reflect official policy or

how individuals would behave in practice. We also use the terms 'government' and 'policy/ decision

makers' loosely throughout, and acknowledge that this relates to different individuals and committees in

each country, due to the multifaceted political structures. A closer examination of these roles is a point for

future work[40]. The wider political, economic, and cultural factors experienced by different government

actors may also influence the uptake of scientific evidence, but this was outside the scope of our article.

We acknowledge that state-level and regional (multi-country) modelling were outside the scope of this

study, as were case study countries where modelling *did not* contribute to national COVID-19 response.

State-level modelling played a role in some of the larger countries of our study, such as Colombia, Canada,

and Australia, but these activities are not reported here. Examining countries with little or no use of

modelling in outbreaks is suggested for further research as this could inform key challenges or

recommendations that have not yet been reported.

Conclusions

With the above limitations in mind, our study provides a first evidence base for the use of epidemiological

modelling in different countries during a major recent health crisis (COVID-19). Analysis has enabled a

classification framework and context-specific recommendations to be suggested, from which further

research and testing can be built.

Looking to the future, we advocate for actors to identify the system they are in and look to other countries

reported in this article with a similar setup (similar size and organisation of modelling capacity, similar level

of government involvement - Figure R2) for key actions to consider for strengthening the use of

epidemiological modelling in outbreaks. No optimal modelling-policy system was observed - there were

benefits to both small and large modelling capacity and to in-house vs independent modelling capacity.

However, disjointed modelling capacity may be ineffective, with isolated modellers in Uganda and South

Korea struggling to identify and recruit adequate support for modelling or networks within the discipline.

Data in this article and in the supplementary dataset (S4) can inform how countries similar to yours have

managed data sharing challenges, communication, relationships, etc., as many of the same challenges

were observed by interviewees in similar systems in our study (section 3.2). Although Figure R2 places

countries in one of five categories based on COVID-19, we note that infrastructure can change significantly

over time, across different disease expertise, or in non-crisis times, so it will be important to update these

classifications. For example, post-COVID-19, some of the countries in our study (for example Canada,

Kenya, South Korea) reported that they have been able to maintain or adapt their COVID-19 modelling-

policy systems post-pandemic, but at the time of interview, New Zealand and South Africa had been

unsuccessful in sourcing long-term funding for modelling-policy systems and their formal epidemiological

modelling committees has been discontinued (Supplementary Material S4 dataset). In addition, Australia

has since begun developing a Centre for Disease Control to maintain the relationships built between

communicable disease scientists and actors in COVID-19[41].

From the guidance given in Table R3, we also advocate for strong data access as a universal

recommendation. For example, the Peru Ministry of Health and Peruvian CDC are working to improve their

data sources after citing poor intervention data as a significant barrier in COVID-19 modelling and outbreak

response (S4). For countries with limited epidemiological modelling capacity, for example a single team

with little capacity for self-verification, reproduction, or consensus with multiple models, there is a critical

need to build standing capacity through secure jobs, training opportunities, and where possible support

from international modelling networks or societies. For example, mass training has taken place in Latin

America and the Caribbean through an open online epidemiological modelling training course held fully in

Spanish by and for Latin American actors, after identifying a lack of capacity in modelling in this region[42].

And finally for countries where modelling is independent from government, it is critical to have an

advocate within policymaking spheres to provide a space for modelling, such as was the case with South

Africa's convenor or the UK's policy co-chair for COVID-19 modelling groups. These individuals enabled

modelling to be integrated and interpreted correctly. Developing relationships with government groups

rather than individuals and establishing agreements and memorandums of understanding ahead of time

will also protect against high turnover and enable rapid scale-up in times of crises.

The relative peacetime that some countries are now in offers a unique opportunity for actors to evaluate

and improve their epidemiological modelling-policy systems, making use of the ideas and context-specific

case studies presented in articles like this one.

This study has directly contributed evidence to formal activities to develop best practice, including the

Lancet Commission on 'Strengthening the Use of Epidemiological Modelling of Emerging and Pandemic

Infectious Diseases'[43].

Author contributions

LH proposed the study. LH, SF, OR, and AT were involved in conception and design. LH and SF led country

and participant selection and recruitment. LH carried out the data collection. LH, CR, and AT carried out

analysis and interpretation of results. LH wrote the first draft of the manuscript. All authors provided

feedback on the manuscript and approved the final version for publication.

Acknowledgements

LH acknowledges support from the Wellcome Trust (block grant no. RG92770). SF also acknowledges

support from the Wellcome Trust (grant no. 210758/Z/18/Z). The authors also thank Sungmok Jung, Mircea

Sofonea, and all other interviewees for dedicating their time and experience to the study.

The authors declare no conflicts of interest.

Data availability

The available anonymised data has been presented in this manuscript and any associated manuscripts

where possible.

References

1. Aguas R, White L, Hupert N, Shretta R, Pan-Ngum W, Celhay O, et al. Modelling the COVID-19 pandemic in context: an international participatory approach. BMJ Glob Health. 2020 Dec 23;5(12):e003126.

2. Jit M, Ainslie K, Althaus C, Caetano C, Colizza V, Paolotti D, et al. Reflections On Epidemiological Modeling To Inform Policy During The COVID-19 Pandemic In Western Europe, 2020–23. Health Aff

(Millwood). 2023 Dec;42(12):1630–6.

3.   Owek CJ, Guleid F, Maluni JK, Jepkosgei J, Were V, Sim SY, et al. LESSONS LEARNED FROM COVID-19
MODELLING EFFORTS FOR POLICY DECISION-MAKING IN LOWER- AND MIDDLE-INCOME COUNTRIES.
BMJ Glob Health [Internet]. 2024 Nov 8 [cited 2024 Mar 13]; Available from:
https://doi.org/10.1136/bmjgh-2024-015247

4.   Xiang Y, Jia Y, Chen L, Guo L, Shu B, Long E. COVID-19 epidemic prediction and the impact of public
health interventions: A review of COVID-19 epidemic models. Infect Dis Model. 2021 Jan 1;6:324–42.

5.   van Elsland SL, O'Hare RM, McCabe R, Laydon DJ, Ferguson NM, Cori A, et al. Policy impact of the
Imperial College COVID-19 Response Team: global perspective and United Kingdom case study. Health
Res Policy Syst. 2024 Nov 13;22(1):153.

6.   McBryde ES, Meehan MT, Adegboye OA, Adekunle AI, Caldwell JM, Pak A, et al. Role of modelling in
COVID-19 policy development. Paediatr Respir Rev. 2020 Sep 1;35:57–60.

7.   Li SL, Bjørnstad ON, Ferrari MJ, Mummah R, Runge MC, Fonnesbeck CJ, et al. Essential information:
Uncertainty and optimal control of Ebola outbreaks. Proc Natl Acad Sci. 2017 May 30;114(22):5659–
64.

8.   Kretzschmar ME, Ashby B, Fearon E, Overton CE, Panovska-Griffiths J, Pellis L, et al. Challenges for
modelling interventions for future pandemics. Epidemics. 2022 Mar 1;38:100546.

9.   Woolhouse M. How to make predictions about future infectious disease risks. Philos Trans R Soc B
Biol Sci. 2011 Jul 12;366(1573):2045–54.

10.   Polonsky JA, Baidjoe A, Kamvar ZN, Cori A, Durski K, Edmunds WJ, et al. Outbreak analytics: a
developing data science for informing the response to emerging pathogens. Philos Trans R Soc B Biol
Sci. 2019 May 20;374(1776):20180276.

11.   Kretzschmar M. Disease modeling for public health: added value, challenges, and institutional
constraints. J Public Health Policy. 2020;41(1):39–51.

12.   Jonkmans N, D'Acremont V, Flahault A. Scoping future outbreaks: a scoping review on the outbreak
prediction of the WHO Blueprint list of priority diseases. BMJ Glob Health. 2021 Sep;6(9):e006623.

13.   Hollingsworth TD. Controlling infectious disease outbreaks: Lessons from mathematical modelling. J
Public Health Policy. 2009 Sep 1;30(3):328–41.

14.   Mollison D, Isham V, Dangerfield C, Hollingsworth D. Preface: Challenges for future pandemics.
Epidemics. 2022 Sep 1;40:100621.

15.   Hadley L, Challenor P, Dent C, Isham V, Mollison D, Robertson DA, et al. Challenges on the interaction
of models and policy for pandemic control. Epidemics. 2021 Dec 1;37:100499.

16.   Howerton E, Contamin L, Mullany LC, Qin M, Reich NG, Bents S, et al. Evaluation of the US COVID-19
Scenario Modeling Hub for informing pandemic response under uncertainty. Nat Commun. 2023 Nov
20;14(1):7260.

17.   Sherratt K, Carnegie AC, Kucharski A, Cori A, Pearson CAB, Jarvis CI, et al. Improving modelling for
epidemic responses: reflections from members of the UK infectious disease modelling community on
their experiences during the COVID-19 pandemic. Wellcome Open Res. 2024 Jan 8;9:12.

18.   CMCC Technical Group. Technical Group Report: Model Fitness-for-purpose Assessment Report

[Internet]. 2020 Aug. Available from:
https://strathprints.strath.ac.uk/78648/1/Megiddo_TGCMMCC_2020_Technical_Group_report_mod
el_fitness_for_purpose_assessment_report.pdf

19. CMCC Policy Group. Guidance on Use of Modelling for Policy Responses to COVID-19 [Internet]. 2020
Aug. Available from: https://www.hitap.net/en/documents/178601

20. Oliwa JN, Guleid F, Owek CJ, Maluni J, Jepkosgei J, Sim SY, et al. A FRAMEWORK TO GUIDE THE USE OF
MATHEMATICAL MODELLING IN EVIDENCE-BASED POLICY DECISION-MAKING. 2024.

21. Christen P, Van Elsland S, Saulo D, Cori A, Fitzner J. Workshop Report: Advanced analytics to inform
decision making during public health emergencies [Internet]. WHO Hub for Pandemic and Epidemic
Intelligence, Berlin, Germany: [object Object]; 2023 May [cited 2024 Apr 15]. Available from:
http://spiral.imperial.ac.uk/handle/10044/1/108600

22. Hadley L, Rich C, Tasker A, Restif O, Funk S. Visual preferences for communicating in crises: evidence
from public health decision makers [Internet]. medRxiv; 2024 [cited 2024 Dec 16]. p.
2024.11.05.24316774. Available from:
https://www.medrxiv.org/content/10.1101/2024.11.05.24316774v2

23. Tong A, Sainsbury P, Craig J. Consolidated criteria for reporting qualitative research (COREQ): a 32-
item checklist for interviews and focus groups. Int J Qual Health Care. 2007 Sep 16;19(6):349–57.

24. Hadley L. Examining the use of infectious disease modelling in outbreak response. University of
Cambridge; 2025.

25. Hitchings R, Latham A. Qualitative methods I: On current conventions in interview research. Prog Hum
Geogr. 2020 Apr 1;44(2):389–98.

26. Palinkas LA, Horwitz SM, Green CA, Wisdom JP, Duan N, Hoagwood K. Purposeful Sampling for
Qualitative Data Collection and Analysis in Mixed Method Implementation Research. Adm Policy Ment
Health Ment Health Serv Res. 2015 Sep 1;42(5):533–44.

27. United Nations Department of Economic and Social Affairs, Population Division. World Population
Prospects 2022 - Documentation [Internet]. 2022 [cited 2024 Apr 15]. Available from:
https://population.un.org/wpp/Download/Documentation/Documentation/

28. United Nations Department of Economic and Social Affairs, Population Division. World Population
Prospects - Demographic Indicators [Internet]. 2022 [cited 2024 Apr 15]. Available from:
https://population.un.org/wpp/Download/Standard/MostUsed/

29. NTI, Johns Hopkins Bloomberg School of Public Health. Global Health Security Index: Building
Collective Action and Accountability. 2019.

30. Dabak S, Lancet Commission Policy Working Group members and secretariat, Dietrich J, Hadley L,
Christen P. Institutional arrangements for strengthening the use of epidemiological modelling for
emerging and pandemic infectious diseases. 2025.

31. Sherratt K, Gruson H, Grah R, Johnson H, Niehus R, Prasse B, et al. Predictive performance of multi-
model ensemble forecasts of COVID-19 across European nations. Wesolowski A, Ferguson NM,
Shaman JL, Pei S, editors. eLife. 2023 Apr 21;12:e81916.

32. Braun V, Clarke V. Using thematic analysis in psychology. Qual Res Psychol. 2006 Jan 1;3(2):77–101.

33. Stapley E, O'Keeffe S, Midgley N. Developing Typologies in Qualitative Research: The Use of Ideal-type

Analysis. Int J Qual Methods. 2022 Apr 1;21:16094069221100633.

34. Caulfield J. How to Do Thematic Analysis | Guide & Examples [Internet]. Scribbr. 2022 [cited 2024 Feb
21]. Available from: https://www.scribbr.co.uk/research-methods/thematic-analysis-explained/

35. Attride-Stirling J. Thematic networks: an analytic tool for qualitative research. Qual Res. 2001 Dec
1;1(3):385–405.

36. Gerhardt U. The Use of Weberian Ideal-Type Methodology in Qualitative Data Interpretation: an
Outline for Ideal-Type Analysis. Bull Sociol Methodol Méthodologie Sociol. 1994 Dec 1;45(1):74–126.

37. Swedberg R. How to use Max Weber's ideal type in sociological analysis. J Class Sociol. 2018 Aug
1;18(3):181–96.

38. Kluge S. Empirically Grounded Construction of Types and Typologies in Qualitative Social Research.
Forum Qual Sozialforschung Forum Qual Soc Res [Internet]. 2000 Jan 31 [cited 2025 Apr 2];1(1).
Available from: https://www.qualitative-research.net/index.php/fqs/article/view/1124

39. Ferguson N. UK Covid-19 Inquiry INQ000185337 - Witness Statement of Professor Neil Ferguson
Module 1, dated 11/05/2023 [Internet]. 2023 Jul [cited 2024 May 21]. Available from:
https://covid19.public-inquiry.uk/documents/inq000185337-witness-statement-of-professor-neil-
ferguson-dated-11-05-2023/

40. Hadley L, Ben Miled S, Clapham H, Djaafara B, Dyson L, Funk S, et al. The elusive 'policy maker': How
understanding the diversity of perspectives and systems in governments can increase the impact of
scientific research. 2025.

41. Australian Government Department of Health and Aged Care. Australian Centre for Disease Control
[Internet]. Australian Government Department of Health and Aged Care; 2023 [cited 2025 Apr 4].
Available from: https://www.health.gov.au/our-work/Australian-CDC

42. Gómez-Bermeo L. Implementation of the Epi training kit: Impact report [Internet]. Epiverse TRACE-
LAC, Pontificia Universidad Javeriana; 2025 Feb [cited 2025 Apr 7] p. 13. Available from:
https://hdl.handle.net/10625/64106

43. Jit M. How modelling can better support public health policy making: the Lancet Commission on
Strengthening the Use of Epidemiological Modelling of Emerging and Pandemic Infectious Diseases.
The Lancet [Internet]. 2023 Dec 20 [cited 2024 Feb 16];0(0). Available from:
https://www.thelancet.com/journals/lancet/article/PIIS0140-6736(23)02758-7/fulltext