Title: Preoperative Exposure to Fine Particulate Matter and Risk of Postoperative Complications: A Single Center Observational Cohort Bayesian Analysis

Authors with affiliations

John F. Pearson MD [1,2], Cameron K. Jacobson MS [1], Calvin S. Riss BS [3], Matthew J. Strickland PhD [3], Longyin Lee BS [4], Neng Wan PhD [4], Tabitha M. Benney PhD [5], Nathan L. Pace MD [1], Ben K. Goodrich PhD [6], Jonah S. Gabry MS [6], John V. Pham MD [1], Cade K. Kartchner BS [1], Jake S. Wood BS [7], Michael H. Andreae MD-PhD[1]

[1] Department of Anesthesiology, University of Utah School of Medicine

[2] Global Change and Sustainability Center, University of Utah

[3] Department of Biostatistics, Epidemiology and Environmental Health, University of Nevada Reno School of Public Health

[4] School of Environment, Society and Sustainability, University of Utah

[5] Department of Political Science, University of Utah

[6] Applied Statistics Center, Columbia University

[7] Indiana University School of Medicine

Author Institutional Titles

John F. Pearson MD, Associate Professor of Anesthesiology, johnf.pearson@utah.edu

Cameron K. Jacobson, MS, Associate Director of Information Technology, cameron.jacobson@hsc.utah.edu

Calvin S. Riss, Graduate Research Assistant, criss@unr.edu

Matthew J. Strickland, Professor and Chair of Public Health, mstrickland@unr.edu

Longyin Lee, Graduate Research Assistant, u1467261@utah.edu

Neng Wan, Associate Professor, neng.wan@utah.edu

Tabitha M. Benney, Associate Professor of Political Science, Tabitha.benney@poli-sci.utah.edu

Nathan L. Pace, Professor of Anesthesiology, n.l.pace@utah.edu

Ben K. Goodrich, Associate Research Scholar, Benjamin.goodrich@columbia.edu

Jonah S. Gabry, Senior Research Associate, Jonah.sol.gabry@columbia.edu

John V. Pham, Assistant Professor of Anesthesiology, john.pham@hsc.utah.edu

Cade K. Kartchner, Medical Student Year 2 (MS-2), cade.kartchner@hsc.utah.edu

Jake Wood, Medical Student Year 4 (MS-4), woodjac@iu.edu

Michael H. Andreae, Professor of Anesthesiology, michael.andreae@hsc.utah.edu

Corresponding Author Contact Information:

John F. Pearson, MD

Department of Anesthesiology, University of Utah School of Medicine

30 North Mario Capecchi Dr.

Salt Lake City, 84112

Phone: +1 978-97-3043

Email: johnf.pearson@utah.edu

Prior Presentations: Association of Air Pollution with Perioperative Complications, 18th Annual World Congress of Anaesthesiologists, March 5, 2024, Singapore.

Preprint Server: URL: https://doi.org/10.1101/2024.08.13.24311943 Title: Preoperative Exposure to Fine Particulate Matter and Risk of Postoperative Complications: A Single Center Observational Cohort Bayesian Analysis. Authors: John F. Pearson, Cameron Jacobson, Calvin Riss, Matt Strickland, Longyin Lee, Neng Wan, Tabitha M. Benney, Nathan L. Pace, Ben Goodrich, Jonah Gabry, Cade Kartchner, Michael H. Andreae. *MedRxiv*.

Acknowledgements: The authors would like to acknowledge TELLUS Networked Sensor Solutions, Inc. for the preparation and provision of the data and visualizations in Figure 1.

Word Count: Abstract: 299; Introduction: ; Discussion ; Number of Figures: 3; Number of Tables: 3; Number of Appendices: 2; Number of Supplemental Digital Files: 1;

Abbreviated Title (Running Head): Association of Air Pollution and Complications

Funding Statement: This study was funded from an intramural University of Utah Wilkes Center for Climate Science and Policy seed grant [to JFP and NW] as well as partially supported by the National Institute of Health National Institute of Environmental

Health Sciences and National Cancer Institute (NIH grant 5R01ES029528-05 [to MS], 5R37CA276365-02 [to NW]). Further funding was from the National Science Foundation (NSF grant 2051246 [BG] and 2153019[BG]). The funding sources were not involved in the study design, collection, analysis and interpretation of the data, in writing the report, or in the decision to submit the article for publication. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health, the National Science Foundation, or the University of Utah Wilkes Center for Climate Science and Policy.

Conflicts of Interest: Ben Goodrich and Jonah Gabry are both member-managers of GG Statistics, LLC. The remainder of the authors declare no competing interests.

Use of Artificial Intelligence Tools: During the preparation of this work the authors used OpenAI's ChatGPT for assistance with writing clarity and literature search. After using this service, the authors reviewed and edited the content and take full responsibility for the content of this publication. This information is included in the Methods section of the manuscript.

Author Approval:

All authors have seen and approved the manuscript.

Abstract

Background: While exposure to fine particulate matter air pollution (PM$_{2.5}$) is known to cause adverse health effects, its impact on postoperative outcomes in US adults remains understudied. Perioperative exposure to PM$_{2.5}$ may induce inflammation that interacts insidiously with the surgical stress response, leading to higher postoperative complications.

Methods: We conducted a single center, retrospective cohort study using data from 49,615 surgical patients living along Utah's Wasatch Front and who underwent elective surgical procedures at a single academic medical center from 2016-2018. Patients' addresses were geocoded and linked to daily Census-tract level PM$_{2.5}$ estimates. We hypothesized that elevated PM$_{2.5}$ concentrations in the week prior to surgery would be associated with an increase in a bundle of major postoperative complications. A hierarchical Bayesians regression model was fit adjusting for age, sex, season, neighborhood disadvantage, and the Elixhauser index of comorbidities.

Results: Postoperative complications increased in a dose-dependent manner with higher concentrations of PM$_{2.5}$ exposure, with a relative increase of 8% in the odds of complications (OR=1.082) for every 10ug/m^3 increase in the highest single-day 24-hr PM$_{2.5}$ exposure during the 7 days prior to surgery. For a 30 fold increase in PM$_{2.5}$ (1 ug/m^3 to 30ug/m^3) the odds of complication rose to over 27% (95%CI: 4%-55%). The

association persisted after controlling for comorbidities and confounders; our inferences were robust to modeling choices and sensitivity analysis.

Conclusions: In this large Utah cohort, exposure to elevated $PM_{2.5}$ concentrations in the week before surgery was associated with a dose-dependent increase in postoperative complications, suggesting a potential impact of air pollution on surgical outcomes. These findings merit replication in larger datasets to identify populations at risk and define the interaction and impact of different pollutants. $PM_{2.5}$ exposure is a potential perioperative risk factor and, given the unmitigated air pollution in urban areas, a global health concern.

Introduction

Short-term exposure to air pollution, especially fine particulate matter of less than 2.5 microns in size ($PM_{2.5}$) is a well-established risk factor for cardiovascular, respiratory, metabolic, and neurologic, morbidity, and mortality worldwide.[1-6] The increasing prevalence of wildfires induced by climate change worsens exposure to fine particulate matter in the US,[7] while millions globally continue to suffer under high levels of $PM_{2.5}$[8,9]. There is emerging evidence that air pollution and the stress response to surgery (**SRS**) interact insidiously as there is overlap between the SRS and the oxidative stress-induced injuries of air pollution. SRS results in pulmonary trauma, hemodynamic stress, and proinflammatory cytokines,[10-16] while air pollution similarly results in increased risk of thrombosis, airway inflammation, and alterations in the immune response.[17,18] The mechanism by which air pollution induces systemic inflammation is well established: small airborne pollutants enter through the lungs or skin[19] and disseminate across the body. Such pollutants can be found in every organ system, including the lungs, brain, the heart, and gastrointestinal tract among others[20]. The impact of individual patient air pollution exposure (IPAPE) thus may lead to an inflammatory response that compounds the SRS[13,21], including pulmonary inflammation, endothelial dysfunction, thrombosis, and membranous nephropathy (Appendix Figure 1).[10,11,15,16] Only a small number of studies[22-26] examined the impact of IPAPE on perioperative outcomes and those that did were limited to specialized populations (Appendix Table 1: Rigor of Prior Research). A recent analysis of over 1.3 million patients in China found increased 30-day postoperative mortality was associated with higher preoperative $PM_{2.5}$ concentrations at the city-level on day of surgery, especially among patients with

preexisting cardiopulmonary conditions[23]. Yet no study examined major post-operative medical complications in any U.S. adult surgical population.

The Salt Lake metropolitan region of Northern Utah is an ideal setting to study acute pollution exposures. Due to the various air pollution and weather events common to the region (e.g. inversion, wildfires) which quickly clear when the weather changes, the random exposure of patients to variable air pollution in the days prior to surgery leads to a quasi-natural experiment as patients are scheduled for surgery regardless of air pollution exposure (Figure 1: The rapidly changing pollution levels along the Wasatch Front).

Our objective was to estimate the impact of preoperative particulate matter ($PM_{2.5}$) air pollution on a bundle of post-operative complications in a local cohort of cases in the national perioperative electronic health registry MPOG.org. We hypothesized that exposure to $PM_{2.5}$ in the 7-day period prior to surgery results in an increased risk of a bundle of post-operative complications, after controlling for comorbidities, confounders, and other well documented drivers of post-operative complications[27,28] in a hierarchical Bayesian regression model.

Materials and Methods

Study Design and Data Sources

We performed a single center cohort analysis of the University of Utah local Multi-Center Perioperative Outcomes Group (MPOG)[29] electronic health registry, which was supplemented with data from the University of Utah Health's Epic database for the period January 1, 2016 to December 31, 2018. We adhered to the Strengthening the Reporting of Observational Studies (STROBE) statement and principles[30]. The study was approved as Exemption Category 4 by the Institutional Review Board at the University of Utah, with IRB approval number 00142167. Prior to access of data, we wrote of our intention to analyze the influence of environmental factors and air pollution on postoperative outcomes with the University of Utah IRB. During the preparation of this work the authors used OpenAI's ChatGPT for assistance with writing clarity and literature search. After using this service, the authors reviewed and edited the content and take full responsibility for the content of this publication.

Study Population: Inclusion and Exclusion Criteria

Extracting information from our Epic and local MPOG instances we included elective and non-emergent general anesthesia cases performed at University of Utah Health operating room locations on or after January 1, 2016 and on or before December 31, 2018. We excluded cases with ASA 5 or 6, age <18 years, cases performed without general anesthesia, as well as obstetric, electroconvulsive therapy, and bronchoscopy procedures. We only included cases where geocoding could match to an address. We

limited our study area further to State of Utah counties along the Wasatch Front, as these counties episodically experience some of the worst particulate matter pollution in the US, and globally, due to inversion events where cold air becomes trapped along the mountain valleys, and are also impacted by wildfire smoke in the summer[31,32]. As such, we included only patients residing in the Counties of Salt Lake, Utah, Davis, Weber, Cache, and Box Elder, all of which are served by the University of Utah Health locations in Salt Lake and Davis Counties of Utah. Further exclusions were made based on availability of the PM$_{2.5}$ estimates and ability to assign Elixhauser Comorbidity scores (Figure 2).

Fine Particulate Matter (PM2.5) Exposure

Daily fine particulate matter with diameter <2.5μm (PM2.5) measurements were obtained from a public dataset of machine learning-derived daily PM$_{2.5}$ concentration estimates at the County, Zip Code, and Census Tract for 11 Western States 2008-2018[33]. These estimates utilize a combination of EPA and state-level ground sensors and satellite derived pollution estimates to provide validated and accurate concentrations of PM$_{2.5}$ across the Western US. We then matched these estimates to individual patients at the census tract of their geocoded home address. For our primary analysis, we utilized the maximum value of PM$_{2.5}$ within 7 days of surgery, which included the day of surgery itself, for two reasons: (1) inversion and pollution events in northern Utah tend to be of short duration[32], so as to ensure catchment of short-lived events without obscuring them with use of multiple day means, the maximum value in

the window was used and (2) prior literature suggests pollution effects on inflammation

and thrombosis may manifest over 1-4 weeks, but most acutely within days of the

exposure event[34,35].

Outcome Bundle: Major Post-operative Complications

The primary outcome was a composite of major postoperative complications occurring

during in-hospital stays after surgery as derived from their presence in the discharge

diagnosis codes captured in Epic and our local MPOG database[36-38], including:

pneumonia, surgical site infection, urinary tract infection, sepsis, stroke, myocardial

infarction, or thromboembolic event. We classified the presence of any post-operative

complication into a binary outcome measure, with a positive (yes) if any complication

was present, while negative (no) if no complications were observed.

Covariates

Multivariable models were adjusted for patient age, sex, year, season, County of

residence, neighborhood deprivation (which is an index of poverty, race, education and

income generated by the National Neighborhood Data Archive (NANDA)[39]), and the

Elixhauser comorbidity index[27,28]. Elixhauser comorbidity index was utilized in place of

the American Society of Anesthesiologists Physical Classification Score (ASA-PS) as it

has demonstrated equivalent reliability in predicting post-operative outcomes[40-42], and

as it is a validated model that relies on documented comorbidities that can be calculated

based on Epic data rather than subjective assessments at the bedside. We assigned

Elixhauser comorbidity scores to all patients utilizing their EPIC records following

standard assignment procedures as outlined by Syed et al[43]. We hypothesized that complications would rise with increasing Elixhauser comorbidity index score.

Statistical Analysis

We investigated the impact of $PM_{2.5}$ exposure on postoperative complications (outcome) using a multivariable Bayesian model. We utilized a three-season exposure model to adjust for possible confounding factors, where summer months (Jun-Aug) were considered "fire season", winter months (Dec-Mar) were considered "inversion" season, and other months (Sep-Nov, Apr-May) were considered a baseline period. Thus, all months were included in the model.

Model fit was by hierarchical Bayesian regression methods using Markov chain Monte Carlo algorithms, specifically Hamiltonian Monte Carlo with the No-U-Turn Sampler having more rapid convergence for high-dimensional models. Models were run with six chains, 2000 iterations and a 50% thinning of the initial estimates. For the beta regression parameters, we used two different weakly informative prior distributions to test for sensitivity of parameter estimates. These were the standard normal distribution N(0, 1) and the R2D2M2(0.25, 4, 05) prior[44]. The priors for the variance parameters with the exponential (exp(1)) and the R2D2M2(0.25, 4, 05) priors[44]. County was included in the statistical model as group effect.

Convergence characteristics of the model estimation was assessed using the Gelman-Rubin R-hat statistic, effective sample size (ESS), chain mixing, and chain

autocorrelation. The posterior predictive distribution was used to generate a predictive

accuracy metric as measured by leave-one-out cross-validation. Nested models were

compared by expected-log-predictive-density. Model fits and parameter values were

explored using conditional effects, R^2 coefficient of determination, and Bayesian

hypothesis testing.

Model results are presented as parameter estimates and odds ratio transformed values

using means, medians, standard deviations, and 95% credible intervals (CI). A 95%

credible interval has a 95% probability of containing the true parameter value. Model

coefficients are also presented with forest plots to show the probability of direction.

Analyses were conducted using R Statistical Software v4.4.1, and used two software

packages *built on STAN* (*brms*) and a Hamiltonian MonteCarlo based software to

estimate Bayesian models[45]. Additional data and model description was done in the R

language using the tableone, loo, mcmcplot, posterior, and tidybayes packages.

Results

Study Population Characteristics

Our initial cohort of patients was n=77,685 for the study period of January 1, 2016 to December 31, 2018, as downloaded and filtered from MPOG.org DataDirect. The majority of exclusions occurred from limiting the sample to the Wasatch Front counties (n=23,243), while a small amount of exclusions occurred due to invalid addresses (n=3,230) and missing $PM_{2.5}$ data (n=544), for a final cohort of n=49,615. Median age was 53.22 years (SD=17.67) and a majority were female (52.2%). Most patients had low comorbidity burden, with a mean Elixhauser of 0.89 (SD=1.95). A majority of patients presented from Salt Lake County (n=34,399), which is the most populous county in the state. There was also a slight increase in total cases per year from 15,375 in 2016, to 16,704 in 2018. Table 1 presents baseline characteristics of patients by study year.

Post-Operative Complications

The overall rate of the composite complication outcome was 4.85% (n=2,411 events among total cohort of 49,615); complication rates are presented in Table 2 broken down by population characteristics in a bivariate analysis (yes vs. no complications). The complication rate varied from a low of 4.41% in Davis County to a high of 7.58% in Box Elder County. While female patients dominated the cohort, they were less likely to experience complications than males (3.86% female vs. 5.94% male respectively). As expected, complication rates increased with a greater Elixhauser comorbidity index, with those without complications having a mean index of 0.6 (SD=1.52) while those with complications had a mean index of 5.15 (SD=3.72) (p<0.001). We also observed a

slight downtrend in complications over time, from 5.13% in 2016 to 4.47% in 2018 (p=0.014). Complications did not vary significantly by season (p=0.834). Finally, we observed that when $PM_{2.5}$ was dichotomized by the EPA daily limit of 35ug/m^3, there was an increased complication rate, from 4.8% (n=2332/48468) on days below 35 ug/m^3 $PM_{2.5}$, to 6.2% (n=79/1147) on days above 35 ug/m^3 (SMD=0.089), and when this entered into the full model had a posterior distribution of 93.2%. The exposure to high pollution did not differ greatly between Counties, nor did it vary by season. These results are summarized in table 3.

Multivariable Exposure Analysis of $PM_{2.5}$ impact on outcome bundle

Model estimation satisfied usual criteria (Gelman-Rubin R-hat, ESS, posterior predictive error checks). There was no significant autocorrelation. The posterior density plots for model parameters indicate reasonable unimodal distributions. Our sensitivity analysis exploring both two weakly informative priors did not change the inferences or results. Our findings were robust to model parameters and model specifications.

In our main Bayesian multivariable model, we found an increased risk of post-operative complications with increasing concentrations of $PM_{2.5}$ with a regression coefficient estimate of 0.01 with a 95% CI (0.00-0.01). A Bayesian hypothesis test showed a 99% probability of an increasing probability of complications with increasing $PM_{2.5}$. The Odds Ratio(OR) of a major perioperative complication for a one unit increase in $PM_{2.5}$ is 1.0079, and for a ten unit increase the OR is 1.082. In clinical terms there is an 8%

increase in the chance of complication for every 10ug/m^3 increase in the maximum

PM2.5 observed in the 7-day preoperative period. For a 30 fold increase in PM$_{2.5}$ (1

ug/m^3 to 30ug/m^3) the odds of complication is over 27% (95%CI: 4%-55%). This

increase was noted in a curvilinear fashion, as presented in Figure 3. There was no

apparent change in this relationship across seasons as we defined them for this region

(fire, inversion, baseline), nor did the inclusion of neighborhood disadvantage alter the

findings. The impact of PM$_{2.5}$ was of greater magnitude among patients with a higher

Elixhauser comorbidity score, especially for those with Elixhauser of 3 or greater

compared to those with Elixhauser <1, as shown in Figure 3. Overall the parameters in

the model explained about 1/4th of the variance (R2 = 0.28).

Discussion

Summary of Findings

In our single center cohort of over 49,000 patients undergoing elective surgery at an academic medical center near the Wasatch front in Utah, we found that increased exposure to $PM_{2.5}$ in the 7-days prior to surgery was associated with significantly higher risk of postoperative complications, confirming our primary hypothesis. The findings were statistically highly significant, and the effect size is consistent with risk observed in the air pollution epidemiology literature[46], providing further confirmation of our findings. We observed a dose-response relationship, with over 27% increased risk of complications at the highest $PM_{2.5}$ concentrations compared to low concentrations, with significant increase in risk above an apparent cut-off exposure of 35 ug/m^3 (Table 3). This association and the cutoff were consistent across all patient age groups, with individuals with higher Elixauser Co-morbidity status appearing to have greater susceptibility to elevated $PM_{2.5}$ (Figure 3). Age, sex, County, year of procedure, neighborhood deprivation (a measure of social distress) and season appeared to have minimal influence on the association. We found that the rate of our complication bundle decreased year over year, while the distribution of Elixhauser scores remained consistent, and the exposure to air pollution remained comparable between years.

Implications for clinical practice, research and policy

To our knowledge, this is the first large-scale study demonstrating the acute impact of preoperative IPAPE on postsurgical inpatient outcomes in a large surgical patient cohort in an OECD country[22,26]. The implications are threefold; our study should inform: (1)

Policy, demonstrating a potentially novel and unaccounted for susceptible population (surgery patients), (2) Individual patient scheduling decisions for high-risk patients (elevated Elixhauser index) and targeted mitigation (masks, filters), and (3) Further research into the pathophysiological mechanisms of healing after surgery in human and animal studies.

Policy Implications

While air pollution in some regions of the U.S. have improved dramatically since the 1970s, improvements have often benefitted certain groups over others. For instance, Black Americans are exposed to higher annual concentrations of air pollution containing fine particulate matter than their White counter parts[47]. Minority populations often face greater pollution burdens and may also be more susceptible to their health effects due to their lack of access to health services and lower income and educational status, which reduces accessibility to pollution information[47] and other forms of prevention. This problem tracks globally as well. Today, air pollution is considered a leading causes of health complications and mortality worldwide, especially affecting lower-income groups, who tend to be more exposed and vulnerable[48]. Consequently, the findings presented in this research have important policy implications for public health and contribute to the literature on health disparities.

Climate change is expected to increase the incidence of wildfires by 29% worldwide[49], which will disproportionately burdening urban and developing regions. These regions are already facing a growing severity of pollution impacts; this research suggests that

the interaction of surgery and pollution will also lead to increasing health burdens. For instance, in China where approximately 10% of GDP[50] is spent on the health effects of pollution, a 1% increase of $PM_{2.5}$ currently leads to a 2.942% increase in household healthcare expenditure[51] and these numbers are expected to grow over the next ten years. Subsequently, developing states, global megacities, and urban areas in Asia are already faced with tough and costly choices. Economically, the impact of air pollution on surgery outcomes will further contribute to decreases in labor productivity for patients and their families, and, therefore, a lower tax base for these communities. In addition, poor surgical outcomes lead to inefficient use of hospital resources, increases the costs of insurance for all, and maybe lead to long term welfare costs, especially where loss of life is concerned. When combined with the disproportionate health disparities, the scale of costs are dramatic for all major population areas.

Scheduling and Targeted Mitigation

Targeted mitigation efforts around the perioperative period may reduce complications attributable to $PM_{2.5}$ exposure among susceptible patients. Surgical delay or protection from pollution during high-risk periods through use of indoor HEPA filtration systems could be considered for patients with planned procedures (if they work indoors or from home) during seasonal inversion events or wildfire smoke conditions. Huang et al's study from China[23], though it focused on mortality, found a beneficial economic impact from rescheduling elective surgery cases in at-risk populations during high pollution events. The implications for elective and non-urgent major surgery, especially during wildfire events, merit further investigation with multidisciplinary teams.

Research Implications and Future Directions

Our findings lead to several new research questions. Our ability to identify high risk populations through sub-group analyses was limited due to sample size, and extension of our study to more years and more sites should help elucidate questions regarding the most vulnerable surgical populations. Additional outcomes should be studied to corroborate our findings and refine the impact of air pollution, incorporating outcome measures such as length of stay, mortality, and additional long- and short-term outcomes. Ours and others future epidemiological studies should also guide and be balanced with animal models where more precise biochemical mechanisms can be uncovered. We are also concerned that social determinants of health might confound the association of small particle pollution with post-surgical outcomes and will study this association in subsequent investigations. Finally, while we did identify that the relationship with $PM_{2.5}$ appeared to accelerate with higher concentrations of pollution, formulating this exposure mathematically into a functional relationship will be a challenge for future research.

Strength and Limitations

This study has several strengths:

1. The natural geography of Utah creates a unique strength to this study in that the same population is exposed to low and high concentrations of pollution.

2. We examined a large cohort of over 49,000 patients over a 3-year contemporary period. We utilized precise $PM_{2.5}$ estimates at the small census tract level and

investigated the effects across the full range of PM2$_{2.5}$ concentrations. We tested

different exposure windows and adjusted for clinical comorbidities.

3. Our outcome measure was a previously used and validated composite of serious

complications encompassing major morbidity events. We specifically focused on

major morbidity complications that are meaningful outcomes for quality

improvement and risk mitigation efforts around elective surgery timing as well as

patient optimization and that are plausibly related to IPAPE. We controlled for

individual patient comorbidities with the validated and widely used Elixhauser

comorbidity index.

Our study has some limitations:

1. Exposure: While our study leveraged a large sample size, we estimated IPAPE

based on census tract locations. This model does not account for proximity to

highways, industrial sources, or other sources of PM$_{2.5}$ pollution that may be

more chronic rather than episodic in nature. This model also used daily mean

exposure estimates and lacked elevation in the model. Both factors could be

biased in terms of missing peaks of exposure within 24-hour windows and at

different elevations, which could lead to further bias in exposure estimates.

Additionally, we evaluated PM$_{2.5}$ mass concentration and did not have data on

particulate composition that may influence toxicity[52].

2. Mitigation and social determinants of health: We could not control for in-home filtration or other personal mitigation measures, which may be less available to those of lower incomes, non-English Language speakers, and in more socially vulnerable neighborhoods.

3. Population: Our cohort was predominantly white and treated at a single health system, although population-level variability in $PM_{2.5}$ exposure was leveraged. We did not account for patient reported or EHR recorded race and ethnicity, which may confound our results. Despite the Salt Lake City region's reputation for homogeneity, the metropolitan area is close to the median diversity index for mid-size US cities. As a result, the findings of this study could be generalized to similar metropolitan areas, both larger and smaller, though the region is dominated by white and Hispanic populations.

4. Analysis Type: As a retrospective analysis, unmeasured confounding is also a possibility, as is incorrect inferences due to the ecological fallacy, we discussed in detail elsewhere[53], though the use of residential address mitigates this in part.

Comparison with the literature and proposed mechanistic pathways

Our findings from over 49,000 patients along Utah's heavily polluted Wasatch Front build on limited prior data on air pollution and postoperative outcomes. In the China analysis, city-level $PM_{2.5}$ during the week prior to surgery was associated with 1% higher adjusted 30-day mortality per 10 µg/m^3 increase in $PM_{2.5}$[23]. Their mortality

association was stronger at higher $PM_{2.5}$ levels (>100 µg/m³), similar to our findings.

The few other studies that have examined adverse impacts of air pollution on

perioperative outcomes have been limited in scope, methodologies and patient

populations: A majority of these studies examined only organ transplants[26]: while a

reasonable suspicion given their immunocompromised nature, they are a relatively

small and high-risk patient populations thus limiting generalizability. Spencer-Hwang et

al found that kidney transplants have an increased risk of fatal MI with increasing ozone,

in a dose-dependent manner[54], while studies of lung transplant recipients suggest that

proximity to major roads can increase risk of chronic allograft dysfunction[26]. Recent data

from California also suggests pediatric patients are susceptible to adverse pulmonary

events under anesthesia during wildfire events, especially those with reactive airway

disease[25].

The 7-day $PM_{2.5}$ exposure timeframe corresponds to the period when detrimental

effects of pollution on inflammatory, thrombotic, and immune pathways implicated in

surgical complications may become manifest[55]. For example, $PM_{2.5}$ instigates systemic

inflammation through release of IL-1, IL-6, and CRP as well as reactive oxygen

species[14,56]. Resultant endothelial dysfunction promotes a prothrombotic state over 7-14

days[57]. Surgical trauma induces a similar acute phase response and

immunomodulation[10-12], which combined with the biological impact from recent pollution

exposure could synergistically heighten complication risk, as detailed in Appendix

Figure 1. Taken as a whole, the literature suggests adverse impacts from air pollution,

and most notably fine particulate matter, on outcomes after surgery. Though major gaps

exist on which pollution source may be most harmful and which patient population may be most affected. Further studies should explore possible mitigation strategies, either pre-operatively or post-operatively, for example the offering of patient bedroom air filters or patient masks preoperatively, or the rescheduling of elective high-risk patients during extreme exposure events, like wildfires.

Conclusion

In our single center cohort study, we found that elevated individual patient exposure to small particles in the week prior to surgery was associated with significantly increased postoperative complications. We demonstrated a dose-response relationship in all age groups, regardless of patient co-morbidity (as measured by the Elixhauser index) in a large cohort at an academic medical center in Utah. This clearly demonstrates a statistically significant impact of air pollution exposure on surgical outcomes, with an effect size consistent with the broader pollution literature[46].

Our results cover a wide swath of surgical specialties and thus have implications for much of elective surgery performed in areas suffering from periodic acute pollution episodes, such as inversion events and, much more commonly with climate change, wildfire smoke. At-risk patients may benefit from pollution mitigation and close monitoring in the postoperative period after procedures preceded by heavy pollution exposure. Overall, our findings highlight that limiting particulate matter exposure through clean air policies and practices can have wide-ranging health benefits beyond just cardiopulmonary disease, including reducing complications of surgery.

References

1.   Farhadi Z, Abulghasem Gorgi H, Shabaninejad H, Aghajani Delavar M, Torani S: Association between PM(2.5) and risk of hospitalization for myocardial infarction: a systematic review and a meta-analysis. BMC Public Health 2020; 20: 314

2.   Atkinson RW, Kang S, Anderson HR, Mills IC, Walton HA: Epidemiological time series studies of PM2.5 and daily mortality and hospital admissions: a systematic review and meta-analysis. Thorax 2014; 69: 660-5

3.   Fu P, Guo X, Cheung FMH, Yung KKL: The association between PM(2.5) exposure and neurological disorders: A systematic review and meta-analysis. Sci Total Environ 2019; 655: 1240-1248

4.   Reid CE, Brauer M, Johnston FH, Jerrett M, Balmes JR, Elliott CT: Critical Review of Health Impacts of Wildfire Smoke Exposure. Environ Health Perspect 2016; 124: 1334-43

5.   Shah AS, Lee KK, McAllister DA, Hunter A, Nair H, Whiteley W, Langrish JP, Newby DE, Mills NL: Short term exposure to air pollution and stroke: systematic review and meta-analysis. BMJ 2015; 350: h1295

6.   Pearson JF, Bachireddy C, Shyamprasad S, Goldfine AB, Brownstein JS: Association between fine particulate matter and diabetes prevalence in the U.S. Diabetes Care 2010; 33: 2196-201

7.   Burke M, Childs ML, de la Cuesta B, Qiu M, Li J, Gould CF, Heft-Neal S, Wara M: The contribution of wildfire to PM(2.5) trends in the USA. Nature 2023; 622: 761-766

8.   Marlier ME, Jina AS, Kinney PL, DeFries RS: Extreme Air Pollution in Global Megacities. Current Climate Change Reports 2016; 2: 15-27

9.   Collaborators GBDRF: Global burden of 87 risk factors in 204 countries and territories, 1990-2019: a systematic analysis for the Global Burden of Disease Study 2019. Lancet 2020; 396: 1223-1249

10.   McSorley ST, Watt DG, Horgan PG, McMillan DC: Postoperative Systemic Inflammatory Response, Complication Severity, and Survival Following Surgery for Colorectal Cancer. Ann Surg Oncol 2016; 23: 2832-40

11.   Iwasaki M, Edmondson M, Sakamoto A, Ma D: Anesthesia, surgical stress, and "long-term" outcomes. Acta Anaesthesiol Taiwan 2015; 53: 99-104

12.   Alam A, Hana Z, Jin Z, Suen KC, Ma D: Surgery, neuroinflammation and cognitive impairment. EBioMedicine 2018; 37: 547-556

13.   Takenaka K, Ogawa E, Wada H, Hirata T: Systemic inflammatory response syndrome and surgical stress in thoracic surgery. J Crit Care 2006; 21: 48-53; discussion 53-5

14.   Costa LG, Cole TB, Dao K, Chang YC, Coburn J, Garrick JM: Effects of air pollution on the nervous system and its possible role in neurodevelopmental and neurodegenerative disorders. Pharmacol Ther 2020; 210: 107523

15.   Shkirkova K, Lamorie-Foote K, Connor M, Patel A, Barisano G, Baertsch H, Liu Q, Morgan TE, Sioutas C, Mack WJ: Effects of ambient particulate matter on vascular tissue: a review. J Toxicol Environ Health B Crit Rev 2020; 23: 319-350

16.   Xu X, Nie S, Ding H, Hou FF: Environmental pollution and kidney diseases. Nat Rev Nephrol 2018; 14: 313-324

17.   Glencross DA, Ho TR, Camina N, Hawrylowicz CM, Pfeffer PE: Air pollution and its effects on the immune system. Free Radic Biol Med 2020; 151: 56-68

18.   Lodovici M, Bigagli E: Oxidative stress and air pollution exposure. J Toxicol 2011; 2011: 487074

19.   Weschler CJ, Beko G, Koch HM, Salthammer T, Schripp T, Toftum J, Clausen G: Transdermal Uptake of Diethyl Phthalate and Di(n-butyl) Phthalate Directly from Air: Experimental Verification. Environ Health Perspect 2015; 123: 928-34

20.   Li T, Yu Y, Sun Z, Duan J: A comprehensive understanding of ambient particulate matter and its components on the adverse health effects based from epidemiological and laboratory evidence. Part Fibre Toxicol 2022; 19: 67

21.   Cusack B, Buggy DJ: Anaesthesia, analgesia, and the surgical stress response. BJA Educ 2020; 20: 321-328

22.   Kovtun R, Ha L, Mendoza D, Pearson J: Influence of Air Pollution on Perioperative Outcomes & Potential for Big Data Driven Discoveries. Anesthesia and Analgesia 2021; 132: 49-51

23.   Huang JL, Xing JW, Zou EY: (Re)scheduling pollution exposure: The case of surgery schedules. Journal of Public Economics 2023; 219

24.   Oh TK, Song IA: Mortality Risk After a Major Cancer Surgery Is Associated With Preoperative Exposure to Air Pollution. Journal of Occupational and Environmental Medicine 2024; 66: 192-197

25.   Marsh BJ, Kolodzie K, Robinowitz D, Jacobson A, Ferschl M: Wildfire Smoke Exposure Is Associated with Adverse Respiratory Events under General Anesthesia in At-risk Pediatric Patients. Anesthesiology 2022; 137: 543-554

26.   Supphapipat K, Leurcharusmee P, Chattipakorn N, Chattipakorn SC: Impact of air pollution on postoperative outcomes following organ transplantation: Evidence from clinical investigations. Clin Transplant 2024; 38: e15180

27.   van Walraven C, Austin PC, Jennings A, Quan H, Forster AJ: A modification of the Elixhauser comorbidity measures into a point system for hospital death using administrative data. Med Care 2009; 47: 626-33

28.   Austin SR, Wong YN, Uzzo RG, Beck JR, Egleston BL: Why Summary Comorbidity Measures Such As the Charlson Comorbidity Index and Elixhauser Score Work. Med Care 2015; 53: e65-72

29.   Kheterpal S: Clinical research using an information system: the multicenter perioperative outcomes group. Anesthesiol Clin 2011; 29: 377-88

30.   von Elm E, Altman DG, Egger M, Pocock SJ, Gotzsche PC, Vandenbroucke JP, Initiative S: The Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) Statement: guidelines for reporting observational studies. Int J Surg 2014; 12: 1495-9

31.   Wilmot TY, Hallar AG, Lin JC, Mallia DV: Expanding number of Western US urban centers face declining summertime air quality due to enhanced wildland fire activity. Environmental Research Letters 2021; 16

32.   Mitchell LE, Zajchowski CAB: The History of Air Quality in Utah: A Narrative Review. Sustainability 2022; 14

33.   Reid CE, Considine EM, Maestas MM, Li G: Daily PM(2.5) concentration estimates by county, ZIP code, and census tract in 11 western states 2008-2018. Sci Data 2021; 8: 112

34.   Arias-Perez RD, Taborda NA, Gomez DM, Narvaez JF, Porras J, Hernandez JC: Inflammatory effects of particulate matter air pollution. Environ Sci Pollut Res Int 2020; 27: 42390-42404

35.   Pope CA, 3rd, Ezzati M, Dockery DW: Fine-particulate air pollution and life expectancy in the United States. N Engl J Med 2009; 360: 376-86

36.   Sun E, Mello MM, Rishel CA, Vaughn MT, Kheterpal S, Saager L, Fleisher LA, Damrose EJ, Kadry B, Jena AB, Multicenter Perioperative Outcomes G: Association of Overlapping Surgery With Perioperative Outcomes. JAMA 2019; 321: 762-772

37.   Wang HH, Tejwani R, Zhang H, Wiener JS, Routh JC: Hospital Surgical Volume and Associated Postoperative Complications of Pediatric Urological Surgery in the United States. J Urol 2015; 194: 506-11

38.   Lawson EH, Louie R, Zingmond DS, Brook RH, Hall BL, Han L, Rapp M, Ko CY: A comparison of clinical registry versus administrative claims data for reporting of 30-day surgical complications. Ann Surg 2012; 256: 973-81

39.   Research IfS: The National Neighborhood Data Archive (NaNDA), The Regents of the University of Michigan, 2024

40.   Hinton ZW, Fletcher AN, Ryan SP, Wu CJ, Bolognesi MP, Seyler TM: Body Mass Index, American Society of Anesthesiologists Score, and Elixhauser Comorbidity Index Predict Cost and Delay of Care During Total Knee Arthroplasty. Journal of Arthroplasty 2021; 36: 1621-1625

41.   Maron SZ, Neifert SN, Ranson WA, Nistal DA, Rothrock RJ, Cooke P, Lamb CD, Cho SK, Caridi JM: Elixhauser Comorbidity Measure is Superior to Charlson Comorbidity Index In - Predicting Hospital Complications Following Elective Posterior Cervical Decompression and Fusion. World Neurosurgery 2020; 138: E26-E34

42.   Menendez ME, Neuhaus V, van Dijk CN, Ring D: The Elixhauser comorbidity method outperforms the Charlson index in predicting inpatient death after orthopaedic surgery. Clin Orthop Relat Res 2014; 472: 2878-86

43.   Syed S, Baghal A, Prior F, Zozus M, Al-Shukri S, Syeda HB, Garza M, Begum S, Gates K, Syed M, Sexton KW: Toolkit to Compute Time-Based Elixhauser Comorbidity Indices and Extension to Common Data Models. Healthcare Informatics Research 2020; 26: 193-200

44.   Aguilar JE, Bürkner PC: Intuitive joint priors for Bayesian linear multilevel models: The R2D2M2 prior. Electronic Journal of Statistics 2023; 17: 1711-1767

45.   Bürkner PC: brms: An R Package for Bayesian Multilevel Models Using Stan. Journal of Statistical Software 2017; 80: 1-28

46.   Lee D, Walton H, Evangelopoulos D, Katsouyanni K, Gowers AM, Shaddick G, Mitsakou C: Health impact assessment for air pollution in the presence of regional variation in effect sizes: The implications of using different meta-analytic approaches. Environ Pollut 2023; 336: 122465

47.   Josey KP, Delaney SW, Wu X, Nethery RC, DeSouza P, Braun D, Dominici F: Air Pollution and Mortality at the Intersection of Race and Social Class. N Engl J Med 2023; 388: 1396-1404

48.   Rentschler J, Leonova N: Global air pollution exposure and poverty. Nat Commun 2023; 14: 4432

49.   Senande-Rivera M, Insua-Costa D, Miguez-Macho G: Spatial and temporal expansion of global wildland fire activity in response to climate change. Nat Commun 2022; 13: 1208

50.   Wang XP, Mauzerall DL: Evaluating impacts of air pollution in China on public health: Implications for future air pollution and energy policies. Atmospheric Environment 2006; 40: 1706-1721

51.   Yang J, Zhang B: Air pollution and healthcare expenditure: Implication for the benefit of air pollution control in China. Environment International 2018; 120: 443-455

52.   Park M, Joo HS, Lee K, Jang M, Kim SD, Kim I, Borlaza LJS, Lim H, Shin H, Hyuckchung K, Choi YH, Park SG, Bae MS, Lee J, Song H, Park K: Differential toxicities of fine particulate matters from various sources. Scientific Reports 2018; 8

53.   Pearson J, Jacobson C, Ugochukwu N, Asare E, Kan K, Pace N, Han J, Wan N, Schonberger R, Andreae M: Geospatial analysis of patients' social determinants of health for health systems science and disparity research. Int Anesthesiol Clin 2023; 61: 49-62

54.   Spencer-Hwang R, Knutsen SF, Soret S, Ghamsary M, Beeson WL, Oda K, Shavlik D, Jaipaul N: Ambient air pollutants and risk of fatal coronary heart disease among kidney transplant recipients. Am J Kidney Dis 2011; 58: 608-16

55.   Robertson S, Miller MR: Ambient air pollution and thrombosis. Particle and Fibre Toxicology 2018; 15

56.   Yan YH, Chou CC, Lee CT, Liu JY, Cheng TJ: Enhanced insulin resistance in diet-induced obese rats exposed to fine particles by instillation. Inhal Toxicol 2011; 23: 507-19

57.   Nemmar A, Al-Salam S, Beegam S, Yuvaraju P, Ali BH: The acute pulmonary and thrombotic effects of cerium oxide nanoparticles after intratracheal instillation in mice. Int J Nanomedicine 2017; 12: 2913-2922

58.   Ou J, Pirozzi CS, Horne BD, Hanson HA, Kirchhoff AC, Mitchell LE, Coleman NC, Pope CA: Historic and Modern Air Pollution Studies Conducted in Utah. Atmosphere 2020; 11

59.   Errigo IM, Abbott BW, Mendoza DL, Mitchell L, Sayedi SS, Glenn J, Kelly KE, Beard JD, Bratsman S, Carter T, Chaney RA, Follett A, Freeman A, Frei RJ, Greenhalgh M, Holmes HA, Howe PD, Johnston JD, Lange L, Martin R, Stacey A, Tran T, Wilson D: Human Health and Economic Costs of Air Pollution in Utah: An Expert Assessment. Atmosphere 2020; 11

Figures

Figure 1: Rapidly changing pollution levels along the Wasatch front

   Legend

Figure 1 illustrates the rapidly changing pollution levels along the "Wasatch Front" mountain range in Salt Lake City, UT. The unique mountain geography of the "Wasatch Front," along with five oil refineries, multiple industrial facilities, 2.2 million people, the intersection of two major interstate highways, and a national train depot, work together to produce the increased occurrence of extreme pollution events in the region, which dramatically impact ambient $PM_{2.5}$ concentrations[32,58,59]. This results in inversion conditions during the winter, with warm air aloft the valley trapping cold air and pollutants in the densely populated valley below. Similarly, in the summer when wildfires throughout the American West dominate pollution exposures, the geological bowl produced by the intersection of multiple mountain ranges that comprise the Wasatch Front act as a shield that accumulates wildfire smoke in the metropolitan region. In both instances, low pressure systems tend to rapidly clear the area of pollutants, at random. As a result, patients are quasi-randomly exposed to high or low pollution levels, while their surgery scheduling is disregarding this exposure, leading to a natural experiment. The left map illustrates high observed pollution levels (red/purple for unhealthy high pm 2.5 levels) due to a recent wildfire event on September 11-12, 2024 while the right map demonstrates how a change in weather patterns cleared the air, leading to suddenly much lower pollution exposures (blue). The graph demonstrates this timeline of rapidly changing 2.5 small particle pollution levels, with markers for the dates of the maps.

Figure 2: Consort Flow Diagram:

Legend

This figure details our initisal cohort downloaded from Data Direct from MPOG.org and the process of exclusion criteria applied to generate our final cohort. Of note, the greatest number of patients eliminated was when excluding those who lived outside of the Wasatch Front study area.

Cases Filtered Using Data Direct
(e.g. 2016-2018, Adults, Procedure Type, etc.)
(n = 77685)

Invalid Addresses
(n = 3230)

Geocoded to Valid Addresses
(n = 74360)

Outside Wasatch Front
(n = 23243)

Wasatch Front
(n = 51117)

No UoC PM2.5 Available (n = 544)

University of Colorado
PM2.5 Estimates by Tract
(n = 50573)

Missing or Negative PM2.5
n=953

Positive PM2.5
(n = 49620)

No Elixhauser Comorbidity Scores
(n = 0)

Elixhauser Comorbidity Scores
(n = 49620)

No NaNDA Data
(n = 5)

NaNDA Social Determinants of Health
(n = 49615)

Final Data Set
(n = 49615)

Figure 3: Maximum PM2.5 in the 7 days pre-operatively (Lag0-Lag6) vs. Complication Rate.

Legend:

Panel A (top) details Complications Rate vs. Maximum PM2.5 of the 7-day lag period, while Panel B (bottom) details the same broken down by Elixhuaser Comorbidity Index. Complication rate is shown as absolute value, so here 0.04 = 4% complication rate. The PM2.5 values are the maximum observed in the 7-day preop window, in ug/m^3, Lag 0 is day of surgery. In Panel B as noted, the overall complication rate for those with higher Elixhauser comorbidity index is higher at baseline, but rose more quickly with elevations in PM2.5. This response to PM2.5 continued to increase in magnitude as Elixhauser index increased. Note that the effect of PM2.5 was greater among this with Elixhauser of 5, but that the baseline complication rate was also higher.

Tables

Table 1: Cohort Characteristics by Year

Overall cohort presented by year. Complications fell marginally year over year, while overall number of cases remained relatively steady. The majority of cases were in Salt Lake County, where the greatest number of University of Utah operating rooms are located.

Factor	Level	Overall	2016	2017	2018	p-test	SMD
Sample Total		49615	15375 (30.9)	17536 (35.3)	16704 (33.6)		
Complications (%)	No	47204 (95.1)	14586 (94.9)	16661 (95.0)	15957 (95.5)	0.014	0.02
	Yes	2411 (4.8)	789 (5.1)	875 (5.0)	747 (4.5)		
Elix (mean (SD))		0.89 (1.95)	0.83 (1.83)	0.91 (2.02)	0.91 (1.99)	<0.001	0.029
Sex (%)	Male	23704 (47.8)	7339 (47.7)	8366 (47.7)	7999 (47.9)	0.939	0.002
	Female	25911 (52.2)	8036 (52.3)	9170 (52.3)	8705 (52.1)		
Season (%)	None	17044 (34.4)	4930 (32.1)	5980 (34.1)	6134 (36.7)	<0.001	0.075
	Fire	16472 (33.2)	5457 (35.5)	5704 (32.5)	5311 (31.8)		
	Cars	16099 (32.4)	4988 (32.4)	5852 (33.4)	5259 (31.5)		
Disadvantage (mean (SD))		0.06 (0.04)	0.06 (0.04)	0.06 (0.04)	0.06 (0.04)	0.214	0.013
Age (mean (SD))		53.22 (17.6)	53.31 (17.61)	53.43 (17.73)	52.92 (17.67)	0.023	0.019
County Name (%)	Box Elder	672 (1.4)	193 (1.3)	246 (1.4)	233 (1.4)	0.029	0.034
	Cache	1098 (2.2)	335 (2.2)	386 (2.2)	377 (2.3)		
	Davis	7047 (14.2)	2055 (13.4)	2526 (14.4)	2466 (14.8)		
	Salt Lake	34399 (69.3)	10827 (70.4)	12113 (69.1)	11459 (68.6)		
	Utah	3847 (7.8)	1179 (7.7)	1386 (7.9)	1282 (7.7)		
	Weber	2552 (5.1)	786 (5.1)	879 (5.0)	887 (5.3)		

Table 2: Cohort characteristics dichotomized by presence of a complication

Cohort dichotomized by presence of complication. The complication rate fell slightly (p=0.014) from 2016 to 2018, with an overall rate of 4.85%. The Elixhauser comorbidity score was markedly higher in for those with complications (5.15) than those without (0.6), while age only varied slightly by those without (53.11) and with (55.28) complications.. Complication rates by County also varied, with Box Elder (7.58%) having the greatest rate and Davis with the lowest (4.41%).

Factor	Level	Overall	No	Yes	Rate	p-test	SMD
n		49615	47204	2411	4.85%		
Complications = Yes (%)		2411 (4.8)	0 (0.0)	2411 (100.0)		<0.001	
Procedure Year Factor (%)	2016	15375 (31.0)	14586 (30.9)	789 (32.7)	5.13%	0.014	0.062
	2017	17536 (35.3)	16661 (35.3)	875 (36.3)	4.98%		
	2018	16704 (33.7)	15957 (33.8)	747 (31.0)	4.47%		
Elix (mean (SD))		0.89 (1.95)	0.6 (1.52)	5.15 (3.72)		<0.001	1.576
Sex = Female (%)		25911 (52.2)	24909 (52.8)	1002 (41.6)	3.86%	<0.001	0.226
Season (%)	None	17044 (34.4)	16207 (34.3)	837 (34.7)	4.91%	0.834	0.013
	Fire	16472 (33.2)	15667 (33.2)	805 (33.4)	4.88%		
	Cars	16099 (32.4)	15330 (32.5)	769 (31.9)	4.77%		
Disadvantage1317 (mean (SD))		0.06 (0.04)	0.06 (0.04)	0.07 (0.04)		<0.001	0.218
Age (mean (SD))		53.22 (17.67)	53.11 (17.72)	55.28 (16.59)		<0.001	0.126
County Name (%)	Box Elder	672 (1.4)	621 (1.3)	51 (2.1)	7.58%	<0.001	0.106
	Cache	1098 (2.2)	1041 (2.2)	57 (2.4)	5.19%		
	Davis	7047 (14.2)	6736 (14.3)	311 (12.9)	4.41%		
	Salt Lake	34399 (69.3)	32775 (69.4)	1624 (67.4)	4.72%		
	Utah	3847 (7.8)	3612 (7.7)	235 (9.7)	6.10%		
	Weber	2552 (5.1)	2419 (5.1)	133 (5.5)	5.21%		

Table 3: Cohort characteristics dichotomized by whether or not exposed to maximum of 35ug/m^3 in pre-operative period.

Table 3 dichotomizes patients based on their exposure to any day of PM$_{2.5}$ greater than 35ug/m^3 in the 7 day pre-operative period. In this instance, the complication rate was higher when patients were exposed to PM$_{2.5}$>35 (SMD=0.089). Notably, 2017 had much higher exposure to elevated PM$_{2.5}$ (546 Cases, 47.6% of total exposures), while disadvantage was not dramatically different between higher (0.07) and lower (0.06). Finally, exposure to fire vs. cars as source of pollution was nearly evenly split, indicating equal exposure to elevated inversion pollution and wildfire smoke.

Factor	Level	Overall	Max PM2.5 > 35	Max PM2.5 <35	SMD
n		49615	1147	48468	
Complications (%)	No	47204 (95.1)	1068 (93.1)	46136 (95.2)	0.089
	Yes	2411 (4.9)	79 (6.9)	2332 (4.8)	
Procedure Year Factor (%)	2016	15375 (31.0)	274 (23.9)	15101 (31.2)	0.260
	2017	17536 (35.3)	546 (47.6)	16990 (35.1)	
	2018	16704 (33.7)	327 (28.5)	16377 (33.8)	
Elix (mean (SD))		0.89 (1.95)	1.02 (2.00)	0.88 (1.95)	0.067
Sex (%)	Male	23704 (47.8)	549 (47.9)	23155 (47.8)	0.002
	Female	25911 (52.2)	598 (52.1)	25313 (52.2)	
Season (%)	None	17044 (34.4)	0 (0.0)	17044 (35.2)	1.042
	Fire	16472 (33.2)	578 (50.4)	15894 (32.8)	
	Cars	16099 (32.4)	569 (49.6)	15530 (32.0)	
Disadvantage (mean (SD))		0.06 (0.04)	0.07 (0.04)	0.06 (0.04)	0.310
Age (mean (SD))		53.22 (17.67)	52.26 (17.80)	53.24 (17.67)	0.056
County Name (%)	Box Elder	672 (1.4)	18 (1.6)	654 (1.3)	0.058
	Cache	1098 (2.2)	29 (2.5)	1069 (2.2)	
	Davis	7047 (14.2)	175 (15.3)	6872 (14.2)	
	Salt Lake	34399 (69.3)	786 (68.5)	33613 (69.4)	
	Utah	3847 (7.8)	77 (6.7)	3770 (7.8)	
	Weber	2552 (5.1)	62 (5.4)	2490 (5.1)	