Gynaecological health patterns and motherhood experiences of female

professional football players

Dimakatso Ramagole,[1,2] Dina C Janse van Rensburg,[2] Charlotte Cowie,[3] Ritan

Mehta,[3] Gopika Ramkilawon,[4] Babette M. Pluim,[1,2] Gino Kerkhoffs,[1,5,6,7], Vincent

Gouttebarge[1,2,5,6,7,8]

1 Amsterdam UMC location University of Amsterdam, Department of Orthopedic

Surgery and Sports Medicine, Meibergdreef 9, Amsterdam, the Netherlands

2 Section Sports Medicine, Faculty of Health Sciences, University of Pretoria,

Pretoria, South Africa

3 The Football Association (England), National Football Centre, St George's Park,

Needwood, DE13 4PD, UK

4 Department of Statistics, Faculty of Natural and Agricultural Sciences, University of

Pretoria

5 Academic Center for Evidence-based Sports Medicine (ACES), Amsterdam, the

Netherlands

6 Amsterdam Movement Sciences, Aging & Vitality, Musculoskeletal Health, Sports,

Amsterdam, The Netherlands

7 Amsterdam Collaboration on Health & Safety in Sports (ACHSS), IOC Research

Center of Excellence, Amsterdam, the Netherlands

8 Football Players Worldwide (FIFPRO), Hoofddorp, the Netherlands

CORRESPONDING AUTHOR

Vincent Gouttebarge

Amsterdam UMC location University of Amsterdam, Department of Orthopedic

Surgery and Sports Medicine, Meibergdreef 9, 1105 AZ Amsterdam, the Netherlands

Email: v.gouttebarge@amsterdamumc.nl

Telephone: +31621547499

ORCHID ID's

Dimakatso Ramagole https://orcid.org/0000-0002-5341-6080

Dina C (Christa) Janse van Rensburg http://orcid.org/0000-0003-1058-6992

Charlotte Cowie https://orcid.org/0009-0002-3417-4335

Ritan Mehta

Gopika Ramkilawon https://orcid.org/0000-0002-2919-1918

Babette Pluim https://orcid.org/ 0000-0003-0655-8980

Gino Kerkhoffs https://orcid.org/ 0000-0001-5916-7049

Vincent Gouttebarge https://orcid.org/0000-0002-0126-4177

Abstract

Aims: To explore health patterns in female professional football players in the

domains of gynaecological health patterns, contraceptive use, body perception and

motherhood experiences, including return to play after childbirth.

Methods: An online questionnaire was emailed to active female professional football

players via email. Participants were asked about their menstrual cycle,contraception

use and motherhood. Validated questionnaires were used to assess body

dissatisfaction (BD) and drive for thinness (DT).

Results: A total of 74 female professional football players were enrolled. The mean

age at menarche was 13.5 years, average cycle length of 26 days and a bleeding

period of 5 days. Cycle irregularities were experienced by 30% of participants, and

menstrual symptoms by 74%. Half of the participants used contraceptives, with 60%

using hormonal contraceptives, primarily oral contraceptive pills (38%), followed by

implants (20%). Participants had a normal BD score, but a higher-than-expected DT

score. The motherhood rate was low (1%), with normal conception, vaginal delivery,

return to training after 6 weeks, and return to competition after 12 weeks.

Conclusion: Cycle irregularities are common in female professional football players,

with a significant number of cycle-related symptoms. The majority on contraceptives

preferred hormonal contraceptives, especially oral contraceptive pills (OCP) followed

by implants, reflecting trends seen in elite athletes gynaecological health. While body

satisfaction scores were normal, there was an unexpectedly high drive-for-thinness

score, similar to that observed in lean or weight-category sports. The rate of

motherhood was low, consistent with previous findings in professional football

players.

Keywords: female professional football, gynaecological health patterns, motherhood

Word count: 244

Data availability statement :

All data relevant to the study are included in the article.

What is already known on this topic

- Elite female athletes have unique challenges related to their menstrual cycle and symptoms.

- Elite athletes use mostly hormonal contraceptives, predominantly the OCP.

- They often have to choose between professional sports and motherhood.

What this study adds

Our findings show some similarities to other to other elite female athletes:

- 30% have menstrual irregularities.

- 74% have negative menstrual symptoms.

- Motherhood is low.

- Our participants have a high drive for thinness.

How this study might affect research, practice or policy

- This descriptive study is a basis for more prospective and epidemiological research in this cohort that has been established.

- More research will be stimulated in the domains of menstrual symptoms and coping strategies, choice of contraception and motherhood.

- This study should also stimulate communications regarding player contracts and maternity benefits, and include after career support for female professional football players.

INTRODUCTION

In recent years, there has been an increase in female participation in sports.

Menstrual irregularities, delayed menarche, secondary amenorrhoea and disordered

eating (DE) are common amongst female athletes and have been associated with

disadvantages in the sporting career, as well as shortening the sporting career of

endurance athletes.[1] The menstrual cycle is an integral part of female athletes, and

some earlier research suggested that early menarche (below age 12 years) has been

associated with a higher risk of breast cancer due to the early onset of ovulatory

cycles.[2]

Several surveys have investigated contraceptive use, including the methods chosen

and the reasons behind their use.[3-4] These surveys showed that more than 50% of

their participants were not on contraceptives at all,[4] and those who were on

contraceptives predominantly used oral contraceptive pills (OCP).[3]

Motherhood experiences (ME) have unique challenges for female athletes, as

highlighted in the article 'We are not superhuman, we're human.'[5] Some of the

concerns raised by athletes included: 'training a new body' postpartum; safe return to

training; breastfeeding and exercise; and managing time between sports and

motherhood. These athletes emphasised the need to develop best practices and

secure funding for a safe return to play (RTP).[5-6]

Planned and well-structured physical activity under the guidance of an obstetrician-

gynaecologist during and after pregnancy has been shown to benefit both physical

and mental well-being. These benefits include a decrease in gestational diabetes,

caesarean and operative vaginal deliveries, and a shorter post-partum recovery

time.[7]

Several surveys highlighted the difficult choice between motherhood and elite sports

participation, the need for careful planning of pregnancy and fertility,[8] concerns about

pregnancy disclosures and fear of discrimination, challenges in training the pregnant

body safely, and the lack of a supportive network and funding during the pre- and

post-natal periods.[5]

These interviews suggest that fewer than 50% of female professional football players

have employment contracts comparable to those of their male counterparts. This lack

of contractual security places female footballers at a disadvantage because they

struggle to sustain themselves financially if they get injured or decide to start a

family.[8]

Based on the considerations mentioned above, this study aimed to achieve two

primary objectives: 1) To explore gynaecological health patterns in female

professional football players, including the menstrual cycle, menstrual symptoms,

contraceptive use and body perception; 2) To examine motherhood experiences,

including methods of conception and delivery, and return-to-play/return to competition

after childbirth.

METHODS

Study designs and ethical considerations

An observational prospective cohort study over a follow-up period of six months was

conducted using the Strengthening the Reporting of Observational Studies in

Epidemiology statement in order to guarantee the quality of reporting.[9] The Medical

Ethics Review Committee of the Amsterdam University Medical Centers (Amsterdam

UMC, location AMC) provided ethical approval for the study (Drake Football Study:

NL69852.018.19 | W19_171#B202169). The study was conducted in accordance

with the Declaration of Helsinki (2013).

Setting

Participants from English and French speaking countries were recruited by Football

Players Worldwide (FIFPRO) and affiliated national unions via email from September

202 to May 2021. Initial data collection was completed in December 2022, and

participants will be followed at 6 months' intervals over a period of 10 years..

Participant selection

The study population consisted of female professional footballers. Inclusion criteria

were: (a) being a professional footballer, (b) being female, (c) being able to read and

comprehend texts in English or French. In our study, the definition for a professional

footballer was that she (i) trains to improve performance, (ii) competes in the highest

or second highest national league, and (iii) has football training and competition as a

major activity (way of living) or focus of personal interest, devoting several hours in all

or most of the days for these activities, and exceeding the time allocated to other

types of professional or leisure activities. Sample size calculation indicated that at

least 40 participants were needed to reach a power of 80% (confidence interval (CI)

of 95% and absolute precision of 7%) under the assumption of an anticipated

population proportion (prevalence) of 5%.[10]

Dependent variables

Various female-specific variables were collected through single questions related to

(history of) pregnancy, fertility (e.g., in vitro fertilisation, sperm donation), miscarriage,

adoption, date of menarche, menstrual cycles, (history of) amenorrhoea, use of

contraceptive measures (e.g., oral, implant), experiences related to motherhood (e.g.,

vaginal delivery, caesarean delivery), return to training (RTT)/return to competition

(RTC) and body perception.

Body perception was measured by using the body dissatisfaction (BD) and drive for

thinness (DT) scores. These were measured with two of the 12 primary scales of the

Eating Disorder Inventory (EDI-3).[11] This validated tool ensures a reliable

interpretation of the psychometric test.[12] BD relates to the disapproval of the overall

shape of one's body and the size of specific body regions of particular concern. DT

consists of perceptual, behavioral, and attitudinal parts and is probably triggered

when there is a discrepancy between actual and ideal body weight that exceeds the

specific idealised preference of cultural thinness and involves body image

dissatisfaction.[11]

The supplementary material presents all the questions in the survey. Annexure A.

Descriptive variables

Participant characteristics and several descriptive variables (e.g., age, height, body

mass index (BMI), player position, and stress fractures) were collected using an

electronic questionnaire.

Procedures and data storage

A baseline and a follow-up electronic questionnaire were set up in English and

French (CastorEDC, CIWIT B.V, Amsterdam, the Netherlands), including all study

outcome measures. In addition, the following descriptive variables were added to the

baseline questionnaire: level of education, parallel activity (e.g., study, work), level of

football, number of seasons as a professional footballer, history of hospitalisation,

history of eating disorders, and smoking status. Information about the study was sent

per email to potential participants by FIFPRO and affiliated national unions, and

procedures were hidden from the principal researcher for privacy reasons. If

interested in the study, all participants gave informed consent and received access to

the baseline questionnaire. The follow-up questionnaire was sent six months later.

Each questionnaire took about 15 minutes to complete. The responses to the

questionnaires were coded and made anonymous for reasons of privacy and

confidentiality. The electronic questionnaires were saved automatically on a secured

electronic server that only the principal researcher could access. Players participated

voluntarily in the study and were not rewarded for their participation.

Statistical analyses

The Statistical software, R (version 4.2.1; http://www.r-project.org) was used to

perform all data analyses. Descriptive analyses (mean, standard deviation (SD),

frequency and proportions) were performed for all variables included in the study,

including Body Mass Index (BMI) calculated as a ratio of weight/height squared.[13]

RESULTS

Participant characteristics

Seventy-four women athletes were recruited, with a mean age of 25 years and an

average BMI of 22. Table 1 depicts demographics, football characteristics, career

level and employment status. The majority of our participants were defenders (34%),

followed by forwards (30%), midfielders (23%) and goalkeepers (14%). Most

participants were did not have another paid job (70%), and most (87%) participated

at the highest national level.

Table 1: Demographics, football characteristics, employment status and career level

Demographics		All Participants (n = 74)
Age (yrs) mean (SD)		24.95 (2.69)
Height (cm) mean (SD)		168.39 (5.38)
Weight (kg) mean (SD)		62.81 (5.4)
BMI (kg/m^2) mean (SD)		22.14 (1.47)
Stress fractures n (%)		11 (15)
Low bone density n (%)		1 (1.35)
Other characteristics		
Field Postion n (%)	defender	25 (34%)
	forward	22 (30%)
	midfield	17 (23%)
	goalkeeper	10 (13%)
Number of seasons as a professional player Mean (SD)		5.74 (3.12)
Employed	yes	22 (30%)
Level of Football n (%)	highest national level	64 (87%)
	second highest national level	6 (8 %)
	other	4 (5%)

Yrs = years; SD = standard deviation; cm = centimetres; kg = kilograms; BMI = Body Mass Index

Gynaecological health patterns and body perception

Table 2 depicts gynaecological health patterns. The mean age at menarche was 13.5

years, with a menstrual cycle of 26 days and duration of menstrual bleeding of 5

days. Most participants reported pain during the bleeding period. Half of the

participants used contraception.

Table 2: Gynaecological health patterns, BD and DT

Variable	Characteristics	All Participants (n = 74)
Age of Menstruation (yrs) Mean (SD)		13.47 (1.3)
Duration of menstruation cycle (days Mean (SD)		25.81 (13.09)
Duration of menstrual bleeding period (days) Mean (SD)		5.16 (3.42)
Description of menstruation cycle n (%)	regular	52 (70%)
	irregular	15 (20%)
	very irregular	7 (10%)
Pain during menstruation cycle n (%)	during bleeding period	40 (54%)
	none	19 (26%)
	before bleeding period	15 (20%)
Contraception Methods n (%)	none	37 (50%)
	pill	14 (38%)
	condom	12 (32%)
	other	8 (22%)
	implant	7 (20%)
	hormone replacement	4 (10%)
	injection	1 (2%)
Body dissatisfaction Mean (SD)		4.88 (5.08)
Drive for thinness Mean (SD)		3.59 (5.04)

SD = standard deviation; BD = Body dissatisfaction; DT = Drive for thinness

Motherhood experiences and RTT/RTC

Only one (1.4%) of the 74 participants had two children, both conceived naturally and

delivered vaginally. Her RTT was 6 weeks, and RTC 12 weeks on both occasions.

DISCUSSION

The findings from this study provide a baseline of active female professional football

players' gynaecological health behaviours, whilst most studies reported on retired

elite athletes,[14] or elite athletes from other sports.[15-16] Our findings were that:1)

hormonal contraceptives, particularly the oral contraceptive pill, were the most

commonly used, 2) cycle irregularities were common, 3) most participants

experienced cycle-related pain, and 4) motherhood experiences were rare, at just

1%.

Menstrual cycle

Mernache is the average age at which menstrual bleeding starts in young females,

and usually, this is between the ages of 12-13 years. It has been reported that the

onset of menarche is usually delayed in athletes participating in lean sports, but no

variations were noted between individual and team sports.[4] Our participants had a

mean menarche age of 13.5 years, comparable to other studies on elite athletes.[14]

Our results showed that 30% of our participants had cycle irregularities. Our findings

indicate a higher prevalence of cycle irregularities compared to Swiss elite athletes

from various sports, where approximately 15% reported such issues.[4] However, our

results are lower than those for elite athletes in Denmark[17] and the United Kingdom,[15]

where 51% of athletes reported cycle irregularities. Retired semi- and professional

football players reported a 40% prevalence of menstrual irregularities, with 22%

experiencing amenorrhoea lasting longer than 3 months.[14]

Menstrual pain was reported by 74% of our participants. Symptoms were present

before the bleeding period in 20% and during the menstrual phase in 54% of

participants. This is comparable to the findings in a UK survey where 77% of elite

athletes reported menstrual pain.[15] Elite rugby players reported menstrual pain in

93% of participants,[16] and 84% of elite Denmark athletes reported negative

symptoms related to their menstrual cycle.[17]

A study among elite rugby and football players on premenstrual pain and coping

strategies suggested a model including avoiding harm, awareness and acceptance,

adjusting energy, self-care, and communication to help maintain well-being and

physical activity during this period.[18] Our study was not designed to investigate

coping strategies, and this will require further investigation.

Several studies have shown that athletes believe that their performance declines

when they experience menstrual-related symptoms.[16-17,19-20] A survey of sub-elite

female football players showed a reduction in endurance performance during the

menstrual phase,[21] while elite Australian football players reported higher perceived

fatigue during this period.[22] Our study did not investigate the impact of the menstrual

cycle on exercise performance, highlighting the need for future research to address

this gap in the literature on female professional football players.

Contraceptive use

Our findings confirm that 50% of our participants were on contraceptives, mainly HCs

(60%), predominantly the OCP (38%), followed by implants (20%) and injectable

contraception (2%). These findings are consistent with findings amongst other elite

athletes, with 50% on HCs, predominantly the OCP (68.%), followed by the implants

(13%), injection (4%) and intra-uterine devices (3%).[15] In elite Australian football

codes, HCs use was 33%, with 81% on the OCP.[3] A survey on Swiss players

reported that 46% were on contraception, and 26% used the OCP,[4] and a study on

Danish players reported 57% use of HCs and all of them on OCP.[17] Amongst retired

professional football players from the United States (US), the trend was similar,

although more participants (64.5%) had used HCs, predominantly the OCP

(94.5%).[14] Some of our participants reported the use of more than one contraceptive

method. Our study was not designed to investigate reasons for the use of different

methods and what 'other methods' were used. Further research is warranted on this

topic.

BD and DT

Our participants had a normal BD score of 5. This finding aligns with findings in

federated Spanish amateur football players, who also reported high positive body

images and a normal BD score of 5.[23] We are not aware of other similar studies in

female professional football players. A systematic review and meta analysis across

different sports and genders highlighted the importance of sports participation in

fostering positive body image. The study found that female athletes had higher BD

scores than males, normal-weight athletes had higher BD scores than underweight

ones, and compared to non-athletes, female athletes tend to have a higher body

image perception.[24]

Our participants had a high mean DT of 4, which was not expected in football. Higher

DT scores are common in aesthetic and weight category sports.[25] We are not aware

of other comparable studies in female professional football players. High DT has

been correlated to susceptibility to musculoskeletal injuries due to low energy

availability in the National Collegiate Athletic Association (NCAA) Division II female

athletes and higher time loss from training.[26] In this NCAA survey, 82% of

participants sustained musculoskeletal injuries in-season. The mean number of

injuries/training days loss was higher in the high DT group (over 2.0 ± 0.3), with 9.8

(2.6) days loss from training, whilst the low DT group (under 2.0 ± 0.3) had 6.9 (2.6)

days loss.[26]

We found that 70% of our participants did not have another paid job, 58% were not

studying, and 13 (17.6%) smoked. These factors, combined with a high DT, warrant

further monitoring of these athletes to promote well-being. As confirmed in one study,

high DT is a risk factor for the development of ED, and personal risk factors for DT

include perfectionism, low self-esteem in sport, unemployment, and not studying[27]

and has been linked to tendencies towards smoking.[28] Although ED was found to be

higher in non-athletes compared to elite female football players,[29] the combination of

risk factors for ED and susceptibility to musculoskeletal injuries warrant further

monitoring and research.

Motherhood experiences

In our study, only one participant (1%) reported having had two pregnancies and she

conceived naturally. Both children were delivered vaginally, and she returned to

training and competition 6 and 12 weeks post-partum, respectively. The US study

reported that 50% of the participants reported trying to fall pregnant, with a 94%

success rate, mostly by intra-uterine insemination (51%), medication (49%) and in-

vitro fertilisation (43%).[14] Our sample is too small to compare. Still, we could assume

that pregnancy during a professional football career remains challenging as players

might be afraid to potentially jeopardise their career due to motherhood.

It has been shown that some professional athletes opt to choose between being

mothers and continuing to participate in sports,[8] but similar to our study, other

athletes have successfully fallen pregnant and managed to RTT/RTC earlier than

non-athletes.[30] Although early RTT has been associated with an increased risk for

stress fractures, high BD and high DT,[30] this has not been confirmed in our study.

FIFPRO reported in 2017 that only 2% of female professional football players were

mothers, and almost 45% expressed that they would have to leave the game early to

start a family.[31] Our study's low percentage of motherhood is comparable to the

FIFPRO findings. The choice between motherhood and professional football was not

canvassed in our study, but it will form a basis for future research.

Strengths and Limitations

The strength of our study lies in its pioneering focus on the gynaecological health

patterns and motherhood experiences of active female professional football players.

Although our study is descriptive, our research provides a valuable cohort for future

prospective and epidemiological studies. We hope this study will inspire further

research in the realm of female professional football players.

Our study has several limitations, including the small sample size and the lack of

investigations into participants' perceptions of exercise performance during the

menstrual cycle, their coping strategies for menstrual symptoms, reasons for

contraceptive choices, and their views on balancing motherhood with professional

sport. We did not investigate any correlation between injuries and the menstrual

cycle. Our findings on motherhood experiences are also limited by the low number of

participants. Further research is needed to address these gaps.

Implications

Women's football is growing globally, and factors unique to women may affect their performance and/or shorten their sporting career. This study highlights the challenges faced by these athletes, including menstrual cycle-related symptoms that may affect performance and sporting career, and the importance of specialised gynaecological care for female elite athletes. Healthcare practitioners, coaches and team managers working with elite female athletes should be trained to improve communication, guide athletes on general trends in gynaecological health patterns, optimise well-being, and advise on safe RTT and RTC after motherhood.

More research, such as athlete perception of menstrual cycle effects on performance, coping strategies when experiencing menstrual pain, choice and reasons for use of contraceptives, and choice of sports versus motherhood, is needed in female professional football players.

CONCLUSION

Cycle irregularities are common in female professional football players, with a significant number of cycle-related symptoms. The majority of those using contraceptives preferred hormonal contraceptives, predominantly oral contraceptive pills followed by implants, reflecting trends similar to other female elite athletes. While body satisfaction scores were normal, there was an unexpectantly high DT score, similar to that observed in lean or weight-category sports. The rate of motherhood was low, consistent with previous findings in professional football players.

PATIENT CONSENT FOR PUBLICATION

Not required.

ACKNOWLEDGEMENTS

The authors would like to thank Football Players Worldwide (FIFPRO) and affiliated

national unions for the support in recruiting professional footballers. The authors are

grateful to all participants involved in the study.

CONTRIBUTORS

All authors were involved in the design of the study. VG and GK were responsible for

data collection. All authors were involved in the data analysis and data interpretation.

DR drafted the manuscript, with critical review provided by all authors. All authors

approved the final version of the manuscript.

FUNDING

The study received seed funding from the Drake Foundation located in London (UK)

and financial support from Mehilainen NEO Hospital located in Turku (Finland), from

Sports Hospital Mehilainen located in Helsinki (Finland) and from Nea International

bv. located in Maastricht (the Netherlands).

COMPETING INTERESTS

None declared.

PATIENT AND PUBLIC PARTNERSHIPS

There are no patient and public partnerships to declare.

ETHICAL APPROVAL

The Medical Ethics Review Committee of the Amsterdam University Medical Centers

(Amsterdam UMC, location AMC) provided ethical approval for the study (Drake

Football Study: NL69852.018.19 | W19_171#B202169). The study was conducted in

accordance with the Declaration of Helsinki (2013).

REFERENCES

1. Ravi S, Valtonen M, Ihalainen JK, Holopainen E, Kosola S, Heinonen S, et al. Eating behaviours, menstrual history and the athletic career: A retrospective survey from adolescence to adulthood in female endurance athletes. BMJ Open Sport Exerc Med. 2023; 9(1):e001489. doi:10.1136/bmjsem-2022-001489

2. Apter D, Vihko R. Early menarche, a risk factor for breast cancer, indicates early onset of ovulatory cycles. J Clin Endocrinol Metab. 1983; 57(1):82-6. doi:10.1210/jcem-57-1-82

3. Clarke AC, Bruinvels G, Julian R, Inge P, Pedlar CR, Govus AD. Hormonal contraceptive use in football codes in australia. Front Sports Act Living. 2021; 3:634866. doi:10.3389/fspor.2021.634866

4. Baumgartner S, Bitterlich N, Geboltsberger S, Neuenschwander M, Matter S, Stute P. Contraception, female cycle disorders and injuries in swiss female elite athletes-a cross sectional study. Front Physiol. 2023; 14:1232656. doi:10.3389/fphys.2023.1232656

5. Davenport MH, Nesdoly A, Ray L, Thornton JS, Khurana R, McHugh TF. Pushing for change: A qualitative study of the experiences of elite athletes during pregnancy. Br J Sports Med. 2022; 56(8):452-7. doi:10.1136/bjsports-2021-104755

6. Davenport MH, Ray L, Nesdoly A, Thornton J, Khurana R, McHugh TF. We're not superhuman, we're human: A qualitative description of elite athletes' experiences of return to sport after childbirth. Sports Med. 2022:1-11. doi:10.1007/s40279-022-01730-y

7. Physical activity and exercise during pregnancy and the postpartum period: Acog committee opinion, number 804. Obstet Gynecol. 2020; 135(4):e178-e88. doi:10.1097/aog.0000000000003772

8. Culvin A, Bowes A. The incompatibility of motherhood and professional women's football in england. Front Sports Act Living. 2021; 3:730151. doi:10.3389/fspor.2021.730151

9. von Elm E, Altman DG, Egger M, Pocock SJ, Gøtzsche PC, Vandenbroucke JP, et al. The strengthening the reporting of observational studies in epidemiology (strobe) statement: Guidelines for reporting observational studies. PLoS medicine. 2007; 4(10):e296-e. doi:10.1371/journal.pmed.0040296

10. Woodward M. Epidemiology: Study design and data analysis, third edition. Woodward M, editor: Chapman & Hall; 2014.

11. Garner DM, Olmstead MP, Polivy J. Development and validation of a multidimensional eating disorder inventory for anorexia nervosa and bulimia. International Journal of Eating Disorders. 1983; 2(2):15-34. doi:https://doi.org/10.1002/1098-108X(198321)2:2<15::AID-EAT2260020203>3.0.CO;2-6

12. Punzi C, Tieri P, Girelli L, Petti M. Network-based validation of the psychometric questionnaire edi-3 for the assessment of eating disorders. Sci Rep. 2023; 13(1):1578. doi:10.1038/s41598-023-28743-5

13. Keys A, Fidanza F, Karvonen MJ, Kimura N, Taylor HL. Indices of relative weight and
obesity. J Chronic Dis. 1972; 25(6):329-43. doi:10.1016/0021-9681(72)90027-6

14. Ling DI, Hannafin JA, Prather H, Skolnik H, Chiaia TA, de Mille P, et al. The women's
soccer health study: From head to toe. Sports Med. 2023; 53(10):2001-10.
doi:10.1007/s40279-023-01860-x

15. Martin D, Sale C, Cooper SB, Elliott-Sale KJ. Period prevalence and perceived side
effects of hormonal contraceptive use and the menstrual cycle in elite athletes. Int J Sports
Physiol Perform. 2018; 13(7):926-32. doi:10.1123/ijspp.2017-0330

16. Findlay RJ, Macrae EHR, Whyte IY, Easton C, Forrest Née Whyte LJ. How the menstrual
cycle and menstruation affect sporting performance: Experiences and perceptions of elite
female rugby players. Br J Sports Med. 2020; 54(18):1108-13. doi:10.1136/bjsports-2019-
101486

17. Oxfeldt M, Dalgaard LB, Jørgensen AA, Hansen M. Hormonal contraceptive use,
menstrual dysfunctions, and self-reported side effects in elite athletes in denmark. Int J Sports
Physiol Perform. 2020; 15(10):1377-84. doi:10.1123/ijspp.2019-0636

18. Modena R, Bisagno E, Schena F, Carazzato S, Vitali F. How do elite female athletes cope
with symptoms of their premenstrual period? A study on rugby union and football players'
perceived physical ability and well-being. Int J Environ Res Public Health. 2022; 19(18)
doi:10.3390/ijerph191811168

19. Armour M, Parry KA, Steel K, Smith CA. Australian female athlete perceptions of the
challenges associated with training and competing when menstrual symptoms are present.
International Journal of Sports Science & Coaching. 2020; 15(3):316-23.

20. Randell RK, Clifford T, Drust B, Moss SL, Unnithan VB, De Ste Croix MBA, et al.
Physiological characteristics of female soccer players and health and performance
considerations: A narrative review. Sports Med. 2021; 51(7):1377-99. doi:10.1007/s40279-
021-01458-1

21. Julian R, Hecksteden A, Fullagar HH, Meyer T. The effects of menstrual cycle phase on
physical performance in female soccer players. PLoS One. 2017; 12(3):e0173951.
doi:10.1371/journal.pone.0173951

22. Carmichael MA, Thomson RL, Moran LJ, Dunstan JR, Nelson MJ, Mathai ML, et al. A
pilot study on the impact of menstrual cycle phase on elite australian football athletes. Int J
Environ Res Public Health. 2021; 18(18) doi:10.3390/ijerph18189591

23. Godoy-Izquierdo D, Díaz I. Inhabiting the body(ies) in female soccer players: The
protective role of positive body image. Front Psychol. 2021; 12:718836.
doi:10.3389/fpsyg.2021.718836

24. Zaccagni L, Gualdi-Russo E. The impact of sports involvement on body image perception
and ideals: A systematic review and meta-analysis. Int J Environ Res Public Health. 2023;
20(6) doi:10.3390/ijerph20065228

25. Joy E, Kussman A, Nattiv A. 2016 update on eating disorders in athletes: A
comprehensive narrative review with a focus on clinical assessment and management. British
Journal of Sports Medicine. 2016; 50(3):154-62. doi:10.1136/bjsports-2015-095735

26. Scheid JL, Stefanik ME. Drive for thinness predicts musculoskeletal injuries in division ii
ncaa female athletes. J Funct Morphol Kinesiol. 2019; 4(3) doi:10.3390/jfmk4030052

27. Maurin J, Labossière S, Pomerleau-Fontaine L, Boudreault V, Brassard S, Dion J, et al.
Personal risk factors and types of sport associated with drive for thinness and drive for
muscularity in nextgen athletes. Front Nutr. 2024; 11:1392064.
doi:10.3389/fnut.2024.1392064

28. Copeland AL, Spears CA, Baillie LE, McVay MA. Fear of fatness and drive for thinness
in predicting smoking status in college women. Addict Behav. 2016; 54:1-6.
doi:10.1016/j.addbeh.2015.11.010

29. Abbott W, Brett A, Brownlee TE, Hammond KM, Harper LD, Naughton RJ, et al. The
prevalence of disordered eating in elite male and female soccer players. Eating and weight
disorders : EWD. 2021; 26(2):491-8. doi:10.1007/s40519-020-00872-0
30. Sundgot-Borgen J, Sundgot-Borgen C, Myklebust G, Sølvberg N, Torstveit MK. Elite
athletes get pregnant, have healthy babies and return to sport early postpartum. BMJ Open
Sport Exerc Med. 2019; 5(1):e000652. doi:10.1136/bmjsem-2019-000652
31. Players'Union FW. 2017 fifpro global employment report. Web: 2017.
506

Gynaecological health patterns and motherhood experiences of female professional football players.

Dimakatso Ramagole,[1,2] Dina C Janse van Rensburg,[2] Charlotte Cowie,[3] Ritan Mehta,[3] Gopika Ramkilawon,[4] Babette M. Pluim,[1,2] Gino Kerkhoffs,[1,5,6,7], Vincent Gouttebarge[1,2,5,6,7,8]

Amsterdam UMC location University of Amsterdam, Department of Orthopedic Surgery and Sports Medicine, Meibergdreef 9, Amsterdam, the Netherlands

Section Sports Medicine, Faculty of Health Sciences, University of Pretoria, Pretoria, South Africa

The Football Association (England), National Football Centre, St George's Park, Needwood, DE13 4PD, UK

Department of Statistics, Faculty of Natural and Agricultural Sciences, University of Pretoria

Academic Center for Evidence-based Sports Medicine (ACES), Amsterdam, the Netherlands

Amsterdam Movement Sciences, Aging & Vitality, Musculoskeletal Health, Sports, Amsterdam, The Netherlands

Amsterdam Collaboration on Health & Safety in Sports (ACHSS), IOC Research Center of Excellence, Amsterdam, the Netherlands

Football Players Worldwide (FIFPRO), Hoofddorp, the Netherlands

CORRESPONDING AUTHOR

Vincent Gouttebarge

Amsterdam UMC location University of Amsterdam, Department of Orthopedic Surgery and Sports Medicine, Meibergdreef 9, 1105 AZ Amsterdam, the Netherlands

Email: v.gouttebarge@amsterdamumc.nl

Telephone: +31621547499

DISCLAIMER: The views expressed in the article are that of the authors based on the findings, and not influenced by the funder.

COMPETING INTERESTS: None declared.

FUNDING

The study received seed funding from the Drake Foundation located in London (UK) and financial support from Mehilainen NEO Hospital located in Turku (Finland), from Sports Hospital Mehilainen located in Helsinki (Finland) and from Nea International bv. located in Maastricht (the Netherlands).

WORD COUNT

Abstract: 242
Main Body excluding tables: 2728
Number of tables: 2

STROBE Statement—checklist of items that should be included in reports of observational studies

	Item No	Recommendation	Page No
Title and abstract	1	(*a*) Indicate the study's design with a commonly used term in the title or the abstract	6
		(*b*) Provide in the abstract an informative and balanced summary of what was done and what was found	2-3
Introduction			
Background/rationale	2	Explain the scientific background and rationale for the investigation being reported	4-5
Objectives	3	State specific objectives, including any prespecified hypotheses	5
Methods			
Study design	4	Present key elements of study design early in the paper	6
Setting	5	Describe the setting, locations, and relevant dates, including periods of recruitment, exposure, follow-up, and data collection	6
Participants	6	(*a*) *Cohort study*—Give the eligibility criteria, and the sources and methods of selection of participants. Describe methods of follow-up *Case-control study*—Give the eligibility criteria, and the sources and methods of case ascertainment and control selection. Give the rationale for the choice of cases and controls *Cross-sectional study*—Give the eligibility criteria, and the sources and methods of selection of participants	6
		(*b*) *Cohort study*—For matched studies, give matching criteria and number of exposed and unexposed *Case-control study*—For matched studies, give matching criteria and the number of controls per case	6-7
Variables	7	Clearly define all outcomes, exposures, predictors, potential confounders, and effect modifiers. Give diagnostic criteria, if applicable	7
Data sources/ measurement	8*	For each variable of interest, give sources of data and details of methods of assessment (measurement). Describe comparability of assessment methods if there is more than one group	7
Bias	9	Describe any efforts to address potential sources of bias	N/A
Study size	10	Explain how the study size was arrived at	6-7
Quantitative variables	11	Explain how quantitative variables were handled in the analyses. If applicable, describe which groupings were chosen and why	N/A
Statistical methods	12	(*a*) Describe all statistical methods, including those used to control for confounding	8
		(*b*) Describe any methods used to examine subgroups and interactions	N/A
		(*c*) Explain how missing data were addressed	N/A
		(*d*) *Cohort study*—If applicable, explain how loss to follow-up was addressed *Case-control study*—If applicable, explain how matching of cases and controls was addressed *Cross-sectional study*—If applicable, describe analytical methods taking account of sampling strategy	N/A
		(*e*) Describe any sensitivity analyses	N/A

Results

Participants	13*	(a) Report numbers of individuals at each stage of study—eg numbers potentially eligible, examined for eligibility, confirmed eligible, included in the study, completing follow-up, and analysed	9
		(b) Give reasons for non-participation at each stage	N/A
		(c) Consider use of a flow diagram	N/A
Descriptive data	14*	(a) Give characteristics of study participants (eg demographic, clinical, social) and information on exposures and potential confounders	9
		(b) Indicate number of participants with missing data for each variable of interest	N/A
		(c) *Cohort study*—Summarise follow-up time (eg, average and total amount)	6
Outcome data	15*	*Cohort study*—Report numbers of outcome events or summary measures over time	10
		Case-control study—Report numbers in each exposure category, or summary measures of exposure	
		Cross-sectional study—Report numbers of outcome events or summary measures	10
Main results	16	(*a*) Give unadjusted estimates and, if applicable, confounder-adjusted estimates and their precision (eg, 95% confidence interval). Make clear which confounders were adjusted for and why they were included	N/A
		(*b*) Report category boundaries when continuous variables were categorized	N/A
		(*c*) If relevant, consider translating estimates of relative risk into absolute risk for a meaningful time period	N/A
Other analyses	17	Report other analyses done—eg analyses of subgroups and interactions, and sensitivity analyses	N/A

Discussion

Key results	18	Summarise key results with reference to study objectives	10-15
Limitations	19	Discuss limitations of the study, taking into account sources of potential bias or imprecision. Discuss both direction and magnitude of any potential bias	15-16
Interpretation	20	Give a cautious overall interpretation of results considering objectives, limitations, multiplicity of analyses, results from similar studies, and other relevant evidence	16
Generalisability	21	Discuss the generalisability (external validity) of the study results	16-17

Other information

Funding	22	Give the source of funding and the role of the funders for the present study and, if applicable, for the original study on which the present article is based	17

*Give information separately for cases and controls in case-control studies and, if applicable, for exposed and unexposed groups in cohort and cross-sectional studies.

Note: An Explanation and Elaboration article discusses each checklist item and gives methodological background and published examples of transparent reporting. The STROBE checklist is best used in conjunction with this article (freely available on the Web sites of PLoS Medicine at http://www.plosmedicine.org/, Annals of Internal Medicine at http://www.annals.org/, and Epidemiology at http://www.epidem.com/). Information on the STROBE Initiative is available at www.strobe-statement.org.

Supplementary material

Tables

Table 1: Demographics, football characteristics, employment status and career level

Demographics		All Participants (n = 74)
Age (yrs) mean (SD)		24.95 (2.69)
Height (cm) mean (SD)		168.39 (5.38)
Weight (kg) mean (SD)		62.81 (5.4)
BMI (kg/m^2) mean (SD)		22.14 (1.47)
Stress fractures n (%)		11 (15)
Low bone density n (%)		1 (1.35)
Other characteristics		
Field Postion n (%)	defender	25 (34%)
	forward	22 (30%)
	midfield	17 (23%)
	goalkeeper	10 (13%)
Number of seasons as a professional player Mean (SD)		5.74 (3.12)
Employed	yes	22 (30%)
Level of Football n (%)	highest national level	64 (87%)
	second highest national level	6 (8 %)
	other	4 (5%)

Yrs = years; SD = standard deviation; cm = centimetres; kg = kilograms; BMI = Body Mass Index

Table 2: Gynaecological health patterns, BD and DT

Variable	Characteristics	All Participants (n = 74)
Age of Menstruation (yrs) Mean (SD)		13.47 (1.3)
Duration of menstruation cycle (days Mean (SD)		25.81 (13.09)
Duration of menstrual bleeding period (days) Mean (SD)		5.16 (3.42)
Description of menstruation cycle n (%)	regular	52 (70%)
	irregular	15 (20%)
	very irregular	7 (10%)
Pain during menstruation cycle n (%)	during bleeding period	40 (54%)
	none	19 (26%)
	before bleeding period	15 (20%)
Contraception Methods n (%)	none	37 (50%)
	pill	14 (38%)
	condom	12 (32%)
	other	8 (22%)
	implant	7 (20%)
	hormone replacement	4 (10%)
	injection	1 (2%)
Body dissatisfaction Mean (SD)		4.88 (5.08)
Drive for thinness Mean (SD)		3.59 (5.04)

SD = standard deviation; BD = Body dissatisfaction; DT = Drive for thinness

What is your age?
[Number]
How tall are you (cm)?
[Number]
How much do you weigh (kg)?
[Number]
What is your percentage of body fat (%)?
[Number]
What is the highest level of education you have completed?
[No school completed; Nursery/Elementary school; High school; Vocational/technical school; College, university or equivalent]
Apart from your professional football career, are you currently studying?
[Yes; No]
If yes, how many hours per week on average do you study?
Apart from your professional football career, do you have currently another paid job?
[Yes; No]
If yes, what other kind of (paid) work do you do?
If yes, how many hours per week on average do you work (apart from your professional football career)?
What is your position on the field?
[Goalkeeper; Defender; Midfield; Forward]
At what level do you play?
[Highest national level/league; Second highest national level/league; Other]
For how many seasons are you a professional footballer?
[Number]
In the past few years, were you admitted to an hospital?
If yes, why?
Have you been diagnosed with a mental health disorder (e.g., depression, anxiety, insomnia)?
[Yes; No; I don't know]
If yes, which mental health disorder(s)?
How many cigarettes (approx.) have you smoked in your entire life?
[Number]
How many days have you smoked cigarettes during the past 30 days?
[Number]

Have you been pregnant?
[Yes; No]
IF YES How many times have you been pregnant?
[Number]
IF YES How did you get pregnant?
[Intercourse; Insemination with partner's sperm; Insemination with sperm donation; I rather not tell]
How many miscarriages did you have?
[Number]
Body dissatisfaction (validated EDI-3 subscale)
I think that my stomach is too big.
I think that my thighs are too large.
I think that my stomach is just the right size.
I feel satisfied with the shape of my body.
I like the shape of my buttocks.
I think that my hips are too big.
I think that my thighs are just the right size.
I think that my buttocks are too large.
I think that my hips are just the right size.
[Always; Usually; Often; Sometimes; Rarely; Never]
Drive for thinness (validated EDI-3 subscale)
I eat sweets and carbohydrates without feeling nervous.
I think about dieting.
I feel extremely guilty after overeating.
I am terrified of gaining weight.

I exaggerate or magnify the importance of weight.
I am preoccupied with the desire to be thinner.
If I gain a pound, I worry that I will keep gaining.
[Always; Usually; Often; Sometimes: Rarely; Never]
At about what age did you begin to menstruate?
[Age]
What is the average duration of your menstrual cycle (i.e., average number of days between the first day of your menstrual cycles)?
[Number of days]
What is the average duration of your menstrual bleeding period?
[Number of days]
How would you describe your menstrual cycle?
[Regular; Irregular, Very irregular]
Since your first menstrual cycle, did your menstrual bleeding period stop for three or more consecutive months (for other reasons than hormonal contraception)?
[No; Yes]
Do you generally experience pain during your menstrual cycle?
[No; Yes, before a bleeding period; Yes, during a bleeding period; Yes, after a bleeding period]
Are you currently using contraceptives?
[No; Yes, cap; Yes, pill; Yes, condom; Yes, contraceptive implant; Yes, contraceptive injection; Yes, contraceptive patch; Yes, diaphragm; Yes, female condom; Yes, namely…]
Are you currently using hormone replacement?
[No; Yes]
Are you a mother?
[No; Yes]
If YES How many children do you have?
[Number]
If YES How did you give birth?
[Vaginal delivery; Caesarean delivery; Adoption]
If DELIVERY How long did it take before you returned to training?
Child 1 Child 2 Child X [Number of weeks]
If DELIVERY How long did it take before you returned to competitive matches?
[Number of weeks]
Have you ever had a stress fracture?
[No; Yes]
If YES please indicate when it occurred and what bone(s).
Have you even been told you have low bone density (e.g., osteopenia or osteoporosis)?
[No; Yes]

How many severe injuries in your LEFT hip have you had so far as professional footballer?
[Number]
How many severe injuries in your RIGHT hip have you had so far as professional footballer?
[Number]
How many severe injuries in your LEFT knee have you had so far as professional footballer?
[Number]
How many severe injuries in your RIGHT knee have you had so far as professional footballer?
[Number]
How many severe injuries in your LEFT ankle have you had so far as professional footballer?
[Number]
How many severe injuries in your RIGHT ankle have you had so far as professional footballer?
[Number]