Rapid and robust identification of sepsis using SeptiCyte RAPID in a heterogenous patient population

Robert Balk [1,*], Annette M. Esper [2], Greg S. Martin [2], Russell R. Miller III [3], Bert K. Lopansri [4,5], John P. Burke [4,5], Mitchell Levy [6], Richard E. Rothman [7], Franco R. D'Alessio [7], Venkataramana K. Sidhaye [7], Neil R. Aggarwal [8], Jared A. Greenberg [1], Mark Yoder [1], Gourang Patel [1], Emily Gilbert [9], Jorge P. Parada [9], Majid Afshar [10], Jordan A. Kempker [2], Tom van der Poll [11], Marcus J. Schultz [12,13], Brendon P. Scicluna [14,15], Peter M. C. Klein Klouwenberg [16], Janice Liebler [17,18], Emily Blodget [17,18], Santhi Kumar [17,18], Winnie Mae [19], Krupa Navalkar [20], Thomas D. Yager [20,*], Dayle Sampson [20], James T. Kirk [20], Silvia Cermelli [20], Roy F. Davis [20] and Richard B. Brandon [20,*]

Rush Medical College and Rush University Medical Center, Chicago, IL 60612, USA; jared_greenberg@rush.edu (J.A.G.); mark_a_yoder@rush.edu (M.Y.); gourang_p_patel@rush.edu (G.P.)

Grady Memorial Hospital and Emory University School of Medicine, Atlanta, GA 30322, USA; aesper@emory.edu (A.M.E.); greg.martin@emory.edu (G.S.M.); jkempke@emory.edu (J.A.K.)

FirstHealth of the Carolinas, Pinehurst, NC 28374, USA; rmiller@firsthealt.org

Intermountain Medical Center, Murray, UT 84107, USA; bert.lopansri@imail.org (B.K.L.); john.burke@imail.org (J.P.B.)

School of Medicine, University of Utah, Salt Lake City, UT 84132, USA

Warren Alpert Medical School, Brown University, Providence, RI 02912, USA; mitchell_levy@brown.edu (M.L.)

School of Medicine, Johns Hopkins University, Baltimore, MD 21205, USA; rrothma1@jhmi.edu (R.E.R.); fdaless2@jhmi.edu (F.R.D.); vsidhay1@jhmi.edu (V.K.S.)

Anschutz Medical Campus, University of Colorado, Denver, CO 80045, USA; neil.aggarwal@cuanschuj.edu

Loyola University Medical Center, Maywood, IL 60153, USA; emgilbert@lumc.edu (E.G.); jparada@lumc.edu (J.P.P.)

School of Medicine and Public Health, University of Wisconsin, Madison, WI 53705, USA; mafshar@medicine.wisc.edu

Amsterdam UMC, University of Amsterdam, 1105 AZ Amsterdam, The Netherlands; t.vanderpoll@amsterdamumc.nl (T.v.d.P.)

Division of Cardiothoracic and Vascular Anesthesia and Intensive Care Medicine, Department of Anesthesia, General Intensive Care, and Pain Management, Medical University of Vienna, Vienna, Austria; marcus.j.schultz@gmail.com

Nuffield Department of Medicine, University of Oxford, Oxford, UK

Centre for Molecular Medicine and Biobanking, University of Malta, Msida MSD 2080, Malta; brendon.scicluna@um.edu.mt

Department of Applied Biomedical Science, Faculty of Health Sciences, Mater Dei Hospital, University of Malta Msida MSD 2080 Malta;

Fundashon Mariadal, Kralendijk, Kralendijk, Bonaire, Netherlands Antilles; pkleinklouwenberg@fundashonmariadal.org

Keck Hospital of University of Southern California (USC), Los Angeles, CA 90033, USA; liebler@usc.edu (J.L.); eblodge1@hs.uci.edu (E.B.); santhi.kumar@med.usc.edu (S.K.)

Los Angeles General Medical Center, Los Angeles, CA 90033, USA

Princeton Pharmatech, Princeton, NJ 08540, USA; winnie.mei@princetonpharmatech.com

Immunexpress Inc., Seattle, WA 98109, USA; krupa.n@immunexpress.com (K.N.); dayle.s@immunexpress.com (D.S.); james.k@immunexpress.com (J.T.K.); silvia.c@immunexpress.com (S.C.); roy.d@immunexpress.com (R.F.D.)

*Correspondence: robert_balk@rush.edu (R.B.); thomas.y@immunexpress.com (T.D.Y.); richard.b@immunexpress.com (R.B.B.)

ABSTRACT

Background: SeptiCyte RAPID is a transcriptional host response assay that discriminates between sepsis and non-infectious systemic inflammation (SIRS) with a one-hour turnaround time. The overall performance of this test in a cohort of 419 patients has recently been described [Balk et al., J Clin Med 2024, 13, 1194]. In this study we present results from a detailed stratification analysis in which SeptiCyte RAPID performance was evaluated in the same cohort across patient groups and subgroups encompassing different demographics, comorbidities and disease, sources and types of pathogens, interventional treatments, and clinically defined phenotypes. The aims were to identify variables that might affect the ability of SeptiCyte RAPID to discriminate between sepsis and SIRS, and to determine if any patient subgroups appeared to present a diagnostic challenge for the test. **Methods:** 1) Subgroup analysis, with subgroups

defined by individual demographic or clinical variables, using conventional statistical comparison tests. 2) Principal component analysis and k-means clustering analysis, to investigate phenotypic subgroups defined by unique combinations of demographic and clinical variables. **Results:** No significant differences in SeptiCyte RAPID performance were observed between most groups and subgroups. One notable exception involved an enhanced SeptiCyte RAPID performance for a phenotypic subgroup defined by a combination of clinical variables suggesting a septic shock response. **Conclusions:** We conclude that for this patient cohort SeptiCyte RAPID performance was largely unaffected by key variables associated with heterogeneity in patients suspected of sepsis.

Keywords: sepsis, SIRS, sepsis likelihood, SeptiCyte, host immune response, stratification, phenotype

INTRODUCTION

Accurate and rapid identification of sepsis is often clinically challenging, in part due to non-specific presenting clinical signs (Liu, 2021; Lengquist, 2023), patient heterogeneity (Seymour, 2019; Fohner, 2019; Reddy, 2020) and a lack of timely information (Lopansri et al., 2019). Clinical signs of sepsis are often vague and can include dyspnea, weakness, altered mental status, pain and cough, which are signs commonly associated with other disease conditions including heart failure, stroke and respiratory failure (Liu, 2021; Lengquist, 2023).

To enable better identification and personalized treatment for sepsis, efforts have been made to identify subclasses of patients based on "phenotypes" (Seymour, 2019; Bhavani et al., 2022; Sinha et al., 2023) or "endotypes" (Davenport et al., 2016; Scicluna et al., 2017, Sweeney et al., 2018). For example, Seymour et al. (2019) described four phenotypes differentiated by clinical parameters such as vasopressor requirement, age, chronic illness, renal dysfunction, inflammation, pulmonary dysfunction, shock and liver dysfunction. As another example, Sinha et al. (2023) showed that the "hyperinflammatory" and "hypoinflammatory" phenotypes previously identified in ARDS patients could also be applied to septic shock patients. Categorizing sepsis patients into

clinically relevant phenotypes could potentially lead to personalized treatments (precision medicine) and better outcomes (De Merle, 2021). However, successful application of precision medicine in sepsis assumes that this condition can be identified early and accurately in the first place. Traditional sepsis diagnostics that rely on the isolation and/or identification of causative pathogens, such as blood culture, have been shown to lack sensitivity, timeliness and in general have not taken patient heterogeneity into account (Coburn, 2012). More recent efforts to improve sepsis diagnosis have included identification of host immune response biomarkers (Vincent et al., 2020).

SeptiCyte RAPID is a host immune response test which measures mRNA expression levels of two genes, PLAC8 and PLA2G7, using a small peripheral blood sample (Miller et al., 2018; Balk et al, 2024). The discovery of these biomarkers was achieved via machine learning on a heterogenous patient dataset (McHugh et al., 2015). Results are reported as a "SeptiScore", on a scale of 0-15 and in four "Bands", with increasing likelihood of sepsis associated with higher SeptiScores and Bands.

In the present study, we investigated the performance of SeptiCyte RAPID in a heterogeneous, critically ill adult patient cohort stratified by demographics, comorbidities and diseases, sources and types of infecting pathogen, therapeutic interventions, and clinical phenotypes. The purpose was to determine if the performance of SeptiCyte RAPID for differentiating sepsis from infection-negative systemic inflammatory response syndrome (SIRS) is generalizable across different patient subgroups. The subgroups examined included patients with conditions that 1) present with overlapping clinical signs of sepsis, 2) may predispose to sepsis, and 3) could affect performance of a host immune response assay.

METHODS

Dataset: The dataset for this analysis (N=419 patients) was from the MARS, VENUS and NEPTUNE studies as described in Balk et al. (2024). Stratifications were performed according to sex, age, race/ethnicity, comorbidities and diseases, therapeutic interventions, and clinically defined phenotypes.

SeptiCyte RAPID Performance vs. Comparator: The performance of SeptiCyte RAPID for discrimination of sepsis vs. SIRS was evaluated using Retrospective Physician Diagnosis (RPD) as the comparator. The RPD process is a clinical evaluation by a panel of three expert clinicians not involved in the care of the patients, described in Miller et al. (2018) and Balk et al. (2024). In this study only "forced" RPD was used. That is, if a patient was initially called "indeterminate" by the RPD panelists, the panelists were then forced to make a consensus or unanimous call of sepsis or SIRS.

Statistics: Statistical tests were mainly conducted with R packages or with Medcalc (medcalc.org). Additional details and cross-checks were as follows. p-values for two-group comparisons were calculated with the Wilcoxon-Mann-Whitney test as implemented in the R 'stats' package (R core team, 2013), unless otherwise noted. Some p-value calculations for small N strata were conducted with Student's t-test as implemented in Microsoft Excel, and also with the Mann-Whitney U test as implemented in Medcalc (medcalc.org) and cross-checked with the web applet at www.socscistatistics.com/tests/mannwhitney/default2.aspx. Proportions tests were conducted with the R 'stats' package and cross-checked with Medcalc. One-way ANOVA was conducted with the R 'stats' package. Cohen's kappa was calculated with the web applet at http://vassarstats.net/kappa.html.

Receiver Operating Characteristic (ROC) curve analysis and calculation of area under curve (AUC) values was performed using the pROC library in R (Robin et al., 2011), and cross-checked with JROCFIT [http://www.rad.jhmi.edu/jeng/javarad/roc/JROCFITi.html] and Medcalc. Confidence intervals for AUC were calculated by the Binomial Exact method, the Hanley & McNeil (1982) method, or the method of DeLong et al. (1988) as implemented in Medcalc, or by bootstrapping as implemented with the web applet of Skalsk´a & Freylich (2006) at http://www.freccom.cz/stomo/input.php. The Hanley & McNeil CI values were cross-checked with the web applet at https://riskcalc.org/ci/. AUC comparisons were performed with either DeLong's test (DeLong et al., 1988) or the bootstrap method, both implemented in R. Note that the ROC analysis of small sample sets (AUC and 95% CI) relies on the equivalence relation AUC $= U/(n_1 * n_2)$ where U = the Mann-Whitney U statistic and n_1, n_2 are the sizes of the SIRS and

sepsis groups (see Mason & Graham 2002; Xu et al. 2013; and Supplementary Material, Section 6).

Principal components analysis (PCA) was performed using the FactoMineR package in R [Le et al., 2008]. The following 16 quantitative clinical variables measured within the first 24 hours of ICU admission defined the dimensions (vectors) by which patients were separated in n-dimensional space: temperature (min & max), heart rate (HR, min & max), respiratory rate (RR, min & max), mean arterial pressure (MAP, min & max), glucose (min & max), white blood cells (WBC, min & max), platelets (min), procalcitonin (PCT), lactate and age. Each of these variables had <=33% missing data, and missing values were replaced with mean values before conducting the PCA. SeptiScore and RPD category were not used in the analysis.

The following qualitative or ordinal variables (not used in constructing the PCA dimensions) were also mapped onto the PCA plot: sex, vasopressor use, mechanical ventilation, pathogen type (bacterial, viral, fungal) and source of infection (blood, urine, sputum, or other), SOFA score, SOFA component scores, and qSOFA & component scores. Missing values for the ordinal variables SOFA, SOFA components and qSOFA were replaced with mean values, as for the quantitative variables.

Hierarchical clustering (HC) was then performed upon the PCA according to the method described by Yang et al. (2015), Maugeri et al. (2021), and Koh et al. (2022). This method allowed the PCA-based separation of patients to be further stratified into subgroups, based on combinations of all the variables (quantitative, qualitative, and ordinal) defined above.

k-means clustering: The k-means clustering algorithm, implemented in the stats package in R, was applied to either the sepsis group (N=176; results in the main text), or to the entire cohort (Sepsis+SIRS) (N=419; results in the Supplementary Material). Many of the same variables used for the PCA analysis were also used for the k-means, including: temperature (min & max within 24 hours), heart rate (HR, min & max), respiratory rate (RR, min & max), mean arterial pressure (MAP, min & max), glucose (min & max), white blood cell count (WBC, min & max), platelets (min), procalcitonin (PCT), lactate, SOFA score, individual SOFA component scores, qSOFA

score, age, sex, vasopressor use (Y/N), mechanical ventilation (Y/N), culture/PCR result (+/-) for bacterial, viral or fungal pathogen, source of positive culture/PCR result (urine, blood, sputum, other). To impute missing data for each variable, the overall mean value for that variable was used. Prior to clustering, the continuous variables were transformed to have a mean of zero and a standard deviation of one. SeptiScore and RPD category were not used as input variables.

RESULTS

1. Demographics

Patient stratification was performed according to sex, age, and race/ethnicity. Patient ages ranged from 18 – 90 years and were binarized into subgroups <60 years or ≥ 60 years of age. The performance of SeptiCyte RAPID for discriminating sepsis vs. SIRS never fell below AUC 0.80 for any subgroup. For each demographic comparison, the p-value for discriminating sepsis vs. SIRS was significant at p<0.05. When considered together in a multiple comparison context, the p-values all remained significant (p<0.05/8) after applying a Bonferroni correction. No significant AUC differences were observed between subgroups except for a marginal difference for White vs. Hispanic (p = 0.03) (**Table 1**).

Upon further analysis, a significant White vs. Black difference in SeptiScore performance (p<0.003) was observed for the discrimination of *septic shock* vs. SIRS. For this stratification, the White demographic subgroup gave AUC = 0.83, while the Black subgroup gave AUC = 0.96. See Supplementary Material, Section 1.

Table 1 – SeptiCyte RAPID performance for sepsis / SIRS discrimination for patients stratified by sex, age, or race/ethnicity. Performance quantified by AUC and p-value using forced adjudication. Comparison of AUC values by DeLong's test. AUC confidence intervals (CI) by formula of Hanley & McNeil (1982), as computed by the applet at https://riskcalc.org/ci/.

Condition	N sepsis	N SIRS	AUC	AUC 95% CI	p-value	DeLong's p-value
male (M)	95	137	0.81	0.75 - 0.87	1.4 E-15	M vs. F: 0.52
female (F)	81	106	0.84	0.78 - 0.90	1.6 E-15	
age < 60 years	77	138	0.83	0.77 - 0.89	3.9 E-14	Age (< 60) vs. (≥ 60): 0.82
age ≥ 60 years	99	105	0.82	0.76 - 0.88	5.7 E-16	
Black (B)*	45	70	0.85	0.77 - 0.93	5.9 E-11	B vs. W: 0.30
White (W)*	108	146	0.80	0.74 - 0.86	7.2 E-16	B vs. H: 0.21 B vs. A: 0.62
Hispanic (H)*	10	12	0.93	0.81 - 1.05	0.00014	W vs. H: 0.03
Asian (A)*	10	11	0.89	0.74 - 1.04	0.0013	W vs. A: 0.23 H vs. A: 0.63

* Seven patients were unclassified as to race/ethnicity, so were left out of this analysis.

2. Comorbidities and disease

Patients were stratified by the presence or absence of different comorbidities or diseases, specifically hyperglycemia, impaired immunity, hypertension, cardiovascular disease, kidney disease and obesity. These comorbidities and diseases were chosen for analysis based on available patient numbers, presumed influence on sepsis predisposition, and potential for presenting clinical signs that overlap with those of SIRS or sepsis. The 'impaired immunity' category included patients with organic immune deficiencies, cancer patients on immunosuppressant drugs, and patients that received immunomodulators such as glucocorticoids; some patients fell in more than one sub-category. Patients for whom a particular co-morbidity or disease was not mentioned in the physician notes were assumed to not have the condition. Point-estimate AUCs ranged from 0.81-0.86 with the only exceptions being AUC 0.79 for patients categorized as hypertensive, and AUC 0.75 for patients with diabetic hyperglycemia (**Table 2**). Box and whisker plots corresponding to the entries of Table 2 are shown in **Figure 1 (A-F).** By DeLong's test there were no significant AUC differences in SeptiCyte performance between patients with vs. without the stated condition.

We note several limitations to this analysis, because of small sample sizes. The AUC value of 1.0 for patients with acute cardiovascular disease should be considered imprecise, because of low N. Additionally, while the uncorrected p-value for each individual comparison falls below the conventional cutoff for significance (p~0.05), if we consider as a group the nine comorbidity/disease comparisons with low numbers, then upon applying a Bonferroni correction, the p-values for diabetes and obesity now fall above the adjusted significance cutoff (p~0.05/9) and the p-value for chronic CVD becomes borderline significant.

Table 2 - SeptiCyte RAPID performance stratified by comorbidities and disease. AUC confidence intervals (CI) by formula of Hanley & McNeil (1982), as computed by applet: https://riskcalc.org/ci/. Abbreviations: HG, hyperglycemia; II, impaired immunity; HT, hypertension; CVD, cardiovascular disease; KD, kidney disease. For small sample comparisons, the p-value was calculated using both the t-test (T) and the Mann-Whitney U test (U).

Stratification	N sepsis	N SIRS	AUC	AUC 95% CI	p-value	DeLong's p-value
hyperglycemia (HG)	50	78	0.83	0.76 - 0.92	5.3 E-10	HG vs. no HG noted: 0.76
diabetic hyperglycemia	10	16	0.75	0.51 - 0.93	0.035 (T) 0.038 (U)	
diabetic hyperglycemia not noted	40	62	0.85	0.77 - 0.93	3.7 E-09	
no hyperglycemia noted (no HG)	126	165	0.82	0.76 - 0.86	5.8 E-21	
impaired immunity (II)	27	32	0.83	0.72 - 0.94	6.8 E-06	II vs. No II noted: 0.84
no impaired immunity noted	149	211	0.82	0.77 - 0.87	4.9 E-25	
hypertension (HT)	24	26	0.79	0.66 - 0.92	1.8 E-04	HT vs. no HT noted: 0.59
no hypertension noted	152	217	0.83	0.78 - 0.88	1.03 E-26	
cardiovascular disease (CVD)	16	23	0.85	0.72 - 0.98	1.7 E-04	CVD vs. no CVD noted: 0.63
cardiovascular disease (acute)	7	6	1.0	0.75 – 1.0	0.0031 (T) 0.0034 (U)	
cardiovascular disease (chronic)	9	17	0.81	0.62 - 1.00	0.0046 (T) 0.0105 (U)	
no cardiovascular disease noted	160	220	0.82	0.78 - 0.86	1.9 E-26	
kidney disease (KD)	33	34	0.83	0.73 - 0.93	5.6 E-07	
kidney disease (acute)	13	13	0.86	0.70 -1.00	0.0012 (T) 0.0023 (U)	KD vs. no KD noted: 0.89
kidney disease (chronic)	20	21	0.83	0.70 - 0.96	0.0001 T) 0.0004 (U)	
no kidney disease noted	143	209	0.82	0.77 - 0.87	7.4 E-24	

Stratification	N sepsis	N SIRS	AUC	AUC 95% CI	p-value	DeLong's p-value
obesity (BMI >= 30)	6	14	0.86	0.66 - 1.06	0.0141 (T) 0.0135 (U)	Obesity vs. no obesity noted: 0.63
no obesity noted	170	229	0.82	0.78 - 0.86	1.8 E-28	

Figure 1: Box and whisker plots of SeptiCyte RAPID performance in patients with (A) hyperglycemia, (B) impaired immunity, (C) hypertension, (D) cardiovascular disease, (E) kidney disease, (F) obesity.

E. Kidney disease

F. Obesity

Hyperglycemia - The cohort was stratified based on whether or not patients had hyperglycemia (>=200 mg/dl) (**Figure 1A**). A few patients with hypoglycemia were also noted, but their numbers were too small (14 sepsis, 3 SIRS) for reliable statistical analysis. Hyperglycemia included those patients with and without diabetes (as indicated in the physician notes). For patients without diabetes indicated, data were based on "Glucose.Min" or "Glucose.Max" values recorded over a 24-hour period. The SeptiCyte RAPID performance for patients with diabetic hyperglycemia (AUC 0.73) appeared marginally lower than for all hyperglycemic patients (AUC 0.83) or for non-hyperglycemic patients (AUC 0.82). Note that, although the p-value for sepsis vs. SIRS discrimination in the diabetic hyperglycemia subgroup ($p \approx 0.04$) falls below the conventional cutoff (p~0.05), after a Bonferroni correction this p-value now falls above the adjusted significance cutoff (p~0.05/9).

Impaired immunity - The cohort was stratified based on whether or not patients could be considered to have impaired immunity (**Figure 1B**). The category of impaired immunity included use of immunosuppressants (including corticosteroids), adrenal insufficiency, splenectomy, asplenia, HIV/AIDS and cancer. Of the 13 patients with cancer, no consideration was given to the type of cancer, or whether the patients were being treated with cancer therapy, or the duration of cancer therapy. White blood cell counts (WBC, min) in patients with impaired immunity as defined above ranged from 300 – 37,000 cells / uL, and for patients with no such impaired immunity the range was 300 – 52,000 cells / uL. SeptiCyte RAPID AUCs were 0.83 and 0.82 for patients with impaired immunity vs immunocompetent respectively.

Hypertension - The cohort was stratified based on whether hypertension was noted in the physician comments (**Figure 1C**). The AUC for differentiating sepsis from SIRS was 0.79 for hypertensive patients and 0.83 for those without hypertension noted, a difference that was not significant according to DeLong's test (p=0.59).

Cardiovascular disease (CVD) - The cohort was stratified based on whether patients had been diagnosed with CVD (**Figure 1D**). CVD patients were further subdivided into those with acute (cardiac arrest) vs. chronic disease. The chronic CVD patients included those with congestive heart failure, aortic valve replacement, and bradycardia with cardiogenic shock. Of the patients

with CVD, there were 16 with sepsis, and 23 with SIRS. The remainder of the cohort (without CVD) consisted of 160 patients with sepsis and 220 with SIRS. We calculated AUC 0.85 for patients with CVD as opposed to AUC 0.82 for patients without CVD. This difference was deemed not significant by DeLong's test (p = 0.63).

Kidney disease – the cohort was stratified into those with kidney disease (acute or chronic) vs. no kidney disease noted (**Figure 1E**). Kidney disease included patients with acute renal failure/injury vs. patients with chronic conditions (renal insufficiency/ disease, end stage renal disease, renal cell carcinoma, nephrolithiasis with bilateral ureteral stents, or chronic dialysis). AUCs for differentiating sepsis from SIRS were 0.82 (no kidney disease noted) and 0.83 for those with chronic kidney disease, a difference that was not significant.

Obesity - the cohort was stratified into obese (BMI>=30 or obesity noted in physician comments), vs. BMI<30 or no obesity noted (**Figure 1F**). Obesity is a known risk factor for sepsis (Hu et al., 2023; Wang et al., 2023). AUCs for differentiating sepsis from SIRS were 0.86 (obese patients) and 0.82 for non-obese patients; DeLong's test indicates this AUC difference is not significant. Note that, while the p-value for sepsis vs. SIRS discrimination in the obese subgroup (p ≈ 0.014) falls below the conventional cutoff (p~0.05), after a Bonferroni correction this p-value now falls above the adjusted significance cutoff (p~0.05/9).

3. Source and Type of Infection

Infection source – Of the 176 sepsis patients in the cohort, 150 had an identified source of infection, specifically pulmonary (n=59), abdominal (n=30), blood (n=17), central nervous system (CNS) (n=6), urinary tract (UTI) (n= 24) and "other source" (n=14, as detailed in legend of Figure 2). There were 26 sepsis patients for whom an initial source of infection could not be identified (NI). In the cohort, 243 patients were retrospectively determined to have SIRS (deemed non-infectious), however only 215 did not have a source of infection identified. **Figure 2** presents box and whisker plots for subgroups of septic patients vs. the SIRS patients, with AUCs and p values indicated. SIRS patients with an identified source of infection (4 abdominal, 5 CNS, 15 pulmonary, 2 urinary, 2 other) were excluded from the analysis. For all sepsis subgroups with

identified sources of infection, the median SeptiScore was in Band 4 (highest sepsis probability), as defined in the banding scheme of Balk et al. (2024). For septic patients without an identified source of infection, the median SeptiScore was in Band 3. For the subgroups with sufficient numbers to allow for a reliable AUC comparison, DeLong's test showed no significant differences (pulmonary vs. NI, p=0.08; abdominal vs. NI, p=0.21; UTI vs. NI, p=0.35).

Figure 2 –SeptiCyte RAPID performance stratified by infection source. The "other" group included the following (N per group): cellulitis (3), Fournier's gangrene (1), hip arthroplasty (1), osteomyelitis (1), toe infection (1), sacral wound (1), post-surgical sternal wound (1), bladder/prostate abscess/ peritonitis from cecum micro perforation (1), influenza (1), tracheitis (1), erysipelas (1), skin or soft tissue necrotizing fasciitis (1). Significance: p<=0.01**, p<=0.001***.

Infection type – We previously determined that SeptiCyte RAPID performance was not affected by the Gram (+/-) category of the sepsis pathogen (see Section 9 of the Supplement to Balk et al., 2024). In the present study, we extended this analysis to determine if different sepsis pathogens were associated with different sources of infection. Results of this analysis are provided in **Supplementary Material, Section 2.** To summarize, viral pathogens were more often detected in patients with pulmonary sepsis, which may reflect a bias in the types of pathogen identification tests ordered for this subgroup. *Staphylococcus aureus* was the most frequently isolated pathogen (19% of all pathogen detection events) and was mostly associated with pulmonary and blood sources of septic infection. Gram negative pathogens, in particular *Escherichia coli* and *Pseudomonas aeruginosa*, were most isolated in urosepsis.

4. Therapeutic Interventions

We examined the performance of SeptiCyte RAPID after stratification along two treatment dimensions: either (+/-) pharmaceuticals (immunosuppressants; antibiotics, inotropes, vasopressors); or (+/-) mechanical ventilation. Pharmaceutical treatments were selected from a list of >400 drugs noted in the patient records. Further detail on the immunosuppressants, antibiotics, anti-neoplastics, inotropes and vasopressors listed in patient records is provided in the **Supplementary Material, Section 3.**

Results in **Table 3** and **Figure 3** show that the use of a broad range of immunosuppressants did not affect the overall performance of SeptiCyte RAPID in our cohort. AUC values in patients treated with immunosuppressants (0.80) or not treated (0.82) were not statistically different (p=0.75 by DeLong's test).

Patients treated with antibiotics (Abx) were subdivided into three groups based on when the antibiotic treatment was started relative to SeptiCyte RAPID blood sampling. In total, 190 patients (110 sepsis, 80 SIRS) were given antibiotics up to one day prior, 76 patients (34 sepsis, 42 SIRS) were given antibiotics on the same day, and 148 (29 sepsis, 119 SIRS) received antibiotics on the day following the blood sampling. SeptiCyte RAPID AUCs were not significantly affected by timing of antibiotics over these time periods, with AUCs ranging from 0.79 to 0.84 (**Figure 4**). The purpose of this comparison was to determine if antibiotics interfered with SeptiCyte RAPID performance in a restricted time window from -1 days to +1 day relative to blood draw at day 0.

We limited the analysis to this relatively narrow time window, because a recent study (Peri et al., 2023) indicates that a reduction in SeptiScore is expected over the period 2-6 days following initiation of antibiotics.

Use of vasopressors or inotropes had no evident effect on SeptiCyte RAPID performance (AUC 0.83 versus AUC 0.81 in the absence of vasopressors or inotropes, p= 0.60 by DeLong's test). Similarly, SeptiScores in patients on mechanical ventilation (AUC 0.80) did not differ significantly from patients not on mechanical ventilation (AUC 0.84), p= 0.36 by DeLong's test.

Table 3 - SeptiCyte RAPID performance stratified by therapeutic interventions. Abbreviations: Abx, antibiotics; I, inotropes; IS, immunosuppressants; MV, mechanical ventilation; V, vasopressors. AUC confidence intervals (CI) by formula of Hanley & McNeil (1982), as computed by applet: https://riskcalc.org/ci/.

Therapeutic Intervention	N sepsis	N SIRS	AUC (95% CI)	sepsis vs. SIRS p-value	DeLong's p-value
Immunosuppressants (IS)	29	31	0.80 (0.67-0.91)	1.6 E-05	IS vs. no IS: 0.75
No immunosuppressants	147	212	0.82 (0.77-0.87)	5.8 E-25	
Antibiotics (Abx) 1 day prior to SeptiScore	110	80	0.79 (0.73-0.85)	7.5 E-13	Abx prior vs. on = 0.46 Abx on vs. after = 0.71 Abx after vs. prior = 0.79
Antibiotics same day as SeptiScore	34	42	0.84 (0.75-0.93)	4.9 E-08	
Antibiotics 1 day after SeptiScore	29	119	0.81 (0.71-0.91)	9.4 E-06	
Vasopressors (V) or inotropes (I)	81	76	0.83 (0.77-0.89)	2.3 E-14	V/I vs. no V/I: 0.60
No vasopressors (V) or inotropes (I)	95	167	0.81 (0.75-0.87)	4.8 E-15	
Mechanical ventilation (MV)	62	91	0.80 (0.72-0.88)	3.1 E-10	MV vs. no MV: 0.36
No mechanical ventilation	114	152	0.84 (0.79-0.89)	5.4 E-21	

Figure 3: Box and whisker plot of SeptiCyte RAPID performance for patients treated vs. not treated with immunosuppressants.

Figure 4: Box and whisker plots of the influence of antibiotic treatment (Abx) initiation time on SeptiCyte RAPID performance. The day of blood draw is defined as day 0. (A) treatment initiated -1 days to 0 days relative to blood draw; (B) antibiotic treatment initiated on the same day as blood draw. (C) antibiotic treatment initiated +1 day after blood draw.

5. Phenotypic Subgrouping

We used two unsupervised machine learning methods - Principal Component Analysis / Hierarchical clustering (PCA / HC) and k-means clustering - to conduct stratification analyses. These methods employed combinations of readily available clinical variables as input.

1. Phenotypic subgrouping of the septic patients (N=176) by PCA / HC

We first conducted a PCA, based on 16 pre-selected clinical variables and excluding SeptiScores and RPD determinations. The first two PCA dimensions captured 15.32% and 12.29%, respectively, of the total variation. A hierarchical clustering (HC) analysis was then performed upon the PCA, which gave a separation into three major subgroups (**Figure 5**). There is additionally a single outlier patient, denoted as "subgroup 4"and represented by the blue point. This patient has a combination of elevated values for Glucose (min & max), lactate, WBC (max), SOFA (Liver component score) & lower values for temperature (min). Of note is that blood culture positive patients did not cluster in any particular subgroup (not shown in figure).

Figure 5 – PCA plot of the sepsis group (N=176) with superimposed HC using 16 phenotypic variables. Each variable used in this analysis had <=33% missing data, and mean values were used to impute the missing data. In the plot, the peripheral points in each subgroup were used to define the cluster boundaries for that subgroup. Sepsis subgroup 1 (black) N=110. Sepsis subgroup 2 (red) N=50. Sepsis subgroup 3 (green) N=15. Sepsis subgroup 4 (blue) N=1.

We next evaluated the performance of SeptiCyte RAPID for discriminating the SIRS group (N=243) from each of the three sepsis subgroups defined by PCA / HC. Results are given in **Table 4**. DeLong's test indicates that SeptiCyte RAPID performance is significantly better for subgroup 3 (AUC 0.93) than for subgroups 1 or 2 (AUC 0.81).

Table 4 – SeptiCyte RAPID performance in the three subgroups defined by PCA + HCC analysis

Comparison	N sepsis	N SIRS	AUC	sepsis vs. SIRS p-value	DeLong's p-value
sepsis PCA subgroup 1 vs. SIRS	110	243	0.81	6.9 E-19	1vs2 p=0.95
sepsis PCA subgroup 2 vs. SIRS	50	243	0.81	4.5 E-09	1vs3 p=0.015
sepsis PCA subgroup 3 vs. SIRS	15	243	0.93	2.2 E-06	2vs3 p=0.035

cluster 4 only had 1 patient, so was not included in this table.

To identify the phenotypic variables that cause the three subgroups to differ - and particularly to identify the variables that make subgroup 3 distinct from subgroups 1 and 2 - we conducted an ANOVA. Phenotypic variables with greatest statistical significance (p<0.01) are presented in

Table 5. This analysis revealed that patients in subgroup 3 appeared the most seriously ill as indicated by relatively higher SOFA, PCT, RR (min & max), HR (max), WBC (max), lactate values and vasopressor use, and relatively lower MAP (min), glucose (min), platelets (min) and temperature (min). Of the 15 patients in subgroup 3, 11 were clinically diagnosed with septic shock and 3 with severe sepsis based on sepsis-2 definition with no viral infections. Notably, we find no significant differences between the three sepsis subgroups with respect to site of infection or type of infecting pathogen.

Table 5 - Phenotypic characteristics of patients in the three subgroups from the PCA / HC analysis). p-values calculated by one-way ANOVA for quantitative variables, and by proportions test for categorical variables.

Characteristic	Missing values (%)	Subgroup 1 (N = 110), median (IQR)	Subgroup 2 (N = 50), median (IQR)	Subgroup 3 (N = 15), median (IQR)	p-value
Vital signs					
Temperature (Min)	24 (14%)	36.3 (35.5-36.7)	36.0 (35.3-36.7)	35.4 (33.5-35.7)	<0.001
HeartRate (Max)	0 (0%)	113 (102-128)	128 (114-138)	138 (122-156)	<0.001
Mean Arterial Pressure (Min)	2 (1%)	62 (55-74)	59 (48-65)	50 (45-57)	0.006
Respiratory Rate (Max)	53 (30%)	27 (24-31)	27 (23-31)	37 (33-43)	<0.001
Respiratory Rate (Min)	56 (32%)	20 (14-24)	17 (12-23)	32 (15-36)	<0.001
Clinical parameters					
Glucose (Min)	5 (3%)	123 (98-160)	129 (108-160)	90 (65-115)	0.009
WBC (Max)	7 (4%)	12 (8-15)	21 (18-29)	23 (18-29)	<0.001
WBC (Min)	5 (3%)	9 (6-12)	18 (15-23)	14 (8-23)	<0.001
PCT	27 (15%)	3 (0-17)	5 (2-13)	36 (14-130)	<0.001
Lactate	47 (27%)	2.0 (1.45-3.15)	2.4 (1.60-3.40)	9.3 (4.13-11.88)	<0.001
Platelets (Min)	52 (30%)	145 (102-212)	309 (238-367)	80 (53-221)	<0.001
SOFA	19 (11%)	6 (4-9)	6 (4-8)	11 (7-14)	<0.001
Vasopressor (Y/N)	-	42 Y (38%)	20 Y (40%)	12 Y (80%)	0.008
Infection Site**					1vs2, 1vs3, 2vs3 (proportions test)
Pulmonary		34 (30.9%)	20 (40%)	4 (26.7%)	0.26, 0.73, 0.35
Abdominal		15 (13.6%)	10 (20%)	5 (33.3%)	0.30, 0.05, 0.29
Blood		12 (10.9%)	4 (8.0%)	1 (6.7%)	0.57, 0.23, 0.33
CNS		6 (5.5%)	0 (0%)	0 (0%)	0.08, 0.33, ND

Characteristic	Missing values (%)	Subgroup 1 (N = 110), median (IQR)	Subgroup 2 (N = 50), median (IQR)	Subgroup 3 (N = 15), median (IQR)	p-value
Urinary Tract		13 (11.8%)	8 (16%)	3 (20%)	0.47, 0.37, 0.13
Other		9 (8.2%)	4 (8.0%)	1 (6.7%)	0.97, 0.84, 0.87
Site not identified**		21 (19.1%)	4 (8.0%)	1 (6.7%)	0.07, 0.24, 0.87
Pathogens*					1vs2, 1vs3, 2vs3 (proportions test)
Viral (including coinfections)		23 (21%)	6 (12%)	0 (0%)	0.17, 0.05, 0.16
Escherichia coli		11 (10%)	5 (10%)	2 (13.3%)	1.00, 0.70, 0.72
Staphylococcus aureus		18 (16.4%)	15 (30%)	2 (13.3%)	0.05, 0.76, 0.20
Streptococcus		6 (5.5%)	3 (6%)	2 (13.3%)	0.90, 0.25, 0.36
Enterococcus		4 (3.6%)	3 (6%)	0 (0%)	0.49, 0.46, 0.34

** site of infection was not apparent or identified at time of enrollment or within 24 hours of ICU admission, according to study investigator(s)
Percentages are calculated column wise.

In the sepsis group, the PCA variables contributing greater than average (>6%) phenotypic variability for dimensions 1 and 2 were WBC (min & max), glucose (min & max), platelets (min), Age, Temperature (min) and MAP (min). We analyzed SeptiCyte RAPID performance in different subgroups defined by either these individual driving variables or other biomarkers used for sepsis adjudication (**Figure 6**). Similar SeptiCyte RAPID performance levels (AUC 0.79-0.85) were observed for most subgroups, indicating that SeptiCyte RAPID should still have utility regardless of the values of these driving variables. The highest SeptiScore AUCs (0.85) were found for patient subgroups having high WBC counts (>12E+06/mL) or normal platelet counts (150,000 - 450,000/uL). The lowest SeptiScore AUC (0.71) was found for the subgroup with PCT <0.5 ng/mL. This low PCT subgroup contained 32 sepsis patients which appeared less seriously ill than the high PCT (>0.5 ng/mL) subgroup, as indicated by lower mean SOFA score, and a lower proportion on mechanical ventilation. The low PCT sepsis subgroup also had fewer patients that were bacterial-culture positive (40.6%), and a higher percentage of patients that were viral positive (31.3%), as compared to the high PCT subgroup which had 66% bacterial-culture positive (p<0.01) and only 15% viral positive (p<0.03). The detailed quantitative analysis corresponding to Figure 6 is presented in the **Supplementary Material, Section 5, Table T5.** We note that the individual clinical parameters driving separation in the PCA were not themselves effective at discriminating sepsis vs. SIRS (AUC 0.5-0.6), consistent with the previous analysis of Balk et al. (2024).

Figure 6: Performance for discriminating sepsis vs. SIRS in different phenotypic subgroups defined by individual driving variables in PCA / HC and other biomarkers used for sepsis adjudication. Red: SeptiScore. Black: driving variable. Additional details corresponding to this figure are presented in Supplement Table T5.

2. Phenotypic subgrouping of the septic patients (N=176) by k-means clustering

We conducted another stratification of the septic patients using a different approach (k-means clustering). Results of this analysis are shown in **Figure 7**. The k-means clustering algorithm indicated that 2 subgroups was an optimal number, and produced a very clear separation of the sepsis patients into the two subgroups. We next conducted an ANOVA to identify which of the phenotypic variables in the analysis provided the most significant discrimination between the two subgroups. Results of the ANOVA are presented in **Table 6.** (For vital signs, clinical chemistry measurements, and interventions only those variables giving p<0.01 are presented.) Interestingly, with the sole exception of viral infections being more prominent in subgroup 1, neither the site of infection, or pathogen type, were significant drivers of the separation between the two k-means subgroups. Also, blood culture positive patients did not cluster in either of the two subgroups (not shown in figure). Compared to subgroup 1, the subgroup 2 appeared more seriously ill, being

distinguished by depressed temperature and MAP; elevated WBC, lactate, PCT, and SOFA; and increased administration of vasopressors and mechanical ventilation. Despite these clinical differences between subgroups, SeptiCyte RAPID AUCs for differentiating subgroup 1 sepsis patients (n=96) and subgroup 2 sepsis patients (n=80) from SIRS patients (n=243) were 0.80 and 0.85, respectively.

Finally, we performed a Venn diagram analysis to determine overlap(s) between the subgroups identified by PCA / HC stratification vs. k-means stratification. We determined that all 15 of the patients classified as subgroup 3 in the PCA / HC analysis - patients resembling septic shock or severe sepsis according to the Sepsis-2 criteria - fell within k-means subgroup 2, as indicated in **Figure** 7.

Figure 7 - Sepsis subgroups 1 (red; N=96) and 2 (blue; N=80) identified by k-means clustering. The small, most seriously ill subgroup 3 from the PCA / HC analysis (black; N=15) is contained entirely within k-means group 2. The large colored symbols indicate the centroids of the subgroups.

Table 6 - Phenotypic variables contributing to separation between the two sepsis subgroups in the k-means analysis of Figure 6. p-values were calculated with 1-way ANOVA for quantitative variables, and with Pearson's Chi-squared test or two-sample proportions test for categorical variables. For vital signs, clinical chemistry measurements and interventions, only those variables giving p<0.01 are presented.

Characteristic	Missing values (%)	Subgroup 1 (N = 96), median (IQR)	Subgroup 2 (N = 80), median (IQR)	p-value*
Vital signs				
Temperature (Min)	24 (14%)	36.4 (36.0-36.8)	35.3 (34.4-35.8)	<0.001
Mean Arterial Pressure (Min)	2 (1%)	67 (57-78)	55 (46-61)	<0.001
Respiratory Rate (Max)	53 (30%)	25 (23-31)	29 (25-35)	0.01
Clinical chemistry				
WBC (Max)	7 (4%)	12 (8-18)	18 (14-25)	<0.001
WBC (Min)	5 (3%)	10 (6-14)	14 (9-19)	<0.001
Lactate	47 (27%)	2.10 (1.45-3.00)	3.00 (1.83-5.85)	<0.001
GCS<15 (qSOFA component)	25 (14%)	18 (18.8%)	42 (52.5%)	<0.001
qSOFA>=2	-	22 (23%)	68 (85%)	<0.001
qSOFA<2	-	62 (65%)	8 (10%)	<0.001
SOFA	19 (11%)	4 (2-6)	9 (7-12)	<0.001
PCT_binary (>0.5 ng/ml)	27 (15%)	52 (64%)	65 (96%)	<0.001
Interventions				
Vasopressors used	-	15 (16%)	60 (75%)	<0.001
Mechanical Ventilation	-	18 (19%)	44 (55%)	<0.001
Infection Site*				

Characteristic	Missing values (%)	Subgroup 1 (N = 96), median (IQR)	Subgroup 2 (N = 80), median (IQR)	p-value*
Pulmonary	-	36 (37.5%)	23 (28.8%)	0.22
Abdominal	-	13 (13.5%)	17 (21.2%)	0.18
Blood	-	13 (13.5%)	4 (5%)	0.06
Central nervous system (CNS)	-	4 (4.2%)	2 (2.5%)	0.54
Other	-	5 (5.2%)	9 (11.2%)	0.14
Urinary tract infection (UTI)	-	9 (9.4%)	15 (18.8%)	0.07
Site not confirmed initially	-	16 (16.7%)	10 (12.5%)	0.44
Pathogens*				
Viral (including coinfections)	-	23 (24%)	6 (7.5%)	0.006
Escherichia coli	-	8 (8.3%)	10 (12.5%)	0.36
Staphylococcus aureus	-	20 (20.8%)	15 (18.8%)	0.74
Streptococcus	-	5 (5.2%)	6 (7.5%)	0.53
Enterococcus	-	4 (4.2%)	3 (3.8%)	0.89

* Percentages are calculated column wise.

DISCUSSION

It is well known that patients with sepsis are heterogenous with respect to clinical signs, response to therapy, and outcome. Factors contributing to this heterogeneity include patient demographics, comorbidities and diseases, infecting pathogen, locus of the infection, concurrent therapies, and disease progression and stage (Papathanakos et al., 2023). For maximum clinical utility, any sepsis diagnostic test should retain a high level of performance when confronted with these sources of variability. Accordingly, we investigated the performance of a commercial gene-expression based sepsis test (SeptiCyte RAPID) across a broad range of patient characteristics. We demonstrated generally consistent and robust diagnostic performance of SeptiCyte RAPID in a heterogenous, adult, critically ill patient population suspected of sepsis.

Demographics: We found no statistically significant differences in AUC for SeptiCyte performance in discriminating sepsis vs. SIRS with respect to age, sex or ethnic/racial group, when appropriate number of cases per group were analyzed (Table 1). Upon closer examination, however, we did observe a race/ethnicity -based difference in SeptiScore performance for distinguishing *septic shock* from SIRS (White (septic shock, n=44) AUC 0.83 vs. Black (septic shock, n=16) AUC 0.96; p<0.003 by DeLong's test); see Supplementary Material, Section 1). Higher SeptiScores for septic shock cases in Black as opposed to White patients could relate to genetics or environment, or a mixture of both. However, a number of studies suggest that genetic differences do not influence sepsis mortality or hospital length of stay once socioeconomic factors, such as number of comorbidities and access to healthcare, are taken into account (Guillamet et al., 2021; DiMeglio et al., 2018). Without further data, to include increased patient numbers and details on socioeconomic variables, it is not possible to definitively identify the factors contributing to the observed higher SeptiScores in Black vs. White patients with septic shock.

Comorbidities and diseases: Risk factors for sepsis include certain comorbidities, for example diabetes (Frydrych et al., 2017; Cary et al, 2018; Ahlberg et al, 2023), hypertension (Ahlberg et al., 2023), chronic kidney disease (Doi et al., 2008; Ou et al., 2022), obesity (Hu et al., 2023; Wang et al., 2023), cancer (Gudiol et al., 2021) and cardiovascular disease (Nakada et al., 2019). Because SeptiCyte RAPID is based on measurement of host immune response, comorbidities that

involve the immune system could affect assay performance. AUC values for subgroups based on comorbidity and disease generally clustered around 0.82-0.83. One of the lowest SeptiCyte AUC values observed in this analysis was for patients with hypertension (AUC 0.79), but this was not significantly different (p=0.59) from the AUC 0.83 observed for patients with no hypertension noted. (**Table 2**). Another low SeptiCyte AUC value was observed for patients with diabetic hyperglycemia (AUC 0.73) – a finding which would benefit from additional study in an expanded patient cohort. Despite the small sample sizes in some patient subgroups, these results suggest that certain patients suspected of sepsis need not be excluded from SeptiCyte RAPID testing due to the specific pre-existing comorbidities and diseases we were able to examine in our study cohort.

Type and site of infection: The immune status of leukocytes varies greatly depending upon the body compartment from which they are derived (Cavillon et al., 2024) and type of infection being responded to (Nguyen et al., 2013). For example, in humans an intravenous lipopolysaccharide (LPS) injection suppresses the *ex vivo* peripheral blood mononuclear cell (PMBC) response to LPS but primes that of alveolar macrophages. Therefore, the primary site of infection in sepsis could be expected to influence the PBMC response and hence SeptiCyte RAPID results. To investigate this further, we compared SeptiCyte RAPID performance in patient subgroups stratified by the primary site of infection, including the categories lung, abdomen, central nervous system, urinary tract, blood (bacteremia), other identified site, and 'site not identified' (i.e. sepsis of unknown origin) (Figure 2). We observed no statistically significant differences in SeptiCyte RAPID performance across a broad range of identified infection sites. AUC did appear lower (0.72) for patients with sepsis of unknown origin, which may relate to uncertainty in the retrospective diagnosis of patients in this subgroup (Lopansri et al., 2019; Roger et al., 2023).

Therapeutic interventions: immunosuppressants - We did not observe any significant effect of immunosuppressant therapy on performance of SeptiCyte RAPID (Table 3). However, this comparison covered a broad range of immunosuppressants, so no firm conclusions on the effect of any individual immunosuppressant on SeptiCyte RAPID can be drawn without performing a larger study with sufficient patient numbers. Regarding the glucocorticoid class specifically, it is known that short-term oral prednisone (4 days, 30mg/day) produces a distinct blood gene

expression signature in COPD patients and that neither PLAC8 nor PLA2G7 were differentially expressed when comparing pre-treatment patients to Day 4 of treatment (Takiguchi et al, 2019). Further, PLAC8 and PLA2G7 are not known to be glucocorticoid regulated genes (Chinenov et al., 2014). This suggests that treatment with glucocorticoids does not affect peripheral blood gene expression of the two biomarkers of SeptiCyte RAPID, and therefore is not expected to affect SeptiCyte RAPID results.

Interventional treatment: antibiotics - Although blood cultures are the 'gold standard' for confirming bacteremia, up to fifty percent of patients suspected of sepsis and admitted to the ICU have negative cultures (Gupta et al., 2016) some of which could be caused by prior use of antibiotics (Hirosawa et al., 2023). Standard practice in patients suspected of sepsis is to take blood cultures prior to antibiotic administration to avoid the potential negative impact on growth of organisms in blood culture. In this study we show that use of prophylactic antibiotics within 1 day prior to a patient presenting to ICU does not affect SeptiCyte RAPID performance. Therefore, a clinician need not withhold antibiotics prior to taking a blood sample for SeptiCyte RAPID analysis. However, treatment with antibiotics outside of this timeframe would be expected to affect SeptiScores as patients recover in response to appropriate antibiotic treatment (Peri et al., 2023).

Phenotypic analysis - We conducted phenotypic analyses along the lines described by Seymour et al. (2019) on our sepsis patient cohort (N=176). We employed two different unsupervised clustering methods (PCA/HC, and k-means) using a selection of readily available clinical variables. We identified three sepsis subgroups by PCA / HC, and two sepsis subgroups by k-means clustering. Key driving variables separating the sepsis subgroups included WBC, platelets, MAP, glucose, platelets, age, temperature.

We also conducted the same type of analyses on the entire sepsis+SIRS dataset (N=419). Similarly to what we observed in the first analyses, these same phenotypic variables also resolved three subgroups in the PCA / HC analysis and two subgroups in the k-means analysis (Supplementary Material). Separation between subgroups in either analysis appeared to be driven at least partly by clinical severity of the sepsis response. Interestingly, neither the infection source or type of

infecting pathogen was a significant factor, except that viral infections appeared to be less severe and more highly associated with sepsis of pulmonary origin.

Individual clinical variables that were identified as driving the PCA separation were assessed for their ability to discriminate sepsis from SIRS. Figure 6 compared the performance of these variables versus SeptiCyte RAPID. The results suggest that, in our study at least, patients with sepsis cannot easily be distinguished from those with SIRS using any of the individual clinical variables. In contrast, in subgroups defined by these driving variables, SeptiCyte RAPID differentiated sepsis from SIRS with AUCs ranging from $0.71 - 0.86$. This finding is consistent with a previous demonstration (Balk et al., 2024) showing that combinations of up to 14 clinical variable do not outperform SeptiCyte RAPID. These results suggest that sepsis diagnostic tools that rely on these clinical variables alone (e.g. qSOFA or early warning scores) will be limited in their capacity to differentiate sepsis and SIRS patients.

The clinical variables we have identified as "driving" the separations in this study can be broadly mapped onto Seymour's phenotypes: α, (systolic BP, limited use of vasopressors); β, (SOFA, chronic illness, renal dysfunction, age); γ (RR, WBC, PCT, platelets inflammation and pulmonary dysfunction); and δ (systolic BP, lactate, shock). Further, Sinha et al. (2023) have identified two subgroups in ARDS and septic shock termed "hypoinflammatory" and "hyperinflammatory". Patients with hyperinflammatory septic shock had higher rates of blood culture positivity, increased inflammatory markers and poor outcomes (increased 28-day mortality and lower ICU-free days). In our study, although not restricted to septic shock patients, we identified a sepsis patient subgroup using PCA / HC analysis (subgroup 3) that is more likely to contain a higher proportion of patients with septic shock, as indicated by higher SOFA, RR, lactate and use of vasopressors and lower MAP. With respect to sepsis/SIRS discrimination, with the possible exception of higher SeptiCyte RAPID performance in septic shock patients, we observed similar SeptiCyte RAPID performance across our patient phenotypes and key driving variables. An important practical result is that SeptiCyte RAPID performance appears relatively unaffected by phenotype, or by key clinical variables used to define the phenotypes (Tables 4, 5, 6).

There are several limitations to this study. A first limitation is that the patients in our study cohort consisted of only 419 patients admitted to ICU and tested on the first day of ICU admission. If the study had included patients who were tested at times outside the first day of ICU admission, the relative performance estimates and factor sensitivities might have been different. A second limitation is that the cohort size (N=419) was not large enough to include quantitative analyses of the effects of other comorbidities and diseases. A third limitation is that all the patients were adults ≥18 years of age. We cannot say whether any differences in SeptiCyte RAPID performance occur in adolescents, children, infants or neonates. A fourth limitation is that, with respect to race/ethnicity, the dataset was sufficient only to analyze the main classes of subjects (White, Black, Asian, Hispanic) from North American and European geographic regions, and did not include other minorities or patients from other geographic regions. Future studies will need to be undertaken to establish performance in other racial/ethnic groups within North American and European populations, as well as racial/ethnic groups more generally from other areas of the world. A fifth limitation is that the metadata in our study was not sufficiently comprehensive or granular to allow a stratification according to socioeconomic factors, which are known to correlate with sepsis comorbidities (Ahlberg et al., 2023).

Clinical identification of sepsis, even retrospectively, is characterized by significant diagnostic uncertainty (Rhee et al., 2016; Lopansri et al., 2019). This motivates the development of objective sepsis diagnostic tests (Opal and Wittebole 2020; Pierrakos et al., 2020). In this study we have shown that SeptiCyte RAPID provides consistent discrimination of sepsis vs. SIRS across many subgroups within a heterogeneous adult, ICU patient population suspected of sepsis. Further studies involving larger numbers of patients will be required to confirm the robustness of SeptiCyte RAPID in broader populations.

CONCLUSIONS

We conducted a stratification analysis of SeptiCyte RAPID data from the pivotal studies MARS, VENUS, NEPTUNE (N=419). SeptiCyte RAPID demonstrated consistent performance for discrimination of sepsis vs. SIRS, using AUC as the performance measure, across a heterogenous

adult, critically ill patient population when stratified by key demographic, clinical, microbiological and interventional parameters. This helps to support a clinical utility claim for SeptiCyte RAPID. Stratification analysis with phenotypic variables may be useful for resolving sepsis patients into subgroups based on clinical severity.

Author Contributions:

Conceptualization: K.N., T.D.Y., R.F.D. and R.B.B.;

Data curation: K.N. and T.D.Y.;

Formal analysis: K.N., T.D.Y., D.S., S.C., W.M., R.F.D. and R.B.B.;

Funding acquisition: T.D.Y., R.B.B.;

Investigation: K.N., T.D.Y., D.S., S.C., R.F.D., W.M. and R.B.B.;

J.A.K., T.v.d.P., M.J.S., P.M.C.K.K., J.L., E.B., S.K., J.T.K. and S.C.;

Methodology: K.N., T.D.Y., D.S. and W.M.;

Project administration: K.N., T.D.Y. and R.B.B.;

Resources: T.D.Y. and R.B.B.;

Software: K.N., T.D.Y., W.M. and D.S.;

Supervision: T.D.Y. and R.B.B.;

Validation: K.N., T.D.Y., W.M. and R.B.B.;

Visualization: K.N. and T.D.Y.

Writing—original draft: K.N., T.D.Y. and R.B.B.;

Writing—review & editing: R.B., G.S.M., R.R.M.III, J.P.B., M.L., R.E.R., V.K.S., N.R.A., J.A.G., M.Y., E.G., J.P.P., M.A., J.A.K., T.v.d.P., M.J.S., B.P.S., P.M.C.K.K., E.B., S.K., K.N., T.D.Y., D.S., J.T.K., S.C., R.F.D., W.M. and R.B.B.

All authors have read and agreed to the published version of the manuscript.

Funding: This work was funded by Immunexpress, Inc.

Institutional Review Board Statement: Ethics approval for the MARS trial was given by the medical ethics committee of AMC, Amsterdam (approval # 10-056C, 16 June 2010). Ethics approvals for the VENUS trial were given by the relevant Institutional Review Boards as follows: Intermountain Medical Center/Latter Day Saints Hospital (approval # 1024931, 21 January 2014);

Johns Hopkins Hospital (approval # IRB00087839, 28 January 2016); Rush University Medical Center (approval # 15111104-IRB01, 11 March 2016); Loyola University Medical Center (approval # 208291, 10 March 2016); Northwell Healthcare (approval #16-02-42-03, 1 April 2016). Ethics approvals for the NEPTUNE trial were given by the relevant Institutional Review Boards as follows: Emory University (approval # IRB00115400, 3 December 2019); Grady Memorial Hospital (approval # 00-115400, 14 January 2020); Rush University Medical Center (approval # 19101603-IRB01, 16 January 2020); University of Southern California Medical Center (approval # HS-19-0884-CR001, 10 February 2020). All methods used in this study were carried out in accordance with the relevant guidelines and regulations. Informed Consent Statement: All subjects, or their legally authorized representatives, gave informed consent for participation in this study.

Data Availability Statement: The datasets used and/or analyzed during the current study are available from the corresponding author upon reasonable request.

Acknowledgments: The authors thank the clinical study coordinators of the NEPTUNE study: Joyce D. Brown (Rush University), Liliacna Jara (University of Southern California), Leona Wells and Maya C. Whaley (Grady Memorial Hospital).

Conflicts of Interest: K.N., T.D.Y., D.S., J.T.K., S.C., R.F.D. and R.B.B. declare they are present or past employees or shareholders of Immunexpress, Inc. W.M. declares that she is a present employee of Princeton Pharmatech, Inc. R.R.M. III discloses that, over the time period of relevance to this work he was paid (as a consultant) an honorarium by Immunexpress for conducting a critical review of a Clinical Evaluation Report of SeptiCyte technology, submitted by Immunexpress to the Therapeutic Goods Administration (TGA) of Australia. N.R.A. declares that he has received grants from the NIH and Department of Defense for non-related work, and that payments from these grants are made to his institution, the University of Colorado. No other competing interests are declared.

REFERENCES

Ahlberg CD, Wallam S, Tirba LA, Itumba SN, Gorman L, Galiatsatos P. Linking Sepsis with chronic arterial hypertension, diabetes mellitus, and socioeconomic factors in the United States: A scoping review. J Crit Care. 2023 Oct;77:154324. doi: 10.1016/j.jcrc.2023.154324. Epub 2023 May 7. PMID: 37159971.

Balk R, Esper AM, Martin GS, Miller RR III, Lopansri BK, Burke JP, Levy M, Opal S, Rothman RE, D'Alessio FR, et al. Validation of SeptiCyte RAPID to Discriminate Sepsis from Non-Infectious Systemic Inflammation. *Journal of Clinical Medicine*. 2024; 13(5):1194. https://doi.org/10.3390/jcm13051194

Bhavani SV, Semler M, Qian ET, Verhoef PA, Robichaux C, Churpek MM, Coopersmith CM. Development and validation of novel sepsis subphenotypes using trajectories of vital signs. Intensive Care Med. 2022 Nov;48(11):1582-1592. doi: 10.1007/s00134-022-06890-z. Epub 2022 Sep 24. PMID: 36152041; PMCID: PMC9510534.

Carey IM, et al. Risk of Infection in Type 1 and Type 2 Diabetes Compared With the General Population: A Matched Cohort Study. Diabetes Care. 2018 Mar;41(3):513-521. doi: 10.2337/dc17-2131. Epub 2018 Jan 12. PMID: 29330152

Cavaillon J-M, Chousterman BG, Skirecki T. Compartmentalization of the inflammatory response during bacterial sepsis and severe COVID-19. J Intensiv Med. 2024. https://doi.org/10.1016/j.jointm.2024.01.001.

Chinenov Y, Coppo M, Gupte R, Sacta MA, Rogatsky I. Glucocorticoid receptor coordinates transcription factor-dominated regulatory network in macrophages. BMC Genomics. 2014 Aug 6;15(1):656. doi: 10.1186/1471-2164-15-656. PMID: 25099603; PMCID: PMC4133603.

Clopper, C. J. & Pearson, E. S. (1934). The use of confidence or fiducial limits illustrated in the case of the binomial.*Biometrika*, **26**, 404–413. doi: 10.2307/2331986.

Coburn B, Morris AM, Tomlinson G, Detsky AS. Does This Adult Patient With Suspected Bacteremia Require Blood Cultures? Jama. 2012;308:502–11.

Davenport EE, Burnham KL, Radhakrishnan J, et al. Genomic landscape of the individual host response and outcomes in sepsis: a prospective cohort study. Lancet Respir Medicine. 2016;4(4):259-271. doi:10.1016/s2213-2600(16)00046-1

Deeks JJ, Altman DG. Diagnostic tests 4: likelihood ratios. BMJ. 2004;329:168–9. DOI: 10.1136/bmj.329.7458.168

DeLong ER, DeLong DM, Clarke-Pearson DL. Comparing the areas under two or more correlated receiver operating characteristic curves: a nonparametric approach. Biometrics. 1988 Sep;44(3):837-45. PMID: 3203132.

DeMerle KM, Angus DC, Baillie JK, Brant E, Calfee CS, Carcillo J, et al. Sepsis Subclasses: A Framework for Development and Interpretation. Crit Care Med. 2021. https://doi.org/10.1097/ccm.0000000000004842.

Doi K, Leelahavanichkul A, Hu X, Sidransky KL, Zhou H, Qin Y, Eisner C, Schnermann J, Yuen PS, Star RA. Pre-existing renal disease promotes sepsis-induced acute kidney injury and worsens outcome. Kidney Int. 2008 Oct;74(8):1017-25. doi: 10.1038/ki.2008.346. Epub 2008 Jul 16. PMID: 18633340; PMCID: PMC2614371.

Feng D, Cortese G, Baumgartner R. A comparison of confidence/credible interval methods for the area under the ROC curve for continuous diagnostic tests with small sample size. Stat Methods Med Res. 2017 Dec;26(6):2603-2621. doi: 10.1177/0962280215602040. Epub 2015 Aug 30. PMID: 26323286.

Fohner AE, Greene JD, Lawson BL, Chen JH, Kipnis P, Escobar GJ, Liu VX. Assessing clinical heterogeneity in sepsis through treatment patterns and machine learning. J Am Med Inform Assoc. 2019 Dec 1;26(12):1466-1477. doi: 10.1093/jamia/ocz106.

Frydrych LM, Fattahi F, He K, Ward PA, Delano MJ. Diabetes and Sepsis: Risk, Recurrence, and Ruination. Front Endocrinol (Lausanne). 2017 Oct 30;8:271. doi: 10.3389/fendo.2017.00271. PMID: 29163354; PMCID: PMC5670360.

Gardner IA, Greiner M. Receiver-operating characteristic curves and likelihood ratios: improvements over traditional methods for the evaluation and application of veterinary clinical pathology tests. Vet Clin Pathol. 2006; 35:8-17. doi: 10.1111/j.1939-165x.2006.tb00082.x. PMID: 16511785

Gudiol C, Albasanz-Puig A, Cuervo G, Carratalà J. Understanding and Managing Sepsis in Patients With Cancer in the Era of Antimicrobial Resistance. Front Med. 2021;8:636547.

Gupta S, Sakhuja A, Kumar G, McGrath E, Nanchal RS, Kashani KB: Culture-Negative Severe Sepsis: Nationwide Trends and Outcomes. Chest 2016, 150(6):1251-1259.
Hanczar B, Hua J, Sima C, Weinstein J, Bittner M, Dougherty ER. Small-sample precision of ROC-related estimates. Bioinformatics. 2010 Mar 15;26(6):822-30. doi: 10.1093/bioinformatics/btq037. Epub 2010 Feb 3. PMID: 20130029.

Hanley JA, McNeil BJ. The meaning and use of the area under a receiver operating characteristic (ROC) curve. Radiology. 1982 Apr;143(1):29-36. doi: 10.1148/radiology.143.1.7063747. PMID: 7063747.

Highland HM, Wojcik GL, Graff M, Nishimura KK, Hodonsky CJ, Baldassari AR, et al. Predicted gene expression in ancestrally diverse populations leads to discovery of susceptibility loci for lifestyle and cardiometabolic traits. Am J Hum Genet. 2022;109(4):669-679. doi: 10.1016/j.ajhg.2022.02.013.

Hu J, Gan Q, Zhou D, Xia X, Xiang W, Xiao R, Tang J, Li J. Evaluating the risk of sepsis attributing to obesity: a two-sample Mendelian randomization study. Postgrad Med J. 2023 Nov 20;99(1178):1266-1271. doi: 10.1093/postmj/qgad072. PMID: 37681245.

Ibarz M, Haas LEM, Ceccato A, Artigas A. The critically ill older patient with sepsis: a narrative review. Ann Intensiv Care. 2024;14:6.

Koh K-Y, Ahmad S, Lee J-i, Suh G-H, Lee C-M. Hierarchical Clustering on Principal Components Analysis to Detect Clusters of Highly Pathogenic Avian Influenza Subtype H5N6 Epidemic across South Korean Poultry Farms. Symmetry. 2022; 14(3):598. https://doi.org/10.3390/sym14030598

Lê S, Josse J, Husson F (2008). "FactoMineR: A Package for Multivariate Analysis." *Journal of Statistical Software*, **25**(1), 1–18. doi:10.18637/jss.v025.i01.

Lengquist M, Varadarajan A, Alestam S, Friberg H, Frigyesi A, Mellhammar L. Sepsis mimics among presumed sepsis patients at intensive care admission: a retrospective observational study. Infection. 2024;:1–13.

Liu VX, Bhimarao M, Greene JD, et al. The Presentation, Pace, and Profile of Infection and Sepsis Patients Hospitalized Through the Emergency Department: An Exploratory Analysis. Critical Care Explor. 2021;3(3):e0344. Doi:10.1097/cce.0000000000000344

Lopansri BK, III RRM, Burke JP, Levy M, Opal S, Rothman RE, et al. Physician agreement on the diagnosis of sepsis in the intensive care unit: estimation of concordance and analysis of underlying factors in a multicenter cohort. J Intensive Care. 2019;7:13

Mason SJ, Graham NE (2002). Areas beneath the relative operating characteristics (ROC) and relative operating levels (ROL) curves: Statistical significance and interpretation. Quarterly Journal of the Royal Meteorological Society. 128 (584): 2145–2166. doi:10.1256/003590002320603584. ISSN 1477-870X.

Maugeri, A., Barchitta, M., Basile, G. et al. Applying a hierarchical clustering on principal components approach to identify different patterns of the SARS-CoV-2 epidemic across Italian regions. Sci Rep 11, 7082 (2021). https://doi.org/10.1038/s41598-021-86703-3

Miller RR 3rd, Lopansri BK, Burke JP, Levy M, Opal S, Rothman RE, et al. Validation of a Host Response Assay, SeptiCyte LAB, for Discriminating Sepsis from Systemic Inflammatory Response Syndrome in the ICU. Am J Respir Crit Care Med. 2018 Oct 1;198(7):903-913. doi: 10.1164/rccm.201712-2472OC. Erratum in: Am J Respir Crit Care Med. 2020 Jul 1;202(1):155. PMID: 29624409; PMCID: PMC6835074.

Mogil LS, Andaleon A, Badalamenti A, Dickinson SP, Guo X, Rotter JI, Johnson WC, Im HK, Liu Y, Wheeler HE. Genetic architecture of gene expression traits across diverse populations. PLoS Genet. 2018 Aug 10;14(8):e1007586. doi: 10.1371/journal.pgen.1007586.

Nakada TA, Takahashi W, Nakada E, Shimada T, Russell JA, Walley KR. Genetic Polymorphisms in Sepsis and Cardiovascular Disease: Do Similar Risk Genes Suggest Similar Drug Targets? Chest. 2019 Jun;155(6):1260-1271. doi: 10.1016/j.chest.2019.01.003. Epub 2019 Jan 17. PMID: 30660782.

Nguyen Q-T, Nguyen T-HT, Ju S-A, Lee Y-S, Han SH, Lee S-C, et al. CD137 Expressed on Neutrophils Plays Dual Roles in Antibacterial Responses against Gram-Positive and Gram-Negative Bacterial Infections. Infect Immun. 2013;81:2168–77.

Opal SM, Wittebole X. Biomarkers of Infection and Sepsis. Crit Care Clin. 2020;36:11–22.

Ou SM, Lee KH, Tsai MT, Tseng WC, Chu YC, Tarng DC. Sepsis and the Risks of Long-Term Renal Adverse Outcomes in Patients With Chronic Kidney Disease. Front Med (Lausanne). 2022 Feb 24;9:809292. doi: 10.3389/fmed.2022.809292. PMID: 35280875; PMCID: PMC8908417.

Papathanakos G, Andrianopoulos I, Xenikakis M, Papathanasiou A, Koulenti D, Blot S, et al. Clinical Sepsis Phenotypes in Critically Ill Patients. Microorganisms. 2023;11:2165

Pari B, Gallucci M, Ghigo A, Brizzi MF. Insight on Infections in Diabetic Setting. Biomedicines. 2023 Mar 21;11(3):971. doi: 10.3390/biomedicines11030971. PMID: 36979949; PMCID: PMC10046483.

Peri AM, Rafiei N, O'Callaghan K, Brischetto A, Graves B, Sinclair H, et al. Host response signature trends in persistent bacteraemia and metastatic infection due to Staphylococcus aureus and Gram-negative bacilli: a prospective multicentre observational study. Infect Dis. 2023; ahead-of-print ahead-of-print:1–9.

Pierrakos C, Velissaris D, Bisdorff M, Marshall JC, Vincent J-L. Biomarkers of sepsis: Time for a Reappraisal. 2020. https://doi.org/10.21203/rs.3.rs-15792/v2).

R Core Team (2013). R: A language and environment for statistical computing. R
  Foundation for Statistical Computing, Vienna, Austria. ISBN 3-900051-07-0, URL
  http://www.R-project.org/.

Reddy K, Sinha P, O'Kane CM, Gordon AC, Calfee CS, McAuley DF. Subphenotypes in critical care: translation into clinical practice. Lancet Respir Med. 2020 Jun;8(6):631-643. doi: 10.1016/S2213-2600(20)30124-7.

Rhee C, Kadri SS, Danner RL, Suffredini AF, Massaro AF, Kitch BT, et al. Diagnosing sepsis is subjective and highly variable: a survey of intensivists using case vignettes. Crit Care. 2016;20:89

Robin X, Turck N, Hainard A, Tiberti N, Lisacek F, Sanchez J, Müller M (2011). "pROC: an open-source package for R and S+ to analyze and compare ROC curves." BMC Bioinformatics, 12, 77.

Roger P-M, Keïta-Perse O, Mainardi J-L. Diagnostic uncertainty in infectious diseases: advocacy for a nosological framework. Infect Dis Now. 2023;:104751.

Scicluna BP, Vught LA van, Zwinderman AH, et al. Classification of patients with sepsis according to blood genomic endotype: a prospective cohort study. Lancet Respir Medicine. 2017;5(10):816-826. doi:10.1016/s2213-2600(17)30294-1

Seymour CW, Kennedy JN, Wang S, et al. Derivation, Validation, and Potential Treatment Implications of Novel Clinical Phenotypes for Sepsis. Jama. 2019;321(20):2003-2017. Doi:10.1001/jama.2019.5791)

Singh U, Hernandez KM, Aronow BJ, Wurtele ES. African Americans and European Americans exhibit distinct gene expression patterns across tissues and tumors associated with immunologic functions and environmental exposures. Sci Rep. 2021;11(1):9905. doi: 10.1038/s41598-021-89224-1.

Sinha P, Kerchberger VE, Willmore A, Chambers J, Zhuo H, Abbott J, Jones C, Wickersham N, Wu N, Neyton L, Langelier CR, Mick E, He J, Jauregui A, Churpek MM, Gomez AD, Hendrickson CM, Kangelaris KN, Sarma A, Leligdowicz A, Delucchi KL, Liu KD, Russell JA, Matthay MA, Walley KR, Ware LB, Calfee CS. Identifying molecular phenotypes in sepsis: an analysis of two prospective observational cohorts and secondary analysis of two randomised controlled trials. Lancet Respir Med. 2023 Nov;11(11):965-974. doi: 10.1016/S2213-2600(23)00237-0. Epub 2023 Aug 23. PMID: 37633303; PMCID: PMC10841178.

Sweeney TE, Azad TD, Donato M, Haynes WA, Perumal TM, Henao R, Bermejo-Martin JF, Almansa R, Tamayo E, Howrylak JA, Choi A, Parnell GP, Tang B, Nichols M, Woods CW, Ginsburg GS, Kingsmore SF, Omberg L, Mangravite LM, Wong HR, Tsalik EL, Langley RJ, Khatri P. Unsupervised Analysis of Transcriptomics in Bacterial Sepsis Across Multiple Datasets Reveals Three Robust Clusters. Crit Care Med. 2018 Jun;46(6):915-925. doi: 10.1097/CCM.0000000000003084. PMID: 29537985; PMCID: PMC5953807.

Skalsk´a H, Freylich V (2006) Web-Bootstrap Estimate of Area Under ROC Curve. Austrian J Statistics 35 (2006), 2&3, 325–330.

Takiguchi H, Chen V, Obeidat M, Hollander Z, FitzGerald JM, McManus BM, Ng RT, Sin DD. Effect of short-term oral prednisone therapy on blood gene expression: a randomised controlled clinical trial. Respir Res. 2019 Aug 5;20(1):176. doi: 10.1186/s12931-019-1147-2. PMID: 31382977; PMCID: PMC6683462.

Tiwari S, Pratyush DD, Gahlot A, Singh SK. Sepsis in diabetes: A bad duo. Diabetes Metab Syndr. 2011 Oct-Dec;5(4):222-7. doi: 10.1016/j.dsx.2012.02.026. Epub 2012 Apr 4. PMID: 25572769.

Tosi M, Coloretti I, Meschiari M, Biasi SD, Girardis M, Busani S. The Interplay between Antibiotics and the Host Immune Response in Sepsis: From Basic Mechanisms to Clinical Considerations: A Comprehensive Narrative Review. Antibiotics. 2024;13:406.

Vigeland MD, Flåm ST, Vigeland MD, Espeland A, Zucknick M, Wigemyr M, Bråten LCH, Gjefsen E, Zwart JA, Storheim K, Pedersen LM, Selmer K, Lie BA, Gervin K, The Aim Study Group. Long-Term Use of Amoxicillin Is Associated with Changes in Gene Expression and DNA Methylation in Patients with Low Back Pain and Modic Changes. Antibiotics (Basel). 2023 Jul 21;12(7):1217. doi: 10.3390/antibiotics12071217. PMID: 37508313; PMCID: PMC10376514.

Wacharasint P, Nakada TA, Boyd JH, Russell JA, Walley KR. Normal-range blood lactate concentration in septic shock is prognostic and predictive. Shock. 2012 Jul;38(1):4-10. doi: 10.1097/SHK.0b013e318254d41a. PMID: 22552014.

Wang J, Hu Y, Zeng J, Li Q, He L, Hao W, Song X, Yan S, Lv C. Exploring the Causality Between Body Mass Index and Sepsis: A Two-Sample Mendelian Randomization Study. Int J Public Health. 2023 May 2;68:1605548. doi: 10.3389/ijph.2023.1605548. PMID: 37205044; PMCID: PMC10186272.

Wen J, Xie M, Rowland B, Rosen JD, Sun Q, Chen J, et al. Transcriptome-Wide Association Study of Blood Cell Traits in African Ancestry and Hispanic/Latino Populations. Genes (Basel). 2021;12(7):1049. doi: 10.3390/genes12071049.

Xu, W., Dai, J., Hung, Y. S., & Wang, Q. (2013). Estimating the area under a receiver operating characteristic (ROC) curve: Parametric and nonparametric ways. *Signal Processing, 93(11), 3111–3123*. doi:10.1016/j.sigpro.2013.05.010

Yang C, Yang K, & Zhou B (2015) A hierarchical Clustering Method Based on PCA-Clusters Merging for Quasi-linear SVM. International Conference on Automation, Mechanical Control and Computational Engineering (AMCCE 2015). DOI: 10.2991/amcce-15.2015.407.