Low-cost generation of clinical-grade layperson-friendly pharmacogenetic passports using oligonucleotide arrays

Authors and affiliations

Pauline Lanting[1,2,3*], Robert Warmerdam[1,3*], Jelle Slager[1], Harm Brugge[1], Taichi

Ochi[4], Marloes Benjamins[1], Esteban Lopera-Maya[1], Soesma Jankipersadsing[1], Jody

Gelderloos-Arends[1], Daphne Teuben[1], Dennis Hendriksen[1,6], Bart Charbon[1,6],

Lennart Johansson[1,6], Thijs Oude Munnink[2], Nienke de Boer-Veger[5], Lifelines NEXT,

LifeLines Cohort Study, Bob Wilffert[2,4 †], Morris Swertz[1,6], Daan Touw[2], Patrick

Deelen[1,3], Nine Knoers[1], Jackie Dekens[1,7#], Lude Franke[1,3#]

[*] Contributed equally

[†] Deceased

[#] To whom correspondence should be addressed

1. Department of Genetics, University Medical Center Groningen, University of

   Groningen, Groningen, The Netherlands

2. Department of Clinical Pharmacy and Pharmacology, University Medical

   Center Groningen, University of Groningen, Groningen, The Netherlands

3. Oncode Institute, Utrecht, The Netherlands

4. Groningen Research Institute of Pharmacy, Unit of PharmacoTherapy,

   Epidemiology & Economics, University of Groningen, Groningen, The

   Netherlands

5. Service Apotheek Boterdiep, Groningen, The Netherlands

6. Genomics Coordination Center, Groningen, The Netherlands

7. Center for Development and Innovation, University Medical Center

   Groningen, University of Groningen, Groningen, The Netherlands

28   **Abstract**

Pharmacogenomic (PGx) information is essential for precision medicine, enabling

drug prescriptions to be personalized according to an individual's genetic

background. Almost all individuals will carry a genetic marker that affects their drug

response, so the ideal drug prescription for these individuals will differ from the

population-level guidelines. Currently, PGx information is often not available at first

prescription, reducing its effectiveness. Pharmacogenetic information is most often

obtained using special assays, making it expensive and time-consuming to generate.

We therefore hypothesized that we could also use genome-wide oligonucleotide

genotyping arrays to generate comprehensive PGx information (PGx passports),

thereby decreasing the cost and time required for PGx testing, and lowering the

barrier to generating PGx information prior to first prescription.

Taking advantage of existing genetic data generated in two biobanks, we developed

and validated Asterix, a low-cost clinical-grade PGx passport pipeline for 12 PGx

genes. In these biobanks we performed and clinically validated genetic variant

calling and statistical phasing and imputation. In addition, we developed and

validated a novel *CYP2D6* copy number variant calling tool, foregoing the need to

use separate PCR-based copy number detection. Ultimately, we returned 1227 PGx

passports to biobank participants via a layperson-friendly app, improving knowledge

of PGx among citizens. Our study demonstrates the feasibility of a low-cost clinical-

grade PGx passport pipeline that could be readily implemented in clinical settings to

enhance personalized healthcare, ensuring that patients receive the most effective

and safe drug therapy based on their unique genetic makeup.

Keywords

Pharmacogenetics, pharmacogenomics, PGx profile, PGx passport, clinical

pharmacogenomics

1. Introduction

There is now extensive knowledge available on pharmacogenes, and clinical

guidelines are available for personalized drug treatment using genetic data (1).

However, the application of pharmacogenetics (PGx) in clinical practice remains

limited (2) and is mostly done reactively to understand why patients reacted poorly to

prescribed drugs. However, a pre-emptive pharmacogenetic passport would allow

results to be stored in (electronic) medical files for future medical use (3), ensuring

that PGx data are available at first prescription, thereby enabling optimized treatment

and preventing adverse effects (4,5). Currently, the major impediments to such pre-

emptive testing are cost and speed: PGx information is most commonly obtained

using dedicated assays that target a limited subset of haplotypes of a single gene.

This makes the generation of a complete, multi-gene profile time-consuming and

costly.

In theory, PGx data can also be obtained from genetic data that is routinely used in

both research and clinical genetics. Single nucleotide polymorphism (SNP)

microarrays in particular have proven essential for detection of Copy Number

Variations (CNVs) in clinical genetics and genome-wide association studies (GWAS)

in research (6,7). However, whether this data can be used to provide a clinical-grade

PGx passport to individuals is largely unknown, and there are a number of

challenges that need to be resolved before this data can be effectively used for PGx.

Firstly, SNP arrays do not consider every variant and do not capture on which

haplotype specific information is located, requiring the use of genotype phasing and

imputation. Secondly, *CYP2D6*, an important gene in pharmacogenomics (8) is

highly polymorphic, and it is also affected by CNVs, making the genotyping of this

gene non-trivial. To deliver on the promise of personalized medicine, we need to

bridge these gaps in order to translate available genetic data to clinically relevant

PGx data.

In parallel with the decreasing cost of genetic testing over the past decade, the

number of large-scale biobank initiatives has also increased. Jointly, these biobanks

have now generated genotype data for millions of individuals (9–12). This provides

an opportunity to repurpose readily available genetic data to develop and validate

the use of SNP array data for generating PGx passports. This would allow

researchers to return clinical-grade PGx results to biobank participants, thereby

decreasing the barriers to applying pre-emptive PGx testing and facilitating

personalized medicine in clinical settings.

Another problem is that the disclosure of PGx information so far has mainly been

limited to researchers, clinicians, and pharmacists. Yet sharing relevant PGx

information with citizens is important to improve knowledge and understanding of

PGx testing and its related health outcomes. Digital health technologies, including

mobile phone applications, have increased citizen engagement in the last 10 years

(13,14), and they allow patients to be engaged in the healthcare decision-making

process. Implementation of digital health technology in optimizing medication

adherence in patients with depression, respiratory diseases, and cancers has been

shown to improve patient outcomes. However, the use of digital health technologies

to provide PGx information to patients has been limited. However, with growing

populations throughout the world, governments are now actively stimulating the

development and use of such tools. For instance, the recent Integral Care

Agreement in the Netherlands highlights the importance of digitalization to promote

self-reliant digital healthcare (15).

In the northern Netherlands, the Lifelines Cohort Study, a multigenerational,

longitudinal, dynamic cohort and biobank, presents a unique opportunity to leverage

available genetic data to facilitate healthy aging (16). This biobank has been

genotyping its participants for multiple years, as has the separate Lifelines NEXT

prospective birth cohort biobank. In this study, we developed and clinically validated

1) an automated software tool for the interpretation of array-based genotype data to

generate reliable PGx profiles, 2) a layperson-friendly translation of Dutch

Pharmacogenetics Working Group (DPWG) guidelines, and 3) a digital infrastructure

to communicate these to biobank participants. By developing this tool, we aim to

improve the lives of patients and the general population, starting with the population

of the northern Netherlands and expanding to more individuals in the future.

2. Methods

2.1 Datasets and quality control

We used four different datasets to develop and validate the automated generation of

PGx passports, divided over two different versions of Illumina Genotyping arrays. We

used 38,030 samples from the UMCG Genetics Lifelines Initiative (UGLI) cohort of

the Lifelines prospective follow-up biobank (17) to develop and validate an

automated PGx passport pipeline using the Infinium Global Screening Array®

MultiEthnic Disease (GSA-MD) version 1.0 array. In parallel, we used three datasets

to develop the pipeline for the GSA-MD version 3.0 array: 2473 samples from our

Genome Diagnostics patient cohort (UMCG), 77 validation samples with known PGx

genotypes, and another 574 samples from the Lifelines NEXT cohort (18).

The samples of the Lifelines UGLI cohort were genotyped using GSA-MD version

1.0. We performed quality control on both samples and markers, including removal of

samples and variants with a low genotyping call rate (<97%), variants showing

deviation from Hardy-Weinberg equilibrium ($p < 1 \times 10^{-6}$), excess of Mendelian errors

in families (>1% of the parent–offspring pairs), and samples with very high or low

heterozygosity (4 standard deviations (SD) above or below the mean for autosomes).

We further revised and removed samples that did not show consistent information

between reported sex and genotypes on the X chromosome, or between reported

familial information and observed identity-by-descent sharing with family members.

All other samples were genotyped using GSA-MD version 3.0 and were quality-

controlled similarly to those in Lifelines UGLI. In total, 36,373 Lifelines UGLI

samples, 470 Lifelines NEXT samples, 2285 validation samples of the Genome

Diagnostics patient cohort, and 76 validation samples with known PGx genotypes

passed quality control.

2.2 Selecting individuals to receive pharmacogenetic results

Quality control measures in research settings are commonly less stringent than in

diagnostics. Since Lifelines UGLI samples were genotyped in a research setting, this

dataset could potentially contain sample swaps. Therefore, we used the Idéfix (19)

select a high-quality subset of 5,765 samples suitable for returning personalized

results. In short, Idéfix selects those samples for which the actual phenotypes

sufficiently match the predicted phenotypes based on genetic predispositions. These

genetic predispositions were determined by calculating polygenic scores (PGSs) per

phenotype. These calculations aggregate the effects of common risk-decreasing or -

increasing alleles identified by GWAS. Here, we reran Idéfix using an updated list of

PGSs based on recent GWAS. Quality control, PGS calculation, and running Idéfix

were performed as described in Warmerdam et al. (19). Finally, we returned PGx

results to 996 active Lifelines participants who gave consent to receive their PGx

passport. For Lifelines NEXT, we were able to manually resolve sample swaps

based on family data and duplicated array runs. We returned PGx results to 231

mothers registered as active Lifelines NEXT participants who gave consent to

receive their PGx passport.

2.3 Variant calling

Variants were called in two separate stages. In general, we used the Opticall

software (20) to do genotype-calling on the majority of variants. However, Opticall is

not suitable for genotyping variants covering frequent CNVs. Since CYP2D6 is

affected by relatively frequent CNVs, it is not trivial to genotype variants within and

near this gene. At the same time, inclusion of CYP2D6 is essential to return a

comprehensive PGx passport because it metabolizes a large proportion of

medication. We therefore developed a novel algorithm for CNV-calling of the

CYP2D6 gene using the data obtained from a SNP array. In addition, this method

also performs genotype-calling of individual variants in the CYP2D6 gene.

2.3.1 CYP2D6

SNP arrays have been used extensively to study both SNPs and CNVs. Modern

SNP arrays genotype at hundreds of thousands of assays (probes designed to

genotype one particular variant) simultaneously by measuring the color and intensity

of light emitted when one of four labeled nucleotides hybridizes at the assays. In

addition to enabling study of which alleles are present in an individual through

comparison of the colors of the emitted light, it also allows study of CNVs by

combining the total signal intensity. We combined these two approaches in our

CYP2D6 genotyping methodology, as described below.

For genotyping *CYP2D6*, we first selected a number of independent, high-quality

variants from the quality-controlled genetic data (missingness < 0.01, minor allele

frequency (MAF) > 0.05, Hardy-Weinberg equilibrium p-value > 0.05, *HLA* excluded,

minimum distance of 100kb between variants, R^2 < 0.4). Using these variants, our

method performs a principal component analysis of the intensities of the selected

variants (summed across the two alleles). As the first principal components (PCs)

likely tag batch effects, we chose to regress out the first 100 PCs from the genomic

region of interest – the *CYP2D6* gene – using ordinary linear regression (OLS). Both

PC loadings and OLS regression parameters can be estimated in one dataset, and

later applied in another. The methodology for correcting SNP array intensities follows

the approach previously established by Lude Franke et al. (21).

After correction for batch effects, one variant is selected to assign naive CNV-calls

(rs1135824 for GSA-MDv1, rs1423323203 for GSA-MDv3). Naive, preliminary CNV

assignments are made by binning the total signal intensity of the selected variant so

that the proportions of CNV genotypes match the expected frequency of CNV-calls.

The expected frequencies (*5 = 0.0295, Duplication = 0.0275) are obtained from

PharmGKB (https://www.pharmgkb.org/page/cyp2d6RefMaterials, European

population). Using the means and SDs of the signal intensities for each bin, a naive,

preliminary CNV call is assigned to every individual based on which CNV bin has the

highest likelihood at the signal intensity of the specific variant. If multiple GSA probes

are available for a variant, a majority voting ensemble classifier is used to obtain a

consensus of the various assays. Next, genotyping is performed using a manually

selected set of variants. These variants are selected based on whether the individual

signal intensities show clusters of genotypes in accordance with what can be

expected for a biallelic variant with possible deletions and duplications (i.e., at least

six visible intensity clusters rather than the three intensity clusters expected when

dealing with a biallelic variant outside a CNV).

For each variant and possible copy number, a nearest-neighbors classifier is used to

obtain a set of highly confident samples for every variant. Then, to perform

genotyping, the theta of every sample is calculated based on its intensities. For this,

the median of the intensities for the samples naively assigned to the group of

homozygous deletions is used as the origin. Thereafter, samples are binned into k

bins, where k corresponds to the number of genotypes that can be expected for a

biallelic variant at each copy number. Bins are assigned genotypes so that they best

match Hardy-Weinberg principles. Per variant, naive genotype clusters are then

iteratively improved in a gaussian mixture model algorithm adapted from *scikit-learn*

(22). The maximization step in the gaussian mixture model algorithm was changed to

also weight genotypes according to Hardy-Weinberg principles, which were

expanded to also take deletions and duplications into account. Per variant, this

procedure is performed while accounting for the intensity measures of all assays

tagging that specific variant.

For each combination of a variant, sample and copy number, a total probability is

calculated by summing the probabilities across the genotypes corresponding to that

copy number. Subsequently, the most likely copy number is assigned to each

variant-sample combination by selecting the copy number with the largest total

probability among the possible copy numbers. The CNV status of *CYP2D6* as a

whole is determined by summing the probabilities of variants over all possible copy

numbers, and rescaling these to sum to one. Genotype probabilities are then

calculated by rescaling the possible genotype probabilities in the assigned copy

number status to one.

Genotypes of samples with 1 or 2 *CYP2D6* copies (across both the maternal and

paternal chromosome) are written so that genotypes of samples with single-gene

deletions are homozygous for the reanalyzed variants. Genotypes for variants of

samples that are homozygous for whole-gene deletions do not exist. In human

genetic datasets, the representation of genotypes for variants of samples with a

CYP2D6 duplication is not trivial, making the imputation and phasing non-trivial.

Therefore, these samples are excluded. *CYP2D6* variants in chromosome 22

genotyped using Opticall are replaced with the newly genotyped *CYP2D6* variants.

This is subsequently used in the phasing and imputation pipeline. The *CYP2D6*

CNV-calling algorithm is included as the first part of the *Asterix* software that we

developed to generate PGx profiles.

2.4 Phasing and imputation pipeline

Before phasing and imputation, PLINK files are lifted from Human genome build 37

to build 38 and alleles of insertions and deletions are recoded to match the notation

in the reference dataset. Then, variants for which the strands do not match the

reference panel are removed, and variants with >5% missingness or a Hardy-

Weinberg p-value $< 1 \times 10^{-6}$ are removed. Phasing and imputation are implemented

using Eagle v2.4.1 (23) and Minimac4 (24)(version 1.0.2). The reference dataset for

phasing and imputation is derived from the 1000 Genomes project 30x coverage

whole genome sequencing dataset (25). Finally, we replaced the imputed genotypes

with the original genotypes for the variants that were used as input in the imputation.

2.5 Crucial backbone

A minimum set of variants that should be included in our genotyping panel was

determined per gene (see Table S1). We used available 'tier 1' recommendations for

CYP2C9, *CYP2C19*, *CYP2D6, CYP3A4, CYP3A5,* and *VKORC1* (26–29). For the

other genes in our PGx passport, we assessed the variants/alleles that were

mentioned in DPWG background information and those that are included in our

validation dataset. We included a variant in the crucial backbone when: 1) the variant

results in an allele known to alter the drug response phenotype, and 2) the MAF in

the (North-Western) European population is > 0.001.

2.6 Validation

We validated GSA-MD versions 1 and 3 and *Asterix* following the clinical genome

diagnostic standards in our center. The validation parameters and corresponding

acceptance criteria are shown in Table 1.

Table 1: Validation parameters and acceptance criteria for the clinical validation of the Infinium Global Screening

Array MultiEthnic Disease versions 1 and 3, and the software tool Asterix. *‡For all variants available in both*

gnomAD-NFE and GoNL datasets, the distribution of mutant allele frequency differences between those two

datasets was recorded. Z represents the standard deviation of that distribution. Variants that deviated more than

3*Z from the gnomAD-NFE mutant allele frequency failed validation.

Topic	Validation parameter	Acceptance criterium
Quality of the array data	Call rate	³0.97
Phasing and imputation quality	Concordance with trio-data	³0.97
	R^2	³0.97
Result interpretation	Absolute number of mutant alleles detected	³5
	Mutant allele frequency	<3*Z‡
	Concordance with known genotypes	100%

We created a benchmark set of haplotypes using data of 621 (GSA-MD v1) and 135

(GSA-MD v3) parent–offspring trios to validate phasing performance. Using

SHAPEIT5 (30) we then compared the results of our statistical phasing with the

benchmark haplotypes and computed the switch error rates (SER) of all genetic

variants that were directly typed. The call rate and imputation quality are validated

based on the output of our quality control procedures. Variants were removed if: 1)

they were not directly genotyped nor imputed, 2) they were not in the crucial

backbone and had a low imputation probability ($R^2 < 0.97$), 3) the mutant allele was

observed < 5 times in the full dataset, 4) the mutant allele frequency deviated more

than 3*Z from the gnomAD-non-Finnish Europeans (NFE) mutant allele frequency (Z

represents the SD of differences in mutant allele frequency between the GoNL and

gnomAD-NFE datasets), 5) the phasing SER was > 0%, or 6) the concordance with

known genotypes of validation samples was < 100%.

When a variant was in the crucial backbone, but the imputation probability was low,

we used haploid-level dosage (HDS) to assess the variant call confidence level in an

individual sample.

In addition to variant calling, we also validated our CNV-calling algorithm. To that

end, we selected samples with a confidence score of > 0.97 (1x zero copies, 14x one

copy, and 16x two copies) and samples with a confidence score of with a CNV-

calling confidence score of > 0.9, and < 0.97 (4x one copy, 3x two copies and 5x

three copies). DNA of these samples was analyzed with a dedicated TaqMan real-

time PCR assay specifically targeting exon 9 of the *CYP2D6* gene (Certe,

Groningen, the Netherlands). It is worth noting that this assay cannot detect gene

duplication or hybrid alleles. However, as we report *CYP2D6* to be not available

when >2 copies of *CYP2D6* are detected, these calls do not currently require

validation.

Finally, we confirmed that the software we developed correctly translated 1) the

detected mutations to the corresponding star alleles and, subsequently, 2) the

combination of haplotypes to the corresponding contraindication (CI) and/or

predicted phenotype. This is described in more detail in section 2.9.3 and in the

Supplementary Information.

2.7 Pruning of allele definition tables

We used gene-specific allele definition tables to assign star alleles. Available tables

were downloaded from PharmVar (v5.2.14 for GSA-MD version 3; v6.0.5 for GSA-

MD version 1) for *CYP2B6, CYP2C9, CYP2C19, CYP2D6, CYP3A4, CYP3A5* and

NUDT15 and from PharmGKB for *ABCG2* and *TPMT*

(https://www.pharmgkb.org/page/pgxGeneRef). Custom tables were constructed for

MTHFR, SLCO1B1 and *VKORC1*. We curated all allele definition tables for the GSA-

MD versions 1 and 3 separately and removed all variants that did not pass

acceptance criteria (Tables S2a-b). We then pruned each table to prevent

ambiguous calls caused by haplotypes that we cannot distinguish based on the

variants included in our pipeline. The resulting allele definition tables are shown in

Tables S3a-x.

2.8 Translation to pharmacogenetic contraindication

To translate the diplotype to the PGx CI used in medication surveillance, we

designed custom translation tables (see Tables S4a-l). Allele function of individual

star alleles and their translation into predicted phenotype and PGx CI are based on

PharmVar, PharmGKB, and/or DPWG background information. All possible

diplotypes based on the pruned allele definition tables are included in the translation

tables. We made sure the PGx CIs correspond to those used in the DPWG

recommendations.

2.9 Generating PGx profiles

We developed the *Asterix* tool to produce PGx profiles from genotyping data from

the GSA-MD versions 1 and 3. *Asterix* is composed of four main tasks: 1)

determining the *CYP2D6* copy number status, 2) determining star alleles, 3)

translating these to predicted phenotypes and PGx CIs, and 4) constructing a

logging file.

2.9.1 Determining star alleles, predicted phenotypes, and PGx CIs

After correction of the *CYP2D6* genotypes based on the CNV status, star alleles for

all included genes are assigned using the gene-specific allele definition tables.

Subsequently, predicted phenotypes and PGx CIs are determined with the custom

translation tables. When determining star alleles present for *CYP2D6*, *Asterix* takes

into account whether whole-gene deletions (*CYP2D6*5*) were detected in CNV-

calling.

2.9.2 Constructing a logging file

To be able to check *Asterix* output, an extensive logging file is constructed that

contains information for several checkpoints in our pipeline. Parameters that are

logged include the estimated phased haploid-level dosage (HDS) when < 0.97 (only

for imputed genotypes), missing variants, and haplotypes that were not described in

translation tables. Identical haplotypes are merged together to simplify the validation

process.

2.9.3 Approval and release of the PGx profiles

The PGx profiles generated are to be released by a registered pharmacist. To

maximize process efficiency, the approval and release of the PGx profiles is

performed on the level of unique combinations of detected mutations and unreliable

calls. When a sample contains only previously approved haplotypes, it is

automatically released. Each decision is logged to maintain an audit trail. The

procedure for approving and releasing the profiles is described in more detail in the

Supplementary Information.

2.10 Building layperson-friendly PGx passports

The output of *Asterix* is a PGx profile that is suitable for application in clinical practice

by healthcare practitioners. However, (future) patients should not be expected to

understand the meaning or implications of such a profile. We therefore developed a

layperson-friendly digital PGx passport to fulfill the informational needs of (future)

patients.

2.10.1 Developing layperson-friendly communication materials

We aimed to develop communication materials that explain the impact of drug–gene

interactions (DGIs) that are understandable for 90% of the study population. We

developed layperson-friendly texts based on the DPWG guidelines for healthcare

practitioners (see Table S5), written at Common European Framework of Reference

Common Reference Level B2 (31). One important difference that we had to address

is that the DPWG guidelines do not contain an explanation for normal metabolizers

because standard-of-care applies for these patients. However, texts for the (future)

patient do need to cover this topic because it is valuable to explain that standard-of-

care applies. The texts are constructed using three building blocks (Figure 1).

Block 1: Predicted influence of your DNA.	Predicted influence of your DNA
The *generic drug name* is used and the nature of the DGI is explained. The <u>PGx result</u>, including possible implications such as increased or decreased chance of ADRs and/or efficacy and recommendations for drug use is explained. A reference to the physician or pharmacist for more information is included for actionable DGIs.	How well amitriptyline is processed by the body varies from person to person. This is, in part, determined by your DNA. Your genes show that *amitriptyline* is processed <u>much more slowly</u> than average by your body. The normal dosage can, therefore, lead to a higher chance of side effects. You will probably need a lower dose of *amitriptyline*. Your physician or pharmacist can tell you more about the appropriate dose.
Block 2: Advice. Statement when sharing the PGx result with the physician or pharmacist would be wise. An explanation that a dose currently in use may already be tailored sufficiently is included, as well as a reminder to never change drug use	**Advice** If you are using or, in the future, get prescribed *amitriptyline*, it would be wise to share the information in your Pharmacogenetic Passport with your physician or pharmacist, so they can take this into account. Your physician and pharmacist also consider factors such as age, sex, and other medication you might be using. Based on that

Figure 1: Building blocks used to construct layperson-friendly texts to explain Drug–Gene interactions (left) and the corresponding text for CYP2D6 PM-amitriptyline (translated from Dutch, right).

2.10.2 Digital infrastructure

We created a DGI table that contains all the CIs produced by *Asterix*, a list of known DGIs currently in the DPWG guidelines, and the layperson-friendly texts associated

to each combination of these. The relevant texts for each participant are selected

based on their PGx profile and known DGIs. For example, the text explaining the

consequences for amitriptyline in Figure 1 will be selected for the individuals who are

listed as CYP2D6 PM in their PGx profile. The PGx profile and corresponding texts

are subsequently unlocked through an Application Programming Interface (API)

following the Health Level Seven Fast Healthcare Interoperability Resources (HL7-

FHIR) format. Participants can view their PGx passport in the mobile application

(app) *Gen en Geneesmiddel,* which we developed with Synappz Digital Healthcare

and which is available for iOS and Android (see Figure 4).

2.10.3 Additional features

Upon installing the app, the user is shown a series of information screens that

explain PGx and the content and use of the app. This information can be reviewed

within the app after completion of this initial onboarding process. Furthermore,

frequently asked questions (FAQ) are included in the app to address several

anticipated questions. The FAQ-section is supplemented by new questions from

users and associated answers.

The layperson-friendly texts displayed in the mobile application are accompanied by

articles that describe the PGx CI that interacts with the drug being viewed (see Table

S5). When multiple PGx CIs are relevant for the drug being viewed, all the relevant

articles are shown. The articles provide interested participants with more detailed

information and facilitate communication between (future) patients and healthcare

practitioners.

Finally, a technical report aimed at healthcare practitioners can be downloaded from

the app. This report contains the PGx profile and accompanying information,

including contact information for a helpdesk and instructions on how to interpret and

process the results.

3. Results

3.1. Pharmacogenomic profiles and passports

To validate the use of a layperson-friendly, clinical-grade PGx passport using

oligonucleotide arrays, we have developed a tool (*Asterix*) to produce PGx profiles

from genotyping data from the GSA-MD versions 1 and 3. As described in Figure 2a,

our tool works by determining the *CYP2D6* copy number, phasing and imputing

genetic variants, determining the star alleles of the samples provided, and translating

diplotypes to PGx CIs. Instructions on the installation and use of *Asterix* are

described on the GitHub repository (https://github.com/molgenis/asterix).

3.2. Validation

A total of 34 variants, in 12 genes, were included in the crucial backbone of the PGx

pipeline, and subsequently underwent validation (Table 2). For *CYP2B6* we initially

also included rs3211371 (*5*). However, since the DPWG withdrew CI definitions

differentiating *CYP2B6*5* from *1*, we removed this SNP from the crucial backbone.

Three variants in *CYP2C9* (rs28371686 (*5*), rs9332131 (*6*) and rs7900194 (*8*))

were initially included in the crucial backbone based on their multiethnic allele

frequency (26). However, since each of these variants had a MAF < 0.001 in the

NFE population, we subsequently excluded them from the crucial backbone for GSA-

MD v1. It is important to note that the Lifelines population is known to be reflective of

Figure 2: Overview of the work in this paper. (a) Schematic of the pharmacogenetic (PGx) pipeline developed to

generate PGx profiles. Data collection involves obtaining DNA samples from participants and genotyping using

an oligonucleotide array. We ascertain CYP2D6 copy numbers (CN) using a custom CNV calling approach. This

method estimates CN status at multiple points within the CYP2D6 gene and integrates these estimates to

determine the overall gene copy number status. Twelve genes contribute to the generation of PGx profiles. The

combined effects of these genes are aggregated into medication-level recommendations, which are

communicated to participants via a layperson-friendly app. (b) Broad overview of the methodology used for

validating the PGx pipeline. (c) Overview of the datasets used for development and validation of the PGx

passport (Development datasets), and those for which we also returned PGx-profiles (Repurposed datasets).

the NFE population. For the same reason, we excluded rs41303343 (*CYP3A5*7*).

For *TPMT* variant rs1800462 (*3A*/*3B*), no mutant alleles were detected on GSA-MD

v3. As we could not ensure the associated GSA probe was functional, *TPMT* was

excluded from PGx profiles generated with GSA-MD v3. All other variants in the

crucial backbone passed validation criteria. In case of an imputation score (R^2) <

0.97, imputation quality was judged from sample-specific HDS. When the HDS <

0.97, a registered pharmacist decided during approval and release whether a result

could be released for the associated gene.

Table 2: Validation results of crucial backbone variants.

[a]rs28371686, rs9332131, and rs7900194 were left out of crucial backbone for GSA-MD v1, based on known

ethnic composition of Lifelines

[b]rs41303343 was left out of the crucial backbone for both GSA-MD v1 and GSA-MD v3, based on known ethnic

composition of Lifelines

[c]Reference allele frequency in non-Finnish Europeans (NFE)

Gene	rsID	Alt. allele frequency[c]	GSA-MD v1		GSA-MD v3		
			method	R^2	method	R^2	failed validation
ABCG2	rs2231142	0.1092	direct		direct		
CYP2B6	rs3745274	0.2372	direct		direct		
CYP2C19	rs4244285	0.1480	direct		imputation	0.99	
	rs4986893	0.0002	direct		direct		
	rs12248560	0.2200	direct		direct		
CYP2C9	rs1799853	0.1266	imputation	0.99	imputation	1.00	
	rs1057910	0.0662	direct		direct		
	rs28371686	0.0000	(*imputation*)[a]	(*0.02*)	direct		
	rs9332131	0.0001	(*imputation*)[a]	(*0.00*)	direct		
	rs7900194	0.0005	(*imputation*)[a]	(*0.01*)	imputation	0.90	
	rs28371685	0.0029	direct		direct		
CYP2D6	rs1135840	0.5588	imputation	0.87	imputation	0.93	
	rs16947	0.3347	imputation	0.89	imputation	0.96	
	rs35742686	0.0170	imputation	0.79	imputation	0.82	
	rs3892097	0.1986	imputation	0.91	imputation	0.92	

Gene	rsID					
	rs5030655	0.0117	imputation	0.61	imputation	0.27
	rs5030656	0.0285	imputation	0.67	imputation	0.70
	rs1065852	0.2225	imputation	0.91	imputation	0.95
	rs28371706	0.0011	direct		imputation	0.86
	rs61736512	0.0006	imputation	0.04	imputation	0.97
	rs59421388	0.0006	imputation	0.04	imputation	0.97
	rs1058164	0.5525	imputation	0.86	imputation	0.93
	rs28371725	0.0955	imputation	0.84	direct	
CYP3A4	rs35599367	0.0492	direct		direct	
	rs776746	0.9311	direct		direct	
CYP3A5	rs10264272	0.0009	direct		direct	
	rs41303343	0.0004	(*imputation*)[b]	(0.96)	*n/a*[b]	
MTHFR	rs1801133	0.3373	direct		direct	
NUDT15	rs116855232	0.0029	direct		direct	
SLCO1B1	rs4149056	0.1587	direct		direct	
TPMT	rs1800462	0.0024	imputation	0.46	direct	<5 mutants detected
	rs1800460	0.0382	imputation	0.96	direct	
	rs1142345	0.0423	imputation	1.00	direct	
VKORC1	rs9923231	0.3779	direct		direct	

We considered 731 and 703 variants outside the crucial backbone for inclusion in the

translation tables of GSA-MD v1 (PharmVar tables v6.0.5) and GSA-MD v3

(PharmVar tables v5.2.14), respectively. Results of the validation process are

summarized in Figure 3 and are fully available in Tables S2a-b. Many variants were

removed due to unavailability on the genotyping arrays and in the imputation panel.

Beyond that, poor imputation score and/or low allele frequency were the leading

reasons for exclusion. Phasing quality was verified by comparison to phasing results

using parent–offspring trios. Poorly phased variants were retained when they were

the only variant of interest in that gene, rendering phasing irrelevant. Importantly, 37

variants were typed in 76 validation samples, using a traditional DNA amplification

genotyping method (Leiden University Medical Center). For all 37 variants, the

 genotypes derived from GSA-MD v3 (19 directly typed, 18 imputed) were a 100%

 match to the traditional assays.

 Figure 3: Validation status of variants from the PharmVar translation tables outside the crucial backbone. The

 UpSet plot displays the frequency of variants that either meet or fail to meet the validation criteria. Columns with

 a single marked circle represent the frequency of variants failing exclusively that criterion, while columns with

 multiple marked circles indicate variants that failed multiple corresponding criteria.

 Within *Asterix*, we embedded a custom CYP2D6 copy number (CN) calling

 algorithm. To validate the *Asterix*-derived *CYP2D6* CN calls, a total of 43 samples

 were selected, representing a CN probability of at least 0.9 and each of the four

 possible CN calls (zero copies, one copy, two copies, and three copies). The CN

 status of the aforementioned samples was determined using a dedicated TaqMan

 assay targeting exon 9. To compare the results, we initially evaluated whether the

 overall CN calls with a moderate probability of over 0.9 matched the calls reported by

 the TaqMan assay (Figure 4a). Of the 43 samples, *Asterix* identified three as having

 a CYP2D6 duplication (three copies), whereas the TaqMan assay reported these

 samples as homozygous reference (two copies). In one instance, *Asterix* reported a

 heterozygous deletion (one copy) while the TaqMan assay returned a homozygous

reference call. However, for these four samples, *Asterix* consistently reports a

homozygous reference call at exon 9, which is consistent with the call reported by

the TaqMan assay. For the majority of the other variants, *Asterix* reports calls

matching that of the overall aggregated estimate (Figure 4b). This suggests that

these samples are likely to possess a hybrid allele or tandem arrangement, resulting

in a distinct CN call at exon 9. It is worth noting that the TaqMan assay is not

capable of differentiating between regular copy numbers and hybrid alleles. At a

probability of at least 0.97, all 31 samples exhibit matching CN calls.

Based on the validated variants, final translation tables were created to first translate

variants to haplotypes (Tables S3a-x) and then to translate diplotypes to CI (Tables

S4a-l). These tables were then used as input for *Asterix* to create preliminary PGx

profiles for 996 Lifelines participants and 470 Lifelines NEXT participants. Approval

(or adjustment) and release by a registered pharmacist resulted in the final PGx

profiles. Haplotype frequencies were compared to reference frequencies in a

European population (source: Clinical Pharmacogenetics Implementation

Consortium (CPIC)) or NFE (source: DPWG/gnomAD) (Figure 5 and Tables S6a-b).

Finally, the technical PGx profiles were shared with the biobank participants in the

smartphone application *Gen en Geneesmiddel* (Figure 4). The app links the

technical information in the PGx profile with associated explanatory texts, yielding a

layperson-friendly PGx passport. Rather than using genes as the basic unit of

information, the app categorizes drugs into five categories (Figure 4A): poor

metabolism ('very slow'), intermediate metabolism ('slow'), normal metabolism

('average'), ultrarapid metabolism ('very fast'), and non-metabolic drug–gene

interactions ('other processes'). By selecting one of the categories, users can see a

list of all the medications in that category, based on their pharmacogenetic profile

(Figure 4B). More information can be found on specific medication (Figure 4C), i.e.,

what is it used for and how is it affected by the user's PGx profile. Interested users

can get more information on the associated CIs by proceeding to a linked article.

Finally, the technical information (PGx profile) can be downloaded and/or shared

directly from the app (Figure 4D).

Based on the PGx profiles we generated using *Asterix*, 92% and 90% of the Lifelines

and Lifelines NEXT populations, respectively, carried at least one actionable PGx CI

(Tables S6a-b list frequencies of PGx haplotypes and CIs). We have returned these

profiles to the biobank participants through the *Gen en Geneesmiddel* app, enabling

them to get optimized (future) drug prescriptions with the help of their physician and

pharmacist.

Figure 4: Overview of the 'Gen en Geneesmiddel' app. a) Main menu, where five categories can be selected:

'poor', 'intermediate', 'normal' and 'ultrarapid' metabolism and 'other processes'. b) A personalized list of

medication in the 'poor metabolism' category. c) Explanation of the predicted metabolism of brexpiprazole. d) The

pharmacogenetic profile can be saved or shared from the app.

Figure 5: Validation results of CYP2D6 CNV-calling (a, b). a) correspondence between a standard CYP2D6 CNV-calling procedure using a TaqMan assay (x-axis) and the CNV-calling results determined using GSA-MD v3 and our specially designed algorithm. Only exon 9 was assessed in the TaqMan assay. The y-axis for the three graphs shows the copy number (CN) status determined from the array for a variant at exon 9 only (left), for the overall CN status at a probability threshold of 0.90 (middle), and for the overall CN status at a probability threshold of 0.97 (right). b) the CN status for the four deviating samples with a probability threshold of 0.90 over all variants included in the CNV-calling algorithm. c) Haplotype frequency differences compared to the frequencies reported by CPIC (left) and contraindication frequency differences compared to the frequencies reported by CPIC (right).

4. Discussion

In this study, we use oligonucleotide genotyping arrays to develop a low-cost PGx passport that repurposes existing genotype data from the Lifelines and Lifelines NEXT cohorts for clinical use. We clinically validated the genotyping method by aligning validation parameters and corresponding acceptance criteria to the local clinical diagnostic center and by comparing genotyping results with results from classical PCR-based assays. Using parent–offspring trios, we could also verify the accuracy of haplotype phasing, information that is unavailable when using PCR-based PGx assays. We showed that we could confidently include most of the recommended 'tier 1' PGx variants for the genes included in our study (26–29), with only a few exceptions. *TPMT* was excluded for the GSA-MDv3 oligonucleotide array because variant rs1800462 did not meet validation criteria. In addition, three *CYP2C9* variants classified as tier 1 (mutant allele frequency > 0.001) were excluded from the crucial backbone on the grounds of their much lower allele frequencies in the European population. To apply our pipeline in a broader context, these variants would need to be validated in a population with higher MAFs.

Furthermore, to address previously identified barriers to use of PGx passports by laypeople, we implemented communication of relevant PGx information through a newly developed smartphone application. As the uptake of the PGx passport increases, support will be provided for healthcare professionals to properly apply PGx in their daily practice.

4.1 Producing PGx profiles

As the generation of clinically valid PGx profiles from array-based genotype data is

the cornerstone of this study, we discuss this first. We did not include *DPYD, FV,*

HLA-A, HLA-B, CYP1A2, and *COMT* in the PGx passport because: 1) *DPYD*

genotyping is already pre-emptively performed in combination with phenotyping in

routine care, 2) the DPWG guidelines for mutations in *FV* were retracted, 3) there is

no allele definition table for determining *HLA* alleles, so we could not use the same

pipeline for this region, and 4) *CYP1A2* and *COMT* are part of DPWG guidelines but

currently do not contain DGIs.

An important limitation in our passport production is that we could not provide a

complete passport for every individual. We used imputation quality (R^2) to select

imputed variants of sufficient quality. If a variant with insufficient imputation quality

($R^2 < 0.97$) was in the crucial backbone, the variant was not removed from the

translation tables, as that would lead to the complete exclusion of the corresponding

gene for all individuals. Instead, we use haploid dosage scores to select a subset of

individuals with imputed genotypes of sufficient quality. We could thereby still report

results for a subset of individuals. When applying our method in a diagnostic setting,

genotypes should be determined by a back-up analysis if no reliable result can be

extracted from the array data.

We confirm that actionable PGx variants are present in >90% of the general

population, comparable to the percentages reported in literature. In this

implementation study, we were unable to evaluate the impact of these actionable

variants on drug efficacy and adverse effects, due to the scale of our study and the

absence of high-quality prescription records for our study populations. In the future,

we aim to further investigate this by linking prescription data from an external

database through the Pharmlines initiative (28).

4.2 Layperson-friendly communication of PGx passports

Engaging patients with PGx testing is not limited to increasing their understanding of

genetic information, it is also meant to enable their participation in prescription

decisions. Patients who are well-informed about the implications for their prescription

management are better equipped to collaborate with their healthcare providers in

developing personalized treatment plans that align with their needs and preferences

(32). Furthermore, this engagement builds a sense of empowerment and ownership

over one's health, which has been shown to improve adherence to treatment

regimens and to lead to better health outcomes (33).

The PGx passport, as a digital health tool, facilitates patient engagement by

providing a platform to exchange PGx information and medication management to

enhance patient understanding. By providing layperson-friendly communication

within the PGx passport, it may in the future be used to support patient education

initiatives to broaden the understanding and implementation of pharmacogenomics

across the Netherlands.

We aimed to develop communication materials that are understandable for 80% of

the general population. Since the Lifelines and Lifelines NEXT cohorts are more

highly educated than the general population, we speculate that the material should

be understandable for around 90% of the study population. In an ongoing

implementation study, we will evaluate the level of comprehension among

participants, so that texts can be better adapted to citizens' needs.

Several practical decisions were made in the development of the smartphone

application. For example, *CYP3A5* non-expressers (*3/*3) were placed in the app

category that represents normal metabolism, while both hetero- and homozygous

expressers are placed in the ultrarapid metabolism category. In the Netherlands,

CYP3A5 non-expressers are most prevalent, so the standard of care applies for

people with this metabolizer status.

4.3 Future perspectives

The completion of the PGx passport and smartphone application represents another

step forward on the road toward true personalized medicine. By demonstrating to

healthcare professionals how incorporating PGx into daily practice works to

streamline prescribing practices through the added value of reduced adverse effects,

the PGx passport aims to lower the barrier to making key prescription decisions.

Although clinical implementation of PGx has picked up pace in the past years, it has

not yet been integrated into routine care. One of the main barriers here is likely to be

the lack of strong evidence that its application effectively improves healthcare and

reduces healthcare costs, which are both prerequisites for reimbursement (34).

Other important potential barriers are the complexities associated with integrating

genetic data into electronic health records and the need to ensure that healthcare

professionals have the requisite expertise and resources to comprehend and utilize

PGx insights (35,36). Previous research at our center has shown that, although

these barriers exist, they can be overcome (2).

In this study, we have demonstrated that PGx passports can be generated much

more cheaply than the currently prevailing methodologies. However, the efficacy and

cost-effectiveness of pre-emptive PGx testing still needs to be more thoroughly

proven, and this will require the inclusion of large numbers of participants from the

general population. Only then will the numbers of drug prescriptions dispensed

based on the information in a PGx passport be sufficient to perform quantitative

analysis of adverse effects and/or treatment effectiveness. As the PGx passport is

currently limited to the Netherlands, the findings on cost-effectiveness would only be

applicable to the Netherlands. However, with the push to harmonize the European

Health Data Space, demonstrating cost-effectiveness in the Netherlands will enrich

the changing digital health space. Lastly, as the initial data was derived from the

Lifelines biobank, incorporating *Asterix* into future investigations in Lifelines would

enable more in-depth research into the interplay between genetic variation,

environment, population health, and pharmacotherapy.

4.4 Concluding remarks

This study provides a method to produce low-cost, clinically validated PGx profiles

for use by healthcare professionals from genotype array data. As more and more

individuals are genotyped, both in academic and healthcare settings, translating

available genetic data towards clinical decision-making will be essential for

personalized care. Furthermore, we have developed a smartphone application to

ensure that the results from our pipeline, *Asterix,* reach those whose health may be

impacted. To break down barriers to patient engagement, the application provides a

novel approach to understandably communicate PGx results to laypeople. The

reception of this information by users will be reported in an upcoming publication.

Future steps include the evaluation of the costs and effectiveness of the pre-emptive

PGx passport in the general population and further research into the interplay

between genetic variation and population health and drug response phenotypes.

References

1. Abdullah-Koolmees H, Keulen AM, Nijenhuis M, Deneer VHM.

Pharmacogenetics Guidelines: Overview and Comparison of the DPWG, CPIC,

CPNDS, and RNPGx Guidelines. Front Pharmacol. 2021 Jan 25;11:595219.

2. Lanting P, Drenth E, Boven L, Hoek A, Hijlkema A, Poot E, et al. Practical

Barriers and Facilitators Experienced by Patients, Pharmacists and Physicians to

the Implementation of Pharmacogenomic Screening in Dutch Outpatient Hospital

Care-An Explorative Pilot Study. J Pers Med. 2020 Dec 21;10(4):293.

3. Duarte JD, Dalton R, Elchynski AL, Smith DM, Cicali EJ, Lee JC, et al. Multisite

investigation of strategies for the clinical implementation of pre-emptive

pharmacogenetic testing. Genet Med. 2021 Dec;23(12):2335–41.

4. Swen JJ, Wouden CH van der, Manson LE, Abdullah-Koolmees H, Blagec K,

Blagus T, et al. A 12-gene pharmacogenetic panel to prevent adverse drug

reactions: an open-label, multicentre, controlled, cluster-randomised crossover

implementation study. The Lancet. 2023 Feb 4;401(10374):347–56.

5. Skokou M, Karamperis K, Koufaki MI, Tsermpini EE, Pandi MT, Siamoglou S, et

al. Clinical implementation of preemptive pharmacogenomics in psychiatry.

eBioMedicine [Internet]. 2024 Mar 1 [cited 2024 Aug 27];101. Available from:

https://www.thelancet.com/journals/ebiom/article/PIIS2352-3964(24)00044-

6/fulltext

6. Mason-Suares H, Kim W, Grimmett L, Williams ES, Horner VL, Kunig D, et al.

Density matters: comparison of array platforms for detection of copy-number

711  variation and copy-neutral abnormalities. Genetics in Medicine. 2013 Sep

712  1;15(9):706–12.

713 7. Momtaz R, Ghanem N m., El-Makky N m., Ismail M a. Integrated analysis of

714  SNP, CNV and gene expression data in genetic association studies. Clinical

715  Genetics. 2018;93(3):557–66.

716 8. Taylor C, Crosby I, Yip V, Maguire P, Pirmohamed M, Turner RM. A Review of

717  the Important Role of CYP2D6 in Pharmacogenomics. Genes. 2020

718  Nov;11(11):1295.

719 9. Leitsalu L, Haller T, Esko T, Tammesoo ML, Alavere H, Snieder H, et al. Cohort

720  Profile: Estonian Biobank of the Estonian Genome Center, University of Tartu.

721  International Journal of Epidemiology. 2015 Aug 1;44(4):1137–47.

722 10. Sudlow C, Gallacher J, Allen N, Beral V, Burton P, Danesh J, et al. UK Biobank:

723  An Open Access Resource for Identifying the Causes of a Wide Range of

724  Complex Diseases of Middle and Old Age. PLOS Medicine. 2015

725  mrt;12(3):e1001779.

726 11. Littlejohns TJ, Sudlow C, Allen NE, Collins R. UK Biobank: opportunities for

727  cardiovascular research. European Heart Journal. 2019 Apr 7;40(14):1158–66.

728 12. Minegishi N, Nishijima I, Nobukuni T, Kudo H, Ishida N, Terakawa T, et al.

729  Biobank Establishment and Sample Management in the Tohoku Medical

730  Megabank Project. The Tohoku Journal of Experimental Medicine.

731  2019;248(1):45–55.

13. Mahajan S, Lu Y, Spatz ES, Nasir K, Krumholz HM. Trends and Predictors of

Use of Digital Health Technology in the United States. Am J Med. 2021

Jan;134(1):129–34.

14. Shen N, Jankowicz D, Strudwick G. Patient and Family Engagement Approaches

for Digital Health Initiatives: Protocol for a Case Study. JMIR Research Protocols.

2021 Jul 21;10(7):e24274.

15. Integraal Zorg Akkoord. 2022 Sep;

16. Sijtsma A, Rienks J, Harst P, Navis G, Rosmalen JGM, Dotinga A. Cohort Profile

Update: Lifelines, a three-generation cohort study and biobank. Int J Epidemiol.

2022 Oct 13;51(5):e295–302.

17. Scholtens S, Smidt N, Swertz MA, Bakker SJ, Dotinga A, Vonk JM, et al. Cohort

Profile: LifeLines, a three-generation cohort study and biobank. International

Journal of Epidemiology. 2015 Aug 1;44(4):1172–80.

18. Warmink-Perdijk WDB, Peters LL, Tigchelaar EF, Dekens JAM, Jankipersadsing

SA, Zhernakova A, et al. Lifelines NEXT: a prospective birth cohort adding the

next generation to the three-generation Lifelines cohort study. Eur J Epidemiol.

2020 Feb;35(2):157–68.

19. Warmerdam R, Lanting P, Lifelines Cohort Study, Deelen P, Franke L. Idéfix:

identifying accidental sample mix-ups in biobanks using polygenic scores.

Bioinformatics. 2022 Feb 15;38(4):1059–66.

20. Shah TS, Liu JZ, Floyd JAB, Morris JA, Wirth N, Barrett JC, et al. optiCall: a

robust genotype-calling algorithm for rare, low-frequency and common variants.

Bioinformatics. 2012 Jun 15;28(12):1598–603.

21. Franke L, el Bannoudi H, Jansen DTSL, Kok K, Trynka G, Diogo D, et al.

Association analysis of copy numbers of FC-gamma receptor genes for

rheumatoid arthritis and other immune-mediated phenotypes. European Journal

of Human Genetics. 2016 Feb;24(2):263–70.

22. Pedregosa F, Varoquaux G, Gramfort A, Michel V, Thirion B, Grisel O, et al.

Scikit-learn: Machine Learning in Python. Journal of Machine Learning Research.

2011;12(85):2825–30.

23. Loh PR, Danecek P, Palamara PF, Fuchsberger C, A Reshef Y, K Finucane H, et

al. Reference-based phasing using the Haplotype Reference Consortium panel.

Nat Genet. 2016 Nov;48(11):1443–8.

24. statgen/Minimac4 [Internet]. Center for Statistical Genetics; 2024 [cited 2024 Aug

27]. Available from: https://github.com/statgen/Minimac4

25. Byrska-Bishop M, Evani US, Zhao X, Basile AO, Abel HJ, Regier AA, et al. High-

coverage whole-genome sequencing of the expanded 1000 Genomes Project

cohort including 602 trios. Cell. 2022 Sep 1;185(18):3426-3440.e19.

26. Pratt VM, Del Tredici AL, Hachad H, Ji Y, Kalman LV, Scott SA, et al.

Recommendations for Clinical CYP2C19 Genotyping Allele Selection. The

Journal of Molecular Diagnostics. 2018 May;20(3):269–76.

27. Pratt VM, Cavallari LH, Del Tredici AL, Hachad H, Ji Y, Kalman LV, et al.

Recommendations for Clinical Warfarin Genotyping Allele Selection. The Journal

of Molecular Diagnostics. 2020 Jul;22(7):847–59.

28. Pratt VM, Cavallari LH, Del Tredici AL, Gaedigk A, Hachad H, Ji Y, et al.

Recommendations for Clinical CYP2D6 Genotyping Allele Selection. The Journal

of Molecular Diagnostics. 2021 Sep;23(9):1047–64.

29. Pratt VM, Cavallari LH, Fulmer ML, Gaedigk A, Hachad H, Ji Y, et al. *CYP3A4*

and *CYP3A5* Genotyping Recommendations. The Journal of Molecular

Diagnostics. 2023 Sep 1;25(9):619–29.

30. Hofmeister RJ, Ribeiro DM, Rubinacci S, Delaneau O. Accurate rare variant

phasing of whole-genome and whole-exome sequencing data in the UK Biobank.

Nat Genet. 2023 Jul;55(7):1243–9.

31. Kennedy S. Fluent in 3 Months. 2023 [cited 2024 Aug 28]. CEFR Levels: What

They Are and How to Test Yourself. Available from:

https://www.fluentin3months.com/cefr-levels/

32. Krist AH, Tong ST, Aycock RA, Longo DR. Engaging patients in decision-making

and behavior change to promote prevention. Stud Health Technol Inform.

2017;240:284–302.

33. Merker VL, Hyde JK, Herbst A, Solch AK, Mohr DC, Gaj L, et al. Evaluating the

Impacts of Patient Engagement on Health Services Research Teams: Lessons

from the Veteran Consulting Network. J Gen Intern Med. 2022 Apr;37(Suppl

1):33–41.

34. Ministerie van Volksgezondheid, Welzijn en Sport. Stand van zaken inzet DNA-

technologie voor zorg en preventie [Internet]. Ministerie van Algemene Zaken;

2023 [cited 2024 Aug 27]. Available from:

https://www.rijksoverheid.nl/documenten/kamerstukken/2023/10/25/kamerbrief-

over-stand-van-zaken-inzet-dna-technologie-voor-zorg-en-preventie

35. Allison M, Patterson J, Burton S, Patel D, Brassil KJ. Digital engagement

preferences of individuals aged 65 and older with cancer. JCO. 2020 Oct

10;38(29_suppl):289–289.

36. Borghouts J, Eikey E, Mark G, De Leon C, Schueller SM, Schneider M, et al.

Barriers to and Facilitators of User Engagement With Digital Mental Health

Interventions: Systematic Review. J Med Internet Res. 2021 Mar

24;23(3):e24387.

Acknowledgements

The authors thank the UMCG Genomics Coordination Center, the UG Center for

Information Technology and their sponsors BBMRI-NL & TarGet for storage and

compute infrastructure. They thank Maaike van der Lee and Jesse Swen from the

department of Clinical Pharmacy and Toxicology of the Leiden University Medical

Center for providing the validation samples with known genotypes. The authors

thank Certe Groningen for their assistance in genotyping the CYP2D6 gene in a

number of cases and the GIC of the KNMP for providing their drug information texts

to generate layperson-friendly information. They also thank past and present

members of the ELSI-team of the UMCG for valuable feedback on the developed

communication materials.

Lifelines Cohort Study

The Lifelines Biobank initiative has been made possible by funding from the Dutch

Ministry of Health, Welfare and Sport, the Dutch Ministry of Economic Affairs, the

University Medical Center Groningen (UMCG the Netherlands), University of

Groningen and the Northern Provinces of the Netherlands. The generation and

management of GWAS genotype data for the Lifelines Cohort Study is supported by

the UMCG Genetics Lifelines Initiative (UGLI). The authors wish to acknowledge the

services of the Lifelines Cohort Study, the contributing research centers delivering

data to Lifelines, and all the study participants.

Raul Aguirre-Gamboa (1), Patrick Deelen (1), Lude Franke (1), Jan A Kuivenhoven

(2), Esteban A Lopera Maya (1), Ilja M Nolte (3), Serena Sanna (1), Harold Snieder

(3), Morris A Swertz (1), Judith M Vonk (3), Cisca Wijmenga (1)

(1) Department of Genetics, University of Groningen, University Medical Center

Groningen, The Netherlands

(2) Department of Pediatrics, University of Groningen, University Medical Center

Groningen, The Netherlands

(3) Department of Epidemiology, University of Groningen, University Medical Center

Groningen, The Netherlands

Funding

This work was supported by SNN (Samenwerkingsverband Noord Nederland) [OPSNN0229 PCH Ecosysteem WP4.4]; by Lifelines NEXT, which has been made possible by funds from a grant from the UMCG Hereditary Metabolic Diseases Fund; by the grants awarded to A.Z. (Netherlands Organization for Scientific Research [NWO] VIDI Grant [016.178.056], a European Research Council [ERC] starting grant [ERC-715772], and Gravitation program EXPOSOME-NL (NWO grant number 024.004.017)), and the Lifelines Biobank initiative, which has been made possible by funds from FES (Fonds Economische Structuurversterking), SNN and REP (Ruimtelijk Economisch Programma).

L.F. is supported by grants from NWO [ZonMW-VIDI 917.14.374, ZonMW-VICI 09150182010019], by an ERC Starting grant [637640 ImmRisk] and through a Senior Investigator grant from the Oncode Institute. P.D. is funded by The Netherlands Organisation for Health Research and Development (ZonMw) [VENI 09150161910057].

Author contributions

PL, BW, MS, NK, JD and LF: conceptualization,

PL, RW, HB, TO, JS, MB, ELM, TOM, PD, JD and LF: methodology,

RW, HB, BC, JS, LJ and LF: software,

PL, RW, HB, TO, JS and MB: validation,

PL, RW, HB, TO, JS, MB, ELM and SJ: formal analysis,

SJ, JGA and DTe: investigation,

LLN, LLCS, PD, NK, JD and LF: resources,

PL, RW, HB, JS, ELM, SJ, BC and LJ: data curation,

PL, RW, HB, TO and JS: manuscript – original draft,

RW and JS: visualization,

BW, DTo, PD, NK, JD and LF: supervision,

PL, JS and JD: project administration,

NK, JD and LF: funding acquisition.

Data availability

Due to privacy reasons the individual-level data cannot be made publicly available

but can be made available upon reasonable request. The individual-level data from

Lifelines that support the findings in this publication were obtained from the Lifelines

biobank under project application number ov21_0355. This request should be

directed to the Lifelines Research Office through email (research@lifelines.nl) or by

using the application form on their website (https://www.lifelines.nl/researcher/how-

to-apply/apply-here). Access to the Lifelines NEXT Project data will be granted to all

qualified researchers and will be governed by the provisions laid out in the LLNEXT

Data Access Agreement: https://groningenmicrobiome.org/?page_id=2598. This

access procedure is in place to ensure that the data is requested solely for research

and scientific purposes, in compliance with the informed consent signed by Lifelines

NEXT participants. Scripts and other software central to this manuscript is available

on GitHub (https://github.com/molgenis/PGx-passport-pilot/ and

https://github.com/molgenis/asterix). Other analyses of genotype data and publicly

available reference data is performed using standard bioinformatics practices, for

which the code is made available upon request.

Conflicts of interest

893 NdBV and DTo are members of the Dutch Pharmacogenetics Working Group

894 (DPWG). All other authors report no conflict of interest.