Title: Integrating a host transcriptomic biomarker with a large language model for diagnosis of lower respiratory tract infection

Authors: [†]Hoang Van Phan[1], [†*]Natasha Spottiswoode[1], Emily C. Lydon[1], Victoria T. Chu[2,3], Adolfo Cuesta[1], Alexander D. Kazberouk[4], Natalie L. Richmond[1], Padmini Deosthale[1], Carolyn S. Calfee[5], Charles R. Langelier[1,3]

[†]equal contributions
*Corresponding author: natasha.spottiswoode@ucsf.edu

Affiliations:
[1] Department of Medicine, Division of Infectious Diseases, University of California San Francisco
[2] Department of Pediatrics, Division of Infectious Diseases and Global Health, University of California San Francisco
[3] Chan Zuckerberg Biohub San Francisco
[4] Department of Medicine, University of California San Francisco
[5] Department of Medicine, Division of Pulmonary, Critical Care, Allergy and Sleep Medicine, University of California San Francisco

Description. We present the novel use of a host transcriptional biomarker combined with artificial intelligence analysis of electronic medical record data to diagnose lower respiratory tract infections in a derivation cohort of critically ill adults, then the validation of this approach in a second, fully independent, cohort. This approach demonstrated high diagnostic accuracy compared to a gold standard of post-hoc multi-physician adjudication.

26 **Abstract**
27

28 **BACKGROUND**
29 Lower respiratory tract infections (LRTIs) are a leading cause of mortality worldwide and can be
30 difficult to diagnose in critically ill patients, as non-infectious causes of respiratory failure can
31 present with similar clinical features.
32

METHODS
We developed a LRTI diagnostic method combining the pulmonary transcriptomic biomarker
FABP4 with electronic medical record (EMR) text assessment using the large language model
Generative Pre-trained Transformer 4 (GPT-4). We evaluated this approach in a prospective
cohort of critically ill adults with acute respiratory failure from whom tracheal aspirate *FABP4*
expression was measured by RNA sequencing. Patients with LRTI or non-infectious conditions
were identified using retrospective, multi-physician clinical adjudication. We then confirmed our
findings by applying this method to an independent validation cohort of 115 adults with acute
respiratory failure.
42

RESULTS
In the derivation cohort, a combined classifier incorporating *FABP4* expression and GPT-4–
assisted EMR analysis achieved an AUC of 0.93 (±0.08) and an accuracy of 84%, outperforming
FABP4 expression alone (AUC 0.84 ± 0.11) and GPT-4–based analysis alone (AUC 0.83 ± 0.07).
By comparison, the primary medical team's admission diagnosis had an accuracy of 72%. In the
validation cohort, the combined classifier yielded an AUC of 0.98 (±0.04) and an accuracy of 96%.
49

50 **CONCLUSIONS**
51 Integrating a host transcriptional biomarker with EMR text analysis using a large language model
52 may offer a promising new approach to improving the diagnosis of LRTIs in critically ill adults.

INTRODUCTION

Lower respiratory tract infections (LRTIs) are a leading cause of death worldwide, yet remain challenging to diagnose[1]. This is especially true in the intensive care unit (ICU), where non-infectious acute respiratory illnesses often have similar clinical manifestations. Further complicating accurate diagnosis is the failure to identify a causative pathogen in most clinically recognized cases of LRTI[2]. The resulting diagnostic uncertainty drives the overuse of empiric antibiotics, leading to adverse outcomes ranging from *Clostridioides difficile* infection to the development of antimicrobial resistance[3,4].

Host transcriptional biomarkers are a promising modality for LRTI diagnosis that overcome several limitations of traditional microbiologic tests[5,6]. By offering a more direct and dynamic measure of the host immune response, they can enable earlier and more accurate identification of infection, and differentiate between bacterial and viral causes of pneumonia, even in cases where pathogen detection is challenging[1,2]. Single gene biomarkers are particularly amenable to clinical translation, as they can be readily incorporated into nucleic acid amplification platforms widely used in healthcare settings.

Pulmonary *FABP4*, for instance, was recently identified as a novel LRTI diagnostic biomarker in critically ill patients with acute respiratory failure, achieving an area under the receiver operating characteristic curve (AUC) of 0.85 ± 0.12 in adults[7]. Despite better performance characteristics than existing clinical protein biomarkers such as C-reactive protein[8] or procalcitonin[9], *FABP4*, like most pneumonia diagnostic biomarkers, may not yet provide the accuracy needed to enable confident clinical decisions regarding antimicrobial use in ICU patients with acute respiratory failure.

Given that large language models (LLMs) such as Generative Pre-trained Transformer 4 (GPT-4) have shown promise in a diversity of medical applications[10], we considered the possibility that they could be leveraged to improve host biomarker-based LRTI diagnosis. LLMs have demonstrated remarkable performance for image interpretation[11], patient risk

stratification[12], and assisting with clinical reasoning[13-15], although their utility for diagnosing LRTI

or other critical illness syndromes has not been assessed. Here, we address this gap by

building a diagnostic classifier combining *FABP4* with GPT-4 analysis of electronic health record

(EHR) data. We find that this combination affords remarkably accurate LRTI diagnosis,

suggesting a promising new approach to improve the care of critically ill patients.

METHODS

COHORTS AND ADJUDICATION OF LRTI STATUS

We studied patients from two prospective observational cohorts of critically ill adults with

acute respiratory failure enrolled at the University of California San Francisco (UCSF) Medical

Center (**Figure 1, Table 1**). All patients were enrolled within 72 hours of intubation under UCSF

Institutional Review Board protocols #10-02701 (derivation cohort[16], N=202; enrolled 10/2013-

01/2019) or #20-30497 and #10-02852 (validation cohort; N=115; enrolled 04/2020-12/2023).

Adjudication of LRTI status was performed retrospectively following ICU discharge by

two or more physicians using all available information in the EMR, and based on the U.S.

Centers for Disease Control and Prevention (CDC) PNEU1 criteria[17]. Patients with a clear

alternative reason for their acute respiratory failure besides pulmonary infection, representing

the clinically relevant control group, were also identified (No LRTI group). Any adjudication

discrepancies were resolved by a third physician, and patients with indeterminate LRTI status

were excluded.

EXTRACTION OF EMR DATA

The primary medical or ICU team's clinical note from the day prior to study enrollment

and the chest X-ray (CXR) read from the day of enrollment were extracted from the EMR. If no

note was written on the day prior to enrollment, a note from two days prior was substituted

(**Table 1**). If no CXR was performed on the day of enrollment, the next closest CXR read prior to

Figure 1. Study flow diagram and overview. Abbreviations: LRTI = lower respiratory tract infection; RNA-seq = RNA sequencing; CXR = chest X ray, FABP4 = gene encoding fatty acid binding protein 4; CDC = U.S. Centers for Disease Control and Prevention; GPT-4 = Generative Pre-trained Transformer 4.

the date of enrollment was used instead. Patients with no clinical notes available prior to study

enrollment were excluded (N=7 derivation cohort, N=12 validation cohort). The clinical treatment

team's ICU admission LRTI diagnosis was extrapolated based on administration of antibiotics

for empiric treatment of LRTI within one day of study enrollment, excluding antibiotics for

established non-pulmonary infections.

RNA SEQUENCING

RNA was extracted from tracheal aspirate collected on the day of enrollment and

underwent rRNA depletion followed by library preparation using the NEBnext Ultra 2 kit on a

Beckman-Coulter Echo liquid handling instrument, as previously described[7]. Finished libraries

underwent paired-end sequencing on an Illumina NovaSeq.

FABP4 **DIAGNOSTIC CLASSIFIER**

FABP4 expression was normalized using the varianceStabilizingTransformation function

from DESeq2 package (v1.42.1)[18], and used to train a logistic regression classifier. In each

iteration of k-fold cross-validation, both training and test sets were filtered to retain only genes

with at least 10 counts across 20% of the samples in the training set. The test fold's *FABP4*

expression level was normalized using varianceStabilizingTransformation and the dispersions of

the training folds, and input to the trained logistic regression classifier to assign LRTI or No LRTI

status for each patient in the test fold. The performance and receiver operating characteristic

(ROC) curve for each of the five folds was evaluated using the package pROC v1.18.5[19]. The

mean AUC and standard deviation were calculated from the average AUC derived from each

test fold. The sensitivity and specificity at the Youden's index were extracted for each test fold

separately using the function coords from the pROC package, and the average and standard

deviation was calculated across the cross-validation folds.

GPT-4 INPUT, SCORING, AND PROMPT ENGINEERING

We used the GPT-4 turbo model with 128k context length and a temperature setting of

0.2, implemented in Versa, a University of California San Francisco (UCSF) Health Insurance

Portability and Accountability Act-compliant model. For each patient, compiled clinical notes and

CXR reads were input into the GPT-4 chat interface. Prompt engineering was initially carried out

by iterative testing on clinical notes and CXR reads from five randomly selected patients in the

derivation cohort, who were excluded from further analyses. We employed a chain-of-thought

prompt strategy[20] that involved asking GPT-4 to analyze the note and CXR step-by-step. The

validation cohort included patients enrolled during the height of the COVID-19 pandemic and

thus we redacted the terms "SARS-CoV-2" or "COVID-19" from their notes to avoid biasing the

GPT-4 analysis. In our final version of the prompt (**Supplement, Appendix 1**), we asked GPT-4

to choose either LRTI or no LRTI, as exemplified in (**Supplement, Appendix 2**). For each

patient, GPT-4 was asked to diagnose LRTI in three separate chat sessions. A per-patient GPT-

4 score was calculated based on the total number of LRTI-positive diagnoses made by GPT-4.

INTEGRATED CLASSIFIER

The integrated classifier's performance was tested using 5-fold cross-validation in the

derivation cohort. Because of the smaller sample size, 3-fold cross-validation was used in the

validation cohort. For each test fold, a logistic regression classifier was trained on the remaining

training folds using both normalized *FABP4* expression and the GPT-4 score. The performance

and ROC curve for each fold was evaluated as described above. The sensitivity, specificity and

accuracy were calculated based on whether the out-of-fold predicted probability of LRTI is

greater than or equal to 50%.

COMPARING GPT-4 TO PHYSICIANS PROVIDED THE SAME DATA

We compared LRTI diagnosis by GPT-4 against LRTI diagnosis made by three

physicians trained in internal medicine (ADK) or additionally subspecializing in infectious

diseases (AC, NLR). The physicians were provided identical information and prompt as GPT-4,

and they were asked to assign each patient as either LRTI or No LRTI. The comparison

physician group score (0 to 3) was calculated based on the total number of LRTI-positive

diagnoses made by the comparison physicians.

 DATA AND CODE AVAILABILITY

     The gene count data are available at https://github.com/infectiousdisease-langelier-

 lab/LRTI_FABP4_GPT4_classifier. The code and required source data are available at

 https://github.com/infectiousdisease-langelier-lab/LRTI_FABP4_GPT4_classifier.

 RESULTS

     We evaluated the performance of four different diagnostic approaches (*FABP4*, GPT-4,

 integrated *FABP4/*GPT-4 classifier, and admission diagnosis by the primary medical team)

 against a gold-standard of retrospective LRTI adjudication performed by two or more

 physicians. In the derivation cohort, this adjudication process identified 42 patients with LRTI

 and 56 with no evidence of infection and a clear alternative explanation for respiratory failure

 (No LRTI group) (**Figure 1**). In the validation cohort, 33 LRTI and 26 No LRTI patients were

 identified.

     We provided GPT-4 with practical clinical summary information from the EMR that would

 be available to a treating physician on the day of ICU care: a CXR radiology report from the day

 of enrollment, and a note written by the medical team from the day prior. In the derivation

 cohort, notes and radiology reports from five patients were utilized for GPT-4 prompt

 engineering and optimization (Methods) and seven lacked a clinical note from the day prior to

 study enrollment, leaving a total of 37 LRTI and 49 No-LRTI cases available for analysis.

     We first compared the accuracy of the primary medical team's ICU admission diagnosis

 against the gold-standard retrospective LRTI adjudication. The medical team correctly identified

 36/37 (97%) of true LRTI cases but incorrectly called LRTI in 23/49 (47%) of patients in the No

 LRTI group, equating to an accuracy of 72% (**Figure 2A, Table 1**). We next assessed the

 diagnostic performance of *FABP4* and found that it achieved an AUC of 0.84 ± 0.11 (mean ±

 standard deviation) by five-fold cross validation (**Figure 2B**). We then assessed the

 performance of GPT-4 to diagnose LRTI, with three independent diagnoses per patient. A

logistic regression classifier based on the GPT-4 score achieved an AUC of 0.83 ± 0.07 (**Figure**

2B).

We then combined *FABP4* and GPT-4 in a single logistic regression model and found

that this integrated classifier achieved an AUC of 0.93 ± 0.08 (**Figure 2B**), outperforming both

FABP4 (P = 0.002, one-sided paired t-test) and GPT-4 alone (P = 0.008, one-sided paired t-

test). Considering an out-of-fold probability of 50% as LRTI-positive, the integrated *FABP4/*GPT-

4 classifier had a sensitivity of 78%, specificity of 88%, and accuracy of 84% (**Figure 2A**).

Assessment of the integrated classifier's performance at the Youden's index within each cross-

validation fold demonstrated an average sensitivity of 86%, specificity of 98%, and accuracy of

93%.

Next, we assessed the validation cohort (**Figure 1**), in which the primary medical team

correctly identified 25/25 (100%) of LRTI cases but unnecessarily treated for LRTI in 7/22 (32%)

of patients in the No LRTI group, equating to an accuracy of 85% **(Figure 2C)**. The integrated

*FABP4/*GPT-4 classifier achieved a sensitivity of 96%, specificity of 95%, and accuracy of 96%,

again outperforming either *FABP4* (accuracy 79%) or GPT-4 alone (accuracy 79%). In the

validation cohort, the integrated classifier achieved an AUC of 0.98 ± 0.04 using 3-fold cross-

validation, as compared to *FABP4* (0.86 ± 0.06) or GPT-4 (0.90 ± 0.01) alone (P = 0.08 and P =

0.02, respectively, one-sided paired t-test) (**Figure 2D).**

To gain insight into how GPT-4 returns diagnoses based on limited information, we

compared the LLM against the decision making of three comparison physicians provided

identical input. From the same limited EMR data and prompt provided to GPT-4, we asked the

comparison physicians to assign a diagnosis of LRTI or no evidence of LRTI for each patient in

the derivation cohort. Considering a threshold of at least one LRTI diagnosis per patient across

the three physicians as LRTI-positive, we found a sensitivity of 78%, specificity of 88%, and

accuracy of 84% **(Figure 3A).**

Figure 2. Performance of *FABP4,* **GPT-4 and integrated LRTI diagnostic classifiers in the derivation and validation cohorts. A)** Confusion matrices for initial ICU diagnosis and the integrated *FABP4/*GPT-4 classifier in the derivation cohort. **B)** Receiver operating characteristic curves from GPT-4 classifier, *FABP4* classifier, and integrated *FABP4/*GPT-4 classifier in the derivation cohort. **C)** Confusion matrices for initial ICU diagnosis and the integrated *FABP4/*GPT-4 classifier in the validation cohort. **D)** Receiver operating characteristic curves from GPT-4 classifier, *FABP4* classifier, and integrated *FABP4/*GPT-4 classifier in the validation cohort. In panels A and C, the classifiers output an LRTI diagnosis if the patients had a predicted out-of-fold LRTI probability of 50% or higher. In panels B and D, the area under the curves (AUCs) are presented as mean ± standard deviation.

Considering a threshold of at least one LRTI diagnosis per patient across the three

physicians as LRTI-positive, we found a sensitivity of 78%, specificity of 88%, and accuracy of

84% **(Figure 3A).** Finally, we sought to identify potential biases in GPT-4 diagnoses by

comparing GPT-4 results to those of the comparison physicians (**Figure 3B**), focusing on cases

with two or more discordant LRTI diagnoses. Of the nine patients more frequently diagnosed

with LRTI by GPT-4 versus the comparison physicians (**Figure 3B**), six had clinical notes with

no mention of LRTI, but explicit concern for LRTI in the CXR report. This suggested that GPT-4

may have placed more weight on CXR reads relative to physicians. Of the two patients

disproportionately diagnosed with LRTI by comparison physicians versus GPT-4 (**Figure 3B**),

one had a final diagnosis of e-cigarette/vaping associated lung injury, and the other had LRTI

attributed to rhinovirus.

Figure 3. Comparison of GPT-4 performance to physicians provided the same EMR data from the LRTI derivation cohort. A) Confusion matrix of diagnosis by three GPT-4 comparison physicians who received the same prompt and data as GPT-4. **B)** Comparison of GPT-4 LRTI scores as compared to physicians. In Panel B, X-axis depicts the number of times GPT-4 diagnosed LRTI out of 3, Y-axis shows the number of times the physicians called LRTI out of 3. Blue boxes indicate instances in which GPT-4 diagnoses were most discordant with comparison physicians (the scores differ by 2 or more).

DISCUSSION

Our findings demonstrate that the combination of a host transcriptomic biomarker with AI

assessment of EMR text data can improve LRTI diagnosis in critically ill patients. We found that

an integrated *FABP4/*GPT-4 classifier achieved higher LRTI diagnostic accuracy than *FABP4*

alone, GPT-4 alone, or the treating medical team. In our study population, we found that the

initial treating physicians unnecessarily prescribed antibiotics in a third to half of patients

ultimately found to have non-infectious causes of acute respiratory failure. Had our integrated

classifier results been theoretically available at time of ICU admission, we estimate that

inappropriate antibiotic use could have been prevented in >90% of No LRTI patients who were

unnecessarily treated. Acute respiratory illness is a leading reason for inappropriate antibiotic

use[21], and our results suggest a potential role for biomarker/AI classifiers in antimicrobial

stewardship, a major goal of the U.S. CDC[22] and the World Health Organization[23].

Previous studies have found that GPT-4 is influenced by the precise language used in a

prompt, leading to a need for prompt engineering[14]. By iterating our prompt on a subset of

patients, and through direct comparison to physicians provided identical EMR data, we identified

possible blind spots of GPT-4 and gained insights that may help guide future optimization of

LLMs for infectious disease diagnosis.

A primary strength of this study is the novel combination of a host transcriptional

biomarker with AI interpretation of EMR text data to advance infectious disease diagnosis. We

address one of the most common and challenging diagnostic dilemmas in the ICU, leverage

deeply characterized cohorts, and employ a rigorous post-hoc LRTI adjudication approach

incorporating multiple physicians. Importantly, clinicians with access to a HIPAA-compliant GPT-

4 interface can readily use our prompt without any prior bioinformatics expertise. Weaknesses

of this study include a relatively small sample size, assessment of a biomarker not yet

commonly used in clinical practice, and restriction to mechanically ventilated patients.

Future work can test whether GPT-4 can improve the marginal performance of widely

available clinical biomarkers such as C-reactive protein, assess *FABP4/*GPT-4 classifier

performance in larger independent cohorts, and evaluate these methods for the diagnosis of

other critical illness syndromes such as sepsis.

Corresponding Author. Natasha Spottiswoode, MD, DPhil (natasha.spottiswoode@ucsf.edu),
Department of Medicine, Division of Infectious Diseases, University of California San Francisco,
513 Parnassus Avenue, Suite S380, San Francisco CA 94143

Author Contributions. Drs. N. Spottiswoode and C. R. Langelier had full access to all of the
data in the study and take responsibility for the integrity of the data and the accuracy of the data
analysis. Drs. H. V. Phan and N. Spottiswoode are co-first authors.
262
Concept and design: Phan, Spottiswoode, Langelier
Acquisition, analysis, or interpretation of data: All authors
Drafting of the manuscript: Phan, Spottiswoode, Langelier
Critical review of the manuscript for important intellectual content: Lydon, Chu, Calfee
Statistical analysis: Phan, Spottiswoode, Langelier
Administrative, technical, or material support: UCSF
Supervision: Calfee*,* Langelier
270
271   **Conflict of Interest Disclosures:** No authors report conflicts of interest.
272
273   **Funding/Support:** This study was supported by NIAID R01AI185511 and the Chan-Zuckerberg
274   Biohub (Dr. Langelier) and NHLBI R35HL140026 (Dr. Calfee).
275
276   **Role of the Funder/Sponsor:** The funding organizations had no role in the design and conduct
277   of the study; collection, management, analysis, and interpretation of the data; preparation,
278   review, or approval of the manuscript; or decision to submit the manuscript for publication.
279
280   **Data Sharing Statement:** The gene count data are available at
281   https://github.com/infectiousdisease-langelier-lab/LRTI_FABP4_GPT4_classifier. The code and
282   required source data are available at https://github.com/infectiousdisease-langelier-
283   lab/LRTI_FABP4_GPT4_classifier.

References

1.   The Top 10 Causes of Death. World Health Organization, 2021.
        (https://www.who.int/news-room/fact-sheets/detail/the-top-10-causes-of-death).

2.   Jain S, Self WH, Wunderink RG, et al. Community-Acquired Pneumonia Requiring
        Hospitalization among U.S. Adults. N Engl J Med 2015;373(5):415-27. DOI:
        10.1056/NEJMoa1500245.

3.   Langford BJ, Soucy JR, Leung V, et al. Antibiotic resistance associated with the COVID-
        19 pandemic: a systematic review and meta-analysis. Clin Microbiol Infect
        2023;29(3):302-309. DOI: 10.1016/j.cmi.2022.12.006.

4.   2022 Special Report: COVID-19 US Impact on Antimicrobial Resistance. Centers for
        Disease Control and Prevention, 2022. (https://stacks.cdc.gov/view/cdc/1190252022).

5.   Tsalik EL, Henao R, Nichols M, et al. Host gene expression classifiers diagnose acute
        respiratory illness etiology. Sci Transl Med 2016;8(322):322ra11. DOI:
        10.1126/scitranslmed.aad6873.

6.   Mick E, Tsitsiklis A, Kamm J, et al. Integrated host/microbe metagenomics enables
        accurate lower respiratory tract infection diagnosis in critically ill children. J Clin Invest
        2023;133(7). DOI: 10.1172/JCI165904.

7.   Lydon EC, Phan HV, Mick E, et al. Pulmonary FABP4 Is an Inverse Biomarker of
        Pneumonia in Critically Ill Children and Adults. Am J Respir Crit Care Med
        2024;210(12):1480-1483. DOI: 10.1164/rccm.202403-0516RL.

8.   van der Meer V, Neven AK, van den Broek PJ, Assendelft WJ. Diagnostic value of C
        reactive protein in infections of the lower respiratory tract: systematic review. BMJ
        2005;331(7507):26. DOI: 10.1136/bmj.38483.478183.EB.

9.   Self WH, Wunderink RG, Jain S, Edwards KM, Grijalva CG, Etiology of Pneumonia in
        the Community Study I. Procalcitonin as a Marker of Etiology in Adults Hospitalized With

Community-Acquired Pneumonia. Clin Infect Dis 2018;66(10):1640-1641. DOI:

10.1093/cid/cix1090.

10.   OpenAI. Introducing ChatGPT. (https://openai.com/index/chatgpt/).

11.   Zhou Y, Ong H, Kennedy P, et al. Evaluating GPT-V4 (GPT-4 with Vision) on Detection of

Radiologic Findings on Chest Radiographs. Radiology 2024;311(2):e233270. DOI:

10.1148/radiol.233270.

12.   Beaulieu-Jones BK, Yuan W, Brat GA, et al. Machine learning for patient risk

stratification: standing on, or looking over, the shoulders of clinicians? NPJ Digit Med

2021;4(1):62. DOI: 10.1038/s41746-021-00426-3.

13.   Maillard A, Micheli G, Lefevre L, et al. Can Chatbot Artificial Intelligence Replace

Infectious Diseases Physicians in the Management of Bloodstream Infections? A

Prospective Cohort Study. Clin Infect Dis 2024;78(4):825-832. DOI: 10.1093/cid/ciad632.

14.   Lee P, Bubeck S, Petro J. Benefits, Limits, and Risks of GPT-4 as an AI Chatbot for

Medicine. Reply. N Engl J Med 2023;388(25):2400. DOI: 10.1056/NEJMc2305286.

15.   Goh E, Gallo R, Hom J, et al. Large Language Model Influence on Diagnostic

Reasoning: A Randomized Clinical Trial. JAMA Netw Open 2024;7(10):e2440969. DOI:

10.1001/jamanetworkopen.2024.40969.

16.   Langelier C, Kalantar KL, Moazed F, et al. Integrating host response and unbiased

microbe detection for lower respiratory tract infection diagnosis in critically ill adults. Proc

Natl Acad Sci U S A 2018;115(52):E12353-E12362. DOI: 10.1073/pnas.1809700115.

17.   CDC/NHSN Surveillance Definition for Specific Types of Infections. Centers for Disease

Control and Prevention,

2021.(https://www.cdc.gov/nhsn/pdfs/pscmanual/17pscnosinfdef_current.pdf)

18.   Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for

RNA-seq data with DESeq2. Genome Biol 2014;15(12):550. DOI: 10.1186/s13059-014-

0550-8.

19.   Robin X, Turck N, Hainard A, et al. pROC: an open-source package for R and S+ to

analyze and compare ROC curves. BMC Bioinformatics 2011;12:77. DOI: 10.1186/1471-

2105-12-77.

20.   Wei J, Wang, X., Schuurmans, D., Bosma, M., Ichter, B., Xia, F., Chi, E., Le, Q.V., Zhou,

D. Chain-of-Thought Prompting Elicits Reasoning in Large Language Models. Advances

in Neural Information Processing Systems 35 (NeurIPS 2022): NeurIPS Proceedings

2022.

21.   Merenstein DJ, Barrett B, Ebell MH. Antibiotics Not Associated with Shorter Duration or

Reduced Severity of Acute Lower Respiratory Tract Infection. J Gen Intern Med 2024.

DOI: 10.1007/s11606-024-08758-y.

22.   Antibiotic Resistance Threats In the United States. U.S. Department of Health and

Human Services, 2019. (https://www.cdc.gov/antimicrobial-resistance/media/pdfs/2019-

ar-threats-report-508.pdf).

23.   Antimicrobial stewardship interventions: a practical guide. World Health Organization,

2021. (https://www.who.int/europe/publications/i/item/9789289056267).

351 **Table 1. Clinical and demographic features of cohorts.**

	Derivation Cohort			Validation Cohort		
	LRTI	**No LRTI**	*P*	**LRTI**	**No LRTI**	*P*
N	37	49		25	22	
Age, years (Median, IQR)	65.0 (51.0 – 75.0)	62.0 (53.0-73.0)	0.81	53.9 (49.4 – 66.9)	58.5 (57.5 - 63.9)	0.69
Female Sex (No., %)	11 (30)	30 (61)	0.0074	12 (48)	8 (36)	0.61
Race (No., %)			0.91			0.10
White	18 (49)	24 (49)	-	5 (20)	10 (46)	-
Black/African American	4 (11)	5 (10)	-	2 (8)	1 (5)	-
Asian	8 (22)	9 (18)	-	3 (12)	4 (18)	-
Native Hawaiian/Pacific Islander	1 (3)	0 (0)	-	1 (4)	2 (9)	-
Other/Unknown	6 (16)	11 (22)	-	14 (56)	5 (23)	-
Hispanic ethnicity (No., %)	5 (14)	11 (22)	0.44	10 (40)	5 (23)	0.40
Comorbidities (No., %)	36 (97)	45 (92)	0.54	24 (96)	17 (77)	0.14
Immunosuppressed (No., %)	8 (22)	6 (12)	0.38	1 (4)	3 (14)	0.51
Microbiologic diagnosis (No., %)						
Bacterial	28 (76)			3 (12)		
Viral	4 (11)			1 (4)		
SARS-CoV-2	0 (0)			13 (52)		
Fungal	1 (3)			0 (0)		
Multiple	4 (11)			8 (32)		
No-LRTI group cause of respiratory failure (No., %)			-			-
Surgery		14 (29)	-		6 (27)	-
Neurologic		12 (25)	-		7 (32)	-
Cardiovascular		8 (16)	-		4 (18)	-
Non-LRTI infection		5 (10)	-		6 (27)	-
Other		11 (22)	-		2 (9)	-
ICU admission diagnosis (No., %)			-			-
LRTI	36 (97)	23 (47)	-	25 (100)	7 (32)	-
No LRTI	1 (3)	26 (53)	-	0 (0)	15 (68)	-
Clinical team writing note (No., %)			0.20			<0.0001
Medicine	15 (41)	9 (18)	-	1 (4)	5 (23)	-
Critical Care	5 (14)	7 (14)	-	24 (96)	6 (27)	-
Neurosurgery	3 (8)	9 (18)	-	0 (0)	4 (18)	-
Cardiology	3 (8)	6 (12)	-	0 (0)	2 (9)	-
Other	11 (30)	18 (37)	-	0 (0)	5 (23)	-
Note to enrollment (No., %)			0.080			0.11
1 day	34 (92)	49 (100)	-	21 (84)	22 (100)	-
2 days	3 (8)	0 (0)	-	4 (16)	0 (0)	-
CXR to enrollment (No., %)			0.24			0.11
0 days	30 (81)	33 (67)	-	19 (76)	11 (50)	-
1 day	7 (19)	13 (27)	-	4 (16)	9 (41)	-
2 days	0 (0)	3 (6)	-	2 (8)	2 (9)	

Chi-squared test used for all categorical variables except time from note to enrollment for which Fisher exact test
was used. Wilcoxon rank-sum test used for continuous variables (age). IQR = interquartile range. One No LRTI
patient in the derivation cohort, and three No LRTI patients in the validation cohort, were adjudicated as having ≥1
etiology of respiratory failure. Four LRTI patients in the derivation cohort were adjudicated as having more than

one microbiologic diagnosis (three with viral-bacterial co-infection, and one with viral-fungal co-infection) and eight

LRTI patients in the validation cohort were adjudicated as having more than one microbiologic diagnosis (all with

SARS-CoV-2-bacterial co-infection).