Adaptive Responses to PARP Inhibition Predict Response to Olaparib and Durvalumab: Multi-omic Analysis of Serial Biopsies in the AMTEC Trial

Zahi I. Mitri[1,2,†], Allison L. Creason[3,4,†], Jayne M. Stommel[3,†], Tugba Y. Ozmen[3,6,†], Matthew J. Rames[3,6], Furkan Ozmen[3,6], Daniel Bottomly[3], Boyoung Jeong[3,6], Jeong Youn Lim[5], Shamilene Sivagnanam[6], Konjit Betre[6], Jinho Lee[3,7], Marilyne Labrie[8-10], Lisa M. Coussens[3,6], Christopher L. Corless[3,8], Shannon K. McWeeney[3,11], Gordon B. Mills[3,6*]

[1]British Columbia Cancer Agency, Vancouver, BC, Canada

[2]Division of Medical Oncology, University of British Columbia, Vancouver, BC, Canada

[3]Knight Cancer Institute, Oregon Health & Science University, Portland, OR, USA

[4]Department of Biomedical Engineering, Oregon Health & Science University, Portland, OR, USA

[5]Biostatistics Shared Resource, Oregon Health & Science University, Portland, OR, USA

[6]Department of Cell, Developmental and Cancer Biology, Oregon Health & Science University, Portland, OR, USA

[7]Knight Diagnostic Laboratories, Oregon Health & Science University, Portland, OR, USA

[8]Department of Immunology and Cell Biology, Faculty of Medicine and Health Sciences, Université de Sherbrooke, Sherbrooke, Canada

[9]Centre de Recherche du Centre Hospitalier de l'Université de Sherbrooke (CRCHUS), Sherbrooke, Canada

[10]Institut de Recherche sur le Cancer de l'Université de Sherbrooke (IRCUS), Sherbrooke, Canada

[11]Division of Oncological Sciences, Oregon Health & Science University, Portland, OR, USA

[†]These authors contributed equally to this work.

[*]Corresponding Author

ABSTRACT

In syngeneic murine breast cancer models, poly(ADP-ribose) polymerase inhibitor (PARPi) and anti-PD-L1 combinations induce deep, sustained responses independent of *BRCA1* or *BRCA2* mutation (BRCAm) status. We therefore investigated this combination in the AMTEC clinical trial, in which a one-month olaparib run-in was followed by combined olaparib and durvalumab in participants with non-BRCAm metastatic triple negative breast cancer. To characterize adaptive responses to olaparib monotherapy, paired biopsies taken before and during the PARPi lead-in were deeply characterized by DNA, RNA, and protein multi-omic analyses, including spatially resolved single-cell proteomics for tumor and immune contexture. We identified multiple potential tumor-intrinsic and microenvironmental biomarkers from pre-treatment and on-olaparib biopsies that robustly predicted participant response to combined olaparib and durvalumab. Notably, the on-olaparib biopsy provided the greatest information content, suggesting that early adaptations of malignant and immune cells to PARPi can serve as a predictor of potential benefit from combined PARPi and anti-PD-L1 therapy.

INTRODUCTION

Triple negative breast cancer (TNBC), comprising 15-20% of all breast cancers, predominantly affects younger women and black and Hispanic ethnic groups. Compared to other breast cancer subtypes, TNBC is characterized by aggressive features, a higher propensity for metastasis, and an overall worse prognosis[1]. Despite advancements in understanding TNBC biology, outcomes for metastatic TNBC (mTNBC) remain poor, with an overall survival of 1-2 years with standard therapies[2-4].

The DNA damage response (DDR) pathway is frequently dysregulated in TNBC, with nearly half of all cases exhibiting aberrations. *BRCA1* or *BRCA2* mutations (BRCAm) are observed in about 15% of TNBC, while another 45% of TNBC with genomic instability are non-BRCAm, suggesting that other aberrations can lead to defects in this pathway[1,5-7]. DDR dysregulation renders TNBC sensitive to DNA damaging therapies such as chemotherapy, radiation therapy, and more recently, poly(ADP-ribose) polymerase (PARP) inhibitors (PARPi). PARPi are of particular interest in mTNBC because of the high frequency of defects in homologous recombination (HR) repair, the high-fidelity repair pathway for double-stranded DNA breaks[8].

In mTNBC, the PARPi olaparib and talazoparib, FDA-approved for patients with germline BRCAm, improve clinical outcomes primarily by exploiting synthetic lethality with defective HR (HRD). However, even in this population, responses to PARPi monotherapy are often short-lived, with most patients eventually experiencing disease progression[9,10]. Moreover, PARPi monotherapy has limited activity in non-BRCAm tumors, which constitute the majority of mTNBC[11], with the VIOLETTE trial showing minimal benefit in these patients[12]. Preclinical and early clinical data suggest that PARPi combination therapies are safe and more effective in non-BRCAm TNBC, highlighting potential opportunities for these treatments for this population[13-15].

Immune checkpoint blockade (ICB) has improved outcomes for various tumor types, especially those with high tumor mutational burden (TMB) and neoantigenic propensity, which can bolster anti-tumor immunity[16,17]. In mTNBC, the combination of the programmed cell death 1 (PD-1) inhibitor pembrolizumab and chemotherapy is FDA-approved for patients with tumors expressing programmed cell death ligand 1 (PD-L1). However, this indication excludes the majority of mTNBCs, which are frequently PD-L1 negative, and clinical benefit is often brief even in PD-L1 positive patients[18]. Additionally, TMB

46 in TNBC is usually below the generally accepted 10 mutations/Mb that predicts ICB efficacy in other
47 cancer types[19].
48
Extensive preclinical data demonstrate that PARPi induce adaptive responses in malignant cells and the
tumor ecosystem, including immune cells, in models of HR-deficient and HR-competent tumors[20-24].
Similar adaptive changes have been observed in human tumors, such as breast cancer, under therapeutic
stress induced by PARPi[14,15,25,26]. Although adaptive responses to PARP inhibition vary among patients,
they appear to involve a limited set of stress response pathways, suggesting that identification of patient-
specific adaptive responses is feasible. Targeting these adaptive responses – for example, by increasing
PARPi-induced stress, blocking protective responses, or capitalizing on the rewiring of malignant cells
and the immune microenvironment[27] – has the potential to improve responses to PARPi, including in non-
BRCAm mTNBC. The identification of the specific adaptive response to target may require collecting
tumor biopsies before and during PARPi monotherapy, then cataloguing drug-induced changes that could
be exploited by a combined treatment regimen.
60
Combining PARPi with ICB is a promising therapeutic approach to leverage PARPi-induced adaptive
changes in immune signaling. PARP inhibition enhances antigen presentation, recruits cytotoxic T
lymphocytes, and activates the stimulator of interferon genes (STING) pathway[21,22,28], all of which could
increase the efficacy of ICB. Notably, PARPi can activate the STING pathway in non-BRCAm syngeneic
cancer models that are DNA repair competent[23], potentially leading to an infiltration of adaptive immune
cells that would recognize neoantigens generated through low fidelity DNA repair. Indeed, clinical trials
combining ICB and PARPi showed activity in both BRCAm and non-BRCAm mTNBC, although
response rates are lower in the wild-type group[14,29-31].
69
Building on these findings, we initiated the Adaptive Molecular Therapy of Evolving Cancers (AMTEC)
trial to evaluate the efficacy of combining the PARPi olaparib with the PD-L1 inhibitor durvalumab in
non-BRCAm mTNBC. This trial includes an olaparib monotherapy phase followed by a combined
olaparib and durvalumab phase, with biopsies collected at baseline and during olaparib monotherapy. Both
biopsies were analyzed with a suite of multi-omic assays to provide an in-depth examination of changes
in malignant cells and the tumor ecosystem in response to PARPi-mediated therapeutic stress. Using this
approach, the AMTEC trial was designed to identify adaptive changes induced by PARPi monotherapy
that could serve as biomarkers for selecting patients likely to benefit from PARPi/ICB combination
therapy in follow-on trials. An additional goal was to discover resistance mechanisms to combined
PARPi/ICB, which could be targeted with alternative PARPi combinations. AMTEC closed to enrollment
in 2024 with n=24 participants enrolled. Here we report on the final clinical and initial translational results
for the AMTEC trial.
82
RESULTS
AMTEC is a single-arm, open-label phase II trial assessing the efficacy of olaparib combined with
durvalumab in participants with non-BRCAm mTNBC (Fig. 1a). Eligible participants undergo a pre-
treatment biopsy upon enrollment, followed by a 4-week olaparib treatment. At the end of this olaparib
lead-in, all participants undergo a second biopsy on olaparib, and then durvalumab is added to the
treatment regimen. The combination of olaparib and durvalumab continues until progression or toxicity-
required termination.
90

91 The primary endpoint of the study is the overall response rate (ORR) for olaparib plus durvalumab.
92 Secondary endpoints include clinical benefit rate (CBR), duration of response, progression-free survival
93 (PFS), and overall survival (OS), as well as safety and tolerability of the treatment.
94
Participant and Disease Characteristics
Twenty-four participants with biopsy-proven mTNBC were enrolled in the AMTEC trial between January
2019 and May 2024. Participant characteristics are summarized in Table 1. All participants had an ECOG
performance status of 0 or 1. Similar numbers of participants had 0, 1, or 2 lines of prior therapy in the
metastatic setting, including seven participants who had received platinum-based regimens. Seven
participants had controlled brain metastases at the time of enrollment.
101
Of the 24 enrolled participants, three participants were excluded, resulting in a set of 21 participants in
the primary endpoint ORR analysis (Extended Data Fig. 1a). An additional participant voluntarily
withdrew during olaparib/durvalumab cycle 3 per personal preference. This participant ("S"; Fig. 1b)
achieved a RECIST best response of partial response (PR)[32] on first restaging and was included in the
ORR analysis but omitted from survival analyses, resulting in a total of 20 participants evaluable for the
survival efficacy endpoints (PFS and OS).
108
Efficacy Endpoints
Among the 21 participants who received the combination therapy and were evaluable for the primary ORR
efficacy endpoint, there were 5 PRs and 1 complete response (CR), for an ORR of 29% (Fig. 1b and
Extended Data Table 1a). Four additional participants had stable disease (SD) ≥6 months, for a CBR of
48%. Two participants had SD as best response on first restaging scan but progressed prior to the 6-month
timeline. Of note, two participants with PR (Participant D) and CR (Participant A) as best response on
study were off therapy for 26 and 19 months respectively (Extended Data Fig. 1b). Participant D remains
without evidence of disease progression and has not required additional systemic therapy, whereas
Participant A eventually recurred and died of progressive metastatic disease. These outcomes mirror
findings from our prior three-patient pilot study with the same design as AMTEC, which included one
prolonged PR, one SD, and one PD[14].
120
The best percentage change from baseline for the sum of all target lesion diameters was calculated for
each efficacy-evaluable participant (Fig. 1c). Among the six participants with SD as best response,
percentage changes ranged from 0% to −27%. The four participants with PR as the best response showed
reductions from −30% to −47%, and the single participant with CR exhibited a −100% reduction. In
contrast, the five participants with progressive disease (PD) demonstrated increases ranging from +23%
to +165%. Participant W lacked target lesion size measurements at the time of progression, and percent
change could not be determined.
128
The median PFS for the entire cohort was 4.8 months (95% CI 2.17 – 8.97 months), and median OS was
15.5 months (95% CI 7.85 – 18.1 months; Fig. 1d,e). Participants with a best response of SD vs. PR did
not show statistically significant differences in PFS or OS (PFS for PR vs. SD, p=0.97; OS, p=0.68).
Therefore, for subsequent analyses, cases with SD, PR, or CR (n= 12) were grouped as participants who
derived clinical benefit from the study therapy ("Responders"; R). These were compared to participants
with PD as best response (n=9), who were considered to have primary resistance to the study therapy
("Non-Responders"; NR).
136

Participants with CR, PR, or SD as RECIST best response had significantly better survival outcomes than those with PD. Specifically, participants with CR/PR/SD had improved PFS (9 vs. 2 months, p<0.0001) and OS (18 vs. 5 months, p=0.00049) compared to PD patients (Fig. 1f,g). The marked survival differences between response groups justified efforts to uncover molecular markers that differentiate responders from non-responders and predict clinical benefit, with the potential to guide biomarker evaluation in subsequent trials.

Clinical outcomes were not associated with the total number of prior therapies (p=0.8, R vs. NR) or with prior platinum treatment in the metastatic setting (PFS, p=0.85; OS, p=0.26). Among participants with a history of CNS metastases, CNS progression occurred concurrently with the progression of extra-cranial systemic disease.

Safety Results

All 24 participants who received at least one dose of study-directed therapy were evaluable for safety endpoints (Extended Data Table 1a,b). Most treatment-related adverse events (AEs) were mild to moderate (Grade 1-2 = 190/210; 90%). Thirteen participants (54%) experienced severe AEs (Grade 3), with 20/41 (49%) considered as possibly related to the study treatment. No Grade 4 or 5 AEs attributed to study interventions or treatment were noted.

Among the Grade 3 AEs, two consisted of immune-related elevated liver enzymes attributed to durvalumab, which resolved following dose hold and supportive medications (Extended Data Table 1b). The most common likely olaparib-related Grade 3 AE was anemia in 6/24 participants (25%), which resolved with a dose hold and adjustment per protocol. The two Grade 5 AEs included a motor vehicle accident ("Investigations – other*") and respiratory failure, both unrelated to treatment.

Clinical and Exploratory Research Analytics

To identify patients likely to benefit and to uncover targetable mechanisms of response and resistance, we conducted a comprehensive exploratory analysis of tumor-intrinsic and microenvironmental features in both the pre-treatment and on-olaparib tumor biopsies. This included a focus on signatures that emerged as adaptive responses to the olaparib monotherapy lead-in. In addition to the 21 participants treated on the AMTEC trial, we included three mTNBC participants from our previously reported pilot study who received the same sequence of biopsies and therapy as in AMTEC[14]. This approach serves to increase our power to identify potential biomarkers predictive of response to the olaparib and durvalumab combination (n=24).

Given the relatively small sample size and the absence of an independent test set with comparable study design and multi-omic profiling, we implemented a conservative analytical approach for the high-dimensional omics data (RNAseq, RPPA, mIHC, DSP). We employed the Boruta algorithm for feature selection, which incorporates Bonferroni correction to control for false discovery rate in multiple hypothesis testing. Predictive performance was then validated using F1 scores calculated across iterative stratified train-test splits, ensuring a balanced precision-recall trade-off and minimizing the risks of artifacts from feature selection (see Methods for details).

The clinical and exploratory analytical methods included: 1) clinical immunohistochemistry (IHC); 2) *GeneTrails®* DNA sequencing of 225 cancer-relevant genes; 3) whole exome DNA sequencing (WES); 4) bulk transcriptomic sequencing (RNAseq) for gene rearrangements and expression levels; 5)

NanoString GeoMx® Digital Spatial Profiling (DSP) of 46 proteins and phospho-proteins; 6) reverse phase protein arrays (RPPA) of 480 proteins, including post-translational modifications; 7) RAD51 foci analysis to measure homologous recombination deficiency, and 8) multiplexed immunohistochemistry (mIHC) using 23 antibodies to quantify infiltrating immune cells and their functional states. Due to sample availability and assay-specific technical or biological requirements, such as minimum tumor content and tissue adherence, not all assays were performed on every sample.

Extended Data Fig. 1c and Extended Data Tables 1c,d summarize the results of these clinical and exploratory research analyses, discussed in detail below. Extended Data Fig. 1c shows clinical metadata and assay results for each sample, noting statistically significant associations with best response (R), PFS (P), OS (O), as well as two paired-biopsy analyses. "B" denotes within-group paired comparisons, showing whether samples from responders and non-responders changed from pre-treatment to on-olaparib when each group is analyzed separately. "D" denotes between-group differences in olaparib-induced changes, testing whether the magnitude of change from pre-treatment to on-olaparib differs between responders and non-responders. Extended Data Tables 1c,d list the features that were statistically significant predictors of response to the olaparib-durvalumab combination, including candidate biomarkers selected by Boruta with high F1 scores within each data type (see Methods for details of the statistical analysis).

Malignant Cell-Intrinsic Analytes
The following measurements did not show statistically significant correlations with participant outcomes: 1) mutations in *TP53*, the HRD pathway, or the MAPK pathway; 2) TMB (no sample exceeded 10 mutations/Mb); 3) percentage of malignant cells expressing PD-L1 or Ki-67 by clinical IHC; 4) biopsy site; or 5) tumor content, although this tended to be higher in participants with PD.

The most frequently mutated gene was *TP53*, observed in 22 out of 24 participants (92%; Extended Data Fig. 1c). Of the genes known to confer sensitivity to PARPi[33,34], pilot study Participant B was PR and had an unusual germline deletion of exons 13-15 of *BRCA1* that was not detected on initial screening. In contrast, Participant V was PD with somatic *BRCA1* copy loss (0.5 copies). Notably, this participant also had copy loss of *PALB2* (0.35 copies), homozygous deletions of *ATM* and *RAD51,* and 15-copy gain of AKT2, which may have contributed to their poor response. No other deleterious mutations in known PARPi-sensitizing genes *BRCA1*, *BRCA2, BARD1, PALB2, RAD51C, or RAD51D* were noted. Alterations in other genes implicated in homologous recombination (HR) included germline *ATM* mutations in Participants C (SD; p.R3008C) and R (PD; p.S99_I105dup) and a *CDK12* mutation in Participant K (SD; p.L123fs*3). Regarding mutations possibly related to durvalumab response, responding Participants I and H had deleterious *B2M* mutations (splice site and p.C45_N62del), and non-responding Participant P had low frequency compound *JAK1* mutations (D1003Y and A1005V) detected only in the on-olaparib biopsy.

Defects in HR are well-established predictive biomarkers for PARPi response, even in the absence of *BRCA1/2* mutations[35-38]. To assess HR deficiency in our predominantly non-BRCAm samples, we used an immunofluorescence-based assay that we developed previously to distinguish HR-competent (HRC) from HR-deficient (HRD) cells[39]. Cells in G2 phase were evaluated for the presence of functional HR repair complexes (RAD51 foci) and double-strand DNA breaks (γH2AX), then samples were categorized as HRC, HRD, or Intermediate (with a mixture of HRC and HRD cells) based on the ratio of HRD to HRC nuclei (Extended Data Fig. 2a-c). Strikingly, all participants with tumors classified as HRD (including participant (B) from the pilot study with an unusual *BRCA1* mutation) achieved a best response of CR, PR

229 or SD, including those that shifted from Intermediate to HRD in the second biopsy (Extended Data Fig1c,
230 2d). In contrast, no samples from participants with PD were classified as HRD in either biopsy.
231
Genomic scarring is a common indicator of HRD due to its strong association with *BRCA1/2* mutations
and other DNA damage mechanisms[40]. Given its association with HRD, we assessed the predictive value
of the MutSig3 signature in the WES from the pre-therapy biopsy. MutSig3 was significantly associated
with OS (p=0.03; Fig. 2a) but not with best response or PFS (note only limited samples had WES).
Interestingly, there was no clear correspondence between the HRD/HRC ratio and the MutSig3 signature
in samples with both assays (Extended Data Fig. 1c). This discrepancy may reflect the different aspects
of DNA repair biology captured by each assay: the HRD immunofluorescence assay measures ongoing
DNA repair deficiency, whereas the MutSig3 signature captures the historical mutational footprint or
"scar" of prior homologous recombination defects.
241
RPPA proteomics data were generated from a subset of samples, including 15 pre-treatment biopsies, 15
on-treatment biopsies, and 12 matched pairs. Protein PARylation (PAR), catalyzed by PARP upon binding
single-strand DNA breaks, was elevated in all pre-treatment biopsies from non-responders but in only 2
responder samples (Participant B and I; Extended Data Fig. 1c). Although PAR levels were significantly
higher at baseline in non-responders (p=0.0023 NR vs. R, Extended Data Fig. 2e), they decreased on
olaparib to levels similar to those in the responders (Pre- vs. On- paired t-test p=0.00091, Fig. 2b; NR vs.
R on-olaparib p=ns, Extended Data Fig. 2e), indicating that lack of clinical benefit was not due to
insufficient PARP inhibition.
250
Androgen Receptor (AR) expression has been shown to suppress immune activity across multiple tumor
types and attenuate the response to ICB in prostate cancer[41-43]. In the AMTEC study, participants with
nuclear AR expression >10% by clinical IHC were similarly less responsive to olaparib/durvalumab, with
AR positivity predictive for shorter OS in on-olaparib biopsies (p=0.036; Fig. 2c). High AR expression
(>70% positive) corresponded perfectly with Luminal Androgen Receptor subtype (LAR; see below) in
non-responders as previously observed[44], but was not predictive of response across the entire cohort,
suggesting the presence of multiple, distinct mechanisms of resistance to this therapeutic combination.
258
PI3K-AKT and Receptor Tyrosine Kinase (RTK)-MAPK pathway activation is associated with
therapeutic resistance, including to PARPi and ICB[15,45,46]. To evaluate whether activation of these
pathways influenced the response to olaparib/durvalumab, we catalogued PI3K-AKT and RTK-MAPK
gene alterations and pathway activation via NanoString DSP phosphoprotein profiling. PI3K-AKT gene
alterations were present in biopsies from 3 of 14 responders (21%) participants and 7 of 10 non-responders
(70%) (Extended Data Fig. 1c). These associations were statistically significant for response and PFS but
not OS (pre-treatment p=0.013, 0.018, and ns; on-olaparib ns; Extended Data Fig. 2f). *PTEN* copy number
loss alone was significantly associated with best response, PFS, and OS (on-olaparib p=0.029, 0.008, and
0.009; Extended Data Fig. 2g), indicating that some PI3K-AKT pathway mutations may be passengers
with negligible impact on therapeutic outcome. RTK-MAPK pathway aberrations were present in only
two participants: one with SD and one with PD as best response.
270
271 Samples with elevated phosphoproteins in the PI3K-AKT signaling pathway (p-AKT1 plus p-GSK3B)
272 frequently harbored concurrent pathway gene alterations: 9 of the 10 pathway-activated samples had a
273 PI3K-AKT pathway gene alteration, including all 9 non-responders (Extended Data Fig. 1c). Notably,
274 however, 8 samples with PI3K–AKT gene aberrations lacked detectable pathway activation,

demonstrating the importance of phosphoprotein profiling in determining relevant downstream effects of gene alterations. Elevated protein phosphorylation accurately predicted poor therapeutic response, particularly in the on-olaparib biopsies: p-GSK3B alone was significantly associated with best response and PFS (p= 0.008 and 0.047; Extended Data Fig. 2h), while PI3K-AKT pathway activation correlated with response, PFS, and OS (p=0.011, 0.026, 0.022; Fig. 2d, Extended Data Fig. 2i)

Although RTK-MAPK pathway activity alone was not associated with clinical outcomes, a combined PI3K-AKT/RTK-MAPK pathway protein signature (AKT, p-GSK3B, p-PRAS40, and p-HER2) was associated with PD and reduced PFS and OS ("Mixed Signaling Pathway"; Fig. 2e, Extended Data Fig. 1c, 2j). Notably, activation of this combined signaling pathway in the on-olaparib biopsy was strongly associated with poor clinical outcomes, including best response, PFS, and OS (on-olaparib p<0.001, p=0.001, and p=0.0003), suggesting these signaling proteins may be activated as an adaptive response to olaparib stress. Indeed, the mixed signaling signature demonstrated high predictive accuracy for identifying non-responders, with a positive predictive value (PPV) of 0.80 in pre-treatment biopsies and 1.00 in on-olaparib samples.

Immune Contexture Analyses
In parallel with our analyses of gene and protein expression within malignant cells, we deeply evaluated the tumor microenvironment to identify features predictive of benefit from the olaparib/durvalumab combination. Previous work has shown that olaparib alters the immune landscape in preclinical models and patient tumors, and that immune contexture is a key determinant of durvalumab activity[47]. We therefore emphasized immune contexture in our microenvironment analyses.

As expected, analysis of the RNAseq, mIHC, and DSP data identified multiple immune features and markers associated with PFS, OS, and RECIST best response, including gene expression signatures, cell phenotypes, and protein markers, (summarized in Extended Data Fig. 1c and Extended Data Table 1d; only features that passed Boruta feature selection with Bonferroni adjustment and showed robust performance based on the harmonic mean of precision/recall (F1 score) are shown). Notably, the specific predictors varied from platform-to-platform, likely reflecting technical and assay-specific differences we present in the Discussion below.

To identify cross-platform response predictors that minimize the technical nuances of each assay platform, we first summarized the overall immune activation state within each platform by combining the most predictive individual markers and signatures (see Methods and below). Then, we integrated these single-platform summaries to generate a multi-omic immune consensus signature across platforms.

For the single platform summaries, the RNAseq summary signature was defined as high enrichment scores in at least 10 of the 17 significant immune RNA signatures ("IMMUNE RNA SUMMARY"; Extended Data Fig. 1c), the mIHC summary signature was defined as high levels of 9 or 10 mIHC cell types ("IMMUNE mIHC SUMMARY"), and the DSP consensus immune signature was defined as having high levels of at least 40% of the Immune DSP proteins ("IMMUNE DSP SUMMARY"; 40% was used due to variability in the number of successful DSP assays per sample). Each of these immune summary signatures was significantly associated with favorable PFS, OS, and RECIST best response, with stronger associations observed in the on-olaparib biopsies across the platforms (Fig. 3a-c; Extended Data Fig. 3a-c). Notably, only the immune DSP summary signature was predictive in untreated samples.

The sample-level summary calls were highly concordant across platforms: 83% for RNA vs. mIHC, 79% for RNA vs. DSP, and 77% for mIHC vs. DSP (Extended Data Fig. 1c: compare "IMMUNE RNA SUMMARY", "IMMUNE mIHC SUMMARY", and "IMMUNE DSP SUMMARY"; see Methods for threshold selection). This demonstrates that combining predictors within each assay platform can effectively compensate for technical differences across modalities, as detailed in Discussion.

Importantly, the multi-omic immune consensus signature – defined as positive for at least one of the single-platform immune summaries – strongly predicted clinical outcomes, including best response (p<0.001), OS (p=0.0005), and PFS (p=0.002) in the on-olaparib biopsy ("MULTI-OMIC IMMUNE CONSENSUS"; Fig. 3d, Extended Data Fig. 3d). The predictive value in the on-therapy biopsy was greater than the pre-therapy biopsy (PPV: 0.91 on- vs. 0.77 pre-); however, the pre-therapy biopsy was still predictive of OS but not best response or PFS. Taken together, evaluation of immune contexture, particularly within the on-olaparib biopsy, provides a compelling ability to predict clinical response to combined olaparib/durvalumab therapy.

The pathways and cell types identified by the combined Boruta/F1 analysis and incorporated into the consensus signatures are presented in Extended Data Figure 3. Several RNA-based GSEA Hallmarks – Inflammatory Response, Allograft Rejection, Interferon Gamma Response, IL6/JAK/STAT3 Signaling, IL2/STAT5 Signaling, and Interferon Alpha Response – were variably associated with RECIST best response, PFS, and OS. Multiple adaptive immune cell populations inferred by GSVA were also variably associated with clinical outcomes. Among T cells, Activated CD8$^+$ T Cells, Effector Memory T Cells, T Helper Cells, and Cytotoxic Cells, were most prominent. Regulatory T Cells, typically viewed as immunosuppressive, were unexpectedly associated with favorable RECIST best response in the on-olaparib biopsy (Extended Data Fig. 3n). B Cells were also predictive of response in this analysis (Extended Data Fig. 3u).

Among innate immune populations, Immature Dendritic Cells and NK CD56^{dim} natural killer cells were identified as potential predictors (Extended Data Fig. 3l,s), suggesting that activation of the innate immune response may contribute to benefit from olaparib/durvalumab. This is particularly noteworthy in participants H and I, who despite having a good response to therapy, had *beta-2 microglobulin (B2M)* mutations that are predicted to significantly impact B2M protein function, potentially rendering tumor cells resistant to adaptive immune cell killing. Indeed, the only significant RNA-based immune cell signature in participant H was NK CD65^{dim} natural killer cells.

Potential biomarkers on the DSP and mIHC platforms were generally consistent with those identified on the mRNA platform, including associations in the on-olaparib but not pre-therapy biopsy. These include Activated CD8$^+$ T Cells, Ki-67$^+$ CD8$^+$ T Cells, FOXP3$^-$ CD4$^+$ T Cells (but not FOXP3$^+$ CD4$^+$ T cells) and CD20$^+$ B Cells on the mIHC platform, and high CD45 and CD3 protein levels in both the tumor and stroma compartments on the DSP platform. Interestingly, CD8 and granzyme A (GZMA) were significant only in the stroma compartment by the Boruta/F1 approach. Whether this reflects differential localization of these populations or a technical challenge with the small number of samples remains to be elucidated.

PD-L1 tumor cell positivity was assessed by IHC using two different cutoffs due to discrepancies in the literature[48]. PD-L1 tumor proportion scores ≥10% were observed in at least one biopsy from 7 of the 24 participants, including 5 of 14 (36%) demonstrating clinical benefit (CR/PR/SD) and 2 of 10 (20%) participants who did not (PD; Extended Data Fig. 1c). Using a lower cutoff of ≥1%, positive PD-L1was

367 observed in at least one biopsy from 13 of the 24 participants, including 8 of 14 (57%) demonstrating
368 benefit (CR/PR/SD) and 5 of 10 (50%) who did not (PD). Although not statistically significant,
369 participants with PD-L1 ≥1% were more likely to achieve CR/PR/SD, aligning with other studies
370 indicating an association between PD-L1 positivity and response to ICB in mTNBC[49,50].
371
372 In contrast, PD-L1 assessed by DSP was significantly predictive of best response, with similar associations
373 in the pre- and on-olaparib biopsies (p= 0.022 and 0.035). The greater predictive value of DSP PD-L1
374 may reflect the platform's higher sensitivity and linear quantitative range compared with IHC, which is
375 reported as a dichotomous variable (positive or negative). Differences may also reflect variability in
376 pathologist-interpreted positivity thresholds, or slight differences in sample sets.
377
378 The only immune-related predictor identified by the combined Boruta/F1 analysis from the RPPA
379 platform was ZAP-70, likely reflecting the presence of T cells in the samples (Extended Data Fig. 3mm).
380 The absence of other immune markers may relate to the bulk nature of the RPPA tissue specimens, which
381 included tumor, immune, and adjacent normal cells, possibly diluting immune-specific signals.
382
Olaparib-induced changes
In the analyses above, biopsies from participants with better responses and longer survival tended to have
higher levels of malignant cell-intrinsic and immune-related features than those from non-responders,
suggesting that outcomes may be influenced by adaptive changes in both tumor and infiltrating immune
cells in response to olaparib. To investigate this further, we examined olaparib-induced changes in
matched pairs of pre- and on-olaparib biopsies to identify statistically significant changes correlated with
RECIST best response.
390
As shown in Fig. 4 and Extended Data Fig. 4, olaparib treatment changed multiple immune-related
markers in matched biopsy pairs. Markers that increased in responders but decreased in non-responders
include Activated $CD8^+$ T Cells RNA, Immune Cells by mIHC, and stroma HLA-DR by DSP (p=0.00086,
p=0.0018, and p<0.001; Fig. 4a). CD3, CD4, and HLA-DR in tumor DSP increased in responders but
remained unchanged in non-responders (p=0.002, p<0.001, and p<0.001; Fig. 4b). Effector Memory T
Cells RNA and Granzyme A (GZMA) in tumor DSP were unchanged in responders but decreased in non-
responders (p=0.0016 and p<0.001; Fig. 4c). Finally, the RPPA protein B7-H3, which suppresses T cell
activation, decreased in responders but increased in non-responders (p<0.001; Fig. 4d). Although not
significant in the comparison of change magnitude between response groups, PD-L1 significantly
increased on olaparib in non-responders but remained unchanged in responders by paired biopsy t-test
(Extended Data Fig. 4g).
402
Protein markers associated with malignant cells also changed differentially between responders and non-
responders. RPPA proteins HSP70, COUP-TFII, and SNAIL increased in responders and decreased in
non-responders (p=0.002, p=0.004, and p<0.001; Fig. 4e), while GLI1 only increased in responders
(p<0.001; Fig 4f). In contrast, p-GSK3β (DSP) and MTCO1 and MMP14 (RPPA) showed the opposite
pattern, decreasing in responders and increasing in non-responders p=0.023, p<0.001, and p<0.001; Fig.
4g.). Similar to Fig. 2, PAR decreased in both groups but showed a significantly greater reduction in non-
responders (p=0.011; Fig. 4h). Taken together, the association of olaparib-induced molecular changes
with patient response underscores the importance of evaluating both pre- and on-treatment samples to
better predict outcomes and guide patient selection for optimal therapy.
412

Analyses using predefined biomarker algorithms

We next applied two TNBC marker classification algorithms derived from independent cohorts to the AMTEC pre- and on-olaparib samples to determine whether they could predict response. TNBC classifiers that use bulk RNAseq data to define luminal, basal, mesenchymal, and immune states have been refined based on our studies to include four subtypes: basal-like immune-activated (BLIA), basal-like immune-suppressed (BLIS), luminal androgen receptor (LAR), and mesenchymal (MES) [51,52]. We have since implemented and validated this updated classification algorithm within the CLIA/CAP certified OHSU Knight Diagnostic Laboratories, adding an indeterminant (IND) category for samples with insufficient confidence to discriminate between BLIA and BLIS subtypes.

In the AMTEC cohort, tumors classified as BLIS or LAR in either the pre-treatment or on-olaparib biopsy had significantly poorer outcomes than those classified as BLIA or IND (non-BLIS/LAR), with decreased PFS and OS and a significant likelihood of RECIST best response of PD (on-olaparib biopsy p=0, p=0.005, and p=0.002; Fig. 5a, Extended Data Fig. 5a). Samples classified as BLIA were strongly associated with benefit: all five BLIA participants achieved a best response of CR/SD/PR, whereas all six BLIS/LAR participants had PD (Fig. 5b,c), indicating a 100% predictive value for these dichotomous predictors. IND samples were present in both PD and CR/PR/SD groups, though more frequently in responders. When the BLIS/LAR subset was compared with the combined BLIA/IND subset (non-BLIS/LAR), only three on-therapy biopsies (all IND) were misclassified as PD, corresponding to an 88% accuracy in predicting response. No MES tumors were present in the cohort.

Among the 19 participants with subtyping performed on both pre- and on-olaparib biopsies, four exhibited a change in TNBC subtype following treatment: two changed from IND to BLIS, one from BLIS to IND, and one from BLIA to IND (Extended Data Fig. 1c). All these changes involved at least one IND classification, reflecting cases where subtype assignment could not be confidently determined by the CLIA assay. Notably, these changes were clinically relevant, as the on-olaparib subtype had a higher predictive value for both survival and best response than the pre-treatment biopsy (Fig. 5a, and Extended Data Fig. 5a).

Given the strong immune activation associated with the BLIA subtype, we investigated whether its defining genes reflected an interferon-driven transcriptional program. A highly ranked subset of BLIA genes (centroid >0.5) was interferon-inducible, with a median fold change >2 based on data from interferome.org (Extended Data Fig. 5b). However, classical interferon-stimulated genes such as IRFs and IFITs were not among the top- ranked genes, suggesting that the BLIA classifier captures a novel interferon-induced gene set rather than the canonical interferon signature.

To determine which cell types express the genes defining the TNBC subtypes, we performed single cell RNAseq on a set of 13 advanced breast cancer specimens collected under an internal observational study[53]. BLIA-associated genes were predominantly expressed in immune cells, including various subtypes of T cells, NK cells, myelomonocytic subsets, and dendritic cells, consisting with their association with antigen presentation, T cell localization, and T cell killing (Extended Data Fig. 5c). Expression was minimal in other cell types, including cancer-associated fibroblasts, B cells, malignant cells, or normal epithelial cells. However, like the interferon-induced genes, classic T cell, macrophage, monocyte, or dendritic cell genes were not part of the BLIA signature.

The second TNBC marker classification algorithm evaluated was mTNBC3s, a biomarker signature we
previously derived from TCGA basal breast cancers that identifies a tumor-infiltrating B cell population
strongly associated with therapeutic response[54]. In our prior analyses of a subset of AMTEC samples,
mTNBC3s demonstrated both predictive and prognostic ability, and resolved the IND subtype within the
TNBC subtype classifier[54]. Applied here to the full AMTEC cohort, (excluding LAR samples, as
mTNBC3s was not developed for this subtype), mTNBC3s correctly classified predicted responders
(pSDPR) and non-responders (pPD) in 95% of pre-treatment samples (18/19) and 89% on-olaparib
samples (17/19), and significantly predicted response in both biopsies (pre-treatment $p<0.001$ and on-
olaparib $p=0.002$; Fig. 5d, Extended Data Fig. 1b). Among all pre-treatment markers evaluated, mTNBC3s
provided the strongest predictive signal for RECIST best response; however, its performance for PFS and
OS was surpassed by the on-olaparib biopsy (PFS pre-treatment $p=0.11$ vs. on-olaparib $p=0.0001$, Fig 5d;
OS pre-treatment $p=0.046$ vs. on-olaparib $p=0$, Extended Data Fig. 5d), indicating that olaparib-induced
immune remodeling adds additional predictive value.
471
A reduced three-protein response signature with clinical translation potential
The potential biomarkers that emerged as the strongest predictors of benefit in the earlier analyses were
derived from resource-intensive research platforms that use RNAseq, multi-gene expression signatures,
multiplexed antibody panels, and complex computational analysis pipelines. While these approaches yield
important biological insights, they are not suitable for the rapid, scalable, and regulated testing needed to
guide treatment decisions in routine clinical practice because they require specialized infrastructure,
technical expertise, long processing times, and most are not CLIA-compliant. To develop a more limited
biomarker approach that could feasibly be implemented in standard workflows, we turned to the immune
proteins quantified in our CLIA-validated DSP dataset to find markers that could be measured by routine
IHC. Using a training/test framework with ANOVA-based feature selection applied to the on-olaparib
samples, we identified a three-protein combination - CD3 (Tumor), CD45 (Tumor), and PD-L1 (Stroma)
- as the strongest discriminator of response and used it to train a logistic regression classifier. This three-
protein predictor achieved 100% accuracy in the on-olaparib cohort (Fisher's exact test $p<0.001$; Fig. 6a;
Extended Data Fig. 1c), and predicted responders (pR) exhibited significantly longer OS and PFS than
predicted non-responders (pNR) ($p=0.006$ and $p=0.0001$; Fig. 6a,b). Pre-treatment samples were classified
with lower, though still substantial, accuracy (84%) and showed weaker survival associations, consistent
with the greater predictive value of on-therapy biopsies and olaparib-induced immune remodeling.
489
The three-protein biomarker set can be measured using a CLIA-compliant assay and could potentially be
adapted to a simplified format suitable for broader clinical use, such as IHC. Importantly, when an on-
therapy biopsy cannot be obtained, this three-protein combination could still provide actionable
information for therapy selection from pre-treatment samples. Further validation in independent cohorts
is needed to determine whether this three-protein signature is broadly prognostic of therapeutic response,
or specifically predictive of patient benefit from olaparib–durvalumab therapy.
496
DISCUSSION
The AMTEC trial demonstrated that the combination of olaparib and durvalumab provides meaningful
clinical benefit in advanced mTNBC, even in patients without *BRCA1/2* mutations or other indicators of
HRD. The study regimen - consisting of one month of olaparib monotherapy followed by combined
olaparib and durvalumab - resulted in a median PFS of 4.8 months and OS of 15.5 months, with an overall
response rate of 29% and a clinical benefit rate of 48%. Notably, participants achieving partial response
or stable disease lasting ≥6 months had comparable PFS, indicating that CBR is a relevant measure of

efficacy for this combination. Treatment was reasonably well-tolerated, with toxicities resolving through
dose holds or reductions.
506
A major challenge in evaluating clinical trials with combination therapies is determining the relative
"contribution of components" to overall efficacy. The AMTEC trial design, which incorporated an initial
olaparib monotherapy window, enabled direct assessment of adaptive responses to PARP inhibition and
their relationship to subsequent benefit from the olaparib-durvalumab combination. Because a subset of
participants experienced marked and durable clinical benefit, we performed deep, integrated multi-omic
analyses of pre-treatment and on-olaparib biopsies to identify potential biomarkers of response, with a
focus on adaptive mechanisms induced by olaparib that might be leveraged by immune checkpoint
blockade.
515
Using this multi-omic DNA, RNA, and protein profiling approach, we identified markers of both primary
and adaptive resistance to PARPi, as well as predictors of clinical response to PARPi combined with anti-
PD-L1 in pre- and on-therapy biopsies. Most markers showed limited predictive value in pre-treatment
samples but were strongly associated with outcomes in on-olaparib biopsies, supporting the concept that
PARPi-induced adaptive responses are key determinants of therapeutic outcome. These findings suggest
that treatment strategies guided by evaluation of therapy-induced adaptive responses may optimize patient
outcomes with this regimen and potentially other drug combinations. To advance clinical translation, we
distilled the olaparib-induced response predictors into a set of three proteins measurable on the CLIA-
compliant DSP platform, potentially enabling broad implementation of these markers in clinical practice.
525
526　Among the markers we analyzed, several were not predictive of response to olaparib/durvalumab, despite
527　initial expectations. These include TMB; the percentage of Ki-67$^+$ cells by IHC; mutations in *TP53*,
528　*BRCA1/2* and other HRD pathway genes, or MAPK pathway genes; CD4$^+$ or CD8$^+$ T cell counts by IHC;
529　and percent tumor content.
530
Notably, PD-L1 expression by clinical IHC also did not significantly predict outcomes. Although higher
percentages of PD-L1$^+$ tumor cells were more often observed in responding participants and less common
in non-responders, this did not reach statistical significance. In contrast, PD-L1 protein levels quantified
by DSP were significantly associated with RECIST best response in both pre- and on-olaparib biopsies.
This improved performance likely reflects several advantages of DSP: (1) it is fully quantitative, reducing
observer variability; (2) it captures protein expression in both tumor and infiltrating immune cells; and (3)
it is sensitive to low-abundance protein, enabling detection of biologically meaningful differences that
may be missed by standard IHC.
539
Although mutations in the HR pathway were not predictive of response, MutSig3, which reflects a
genomic "scar" associated with HRD, was predictive of OS, consistent with HR status contributing to the
response to olaparib plus durvalumab. Interestingly, similar findings were reported in ovarian cancer,
where MutSig3, but not other HRD assays, was associated with outcomes following combined PARPi and
ICB[25]. Using a functional HRD assay that counts RAD51 foci exclusively in G2-phase cells with DNA
damage[39], we found that all HRD samples in our study came from participants who benefited from the
combination therapy, although HRD status did not significantly correlate with outcome, likely due to the
limited number of samples definitively categorized as HRD. Importantly, this assay revealed that many
tumors contained mixed populations of HRC and HRD cells, suggesting that intratumoral heterogeneity

in HR competence may undermine the predictive value of HRD biomarkers and adversely affect clinical
outcomes in PARPi-treated mTNBC patients.
551
In contrast, signaling pathway alterations showed significant associations with clinical outcomes. PI3K-
AKT pathway gene alterations, particularly *PTEN* copy loss, were associated with poor outcomes in both
pre-treatment and on-olaparib biopsies. Importantly, protein evidence of pathway activation through both
PI3K-AKT and RTK-MAPK more strongly predicted non-response than gene alterations, particularly in
the on-olaparib biopsies. These findings support the concept that signaling through these pathways is
activated as an adaptive response to PARP inhibition, thereby limiting the efficacy of olaparib plus
durvalumab. Consistent with this, these pathways have been implicated in responses to olaparib
combination therapies in both preclinical studies and clinical trials[15,26,33,46,55]. Notably, signaling activity
in our study was observed predominantly in samples lacking immune activation, suggesting that these
participants may benefit from PARPi combined with an agent targeting the PI3K or MAPK pathways,
rather than PARPi plus ICB.
563
The strongest predictors of response were related to the immune contexture, particularly immune features
in the on-therapy biopsy. We and others have shown in preclinical studies that PARPi generates double-
stranded DNA fragments that activate STING, leading to interferon production[22-24,28]. In the current study,
although a baseline interferon signature was associated with response, a high "Interferon Gamma
Response" RNA signature in the on-olaparib biopsy showed a stronger correlation with clinical benefit.
Because interferon signals through the JAK-STAT pathway, the association of the "IL6/JAK/STAT3
Signaling" RNA signature with outcomes supports the concept that olaparib-induced interferon signaling
contributes to therapeutic efficacy of the olaparib-durvalumab combination.
572
Interestingly, two participants had tumors with *B2M* mutations predicted to decrease class I HLA
expression and impair antigen presentation. Loss of B2M and HLA expression has been linked to
resistance to immune checkpoint blockade[56]. Despite this, both participants benefited from treatment, with
one showing evidence of constitutive NK cell activation and the other exhibiting an olaparib-induced
immune response consistent with the innate immune activation, in which type 1 interferons are potent
inducers.
579
Many predictive markers were associated with immune status in the on-olaparib biopsy, supporting the
contention that olaparib-mediated alterations in the immune contexture contribute to the response to the
olaparib and durvalumab combination. However, not all analyzed markers showed response-dependent
changes with olaparib treatment, indicating that the immune signature predictors likely reflect specific
changes in immune contexture induced by olaparib rather than sample set differences or differential tumor
heterogeneity between responders and non-responders. Importantly, the potential biomarkers identified in
the AMTEC trial paralleled those that we identified in preclinical studies as well as in prior clinical
trials[14,15,20-26]. This strongly supports the validity and mechanistic relevance of the identified biomarkers.
588
While immune indicators from the RNA, mIHC, and DSP platforms each predicted benefit from olaparib
and durvalumab, the specific markers identified differed across platforms. This is not unexpected, as the
mIHC and DSP platforms used different antibodies and were sometimes performed on different cores with
varying degrees of tissue heterogeneity. Moreover, mIHC measures protein expression at the single-cell
level, whereas DSP and RNA sequencing yield composite data from mixed cell populations within tumor-
rich regions. Given that RNA and protein levels are often poorly correlated, especially for proteins

regulated through post-translational modifications rather than gene expression[57], discrepancies across
platforms are anticipated. The limited number of samples, varying cut-off thresholds for positive versus
negative calls, and lack of consensus on markers used for cell type assignments likely also influenced the
statistical significance of individual markers. These factors highlight the need for further refinement of
platform-specific predictors in larger cohorts. Nevertheless, by combining multiple markers into platform-
specific summaries, our patient-by-patient analysis of immune activation states showed high concordance
across platforms.

We used two RNAseq-based algorithms developed from external data sets to predict therapeutic response
in our cohort: the BLIA/BLIS TNBC subtype classifier and the mTNBC3s outcome classifier[54]. The
BLIA/BLIS algorithm is particularly notable because it has been validated and implemented in a CLIA
environment, enabling its potential use for patient selection. TNBC subtype was a strong predictor of
benefit on the pre-treatment biopsy with BLIA/IND vs. BLIS/LAR p-values of 0.001 and 0.04 for PFS
and OS, respectively. Predictive accuracy further improved in on-olaparib biopsies, with p-values of 0 and
0.005 for PFS and OS, and PPV of pre-therapy biopsy 0.75 and on-olaparib biopsy 0.80. Thus, while not
optimal, pre-treatment TNBC subtyping provides substantial predictive value for selecting patients likely
to benefit from olaparib and durvalumab, especially when an on-treatment sample is unavailable.

However, clinical use of the TNBC subtyping classifier is confounded by the frequency of Indeterminant
results on the CLIA platform. Among participants with Indeterminate subtype calls on the on-olaparib
biopsy, most derived clinical benefit (7/10), though a subset did not (3/10). These findings underscore the
need to refine this algorithm to improve its predictive value in the CLIA environment.

The previously developed mTNBC3s algorithm proved highly effective in predicting response to the
olaparib and durvalumab combination in both pre-treatment and on-olaparib biopsies and, importantly,
was strongly predictive in samples classified as Indeterminate by the TNBC subtype assay. This suggests
that mTNBC3s may resolve a key limitation of the current CLIA-based BLIA/BLIS subtype test. Moving
mTNBC3s into the CLIA environment could therefore substantially improve identification of patients
most likely to benefit from PARPi and immune checkpoint inhibitor combinations. Further, because
mTNBC3s predicted response in baseline biopsies in our study and has previously shown to have
prognostic value[54], this algorithm, pending additional validation, may have utility in predicting benefit
beyond PARPi and immunotherapy.

Finally, we identified three proteins on the DSP platform – CD45, CD3, and PD-L1 – that, when
combined, were highly predictive of response in on-olaparib biopsies. Because DSP is CLIA-compliant,
this assay could be used for patient selection in future trials and warrants evaluation to determine whether
it can predict response to other therapeutic regimens. Recognizing that DSP technology is not yet widely
available, the small size of the analyte set also makes it feasible to implement alternative CLIA-compliant
approaches, such as IHC, which could allow broader application in clinical practice.

Taken together, our analyses of pre- and on-olaparib biopsies provide insight into the relative prognostic
and predictive value of the identified markers. Among the features we identified, those that were strongly
associated with response in the pre-treatment biopsy, such as TNBC subtype and mTNBC3s, may reflect
a mixture of prognostic and predictive information. Their performance before therapy suggests an inherent
prognostic component, while their stronger associations in the on-olaparib biopsy indicate added
predictive value driven by olaparib-induced biological changes. Indeed, both algorithms were developed

using TNBC sample sets from patients treated with a variety of therapeutic regimens that did not include a PARPi or ICB. In contrast, our study design, incorporating both pre-treatment and on-therapy biopsies, allowed us to identify markers, such as the Three-Protein Response Predictor, that were not evident at baseline but became predictive only upon olaparib exposure. The emergence of such on-treatment markers supports the idea that some predictors specifically capture adaptive biological changes induced by PARP inhibition rather than underlying disease biology. Thus, a key strength of this study is its ability to distinguish markers with general prognostic significance from those with predictive value for the olaparib-durvalumab combination.

Despite these promising insights, our study has several important limitations. The foremost is the limited number of participants and the lack of an independent test set. While the Boruta algorithm with a Bonferroni adjustment and an F1 statistic helped mitigate false positives, prospective validation of our potential predictors in larger trials is necessary. In addition, some assays were completed on only a subset of participants due to insufficient material or tumor content, despite obtaining up to six cores per biopsy. Further, DSP data were unavailable in some samples because the assay was upgraded mid-trial, and there was insufficient material to re-run analyses. Finally, several immune characterization assays were not concordant, and as discussed above, there are likely multiple reasons for these discrepancies that require further investigation.

In summary, the AMTEC trial demonstrated marked clinical activity with acceptable toxicity. Importantly, we identified a set of potential signatures that strongly predicted participant outcomes, with a subset already validated in the CLIA environment. While some predictive biomarkers were evident in pre-therapy biopsies, the greatest predictive value was observed in on-therapy samples, consistent with adaptive responses to PARP inhibition contributing to the efficacy of the olaparib–durvalumab combination. Taken together, these results demonstrate that on-therapy biopsies capture adaptive tumor responses that are not present at baseline, underscoring their potential to improve treatment selection and provide optimal patient outcomes.

Acknowledgements

This project was carried out with major support from the Oregon Health & Science University (OHSU) SMMART Clinical Trials Program, AstraZeneca, National Institutes of Health (NIH), National Cancer Institute (NCI) Human Tumor Atlas Network (HTAN) Research Center (U2CCA233280), Breast Cancer Research Foundation, and Prospect Creek Foundation. We thank all members of the SMMART Clinical Trials Program for support with sample collection and data management. We thank the OHSU Knight Cancer Institute BioLibrary for assistance with sample management.

Author Contributions

GBM, ZIM, and CLC conceived and designed the study. ZIM enrolled and treated participants. ALC, JMS, ZIM, DB, TYO, and FO generated visualizations and figures. ALC, JMS, JYL, TYO, and DB performed data and statistical analyses. ALC developed software and pipelines for primary data processing. GBM, JMS, ALC, ZIM, TYO, FO, DB, and SKM interpreted the data. SS, KB, JL, DB, ML, TYO, MJR, FO, and BJ generated data. ALC and JMS managed data collection, cleaning, and curation. GBM, JMS, ZIM, and ALC wrote the manuscript. GBM, CLC, SKM, and LMC supervised the project. GBM acquired funding. All authors reviewed and revised the manuscript.

Ethics Declaration

ZIM has received institutional grants/research support from AstraZeneca, Eli Lilly, and GSK; and received consulting fees / advisory boards from Gilead, Lilly, Merck, Novartis, AstraZeneca, and Daiichi Sankyo. LMC has received reagent support from Cell Signaling Technologies, Syndax Pharmaceuticals, Inc., ZielBio, Inc., and Hibercell, Inc.; holds sponsored research agreements with Syndax Pharmaceuticals, Hibercell, Inc., Prospect Creek Foundation, Lustgarten Foundation for Pancreatic Cancer Research, Susan G. Komen Foundation, and the National Foundation for Cancer Research; is on the Advisory Boards for Syndax Pharmaceuticals, Dispatch Pharmateuticals, Carisma Therapeutics, Inc., CytomX Therapeutics, Inc., Shasqi, Kineta, Inc., Hibercell, Inc., Cell Signaling Technologies, Inc., Alkermes, Inc., NextCure, Guardian Bio, AstraZeneca Partner of Choice Network (OHSU Site Leader), Genenta Sciences, Pio Therapeutics Pty Ltd., and Lustgarten Foundation for Pancreatic Cancer Research Therapeutics Working Group, Inc. CLC serves on the SAB for Cepheid. GBM has licensed technologies to Myriad Genetics and NanoString/Bruker; is on the SAB or is a consultant to Amphista, Astex, AstraZeneca, BlueDot, Ellipses Pharmaceuticals, ImmunoMET, Leapfrog Bio, Bruker/Nanostring, Neophore, Nerviano, Nuvectis, Pangea, PDX Pharmaceuticals, Qureator, Rybodyne, Signalchem Lifesciences, Turbine, Zentalis Pharmaceuticals; has sponsored research from AstraZeneca, Zentalis, Nanostring and Nerviano and has stock/options/financial interests in ImmunoMet and SignalChem; financial interest in Nerviano Medical and Turbine that may have a commercial interest in the results of this research and technology. This potential conflict of interest has been reviewed and managed by OHSU.

Methods

TRIAL OVERSIGHT
All biospecimens and data were collected under Adaptive Molecular Therapy of Evolving Cancers (AMTEC) study, ClinicalTrials.gov identifier NCT03801369. Written informed consent was obtained from all participants prior to enrollment. The Oregon Health & Science University (OHSU) Institutional Review Board approved the trial (IRB #18504). The study adhered to the ethical principles of the Declaration of Helsinki and followed the International Council on Harmonization guidelines on Good Clinical Practice, in compliance with all applicable laws, regulatory requirements, and conditions mandated by regulatory authorities and/or institutional review boards.

STUDY DESIGN
AMTEC was a single-center, single-arm, open-label phase II observational study evaluating the efficacy of olaparib in combination with durvalumab in participants with *BRCA1/2^wt* mTNBC (Fig. 1a). Participants with biopsy-confirmed mTNBC first underwent a pre-treatment biopsy, followed by a 4-week induction phase with olaparib (300 mg orally twice daily). A repeat on-olaparib biopsy was performed at Cycle 1, Day 28 (±7 days). After completion of the induction phase, durvalumab (1500 mg intravenously every 4 weeks) was added to continued olaparib. Treatment was planned for 12 cycles, with continuation allowed beyond 12 cycles in the absence of disease progression or unacceptable toxicity. An optional biopsy was offered at the time of progression. Participants were followed for up to one year after completing protocol-directed therapy.

ELIGIBILITY CRITERIA
Eligible participants must have biopsy-proven mTNBC defined as ER <10%, PR <10%, and HER2 non-amplified per ASCO/CAP guidelines[58]. Up to two prior chemotherapy regimens for metastatic disease were permitted. Additional inclusion criteria included ECOG performance status ≤1, tumors amenable to

serial biopsies, and no prior exposure to PARP inhibitors or immune checkpoint inhibitors in the metastatic setting.

Key exclusion criteria were active autoimmune disease requiring systemic steroids or disease-modifying therapy; history of HIV; active hepatitis B or C infection; and symptomatic central nervous system metastases or carcinomatous meningitis.

Of the 24 enrolled participants, three were excluded from the efficacy endpoint analyses: one did not receive durvalumab due to rapid disease progression, one had an orthopedic procedure unrelated to trial therapy, and one was ineligible based on hormone receptor expression on the on-treatment biopsy. This resulted in a n=21 participants for the primary endpoint ORR analysis, and n=24 for the safety analysis group.

ENDPOINTS
The primary endpoint was overall response rate (ORR), defined as the proportion of participants achieving complete response (CR) or partial response (PR) per RECIST v1.1 criteria[32]. Secondary efficacy endpoints included clinical benefit rate (CBR; CR, PR, or stable disease [SD] lasting ≥6 months), duration of response (DOR), progression-free survival (PFS), and overall survival (OS). Safety endpoints included the incidence of grade ≥3 adverse events as defined by CTCAE v5.0.

STATISTICAL ANALYSIS
This study used a Simon 2-stage Minimax design[59]. The null (ICB alone) and alternative (Olaparib + Durvalumab) hypotheses were: H_0: $\pi = 0.15$ and H_a: $\pi = 0.35$. For the primary endpoint, a total sample size of 28 participants will achieve 80% power to detect the ORR difference of 0.20 using the 2-stage Minimax design, with one-sided type I error =0.05.

In Stage 1, 15 evaluable participants (defined as those who received at least one dose of combination therapy) were enrolled (Extended Data Fig. 1a). The study would stop for futility if ≤2 responses were observed; otherwise, it would proceed to Stage 2. The regimen would be considered promising if ≥8 responses were observed among 28 participants. All Stage 1 participants were required to complete at least one post-baseline disease assessment prior to interim analysis.

AMTEC met criteria to proceed to Stage 2, with 4 of 15 evaluable participants achieving a PR. Ultimately, 24 participants were enrolled before the trial closed to accrual in August 2024 due to recruitment challenges and shifting strategic priorities of the pharmaceutical sponsor. With no additional enrollment planned, the current report summarizes outcomes for the full trial cohort.

SURVIVAL ANALYSIS
Survival curves for both progression-free survival (PFS) and overall survival (OS) were calculated using the Kaplan-Meier estimator and plain or log 95% confidence intervals. Cox proportional hazards models were fitted to estimate hazard ratios (HR). Reported unadjusted p-values are indicated for both the logrank score test and Cox regression model test to evaluate statistical differences between groups. When evaluating OS, data were right-censored for events that had not occurred before completion of the study and are indicated on OS curves (+). For all survival analyses, living participants were censored at the data cutoff of April 15, 2025.

 SAMPLE AVAILABILITY
Pre-treatment and on-olaparib biopsies were obtained from the 21 participants evaluable for the primary endpoints. Samples were placed in appropriate preservative (FFPE or flash frozen for RPPA) within 3 minutes to preserve post-translational modification, particularly phosphoproteins[60]. Assay success varied across samples and platforms; Extended Data Fig. 1c summarizes which assays were completed for each sample. Neither the pre-treatment nor the on-olaparib biopsy for Participant C had adequate tumor content for analysis, so Genetrails and IHC data from a biopsy obtained 90 days previously was used for analysis. The pre-treatment biopsy from Participant N contained insufficient tumor for assays other than IHC, and RNAseq failed for both samples from this participant. NanoString DSP was successful on most samples, as it requires limited tumor content. mIHC failed in a subset of samples that became detached from the matrix, a known problem with these assay platforms[61]. RPPA, which requires larger amounts of material, was performed only on a subset of participant samples.

To increase statistical power for marker and feature analyses, samples from a three-participant pilot study were also included[14]. This IRB-approved study (OHSU IRB #18239; NCT03544125) served as a precursor to the AMTEC trial and followed the identical study design, with samples collected at the same time points and participants receiving the same drugs and dosing. The participant IDs from this pilot study were B, J, and U (Extended Data Fig. 1c).

FEATURE SELECTION
Feature selection was conducted using two approaches for discrete vs. continuous data. For continuous values, the Boruta algorithm, a method built around random forests to compare feature importance against randomly permuted shadow features[62]. Unlike alternative feature selection methods that identify a minimal-optimal feature subset, Boruta identifies all features relevant for predicting outcome, which is useful for exploratory biomarker discovery in small sample sizes. This approach provides control against false positives while maintaining sensitivity to detect true associations. For each data type, Boruta was applied with Bonferroni multiple testing correction (mcAdj=TRUE) to adjust for the number of features tested at a significance threshold of $p < 0.05$. To reduce stochasticity, an iterative approach was implemented where Boruta was executed 50 times with different random seeds (seeds 1-50) at a more stringent threshold ($p < 0.01$) for both pre-treatment and on-olaparib biopsy timepoints. Features identified as "Confirmed" or "Tentative" across iterations were retained, and a final Boruta analysis (seed=42, $p < 0.05$) was performed to determine the final feature set.

For data with discrete labels, predictive performance was assessed using F1 score through repeated stratified train-test splits. For continuous data that was subsequently discretized, only features that passed the initial Boruta analysis were considered in the F1 score evaluation. Prior to analysis, optimal label mapping was determined by evaluating which assignment (e.g., Positive=Responder vs. Negative=Responder) maximized F1 score. For each feature, 20 independent splits were performed with a 70:30 train-test ratio using stratified sampling to maintain class balance. Performance metrics including F1 score, precision, recall, and accuracy were calculated for both training and test sets across all splits. A final independent test was conducted using a 60:40 split. Median F1 scores across splits were used to rank feature predictive power. Only features with an F1 score ≥70 are shown.

STASTICAL TESTING
Following feature selection with Boruta and F1 scores, statistical testing was conducted using student's t-test for continuous datasets or Fisher's exact tests for discrete datasets. Comparisons of continuous

824 variables between responder (R) and non-responder (NR) groups were performed using two-sided t-tests,
825 where non-responders were used as the reference group. Associations between categorical variables and
826 treatment response were evaluated using Fisher's exact test. Unadjusted p-values were reported with
827 adaptive precision for up to 4 decimal places.
828
CLINICAL IMMUNOHISTOCHEMISTRY (IHC)
Tissue was formalin-fixed for ~12 h based on ASCO/CAP guidelines[58]. ER, PR, AR, HER2, and Ki-67
IHC stains were performed in the OHSU Knight Diagnostic Laboratories (KDL) on formalin-fixed,
paraffin-embedded tissue, using pre-diluted antibodies intended for diagnostic use in a biotin-free protocol
on a Ventana Benchmark instrument. All antibodies were sourced from Roche. Staining was interpreted
and scored by a pathologist. ER, PR, and AR: Antibodies include anti-ER (CONFIRM clone SP1), #790-
4324; anti-PR (CONFIRM clone 1E2), #790-2223; and anti-AR (Cell Marque clone SP107), # 760-4605.
ER, PR, and AR stains are considered negative if there is nuclear staining in < 10% of the tumor cells.
HER2: Staining was performed with Anti-HER2/neu (PATHWAY clone 4B5), #790-2991. HER2 scoring
and interpretation are per the Ventana scoring guide as expanded by the ASCO/CAP 2013 guidelines. Ki-
67: Staining was performed with anti-Ki-67 (CONFIRM clone 30-9) and reported as percent positive cells.
PD-L1: Anti-PD-L1 staining with clone 22C3 was performed by PhenoPath Laboratories (Seattle, WA)
using pharmDx (Agilent) or by KDL using Dako #M3653. Positive cells were scored by Tumor Proportion
Score. Staining with clone SP263 (Roche Ventana #790-4905) was used when clone 22C3 staining was
unavailable (Participants A, D, T, V).
844
NGS: GENETRAILS® COMPREHENSIVE SOLID TUMOR PANEL
Targeted DNA sequencing was conducted at the CLIA-certified and CAP-accredited OHSU Knight
Diagnostic Laboratories using the GeneTrails Solid Tumor Panel assay. DNA was extracted from tumor-
rich regions that were macro-dissected from FFPE tissue. Library preparation utilized custom QiaSeq
chemistry (QIAGEN) with multiplexed PCR, and sequencing was performed on an Illumina NextSeq
500/550. The DNA library was created with 9,228 custom-designed primer extension assays, covering
613,343 base pairs across 125 to 225 cancer-related genes. The panel is routinely sequenced at an average
read depth exceeding 2,000 to ensure highly sensitive detection of SNVs, short insertions/deletions, and
copy number alterations.
854
PI3K-AKT pathway genes include the following genes from the GeneTrails panel: *AKT1, AKT2, PIK3CA,*
PIK3CB, PIK3R1, or *PTEN*. RTK-MAPK pathway genes include the following genes: *HRAS, KRAS,*
NRAS, ARAF, BRAF, RAF1, MAP2K1, MAP2K2, MAPK1, EGFR, ERBB2, ERBB3, or *MET*. HR genes
include the following*: ARID1A, ATM, ATR, ATRX, BAP1, BARD1, BRCA1, BRCA2, BRIP1, CDK12,*
CHEK1, CHEK2, FAM175A, FANCA, FANCC, FANCD2, FANCE, FANCF, FANCG, FANCM, MRE11A,
NBN, PALB2, RAD50, RAD51, RAD51B, RAD51C, RAD51D, RAD52, RAD54L, and *RECQL*. Immune
genes include the following: *B2M, CD274, HLA-A, HLA-B, HLA-C, IDO1, IDO2, IFNGR1, IFNGR2,*
IRF1, JAK1, JAK2, and *PDCD1LG*.
863
864 Only genomic alterations classified as Tier 1 (Strong Clinical Significance) or Tier 2 (Potential Clinical
865 Significance), or gene copy number gains ≥7 or losses ≤0.5, were included in this study. For samples
866 analyzed using earlier versions of the GeneTrails panel that lacked full coverage of these genes, whole-
867 exome sequencing (WES) was used to determine gene status.
868
WHOLE EXOME SEQUENCING (WES)

Preparation of DNA was described as above at the Knight Diagnostic Laboratories. Total nucleic acid
from FFPE or buffy coat was sent to Tempus Labs Inc. (Tempus Labs, Inc., Chicago, IL, USA) for exome
sequencing using the Tempus xE panel. Raw sequence reads were reprocessed at OHSU following the
GATK Best Practices for Somatic Variant Calling. Briefly, raw sequence FASTQ reads were aligned to
the UCSC GRCh37/hg19 human genome build using BWA-MEM (0.7.12, GATK, Broad Institute) [63,64],
followed by marking duplicate reads (Mark Duplicates, GATK) and base recalibration (BQSR, GATK).
Somatic variants were called using MuTect2 (4.0.4.0, GATK, Broad Institute) between the FFPE (tumor)
and the participant's matched normal from buffy coat. A panel of normal (PON) and the gnomAD (2.0.1)
[65] germline reference resource were used to filter out technical sequencing artifacts and common
polymorphisms, respectively. All analysis tools were run using an OHSU Galaxy instance (v19.01) [66]. On-
olaparib WES was not performed for most participants, given the short duration between biopsies.
881
882 *MUTSIG3 ANALYSIS*
883 COSMIC mutation signatures (v2) were calculated from the somatic variants (VCF) from WES using the
884 MutationalPatterns R package with the hg19 genome reference (BSgenome.Hsapiens.UCSC.hg19) [40,67].
885 The Signature 3 (MutSig3) scores were discretized into High or Low using a threshold of 40.
886
HR DETECTION ASSAY
HR efficiency was assessed as in Ozmen et al.[39], whereby RAD51 foci, which represent functional HR
repair complexes, are assessed in cells that are both Geminin positive (G2 phase) and γH2AX foci positive
(double strand DNA breaks). For IF staining of formalin-fixed, paraffin-embedded (FFPE) tissue samples,
5 μm sections were deparaffinized using xylene twice for 10 minutes, 100% ethanol twice for 10 minutes,
95% ethanol for 5 minutes, 70% ethanol for 5 minutes, and 50% ethanol for 5 minutes, and then left in
PBS. Tissues were antigen-retrieved in a pressure cooker using Citrate (pH 6) and DAKO antigen retrieval
(pH 9) buffers. Tissues were washed twice in PBS and then were further permeabilized with 0.4% Triton
X-100 in PBS for 1 hour. The excess permeabilization solution was removed by washing three times in
PBS, and then a hydrophobic barrier was placed around the mounted tissue. Tissues were blocked with
3% BSA in PBS prior to antibody labeling. The tissues were then incubated with primary antibodies Rad51
(1:1500; Abcam #ab133534), Geminin (1:15, #NCL-L, Leica), and γH2AX (1:400; EMD #05-636) on a
rocker for 2 hours at room temperature in a humidity chamber. After three PBS washes, secondary
antibodies (1:500, Invitrogen #A21428, 1:500, Biotium #20253-1, 1:1000, Invitrogen #A21131) were
applied. After three final PBS washes, all tissue samples were postfixed with 4% PFA for 10 minutes at
room temperature. Hoechst staining solution (1:5000) was used and mounted in Prolong Gold Antifade
mounting media. Sample images were acquired using a spinning disk confocal microscope with a 40x dry
objective. Raw image files from Zen (.czi) were imported into custom ImageJ macro software for image
segmentation. WEKA machine learning plugin was used for both nuclei and foci segmentation. HRC
nuclei were defined as having $\geq$5 RAD51 foci, while HRD cells were defined as RAD51 negative nuclei
with $\geq$3 γH2AX+ foci. The relative contributions of both HRC and HRD nuclei were used to stratify
participants.
909
RNA TRANSCRIPTOME SEQUENCING
Preparation of RNA and transcriptome sequencing was performed at the Knight Diagnostic Laboratories.
Total nucleic acid was extracted and purified from macrodissected, tumor-rich areas of FFPE sections.
Libraries were prepared using a TruSeq RNA Access library preparation kit and sequenced on an Illumina
NextSeq500/550. Approximately 100 million reads were generated per sample. Gene expression was
quantified relative to Gencode v24 transcripts using Kallisto[68,69].

GENE SET VARIATION ANALYSIS (GSVA)

GSVA, a non-parametric unsupervised method to calculate gene set or pathway scores on a per-sample basis, was conducted on the RNAseq data using the GSVA package (v1.44.5) with the 1) Molecular Signatures DataBase (MSigDB) Cancer Hallmarks (v7.5.1) and 2) 16 immune cell types gene sets[70-72]. GSVA scores values were discretized into High or Low using a threshold of 0.

MULTIPLEX IMMUNOHISTOCHEMISTRY (mIHC)

Multiple immunohistochemistry (mIHC) was performed as previously described[73] and following (https://www.protocols.io/view/mihc-staining-ohsu-coussens-39-lab-sop-3i6gkhe). Antibody information is listed in Supplementary Table 1. Briefly, FFPE human tissues were sectioned at 5 microns and mounted on positively charged slides (Tanner Adhesive Slides, Mercedes Medical, TNR WHT45AD). Slides were baked overnight at 55°C followed by an additional baking step for 30 minutes at 58-60°C. They were then deparaffinized in xylene and graded ethanol (Xylene 2 x 5 min, 100% EtOH 2 x 2 min, 95% EtOH 2 x 2 min, 70% EtOH 2 x 2 min, 50% EtOH 1 x 2 min, diH2O 2 x 2 min). Slides were counterstained in Hematoxylin (Dako S3301), underwent heat-mediated antigen retrieval in pH 6.0 Citra solution (BioGenex, Fremont, CA, USA), blocked in Dako Dual Endogenous Enzyme Block (S2003, Dako, Santa Clara, CA, USA), and subjected to protein blocking with 5% normal goat serum and 2.5% bovine serum albumin (BSA) in tris-buffered saline-0.1% Tween (TBST). Slides were incubated with primary antibodies for 1 hour at room temperature or 16–17 hours at 4°C. Primary antibody was washed off in TBST, and either anti-rat, anti-mouse, or anti-rabbit Histofine Simple Stain MAX PO horseradish peroxidase-conjugated polymer (Nichirei Biosciences, Tokyo, Japan) was applied for 30 min at room temperature, followed by AEC chromogen (Vector Laboratories, Burlingame, California, USA).

Images were acquired using the Aperio ImageScope AT (Leica Biosystems) at ×20 magnification. Following acquisition, coverslips were gently removed in 1×TBST while agitating. Removal of AEC and HRP inactivation was accomplished by incubating the slides in 0.6% fresh H_2O_2 in methanol for 15 minutes. AEC removal and stripping of antibodies was accomplished by ethanol gradient incubation and heat-mediated antigen retrieval such as described above between cycles. After washing and protein blocking, samples were subjected to the next round of staining. Some samples failed due to tissue loss during staining and image acquisition. Cell phenotypes were assigned with hierarchically gating (see Supplementary Table 2 for gating strategy) and quantified to cell densities (cells/mm^2). Cell phenotype densities were log2 transformed followed by scaling and centering (i.e. standard scaled or z-scored) as continuous variables and discretized as "High" for samples ≥ 0 and "Low" for <0.

NANOSTRING GEOMX DIGITAL SPATIAL PROFILING (DSP)

DSP was performed by the OHSU Knight Diagnostic Laboratories. A single FFPE slide per sample was used for DSP protein panel analysis. FFPE sections (4-5 micron) were deparaffinized and subjected to antigen retrieval in Citrate buffer, pH 6.0 for 15 minutes using a pressure cooker. The slides were then incubated with a cocktail of oligonucleotide-tagged antibodies together with two fluorescent-tagged morphological markers, pan-CK (Alexa 532nm) and CD45 (Alexa 594nm) at 4°C overnight, followed by blocking for 1 hour with Buffer W. After antibody incubation, slides were washed with TBS-T and post-fixed with 10% neutral buffered formalin for 30 minutes. Following a brief wash with TBS-T, the sections were incubated with SYTO-13 (FITC 525nm) for 30 minutes and then placed on the DSP instrument for sample collection. Slides were scanned and tumor regions were identified based on nuclei staining and visualization markers.

A minimum of three regions of interest (ROIs; 660 micron diameter) were selected for each sample under the guidance of a board-certified pathologist in accordance with recommended practices[74]. ROIs were chosen with the goal of maximizing tumor cell content, avoiding any areas of necrosis or sectioning artifacts on the slide. UV-released oligonucleotide barcodes were collected automatically in a 96-well plate, then hybridized with GeoMx hyb codesets at 67°C for 16 hours and quantitated using the MAX nCounter system.

Assays were run on two versions of the assay platform: 1.2 and 2.0. Some antibodies differed between platform versions (see Supplementary Table 3 for DSP antibody list), but the same analysis pipeline was used for all. A control tissue microarray was included in every run to correct for batch effects. Based on comparison with normalized signals from a 19-sample mixed breast cancer cohort, sample values were discretized with z-score percentiles. To determine appropriate thresholds for DSP protein markers, we performed a density-based bimodal distribution analysis. Density curves were computed for each protein marker using a Gaussian kernel density function where the majority of DSP protein markers followed a bimodal distribution. Using a rolling window approach, the local maximums and minimums were calculated for each marker where peaks were defined as local maximums exceeding both the leading and trailing edges of the window and valleys defined as the minimum density value between the first two identified peaks. The median valley across all markers was defined as the global threshold for either signaling associated or immune markers as density distributions for tumor signaling associated markers tended to be shifted higher. The final thresholds were set to 60 for signaling markers and 50 for immune markers.

IMMUNE CONSENSUS SIGNATURES

For the RNAseq-based Immune Consensus ("IMMUNE RNA SUMMARY"), a sample was classified as "Positive" if at least 10 of 17 High Immune RNA Signatures (Extended Data Fig. 1c) were present. These 17 RNA signatures were identified as predictive of clinical outcomes ($p < 0.05$), selected using the Boruta feature-selection algorithm, and demonstrated an F1 score ≥ 70. The signatures included: Activated CD8$^+$ T Cells, Allograft Rejection, B Cells, Complement, Cytotoxic Cells, Gamma Delta T Cells, Immature Dendritic Cells, IL2/STAT5 Signaling, IL6/JAK/STAT3 Signaling, Inflammatory Response, Interferon Alpha Response, Interferon Gamma Response, NK CD56^{dim} Cells, Regulatory T Cells, T Helper Cells, Central Memory T Cells (TCM), and Effector Memory T Cells (TEM).

For the mIHC-based Immune Consensus ("IMMUNE mIHC SUMMARY"), a sample was classified as "Positive" if at least 9 of 10 High Immune RNA Signatures (Extended Data Fig. 1c) were present. These 10 mIHC Cell Types were identified as predictive of clinical outcomes ($p < 0.05$), selected using the Boruta feature-selection algorithm, and demonstrated an F1 score ≥ 70. The signatures included: CD20$^+$ B Cells, CD4$^+$ T Cells, CD8$^+$ T Cells, Immune Cells, Ki-67$^+$ CD8$^+$ T Cells, Ki-67$^+$ Immune Cells, Ki-67$^+$ TBET/EOMES/PD-1$^-$ CD8$^+$ T Cells, Other Immune Cells, TBET$^-$ FOXP3$^-$ CD4$^+$ T Cells, and TBET/EOMES/PD-1$^-$ CD8$^+$ T Cells.

For the DSP-based Immune Consensus ("IMMUNE DSP SUMMARY"), a sample was classified as "Positive" if at least 40% High Immune DSP Proteins (Extended Data Fig. 1c) were present. For samples in which the stroma compartment was not assayed, only tumor compartment proteins were included. Eleven Immune DSP Proteins were identified as predictive of clinical outcomes ($p < 0.05$), selected using the Boruta feature-selection algorithm, and demonstrated an F1 score ≥ 70. The proteins included: CD3

(Stroma), CD3 (Tumor), CD4 (Tumor), CD45 (Stroma), CD45 (Tumor), CD8 (Stroma), GZMA (Stroma), GZMA (Tumor), HLA-DR (Stroma), HLA-DR (Tumor), and PD-L1 (Tumor).

The MULTI-OMIC IMMUNE CONSENSUS was defined by integrating RNAseq, mIHC, and DSP immune consensus results. A sample was classified as "Positive" if it was positive in at least one of the individual measures: IMMUNE RNA SUMMARY, IMMUNE mIHC SUMMARY, or IMMUNE DSP SUMMARY.

REVERSE PHASE PROTEIN ARRAY (RPPA)
Protein extracts from tumor samples were analyzed with a panel of 480 proteins and phosphoproteins as previously described[75,76]. To scale the protein expression values, the RPPA data from each participant sample was merged within the TCGA primary breast cancer and SMMART-program metastatic breast cancer RPPA datasets, using the replicate-based normalization method[57,77]. The protein expression values were then *z*-scored by the Median Absolute Deviation (MAD) from a Basal classified subset of TCGA samples.

TNBC SUBTYPE CALLING
CLIA TNBC subtypes were computed using a refined list of 77 genes using a novel algorithm[51,52]. Sample gene expression was batch corrected using Removal of Unwanted Variation (RUVg) [78]. Spearman's correlation was computed between the gene expression values and each centroid. Patient samples were classified as immune activated (BLIA), basal immune suppressed (BLIS), luminal androgen receptor (LAR), or mesenchymal (MES) subtypes based correlation to the corresponding centroids. In cases where the difference between the top two centroids was less than 0.1, samples were considered indeterminate (IND).

SINGLE-CELL RNA SEQUENCING (scRNAseq)
Thirteen biospecimens were collected from participants with metastatic breast cancer under an IRB-approved MMTERT observational study[53] (OHSU IRB #16113). Fresh core needle biopsies were disaggregated using gentleMACS kits (Miltenyi Biotec, #130-095-929) following the manufacturer's instructions and filtered through a 70-μm cell strainer. Cells were collected by centrifuging (300g for 7 min at 4°C) and resuspended at 700–1200 cells/μl. Live cells were isolated with a EasySep Dead Cell Removal (Annexin V) Kit (STEMCELL Technologies, #17899). Single cell suspensions of viable cells were processed using a Chromium Next GEM Single Cell 3' Kit v3.1 (10xGenomics) following the manufacturer's protocol. The entire mixed cell population was analyzed without sorting or enriching specific cell subtypes. Briefly, viable single cells were encapsulated into GEMs containing barcoded oligonucleotides. 5,000-10,000 GEMs were then subjected to cell lysis, reverse transcription, and cDNA amplification. Libraries were constructed from the amplified cDNA, and sequencing was performed on the Illumina NovaSeq platform. Raw sequencing data were aligned to GRCh38 genome reference using 10X software CellRanger (Version 6.1.2) with default parameters.

To ensure high-quality data, we used Scanpy[79] to implement three quality control measures on the raw gene-cell-barcode matrix for each cell: 1) the proportion of mitochondrial genes (≤20%), 2) unique molecular identifiers (UMIs, ranging from 400 to 120,000), and 3) gene count (ranging from 200 to 10,000). Doublets were identified and removed from each sample using Scrublet[80]. Normalization of total counts per cell was performed using the normalize_total function in Scanpy, followed by log-

normalization with the log1p function. Clustering was conducted using the Leiden algorithm at a resolution of 1 in Scanpy.

Marker genes for each cluster were identified using a t-test implemented in Scanpy. These marker genes were then used to annotate each cluster using publicly available databases including CellMarker[81] and PanglaoDB[82] to determine cell identities. These annotations were later refined using the CellTypist[83] package with the celltypist.annotate function. The model 'Immune_All_High.pkl' was specified, with majority voting enabled. Through this process, seven major cell type clusters were annotated: epithelial cells, natural killer cells, myeloid cells, T cells, B cells, fibroblasts, and endothelial cells. To distinguish malignant cells, we calculated and identified large-scale chromosomal copy number variation (CNV) using inferCNV[84] and CaSpER[85] tools for each sample based on transcriptomes. T cells and myeloid cells were used as reference cells, and epithelial cells with differing CNV patterns and higher CNV score were annotated as malignant cells.

Datasets were merged across different samples using SCVI[86] based integration. The top 4,000 highly variable genes were used to train the VAE models, with each biopsy as covariate key. After training the initial VAE model, the annotated cell types were used to build an extended model with scANVI[87] for better integration. After integration, NK cells, myeloid cells, T cells, B cells, and fibroblasts were further classified using the following publicly available scRNAseq datasets: NK cells (GSE212890[88]); T cells, myeloid cells, and B cells (GSE169426[89]) and fibroblasts (GSE103322, GSE132465, GSE154778, and GSE212966[90]). The label transfer method was performed using the scArches[91] algorithm, following the best practices described previously[93]. Dot plots were generated using the Scanpy DotPlot function to visualize the expression patterns of a panel of previously defined TNBC subtype markers[52] across various cell types.

mTNBC3s OUTCOME CLASSIFIER
A 34 gene classifier, mTNBC3s, designed to improve upon the CLIA TNBC BLIA/BLIS subtype classifier by reducing the number of indeterminate calls was previously developed and applied to a subset of the AMTEC cohort (see Bottomly et al. [54] for detailed methods). The same classifier was applied to the additional AMTEC samples in this paper. Because this model was trained on datasets that did not include LAR tumors, LAR-classified samples were excluded (Participants P and U).

THREE-PROTEIN RESPONSE PREDICTOR
Using the normalized GeoMx DSP (z-score percentiles), the on-olaparib samples were randomly split into a 65/35 train-test set for feature selection and model development. Using the training set, combinatorial feature selection with ANOVA was used to identify the set of protein markers (continuous measurements) with the highest discriminatory power for predicting best response (categorical outcome). Combinations of 2, 3, and 4 proteins were systematically evaluated and ranked using ANOVA F-statistics (f_classif) with the top-performing feature combinations retained for model building.

A logistic regression classifier with L2 (Ridge) regularization (C=1.0, lbfgs solver) was trained using the same training set as above and the top-performing protein marker combination, including CD3 (Tumor), CD45 (Tumor), and PD-L1 (Stroma). Since feature selection was previously performed with ANOVA, L2 regularization was selected over L1 (LASSO) to retain protein features with weighted coefficients. The classifier was implemented with a fixed random seed (seed=42) and evaluated on the independent test set with performance assessed by precision, recall, F1 score, and accuracy.

1099

1100 *VISUALIZATIONS*
1101 Boxplots, bar plots, and Kaplan-Meier curves were plotted with R using ggplot2, ggpubr, and ggstatsplot.
1102 Swimmer plots were generated in Python using matplotlib. The CoMut plot was generated using the
1103 Python library CoMut[92].

1104

1105 *DATA AVAILABILITY*
1106 All data are available through the HTAN Data Portal as part of the HTAN OHSU Atlas
1107 (https://data.humantumoratlas.org/). Raw sequencing data have been deposited to dbGAP (Project
1108 phs002371.v1.p1). Images can be viewed at the Imaging Data Commons (IDC)
1109 https://portal.imaging.datacommons.cancer.gov/.

1110

1111 *CODE AVAILABILITY*
1112 No new tools were developed for the analyses in this study.

1113
1114

 FIGURE LEGENDS

1116
1117 **MAIN**
1118
1119 **Table 1. Summary of clinical features of AMTEC trial participants at baseline.**

1120
1121 **Figure 1. Overview of the AMTEC trial and clinical outcomes.**

1122
1123 **a)** *Trial design.* Schematic of the AMTEC trial showing key eligibility criteria, treatment schedule, biopsy
1124 timing, treatment discontinuation, and post-treatment follow-up. Created in BioRender. Yildiran Ozmen,
1125 T. (2025) https://BioRender.com/vjju338

1126
b) *Progression free survival (PFS).* Swimmer plot for 21 efficacy-evaluable participants (see Extended
Data Fig. 1a). PFS measured from the start of olaparib treatment until the end of olaparib plus durvalumab.
Best response by RECIST v1.1[32] is adjacent to each treatment duration bar. Orange bars: participants with
Complete Response (CR), Partial Response (PR), or Stable Disease (SD); purple bars: Progressive Disease
(PD). * = passed away from a trial-unrelated accident; ** = voluntarily withdrew from the study. Arrows
= alive at time of data cutoff.

1133
1134 **c)** *Best percentage change from baseline.* Waterfall plot showing best change in the sum of all target
1135 lesions for 20 efficacy-evaluable participants. Lesions from Participant W were not evaluable. Dashed
1136 lines indicate RECIST thresholds for PD (+20%) and PR (−30%). * = new lesions.

1137
d-g) *Kaplan-Meier analyses.* PFS and Overall Survival (OS) for 20 survival-evaluable participants (see
Extended Data Fig. 1a). Shaded areas indicate 95% confidence intervals (CI); median time shown by
dashed line. PFS measured from start of olaparib until end of olaparib plus durvalumab. OS measured
from olaparib start until death or data cutoff; right-censoring denoted by "+".
d) PFS, all.
e) OS, all.
f) PFS, by RECIST best response. Non-responders (NR; purple) = PD. Responders (R; orange) =
CR/PR/SD. P-values from log-rank tests.
g) OS, by RECIST best response.

1147
1148 **Figure 2. Malignant cell-intrinsic signatures of response to olaparib and durvalumab.**

1149
1150 **a)** *Mutation Signature 3 (MutSig3).* Kaplan-Meier analysis of Overall Survival (OS) by High (red) vs.
1151 Low (blue) MutSig3 status from whole-exome sequencing of pre-treatment biopsies. Hazard ratios (HRs)
1152 and p-values from Cox proportional hazards (CoxPH) and log-rank tests are shown. Shaded areas indicate
1153 95% confidence intervals (CI) for the log survival probability. OS was measured from the start of olaparib
1154 until death.

1155
1156 **b)** *PAR protein levels.* Boxplots comparing PAR protein levels from RPPA in Pre-Treatment (Pre-) and
1157 On-Olaparib (On-) biopsies, separated by RECIST non-responders (NR; purple) and responders (R;
1158 orange). Lines connect paired Pre- and On- samples from the same participant. P-values from paired
1159 Student's t-test comparing Pre- versus On- biopsies within each group.
1160

c) *Androgen Receptor (AR) IHC.* Kaplan-Meier analysis of OS stratified by AR IHC ($\geq$10% positive cells, blue; <10%, red) in Pre-Treatment (left) and On-Olaparib (right) biopsies. OS measured from olaparib start until death or data cutoff; right-censoring denoted by "+".

d-e) *Signaling pathway activity.* Kaplan-Meier analysis of OS stratified by High (z-score percentile $\geq$50, red) or Low (<50, blue) DSP signaling pathway protein expression. Insets: stacked bar plots of percent NR (purple) and R (orange) in High (left) and Low (right) groups; p-values from Fisher's exact test for High vs. Low.
d) PI3K-AKT pathway activity. Positive: p-AKT1 and p-GSK3B $\geq$50 percentile.
e) Mixed signaling pathway activity. Positive: at least 3 of p-GSK3B, p-PRAS40, AKT, or p-HER2 $\geq$50 percentile.

Figure 3. Microenvironment signatures of response to olaparib and durvalumab: Immune Consensus signatures.

a-d) *Multi-omic immune summary signatures.* Kaplan-Meier OS in Pre-Treatment (left) and On-Olaparib (right) biopsies. Samples classified as Positive (blue) or Negative (red) based on predefined signature thresholds. HRs and p-values from CoxPH and log-rank tests are shown; shaded areas indicate 95% CI for the log survival probability. OS measured from olaparib start until death or data cutoff; right-censoring denoted by "+". Insets: stacked bar plots of percent NR (purple) and R (orange) in Negative (left) and Positive (right) groups; p-values from Fisher's exact test for Negative vs. Positive. Features shown achieved F1 >0.7 from random train/test split analysis and were significant (p <0.05) in pre-treatment and/or on-olaparib analyses.
a) Immune RNA summary signature. Positive: $\geq$10 of 17 Immune RNA Signatures (Extended Data Figure 1c).
b) Immune mIHC summary signature. Positive: $\geq$9 of 10 mIHC Cell Types (Extended Data Figure 1c).
c) Immune DSP summary signature. Positive: $\geq$40% of Immune DSP proteins (Extended Data Figure 1c).
d) Multi-Omic Immune Consensus signature. Positive: at least one positive summary signature (Extended Data Figure 1c).

Figure 4. Olaparib-induced response markers and signatures.

a-h) *Changes following olaparib treatment.* Boxplots showing changes in feature expression from Pre-Treatment (Pre-) to On-Olaparib (On-) for paired samples from the same participant. Positive values = higher expression on-olaparib; negative values = lower expression. NR (purple) and R (orange) changes were compared using Student's t-test. Features shown were identified by Boruta analyses for each dataset: RNAseq, mIHC, DSP, and RPPA. All features significant for NR vs. R (p <0.05).
a) Immune markers increased in responders, decreased in non-responders.
b) Immune markers increased in responders, unchanged in non-responders.
c) Immune markers unchanged in responders, decreased in non-responders.
d) Immune markers decreased in responders, increased in non-responders.
e) Malignant cell markers increased in responders, decreased in non-responders.
f) Malignant cell markers increased in responders, unchanged in non-responders.
g) Malignant cell markers decreased in responders, increased in non-responders.
h) Malignant cell markers decreased in both responders and non-responders.

Figure 5. TNBC subtype correlates of response to olaparib and durvalumab.

a) *TNBC subtypes*. Kaplan-Meier PFS by TNBC RNA subtype (BLIS/LAR, red; Non-BLIS/LAR, blue) in Pre-Treatment (left) and On-Olaparib (right) biopsies. HR and p-values from CoxPH and log-rank tests are shown; shaded areas indicate 95% CI. PFS measured from start of olaparib treatment until end of olaparib plus durvalumab. Insets: stacked bar plots of percent NR (purple) and R (orange) in BLIS/LAR (left) and Non-BLIS/LAR (right) groups; p-values from Fisher's exact test for BLIS/LAR vs. Non-BLIS/LAR.

b) *Progression free survival with TNBC subtypes.* Swimmer plot showing PFS (time on olaparib ± durvalumab) for 24 participants (21 efficacy + 3 pilot; Extended Data Figure 1c). TNBC subtype and RECIST best response are annotated. Subtyping was from the on-olaparib biopsy except for Participants A, C, and U (Pre-Treatment) and N (no RNA available). Arrows = alive at time of data cutoff.

c) *Overall survival with TNBC subtypes.* Swimmer plot showing OS from treatment start until death or data cutoff for the same participants as in (b). Black squares = treatment end.

d) *mTNBC3s gene signature.* Kaplan-Meier PFS stratified by mTNBC3s[54] predicted non-responders (pPD, red) and predicted responders (pSDPR, blue). Insets: stacked bar plots of percent NR (purple) and R (orange) in pPD (left) and pSDPR (right) groups; p-values from Fisher's exact test for pPD vs. pSDPR.

Figure 6. Three-Protein Response Predictor correlates of response to olaparib and durvalumab.

a-b) *Survival and response correlates*. Kaplan-Meier OS and PFS stratified by combined weighted expression scores from three immune DSP proteins (CD45 tumor, CD3 tumor, and PD-L1 stroma): predicted non-responders (pNR score <0; red) and responders (pR ≥0; blue). HR and p-values from CoxPH and log-rank tests are shown; shaded areas indicate 95% CI. OS measured from olaparib start until death or data cutoff; right-censoring denoted by "+". PFS measured from start of olaparib until end of olaparib plus durvalumab. Insets: stacked bar plots of percent NR (purple) and R (orange) in pNR (left) and pR (right) groups; p-values from Fisher's exact test for pNR vs. pR.
a) OS and best response distributions.
b) PFS.

EXTENDED DATA

Extended Data Table 1. Clinical summary of AMTEC trial participants.

a) *Adverse events.* Safety analysis performed on data from 24 participants who received at least one dose of either drug. Most treatment-related adverse events (89%) were Grade 1-2. Of the 45 Grade ≥3 events, 20 (44%) were treatment-related.

b) *Grades 3-5 adverse events.* Data are from 15 of 24 participants. The two immune-related events were resolved with a dose hold. The two Grade 5 events were unrelated to protocol therapy: one due to respiratory failure and one (Investigations – other*) resulting from a motor vehicle accident.

c) *Response-associated features.* Summary of statistically significant features correlating with response to olaparib and durvalumab. The biopsy in which each feature was observed (Pre-Treatment, Pre-; On-Olaparib, On-) is indicated for significant associations with Best Response (BR), Progression Free Survival (PFS), and Overall Survival (OS). Features identified by Boruta feature selection are noted. Median F1 scores for random training/test split analysis are reported. ns = not significant (p-value >0.05); N/A = not available.

d) *Olaparib-induced features.* Summary of features significantly changed in olaparib-treated samples. All features were identified by Boruta feature selection performed separately for each dataset: Inferred Immune Cell Type gene sets, mIHC Cell Phenotypes, DSP Signaling or Immune Protein Expression, and RPPA Protein Expression.

Extended Data Figure 1. Overview of the AMTEC trial and clinical outcomes.

a) *Trial design and enrollment.* CONSORT flow diagram summarizing participant enrollment, exclusions, and analyses in the AMTEC trial.

b) *Overall Survival (OS).* Swimmer plot for 21 efficacy-evaluable participants. Best response by RECIST v1.1[33], is indicated adjacent to each treatment duration bar. OS measured from olaparib start until death or data cutoff. Orange bars: participants with Complete Response (CR), Partial Response (PR), or Stable Disease (SD); purple bars: Progressive Disease (PD). * = passed away from a trial-unrelated accident; ** = voluntarily withdrew from the study. Black squares = treatment end. Arrows = alive at time of data cutoff.

c) *Multi-omic clinical features and signatures from serial biopsies.* Summary analysis results for all clinical and molecular features in each sample. Columns are sample identifiers (participant identifier appended to biopsy number; Bx1 = pre-treatment biopsy; Bx2 = on-olaparib biopsy). Rows correspond to clinical or molecular feature and signature data. White boxes denote missing data. Features with parenthetical notations were found to be significant in one or more analyses: R = RECIST best response t-test; P = Kaplan-Meier analysis of Progression Free Survival (PFS); O = Kaplan-Meier analysis of Overall Survival (OS); B = paired biopsy t-test for responders and non-responders; D = pre-treatment to on-olaparib change t-test for responders vs non-responders. * p <0.05; ** p <0.005; *** p <0.0005; ns = not significant. See also Extended Data Tables 1c,d for feature significance summaries.

Best Response: determined by RECIST v1.1. CR = Complete Response; PR = Partial Response; SD = Stable Disease; PD = Progressive Disease.

Participant Cohort: AMTEC trial or Pilot study enrollment. Includes all efficacy-evaluable AMTEC participants (see Extended Data Fig. 1a).

PFS: measured from start of olaparib until end of olaparib plus durvalumab (see Fig. 1d).

OS: measured from olaparib start until death or data cutoff (see Fig 1e).

Anatomical Location: organ site of the metastatic lesion from which the biopsy was obtained.

Tumor Content: pathologist-estimated percentage of tumor cells after microdissection of tumor-rich regions from FFPE slides.

Ki-67, **PD-L1**, and **AR IHC:** assayed clinically and defined as defined as follows: Ki-67 = % positive tumor cells; PD-L1 = tumor proportion score (TPS) ≥1% or ≥10%; AR = nuclear staining tumor cells ≥10% or ≥70% (see Fig. 2c).

PAR RPPA: PAR protein levels measured by RPPA. High = median absolute deviation >0 relative to a Basal-classified subset of TCGA samples (see Fig. 2b, Extended Data Fig. 2e, Fig. 4h).

HRC/HRC Ratio: the ratio of HRC to HRD nuclei. Ratios <15% = HRD, >65% = HRC, 15–65% = Intermediate (see Extended Data Fig. 2a–c).

MutSig3: COSMIC Mutational Signature 3, discretized into High or Low using a threshold of 40 (see Fig. 2a).

HR & Immune Genes and **PI3K-AKT & RTK-MAPK Genes:** variants reported clinically by GeneTrails. Copy Gain is ≥7 copies and Copy Loss is ≤0.5 copies. Mutations include missense mutations, nonsense mutations, indels, and splice site substitutions. See *Methods* for pathway gene inclusions.

Signaling Pathway Gene Alteration: the presence of least one DNA alteration among the PI3K-AKT or the RTK-MAPK pathway genes (see Extended Data Fig. 2f). Note that participants K and W had only one of two biopsies with clinically reported *PTEN* loss, possibly due to tumor content variability constraining copy number loss detection.

DSP Signaling Proteins: phosphoprotein levels within tumor ROIs measured by NanoString DSP. High = z-score percentile ≥50 relative to a mixed breast cancer cohort (see Extended Data Fig. 2h, Fig. 4g).

DSP Signaling Pathway Activation: PI3K-AKT PATHWAY DSP activation = both p-AKT1 and p-GSK3B high. MIXED SIGNALING PATHWAY DSP activation is ≥3 high among p-GSK3B, p-PRAS40, AKT, or p-HER2 (see Fig. 2d,e and Extended Data Fig. 2i,j).

ZAP-70 RPPA: ZAP-70 protein levels measured by RPPA. High = median absolute deviation >0 relative to a Basal-classified subset of TCGA samples (see Extended Data Fig 3mm).

Immune RNA Signature: from GSVA scores of MSigDB Cancer Hallmarks and Inferred Immune Cell Type gene sets. Discretized into High or Low using a threshold of zero (see Extended Data Fig. 3e-u, Fig. 4a,c).

IMMUNE RNA SUMMARY: the number of High and Low Immune RNA Signatures per sample. Positive is ≥10 High signatures (see Fig. 3a and Extended Data Fig. 3a).

mIHC Cell Type: immune cell phenotypes assigned by hierarchical gating of protein co-expression and quantified to cell densities. High is logged/scaled z-score ≥0 (see Extended Data Fig. 3v-ee, Fig. 4a, Extended Data Fig. 4a).

IMMUNE mIHC SUMMARY: the number of High and Low mIHC Cell Types per sample. Positive is ≥9 High cell types (see Fig. 3b and Extended Data Fig. 3b).

Immune DSP: levels of immune proteins measured within tumor and stroma ROIs by NanoString DSP. High is z-scores percentile ≥60 (see Extended Data Fig. 3ff-ll, Fig. 4a-c, Extended Data Fig. 4a-c,g).

IMMUNE DSP SUMMARY: the number of High Immune DSP proteins per sample. Positive is ≥40% High proteins. Stroma compartment was unmeasurable for some samples (see Fig. 3c and Extended Data Fig. 3c).

MULTI-OMIC IMMUNE CONSENSUS: the overall immune status in a combined three-platform signature. Positive is at least one High measurement was observed for IMMUNE RNA SUMMARY, IMMUNE mIHC SUMMARY, or IMMUNE DSP SUMMARY (see Fig. 3d and Extended Data Fig. 3d).

Change from Bx1 to Bx2: the difference in feature score between the pre-treatment and on-olaparib biopsies (see Fig. 4). For RNA, mIHC, and RPPA features, Strong Increase ≥ +0.25; Mild Increase > +0.05 and < +0.25; Strong Decrease ≤ -0.25; Mild Decrease < -0.05 and > -0.25; No Change > -0.05 and < +0.05. For DSP proteins, Strong Increase is z-score percentile change ≥ +25; Mild Increase > +5 and < +25; Strong Decrease ≤ -25; Mild Decrease < -5 and > -25; No Change > -5 and < +5.

TNBC Subtype: Basal-Like Immune Activated (BLIA), Basal-Like Immune Suppressed (BLIS), Luminal Androgen Receptor (LAR), or Indeterminate (IND), assigned from RNAseq using a revised 77-gene classifier (see Fig. 5a-c, Extended Data Fig. 5a).

mTNBC3s: predicted responder (pSDPR) or non-responder (pPD), assigned from RNAseq analysis of a 34-gene set predictive of response to therapy in a preliminary study of 13 AMTEC trial participants[54] (see Fig. 5d, Extended Data Fig. 5d).

THREE-PROTEIN RESPONSE PREDICTOR: predicted responder (pR) or predicted non-responder (pNR), assigned from NanoString DSP combined protein expression of CD45 (tumor ROI), CD3 (tumor ROI), and PD-L1 (stroma ROI; see Fig. 6).

Extended Data Figure 2. Malignant cell-intrinsic signatures of response to olaparib and durvalumab.

1378

a-d) *Homologous recombination (HR) repair proficiency.* Quantification of DNA repair complexes and
HR status in cycling cells by immunocytochemistry.

a) RAD51, γH2AX, and Geminin nuclear co-staining. Representative images from three samples: HR
deficient (HRD; Participant S), HR competent (HRC; Participant X), and intermediate (Participant K).
RAD51 and γH2AX foci were counted in Geminin-positive nuclei. HRD: <5 RAD51 foci and >3 γH2AX
foci; HRC >5 RAD51 foci. Scale bar is shown as indicated.

b) Machine learning-based HR classification. Custom machine learning segmentation results from the
samples shown in (d). Sample HR classifications were defined using the ratio of HRC to HRD. Deficient
is <15%; Competent is >65%, and Intermediate is 15-65%. RECIST best response is shown in orange for
responders (PR, SD) and purple for the non-responder (PD).

c) γH2AX positivity ratios. γH2AX positivity ratios calculated in the samples in (d) are represented as
horizontal black bars. Samples falling within the green area are HR Deficient (>95% γH2AX$^+$/Geminin$^+$
co-positive nuclei) and in the red area are HR Competent (<55% co-positive nuclei). Samples falling
within the blue area were Intermediate, tending toward higher γH2AX positivity and a mixture of HRC
and HRD nuclei.

d) *Progression-free survival with HRD status.* Swimmer plot showing PFS (time on olaparib ±
durvalumab) for 24 participants (21 efficacy + 3 pilot; Extended Data Figure 1c). HRD/HRC ratio score
is annotated, except for participant B who was germline BRCAm. Scoring was from the on-olaparib
biopsy except for Participants C, D, J, P, and Q, where Pre-Treatment was used (no on-olaparib sample
available).

1399

1400 **e)** *PAR protein.* Boxplots of PAR protein levels measured from RPPA in Pre-Treatment (left) and On-
1401 Olaparib (right) biopsies from RECIST non-responders (NR, purple) and responders (R, orange). P-values
1402 from Student's t-test compare NR and R protein levels at each timepoint.

1403

f-g) *Gene alterations.* Kaplan-Meier analyses of Overall Survival (OS) and Progression-Free Survival
(PFS) stratified by presence of gene alterations in Pre-Treatment (left) and On-Olaparib (right) biopsies.
Hazard Ratios (HR) and p-values from Cox proportional hazards (CoxPH) and log-rank tests are shown;
shaded areas indicate 95% confidence intervals (CI) for the log survival probability. OS measured from
olaparib start until death or data cutoff; right-censoring denoted by "+"; PFS measured from start of
olaparib until end of olaparib plus durvalumab. Insets: stacked bar plots of percent NR (purple) and R
(orange) in groups with or without gene alterations; p-values from Fisher's exact test comparing each
group.

f) PI3K-AKT gene alterations. Positive is ≥1 PI3K-AKT pathway gene alteration (copy gain ≥7, copy loss
≤0.5, missense mutations, nonsense mutations, indels, or splice site substitutions).

g) PTEN gene status. Copy loss: *PTEN* DNA copy number ≤0.5.

1415

h-j) *Signaling proteins.* Kaplan-Meier analysis of PFS stratified by High (z-score percentile ≥50, red) or
Low (<50, blue) DSP signaling pathway protein expression.

h) p-GSK3B protein levels. Insets: boxplots of protein levels from NR (purple) and R (orange); P-values
from Student's t-test for NR vs. R.

i) PI3K-AKT pathway activity. Positive: p-AKT1 and p-GSK3B ≥50 percentile.

j) Mixed signaling pathway activity. Positive: at least 3 of p-GSK3B, p-PRAS40, AKT, or p-HER2 ≥50
percentile.

1423

Extended Data Figure 3. Microenvironment signatures of response to olaparib and durvalumab.

a-d) *Multi-omic immune summary signatures*. Kaplan-Meier PFS in Pre-Treatment (left) and On-Olaparib (right) biopsies. Samples classified as Positive (blue) or Negative (red) based on predefined signature thresholds. HRs and p-values from CoxPH and log-rank tests are shown; shaded areas indicate 95% CI for the log survival probability. PFS measured from start of olaparib until end of olaparib plus durvalumab. Features shown achieved an F1 score >0.7 from random train/test split analysis and were significant (p <0.05) in pre-treatment and/or on-treatment samples (see Extended Data Table 1c).
a) Immune RNA summary signature. Positive: ≥10 of 17 Immune RNA Signatures (Extended Data Figure 1c).
b) Immune mIHC summary signature. Positive: ≥9 of 10 mIHC Cell Types (Extended Data Figure 1c).
c) Immune DSP summary signature. Positive: ≥40% of Immune DSP proteins (Extended Data Figure 1c).
d) Multi-Omic Immune Consensus signature. Positive: at least one positive summary signature (Extended Data Figure 1c).

e-u) *Immune gene sets and signatures*. Kaplan-Meier OS and PFS in Pre-Treatment (left) and On-Olaparib (right) biopsies. Enrichment scores (treated as continuous variables) were categorized as High (≥0, red) and Low (<0, blue). HRs and p-values from CoxPH and log-rank tests are shown; shaded areas indicate 95% CI for the log survival probability. OS measured from olaparib start until death or data cutoff; right-censoring denoted by "+".
PFS as above. Boxplots: enrichment scores from from NR (purple) and R (orange); p-values from Student's t-test. Features shown were identified by Boruta feature selection within MSigDB Cancer Hallmark or Inferred Immune Cell Type gene sets, achieved an F1 score >0.7 from random train/test split analysis, and were significant (p <0.05) in pre-treatment and/or on-treatment samples (see Extended Data Table 1c).
e) Inflammatory Response (MSigDB Cancer Hallmark). Kaplan-Meier OS and PFS, and boxplots.
f) Interferon Alpha Response (MSigDB Cancer Hallmark). Kaplan-Meier OS and PFS, and boxplots.
g) Interferon Gamma Response (MSigDB Cancer Hallmark). Kaplan-Meier OS and PFS, and boxplots.
h) Allograft Rejection (MSigDB Cancer Hallmark). Kaplan-Meier OS and PFS, and boxplots.
i) IL2/STAT5 Signaling (MSigDB Cancer Hallmark). Kaplan-Meier OS and PFS, and boxplots.
j) IL6/JAK/STAT3 Signaling (MSigDB Cancer Hallmark). Kaplan-Meier OS and PFS, and boxplots.
k) Complement (MSigDB Cancer Hallmark). Kaplan-Meier OS and PFS, and boxplots.
l) Immature Dendritic Cells (Inferred Immune Cell Type). Kaplan-Meier PFS, and boxplots.
m) T Helper Cells (Inferred Immune Cell Type). Kaplan-Meier OS and PFS, and boxplots.
n) Regulatory T Cells. Inferred Immune Cell Type). Boxplots.
o) Central Memory T Cells (TCM; Inferred Immune Cell Type). Boxplots.
p) Effector Memory T Cells (TEM; Inferred Immune Cell Type). Kaplan-Meier OS and PFS, and boxplots.
q) Activated CD8$^+$ T Cells (Inferred Immune Cell Type). Kaplan-Meier OS and PFS, and boxplots.
r) Cytotoxic Cells (Inferred Immune Cell Type). Kaplan-Meier OS and PFS, and boxplots.
s) NK CD56^{dim} Cells (Inferred Immune Cell Type). Kaplan-Meier OS and PFS, and boxplots.
t) Gamma Delta T Cells (Inferred Immune Cell Type). Kaplan-Meier OS and PFS, and boxplots.
u) B Cells (Inferred Immune Cell Type). Boxplots.

v-ee) *mIHC immune cell lineages and states*. Kaplan-Meier OS and PFS in Pre-Treatment (left) and On-Olaparib (right) biopsies. Cell phenotype densities (treated as continuous variables) discretized as High

(≥ 0, red) and Low (< 0, blue). HRs and p-values from CoxPH and log-rank tests are shown; shaded
areas indicate 95% CI for the log survival probability. OS and PFS were measured as above. Boxplots:
cell phenotype densities in biopsies from NR (purple) and R (orange); p-values from Student's t-test.
Features shown were identified by Boruta feature selection, achieved an F1 score >0.7 from random
train/test split analysis, and were significant (p <0.05) in pre-treatment and/or on-treatment samples (see
Extended Data Table 1c).
v) Immune Cells. Boxplots.
w) Ki-67$^+$ Immune Cells. Kaplan-Meier OS and PFS and boxplots.
x) CD4$^+$ T Cells. Boxplots.
y) TBET$^-$ FOXP3$^-$ CD4$^+$ T Cells. Boxplots.
z) CD8$^+$ T Cells. Kaplan-Meier OS and boxplots.
aa) Ki-67$^+$ CD8$^+$ T Cells. Kaplan-Meier OS and boxplots.
bb) TBET/EOMES/PD-1$^-$ CD8$^+$ T Cells. Boxplots.
cc) Ki-67$^+$ TBET/EOMES/PD-1$^-$ CD8$^+$ T Cells. Kaplan-Meier OS and PFS and boxplots.
dd CD20$^+$ B Cells. Boxplots.
ee) Other Immune Cells. Kaplan-Meier OS and PFS and boxplots.
1486
ff-ll) *DSP immune protein expression.* Kaplan-Meier OS and PFS in Pre-Treatment (left) and On-
Olaparib (right) biopsies. Protein levels in both tumor and stroma regions of interest categorized as High
(≥60 z-score percentile, red) and Low (<60 percentile, blue). HR and p-values from CoxPH and log-rank
tests are shown; shaded areas indicate 95% CI for the log survival probability. OS and PFS were
measured as above. Boxplots: protein levels in biopsies from NR (purple) and R (orange); p-values from
Student's t-test. Features shown were identified by Boruta feature selection, achieved an F1 score >0.7
from random train/test split analysis, and were significant (p <0.05) in pre-treatment and/or on-treatment
samples (see Extended Data Table 1c).
ff) CD45 Tumor. Kaplan-Meier OS and PFS and boxplots.
gg) CD45 Stroma. Kaplan-Meier PFS and boxplots.
hh) CD3 Tumor. Kaplan-Meier OS and PFS and boxplots.
ii) CD3 Stroma. Kaplan-Meier PFS and boxplots.
jj) CD8 Stroma. Kaplan-Meier PFS and boxplots.
kk) GZMA Stroma. Boxplots.
ll) PD-L1 Tumor. Boxplots.
1502
mm) *ZAP-70 protein expression from RPPA.* Kaplan-Meier OS and PFS in Pre-Treatment (left) and On-
Olaparib (right) biopsies. Scaled and normalized immune protein expression categorized as High (MAD
z-score ≥0, red) and Low (MAD z-score <0, blue). HRs and p-values from CoxPH and log-rank tests are
shown; shaded areas indicate 95% CI for the log survival probability. OS and PFS were measured as
above. Boxplots: protein expression in biopsies from NR (purple) and R (orange); p-values from Student's
t-test.
1509
1510 **Extended Data Figure 4. Olaparib-induced response markers and signatures.**
1511
1512 **a-g)** *Changes following olaparib treatment.* Boxplots comparing feature scores in Pre-Treatment (Pre-)
1513 and On-Olaparib (On-) biopsies, separated by non-responders (NR, purple, left) and responders (R,

orange, right). Lines connect paired Pre- and On- samples from the same participant. P-values from paired Student's t-test for Pre- vs. On- biopsies. Markers from Fig. 4 and p<0.05 are shown.

a) Immune markers increased in responders, decreased in non-responders (see Fig. 4a).

b) Immune markers increased in responders, unchanged in non-responders (see Fig. 4b).

c) Immune markers unchanged in responders, decreased in non-responders (see Fig. 4c).

d) Malignant cell markers increased in responders, decreased in non-responders (see Fig. 4e).

e) Malignant cell markers increased in responders, unchanged in non-responders (see Fig. 4f).

f) Malignant cell markers decreased in responders, increased in non-responders (see Fig. 4g).

g) PD-L1 Tumor DSP.

Extended Data Figure 5. TNBC subtype correlates of response to olaparib and durvalumab.

a) *TNBC subtypes.* Kaplan-Meier OS stratified by TNBC RNA subtype (BLIS/LAR, red; Non-BLIS/LAR, blue) in Pre-Treatment (left) and On-Olaparib (right) biopsies. HR and p-values from CoxPH and log-rank tests are shown; shaded areas indicate 95% CI for the log survival probability. OS measured from olaparib start until death or data cutoff; right-censoring denoted by "+".

b) *BLIA interferon-regulated genes.* The Interferome web tool (http://www.interferome.org), a database of interferon-regulated genes, was used to identify a subset of highly ranked genes (median Interferome gene change >2, dashed lines, Y-axis) associated with the BLIA subtype (>0.5, X-axis).

c) *TNBC subtype genes within cell types.* Dot plot visualizing expression of representative marker genes (x-axis, bottom) within TNBC subtypes (x-axis, top) across distinct cell type clusters (y-axis). Data are from a single-cell RNAseq breast cancer dataset (see *Methods*). Relative gene expression levels are represented by the color scale, and the fraction of cells within each cell type expressing the gene is represented by the dot size. The plot highlights differential expression patterns of key TNBC markers within each cell cluster, providing insight into the heterogeneity and distribution of TNBC-related gene expression profiles at the single-cell level.

d) *mTNBC3s gene signature.* Kaplan-Meier OS stratified by mTNBC3s[54] predicted non-responders (pPD, red) and predicted responders (pSDPR, blue). OS was measured as above.

References

Foulkes, W. D., Smith, I. E. & Reis-Filho, J. S. Triple-negative breast cancer. *N Engl J Med* **363**, 1938-1948 (2010). https://doi.org:10.1056/NEJMra1001389

Bardia, A. *et al.* Sacituzumab Govitecan in Metastatic Triple-Negative Breast Cancer. *N Engl J Med* **384**, 1529-1541 (2021). https://doi.org:10.1056/NEJMoa2028485

Cortés, J. *et al.* LBA16 KEYNOTE-355: Final results from a randomized, double-blind phase III study of first-line pembrolizumab + chemotherapy vs placebo + chemotherapy for metastatic TNBC. *Annals of Oncology* **32**, S1289-S1290 (2021). https://doi.org:10.1016/j.annonc.2021.08.2089

Modi, S. *et al.* Trastuzumab deruxtecan (T-DXd) versus treatment of physician's choice (TPC) in patients (pts) with HER2-low unresectable and/or metastatic breast cancer (mBC): Results of DESTINY-Breast04, a randomized, phase 3 study. *Journal of Clinical Oncology* **40**, LBA3-LBA3 (2022). https://doi.org:10.1200/JCO.2022.40.17_suppl.LBA3

Glodzik, D. *et al.* Comprehensive molecular comparison of BRCA1 hypermethylated and BRCA1 mutated triple negative breast cancers. *Nat Commun* **11**, 3747 (2020). https://doi.org:10.1038/s41467-020-17537-2

Shimelis, H. *et al.* Triple-Negative Breast Cancer Risk Genes Identified by Multigene Hereditary Cancer Panel Testing. *J Natl Cancer Inst* **110**, 855-862 (2018). https://doi.org:10.1093/jnci/djy106

Winter, C. *et al.* Targeted sequencing of BRCA1 and BRCA2 across a large unselected breast cancer cohort suggests that one-third of mutations are somatic. *Ann Oncol* **27**, 1532-1538 (2016). https://doi.org:10.1093/annonc/mdw209

Morganti, S. *et al.* PARP Inhibitors for Breast Cancer Treatment: A Review. *JAMA Oncol* **10**, 658-670 (2024). https://doi.org:10.1001/jamaoncol.2023.7322

Litton, J. K. *et al.* Talazoparib in Patients with Advanced Breast Cancer and a Germline BRCA Mutation. *N Engl J Med* **379**, 753-763 (2018). https://doi.org:10.1056/NEJMoa1802905

Robson, M. *et al.* Olaparib for Metastatic Breast Cancer in Patients with a Germline BRCA Mutation. *N Engl J Med* **377**, 523-533 (2017). https://doi.org:10.1056/NEJMoa1706450

Lynce, F. & Robson, M. Clinical Use of PARP Inhibitors in BRCA Mutant and Non-BRCA Mutant Breast Cancer. *Cancer Treat Res* **186**, 91-102 (2023). https://doi.org:10.1007/978-3-031-30065-3_6

Tutt, A. *et al.* 161O VIOLETTE: Randomised phase II study of olaparib (ola) + ceralasertib (cer) or adavosertib (ada) vs ola alone in patients (pts) with metastatic triple-negative breast cancer (mTNBC). *Annals of Oncology* **33**, S194-S195 (2022). https://doi.org:10.1016/j.annonc.2022.03.180

Konstantinopoulos, P. A. *et al.* Single-Arm Phases 1 and 2 Trial of Niraparib in Combination With Pembrolizumab in Patients With Recurrent Platinum-Resistant Ovarian Carcinoma. *JAMA Oncol* **5**, 1141-1149 (2019). https://doi.org:10.1001/jamaoncol.2019.1048

Labrie, M. *et al.* Multiomics analysis of serial PARP inhibitor treated metastatic TNBC inform on rational combination therapies. *NPJ Precis Oncol* **5**, 92 (2021). https://doi.org:10.1038/s41698-021-00232-w

Westin, S. N. *et al.* Phase Ib Dose Expansion and Translational Analyses of Olaparib in Combination with Capivasertib in Recurrent Endometrial, Triple-Negative Breast, and Ovarian Cancer. *Clin Cancer Res* **27**, 6354-6365 (2021). https://doi.org:10.1158/1078-0432.CCR-21-1656

16 Morad, G., Helmink, B. A., Sharma, P. & Wargo, J. A. Hallmarks of response, resistance, and
toxicity to immune checkpoint blockade. *Cell* **184**, 5309-5337 (2021).
https://doi.org:10.1016/j.cell.2021.09.020

17 Sun, Q. *et al.* Immune checkpoint therapy for solid tumours: clinical dilemmas and future trends.
Signal Transduct Target Ther **8**, 320 (2023). https://doi.org:10.1038/s41392-023-01522-4

18 Cortes, J. *et al.* Pembrolizumab plus Chemotherapy in Advanced Triple-Negative Breast Cancer.
N Engl J Med **387**, 217-226 (2022). https://doi.org:10.1056/NEJMoa2202809

19 Haricharan, S., Bainbridge, M. N., Scheet, P. & Brown, P. H. Somatic mutation load of estrogen
receptor-positive breast tumors predicts overall survival: an analysis of genome sequence data.
Breast Cancer Res Treat **146**, 211-220 (2014). https://doi.org:10.1007/s10549-014-2991-x

20 Fang, Y. *et al.* Sequential Therapy with PARP and WEE1 Inhibitors Minimizes Toxicity while
Maintaining Efficacy. *Cancer Cell* **35**, 851-867 e857 (2019).
https://doi.org:10.1016/j.ccell.2019.05.001

21 Huang, J. *et al.* The PARP1 inhibitor BMN 673 exhibits immunoregulatory effects in a Brca1(-/-
) murine model of ovarian cancer. *Biochem Biophys Res Commun* **463**, 551-556 (2015).
https://doi.org:10.1016/j.bbrc.2015.05.083

22 Pantelidou, C. *et al.* PARP Inhibitor Efficacy Depends on CD8(+) T-cell Recruitment via
Intratumoral STING Pathway Activation in BRCA-Deficient Models of Triple-Negative Breast
Cancer. *Cancer Discov* **9**, 722-737 (2019). https://doi.org:10.1158/2159-8290.CD-18-1218

23 Shen, J. *et al.* PARPi Triggers the STING-Dependent Immune Response and Enhances the
Therapeutic Efficacy of Immune Checkpoint Blockade Independent of BRCAness. *Cancer Res*
79, 311-319 (2019). https://doi.org:10.1158/0008-5472.CAN-18-1003

24 Wang, Z. *et al.* Niraparib activates interferon signaling and potentiates anti-PD-1 antibody
efficacy in tumor models. *Sci Rep* **9**, 1853 (2019). https://doi.org:10.1038/s41598-019-38534-6

25 Farkkila, A. *et al.* Immunogenomic profiling determines responses to combined PARP and PD-1
inhibition in ovarian cancer. *Nat Commun* **11**, 1459 (2020). https://doi.org:10.1038/s41467-020-
15315-8

26 Labrie, M. *et al.* Adaptive responses in a PARP inhibitor window of opportunity trial illustrate
limited functional interlesional heterogeneity and potential combination therapy options.
Oncotarget **10**, 3533-3546 (2019). https://doi.org:10.18632/oncotarget.26947

27 Labrie, M., Brugge, J. S., Mills, G. B. & Zervantonakis, I. K. Therapy resistance: opportunities
created by adaptive responses to targeted therapies in cancer. *Nat Rev Cancer* **22**, 323-339
(2022). https://doi.org:10.1038/s41568-022-00454-5

28 Ding, L. *et al.* PARP Inhibition Elicits STING-Dependent Antitumor Immunity in Brca1-
Deficient Ovarian Cancer. *Cell Rep* **25**, 2972-2980 e2975 (2018).
https://doi.org:10.1016/j.celrep.2018.11.054

29 Domchek, S. M. *et al.* Olaparib and durvalumab in patients with germline BRCA-mutated
metastatic breast cancer (MEDIOLA): an open-label, multicentre, phase 1/2, basket study.
Lancet Oncol **21**, 1155-1164 (2020). https://doi.org:10.1016/S1470-2045(20)30324-7

30 Vinayak, S. *et al.* Open-label Clinical Trial of Niraparib Combined With Pembrolizumab for
Treatment of Advanced or Metastatic Triple-Negative Breast Cancer. *JAMA Oncol* **5**, 1132-1140
(2019). https://doi.org:10.1001/jamaoncol.2019.1029

31 Yap, T. A. *et al.* Avelumab Plus Talazoparib in Patients With Advanced Solid Tumors: The
JAVELIN PARP Medley Nonrandomized Controlled Trial. *JAMA Oncol* **9**, 40-50 (2023).
https://doi.org:10.1001/jamaoncol.2022.5228

32   Eisenhauer, E. A. *et al.* New response evaluation criteria in solid tumours: revised RECIST
guideline (version 1.1). *Eur J Cancer* **45**, 228-247 (2009).
https://doi.org:10.1016/j.ejca.2008.10.026
33   Swisher, E. M. *et al.* Rucaparib in relapsed, platinum-sensitive high-grade ovarian carcinoma
(ARIEL2 Part 1): an international, multicentre, open-label, phase 2 trial. *Lancet Oncol* **18**, 75-87
(2017). https://doi.org:10.1016/S1470-2045(16)30559-9
34   Tung, N. M. *et al.* TBCRC 048: Phase II Study of Olaparib for Metastatic Breast Cancer and
Mutations in Homologous Recombination-Related Genes. *J Clin Oncol* **38**, 4274-4282 (2020).
https://doi.org:10.1200/JCO.20.02151
35   Castroviejo-Bermejo, M. *et al.* A RAD51 assay feasible in routine tumor samples calls PARP
inhibitor response beyond BRCA mutation. *EMBO Mol Med* **10** (2018).
https://doi.org:10.15252/emmm.201809172
36   Eikesdal, H. P. *et al.* Olaparib monotherapy as primary treatment in unselected triple negative
breast cancer. *Ann Oncol* **32**, 240-249 (2021). https://doi.org:10.1016/j.annonc.2020.11.009
37   Miller, R. E. *et al.* ESMO recommendations on predictive biomarker testing for homologous
recombination deficiency and PARP inhibitor benefit in ovarian cancer. *Ann Oncol* **31**, 1606-
1622 (2020). https://doi.org:10.1016/j.annonc.2020.08.2102
38   Ring, A. *et al.* Olaparib and Ceralasertib (AZD6738) in Patients with Triple-Negative Advanced
Breast Cancer: Results from Cohort E of the plasmaMATCH Trial (CRUK/15/010). *Clin Cancer
Res* **29**, 4751-4759 (2023). https://doi.org:10.1158/1078-0432.CCR-23-1696
39   Ozmen, T. Y. *et al.* Automated machine learning profiling with MAP-HR for quantifying
homologous recombination foci in patient samples. *NAR Cancer* **7**, zcaf025 (2025).
https://doi.org:10.1093/narcan/zcaf025
40   Alexandrov, L. B. *et al.* Signatures of mutational processes in human cancer. *Nature* **500**, 415-
421 (2013). https://doi.org:10.1038/nature12477
41   Guan, X. *et al.* Androgen receptor activity in T cells limits checkpoint blockade efficacy. *Nature*
606, 791-796 (2022). https://doi.org:10.1038/s41586-022-04522-6
42   Hargrove-Wiley, E. & Fingleton, B. Sex Hormones in Breast Cancer Immunity. *Cancer Res* **83**,
12-19 (2023). https://doi.org:10.1158/0008-5472.CAN-22-1829
43   Yang, C. *et al.* Androgen receptor-mediated CD8(+) T cell stemness programs drive sex
differences in antitumor immunity. *Immunity* **55**, 1268-1283 e1269 (2022).
https://doi.org:10.1016/j.immuni.2022.05.012
44   Lim, B. *et al.* Phase 2 study of neoadjuvant enzalutamide and paclitaxel for luminal androgen
receptor-enriched TNBC: Trial results and insights into "ARness". *Cell Rep Med* **5**, 101595
(2024). https://doi.org:10.1016/j.xcrm.2024.101595
45   Sun, C. *et al.* Rational combination therapy with PARP and MEK inhibitors capitalizes on
therapeutic liabilities in RAS mutant cancers. *Sci Transl Med* **9** (2017).
https://doi.org:10.1126/scitranslmed.aal5148
46   Yang, B. *et al.* MEK Inhibition Remodels the Immune Landscape of Mutant KRAS Tumors to
Overcome Resistance to PARP and Immune Checkpoint Inhibitors. *Cancer Res* **81**, 2714-2729
(2021). https://doi.org:10.1158/0008-5472.CAN-20-2370
47   Althammer, S. *et al.* Automated image analysis of NSCLC biopsies to predict response to anti-
PD-L1 therapy. *J Immunother Cancer* **7**, 121 (2019). https://doi.org:10.1186/s40425-019-0589-x
48   Chebib, I. & Mino-Kenudson, M. PD-L1 immunohistochemistry: Clones, cutoffs, and
controversies. *APMIS* **130**, 295-313 (2022). https://doi.org:10.1111/apm.13223

49 Cortes, J. *et al.* Pembrolizumab plus chemotherapy versus placebo plus chemotherapy for
previously untreated locally recurrent inoperable or metastatic triple-negative breast cancer
(KEYNOTE-355): a randomised, placebo-controlled, double-blind, phase 3 clinical trial. *Lancet*
396, 1817-1828 (2020). https://doi.org:10.1016/S0140-6736(20)32531-9

50 Schmid, P. *et al.* Atezolizumab and Nab-Paclitaxel in Advanced Triple-Negative Breast Cancer.
N Engl J Med **379**, 2108-2121 (2018). https://doi.org:10.1056/NEJMoa1809615

51 Burstein, M. D. *et al.* Comprehensive genomic analysis identifies novel subtypes and targets of
triple-negative breast cancer. *Clin Cancer Res* **21**, 1688-1698 (2015).
https://doi.org:10.1158/1078-0432.CCR-14-0432

52 Ding, Y. C. *et al.* Molecular subtypes of triple-negative breast cancer in women of different race
and ethnicity. *Oncotarget* **10**, 198-208 (2019). https://doi.org:10.18632/oncotarget.26559

53 Mitri, Z. I. *et al.* Implementing a comprehensive translational oncology platform: from molecular
testing to actionability. *J Transl Med* **16**, 358 (2018). https://doi.org:10.1186/s12967-018-1733-y

54 Bottomly, D. *et al.* Utilizing cohort-level and individual networks to predict best response in
patients with metastatic triple negative breast cancer. *NPJ Precis Oncol* **9**, 179 (2025).
https://doi.org:10.1038/s41698-025-00959-w

55 Batalini, F. *et al.* Phase 1b Clinical Trial with Alpelisib plus Olaparib for Patients with Advanced
Triple-Negative Breast Cancer. *Clin Cancer Res* **28**, 1493-1499 (2022).
https://doi.org:10.1158/1078-0432.CCR-21-3045

56 Russell, B. L., Sooklal, S. A., Malindisa, S. T., Daka, L. J. & Ntwasa, M. The Tumor
Microenvironment Factors That Promote Resistance to Immune Checkpoint Blockade Therapy.
Front Oncol **11**, 641428 (2021). https://doi.org:10.3389/fonc.2021.641428

57 Akbani, R. *et al.* A pan-cancer proteomic perspective on The Cancer Genome Atlas. *Nat*
Commun **5**, 3887 (2014). https://doi.org:10.1038/ncomms4887

58 Allison, K. H. *et al.* Estrogen and Progesterone Receptor Testing in Breast Cancer: ASCO/CAP
Guideline Update. *J Clin Oncol* **38**, 1346-1366 (2020). https://doi.org:10.1200/JCO.19.02309

59 Simon, R. Optimal two-stage designs for phase II clinical trials. *Control Clin Trials* **10**, 1-10
(1989). https://doi.org:10.1016/0197-2456(89)90015-9

60 Johnson, B. E. *et al.* An omic and multidimensional spatial atlas from serial biopsies of an
evolving metastatic breast cancer. *Cell Rep Med* **3**, 100525 (2022).
https://doi.org:10.1016/j.xcrm.2022.100525

61 Gambella, A. *et al.* Section detachment in immunohistochemistry: causes, troubleshooting, and
problem-solving. *Histochem Cell Biol* **148**, 95-101 (2017). https://doi.org:10.1007/s00418-017-
1558-4

62 Kursa, M. B. & Rudnicki, W. R. Feature Selection with the Boruta Package. *Journal of*
Statistical Software **36**, 1 - 13 (2010). https://doi.org:10.18637/jss.v036.i11

63 Li, H. *et al.* High expression of vinculin predicts poor prognosis and distant metastasis and
associates with influencing tumor-associated NK cell infiltration and epithelial-mesenchymal
transition in gastric cancer. *Aging (Albany NY)* **13**, 5197-5225 (2021).
https://doi.org:10.18632/aging.202440

64 McKenna, A. *et al.* The Genome Analysis Toolkit: a MapReduce framework for analyzing next-
generation DNA sequencing data. *Genome Res* **20**, 1297-1303 (2010).
https://doi.org:10.1101/gr.107524.110

65 Karczewski, K. J. *et al.* The mutational constraint spectrum quantified from variation in 141,456
humans. *Nature* **581**, 434-443 (2020). https://doi.org:10.1038/s41586-020-2308-7

66   Afgan, E. *et al.* The Galaxy platform for accessible, reproducible and collaborative biomedical
analyses: 2018 update. *Nucleic Acids Res* **46**, W537-W544 (2018).
https://doi.org/10.1093/nar/gky379
67   Manders, F. *et al.* MutationalPatterns: the one stop shop for the analysis of mutational processes.
BMC Genomics **23**, 134 (2022). https://doi.org/10.1186/s12864-022-08357-3
68   Bray, N. L., Pimentel, H., Melsted, P. & Pachter, L. Near-optimal probabilistic RNA-seq
quantification. *Nat Biotechnol* **34**, 525-527 (2016). https://doi.org/10.1038/nbt.3519
69   Frankish, A. *et al.* Gencode 2021. *Nucleic Acids Res* **49**, D916-D923 (2021).
https://doi.org/10.1093/nar/gkaa1087
70   Hanzelmann, S., Castelo, R. & Guinney, J. GSVA: gene set variation analysis for microarray and
RNA-seq data. *BMC Bioinformatics* **14**, 7 (2013). https://doi.org/10.1186/1471-2105-14-7
71   Liberzon, A. *et al.* The Molecular Signatures Database (MSigDB) hallmark gene set collection.
Cell Syst **1**, 417-425 (2015). https://doi.org/10.1016/j.cels.2015.12.004
72   Tamborero, D. *et al.* A Pan-cancer Landscape of Interactions between Solid Tumors and
Infiltrating Immune Cell Populations. *Clin Cancer Res* **24**, 3717-3728 (2018).
https://doi.org/10.1158/1078-0432.Ccr-17-3509
73   Banik, G. *et al.* High-dimensional multiplexed immunohistochemical characterization of immune
contexture in human cancers. *Methods Enzymol* **635**, 1-20 (2020).
https://doi.org/10.1016/bs.mie.2019.05.039
74   Bergholtz, H. *et al.* Best Practices for Spatial Profiling for Breast Cancer Research with the
GeoMx((R)) Digital Spatial Profiler. *Cancers (Basel)* **13** (2021).
https://doi.org/10.3390/cancers13174456
75   Labrie, M. *et al.* Exploration of markers of synergistic lethality of PARP and PI3K-akt-mTOR
inhibitors in women's cancers. *Gynecologic Oncology* **154**, 60 (2019).
https://doi.org:https://doi.org/10.1016/j.ygyno.2019.04.143
76   Tibes, R. *et al.* Reverse phase protein array: validation of a novel proteomic technology and
utility for analysis of primary leukemia specimens and hematopoietic stem cells. *Mol Cancer*
Ther **5**, 2512-2521 (2006). https://doi.org/10.1158/1535-7163.Mct-06-0334
77   Shehwana, H. *et al.* RPPA SPACE: an R package for normalization and quantitation of Reverse-
Phase Protein Array data. *Bioinformatics* **38**, 5131-5133 (2022).
https://doi.org/10.1093/bioinformatics/btac665
78   Risso, D., Ngai, J., Speed, T. P. & Dudoit, S. Normalization of RNA-seq data using factor
analysis of control genes or samples. *Nat Biotechnol* **32**, 896-902 (2014).
https://doi.org/10.1038/nbt.2931
79   Wolf, F. A., Angerer, P. & Theis, F. J. SCANPY: large-scale single-cell gene expression data
analysis. *Genome Biol* **19**, 15 (2018). https://doi.org/10.1186/s13059-017-1382-0
80   Wolock, S. L., Lopez, R. & Klein, A. M. Scrublet: Computational Identification of Cell Doublets
in Single-Cell Transcriptomic Data. *Cell Syst* **8**, 281-291 e289 (2019).
https://doi.org/10.1016/j.cels.2018.11.005
81   Hu, C. *et al.* CellMarker 2.0: an updated database of manually curated cell markers in
human/mouse and web tools based on scRNA-seq data. *Nucleic Acids Res* **51**, D870-D876
(2023). https://doi.org/10.1093/nar/gkac947
82   Franzen, O., Gan, L. M. & Bjorkegren, J. L. M. PanglaoDB: a web server for exploration of
mouse and human single-cell RNA sequencing data. *Database (Oxford)* **2019** (2019).
https://doi.org/10.1093/database/baz046

83   Xu, C. *et al.* Automatic cell-type harmonization and integration across Human Cell Atlas
datasets. *Cell* **186**, 5876-5891 e5820 (2023). https://doi.org:10.1016/j.cell.2023.11.026
84   Tickle, T., Tirosh, I., Georgescu, C., Brown, M. & Haas, B.   (Cambridge, MA, 2019).
85   Serin Harmanci, A., Harmanci, A. O. & Zhou, X. CaSpER identifies and visualizes CNV events
by integrative analysis of single-cell or bulk RNA-sequencing data. *Nat Commun* **11**, 89 (2020).
https://doi.org:10.1038/s41467-019-13779-x
86   Gayoso, A. *et al.* A Python library for probabilistic analysis of single-cell omics data. *Nat*
Biotechnol **40**, 163-166 (2022). https://doi.org:10.1038/s41587-021-01206-w
87   Xu, C. *et al.* Probabilistic harmonization and annotation of single-cell transcriptomics data with
deep generative models. *Mol Syst Biol* **17**, e9620 (2021). https://doi.org:10.15252/msb.20209620
88   Tang, F. *et al.* A pan-cancer single-cell panorama of human natural killer cells. *Cell* **186**, 4235-
4251 e4220 (2023). https://doi.org:10.1016/j.cell.2023.07.034
89   Dussiau, C. *et al.* Hematopoietic differentiation is characterized by a transient peak of entropy at
a single-cell level. *BMC Biol* **20**, 60 (2022). https://doi.org:10.1186/s12915-022-01264-9
90   Cords, L. *et al.* Cancer-associated fibroblast classification in single-cell and spatial proteomics
data. *Nat Commun* **14**, 4294 (2023). https://doi.org:10.1038/s41467-023-39762-1
91   Lotfollahi, M. *et al.* Mapping single-cell data to reference atlases by transfer learning. *Nat*
Biotechnol **40**, 121-130 (2022). https://doi.org:10.1038/s41587-021-01001-7
92   Crowdis, J., He, M. X., Reardon, B. & Van Allen, E. M. CoMut: visualizing integrated
molecular information with comutation plots. *Bioinformatics* **36**, 4348-4349 (2020).
https://doi.org:10.1093/bioinformatics/btaa554
1794

Table 1

		N = 24 (%)
Median age (range)	54.0 (25.3 – 78.9)	
ECOG performance status	0	15 (62.5%)
	1	9 (37.5%)
Race	Caucasian	21 (87.5%)
	African American	1 (4.2%)
	Undisclosed	2 (8.3%)
Menopausal status	Pre-menopausal	8 (33.3%)
	Post-menopausal	16 (66.7%)
Lines of therapy in metastatic setting	0	8 (33.3%)
	1	10 (41.7%)
	2	6 (25.0%)
Prior platinum chemotherapy in metastatic setting	Yes	7 (29.2%)
	No	17 (70.8%)
History of brain metastases	Yes	7 (29.2%)
	No	17 (70.8%)

Figure 1

Figure 2

Figure 3 – Microenvironment Signatures

Figure 4 – Olaparib-Induced Response Markers and Signatures

Figure 5 – TNBC Subtypes

Figure 6 – Three-Protein Response Predictor

Extended Data Table 1a

	Participants (N=24)	Adverse events (%)	Treatment-related adverse events (N)
Grade 1	23	222 (55%)	129
Grade 2	18	140 (34%)	61
Grade 3	13	41 (10%)	20
Grade 4	2	2 (<1%)	0
Grade 5	2	2 (<1%)	0
Total		407	210

Extended Data Table 1b

Adverse Event	Grade	Events (N)	Participants (N)	Treatment-related (N)	Immune-related (N)
Investigations – other*	5	1	1	0	
Respiratory failure	5	1	1	0	
Sepsis	4	1	1	0	
Urinary retention	4	1	1	0	
Anemia	3	9	6	1	
Alanine aminotransferase increased	3	1	1	1	1
Alkaline phosphatase increased	3	1	1	0	
Anxiety	3	2	2	0	
Aspartate aminotransferase increased	3	3	2	1	1
Blood and lymphatic system disorders – other	3	1	1	1	
Blood bilirubin increased	3	1	1	0	
Diarrhea	3	1	1	1	
Depression	3	1	1	0	
Dysphagia	3	1	1	0	
Dyspnea	3	1	1	1	
Eosinophilia	3	1	1	1	
Generalized muscle weakness	3	2	2	0	
Headache	3	1	1	1	
Hip fracture	3	1	1	0	
Hoarseness	3	1	1	0	
Hypoglycemia	3	1	1	0	
Pain	3	2	2	0	
Pain in extremity	3	1	1	0	
Pelvic pain	3	1	1	0	
Pleural effusion	3	2	1	0	
Sepsis	3	1	1	1	
Skin and subcutaneous tissue disorders – other	3	1	1	0	
Superior vena cava syndrome	3	1	1	0	
Syncope	3	1	1	1	
Thrush	3	1	1	0	
Vomiting	3	1	1	1	

Extended Data Table 1c

Feature	Category	Omic	Assay	Boruta Selected?	F1 Median Pre-	F1 Median On-	BR p-val, Pre-	BR p-val, On-	PFS p-val, Pre-	PFS p-val, On-	OS p-val, Pre-	OS p-val, On-
PAR	Tumor Intrinsic	Protein	RPPA	Yes	0.91	ns	0.0023	ns	ns	ns	ns	ns
MutSig3	Tumor Intrinsic	DNA	WES COSMIC	N/A	N/A	N/A	ns	N/A	ns	N/A	0.03	N/A
AR CLIA IHC >10%	Tumor Intrinsic	Protein	IHC	N/A	ns	0.75	ns	ns	ns	ns	ns	0.036
PI3K-AKT PATHWAY DSP	Tumor Intrinsic	Protein	DSP	N/A	N/A	N/A	ns	0.011	ns	0.026	0.023	0.022
MIXED SIGNALING PATHWAY	Tumor Intrinsic	Protein	DSP	N/A	N/A	N/A	ns	<0.001	0.027	0.001	0.012	0.0003
PTEN Copy Loss	Tumor Intrinsic	DNA	GeneTrails DNAseq	N/A	0.86	0.8	0.018	0.029	0.002	0.008	0.025	0.009
PI3K-AKT Gene Alterations	Tumor Intrinsic	DNA	GeneTrails DNAseq	N/A	0.80	0.78	0.013	ns	0.018	ns	ns	ns
p-GSK3B	Tumor Intrinsic	Protein	DSP	Yes	ns	0.8	ns	0.008	ns	0.047	ns	ns
IMMUNE RNA SUMMARY	Microenvironment	RNA	RNAseq	N/A	ns	0.9	ns	0.001	ns	0.008	ns	0.007
IMMUNE mIHC SUMMARY	Microenvironment	Protein	mIHC	N/A	ns	0.9	ns	0.001	ns	0.012	ns	0.008
IMMUNE DSP SUMMARY	Microenvironment	Protein	DSP	N/A	0.73	0.8	0.029	0.003	0.004	0.015	0.019	0.014
MULTI-OMIC IMMUNE CONSENSUS	Microenvironment	Multi	RNAseq, mIHC, DSP	N/A	0.74	0.9	ns	<0.001	ns	0.002	0.018	0.0005
Inflammatory Response	Microenvironment	RNA	RNAseq	Yes	ns	0.82	ns	<0.001	ns	0.029	ns	0.029
Interferon Alpha Response	Microenvironment	RNA	RNAseq	Yes	0.75	0.78	0.012	<0.001	ns	ns	0.011	ns
Interferon Gamma Response	Microenvironment	RNA	RNAseq	Yes	0.75	0.88	0.028	<0.001	ns	0.04	0.03	0.009
Allograft Rejection	Microenvironment	RNA	RNAseq	Yes	0.71	0.94	ns	<0.001	ns	0.011	ns	0.002
IL2/STAT5 Signaling	Microenvironment	RNA	RNAseq	Yes	ns	0.82	ns	0.007	ns	0.029	ns	0.029
IL6/JAK/STAT3 Signaling	Microenvironment	RNA	RNAseq	Yes	ns	0.82	ns	<0.001	ns	0.029	ns	0.029
Complement	Microenvironment	RNA	RNAseq	Yes	ns	0.82	ns	0.003	ns	0.029	ns	0.029
Immature Dendritic Cells	Microenvironment	RNA	RNAseq	Yes	ns	0.82	ns	0.031	ns	0.042	ns	ns
T Helper Cells	Microenvironment	RNA	RNAseq	Yes	ns	0.80	ns	0.004	ns	0.026	ns	0.02
Regulatory T Cells	Microenvironment	RNA	RNAseq	Yes	0.71	0.78	ns	0.0059	ns	ns	ns	ns
Central Memory T Cells (TCM)	Microenvironment	RNA	RNAseq	Yes	ns	0.71	ns	0.015	ns	ns	ns	ns
Effector Memory T Cells (TEM)	Microenvironment	RNA	RNAseq	Yes	ns	0.82	ns	0.001	ns	0.029	0.02	0.029
Activated CD8$^+$ T Cells	Microenvironment	RNA	RNAseq	Yes	ns	0.82	ns	<0.001	ns	0.021	ns	0.027
Cytotoxic Cells	Microenvironment	RNA	RNAseq	Yes	0.80	0.82	0.033	0.002	ns	0.021	0.009	0.027
NK CD56^{dim} Cells	Microenvironment	RNA	RNAseq	Yes	ns	0.80	ns	0.002	ns	0.013	ns	0.008
Gamma Delta T Cells	Microenvironment	RNA	RNAseq	Yes	ns	0.82	ns	0.004	ns	0.029	ns	0.029
B Cells	Microenvironment	RNA	RNAseq	Yes	0.75	0.75	0.0015	0.0015	ns	ns	ns	ns
Immune Cells	Microenvironment	Protein	mIHC	Yes	ns	0.75	ns	0.0047	ns	ns	ns	ns
Ki-67$^+$ Immune Cells	Microenvironment	Protein	mIHC	Yes	ns	0.92	ns	0.004	ns	0.032	ns	0.019
CD4$^+$ T Cells	Microenvironment	Protein	mIHC	Yes	0.86	0.86	ns	0.0063	ns	ns	ns	ns
TBET$^-$ FOXP3$^-$ CD4$^+$ T Cells	Microenvironment	Protein	mIHC	Yes	ns	0.86	ns	0.01	ns	ns	ns	ns
CD8$^+$ T Cells	Microenvironment	Protein	mIHC	Yes	ns	0.75	ns	0.007	ns	ns	ns	0.028
Ki-67$^+$ CD8$^+$ T Cells	Microenvironment	Protein	mIHC	Yes	ns	0.86	ns	0.001	ns	ns	ns	0.001
TBET/EOMES/PD-1$^-$ CD8$^+$ T Cells	Microenvironment	Protein	mIHC	Yes	ns	0.86	ns	0.0031	ns	ns	ns	ns
Ki-67$^+$ TBET/EOMES/PD-1$^-$ CD8$^+$ T Cells	Microenvironment	Protein	mIHC	Yes	ns	0.86	ns	<0.001	ns	0.032	ns	0.019
CD20$^+$ B Cells	Microenvironment	Protein	mIHC	Yes	0.86	0.83	0.0091	0.0077	ns	ns	ns	ns
Other Immune Cells	Microenvironment	Protein	mIHC	Yes	0.71	0.92	ns	0.009	ns	0.041	ns	0.027
CD45 Tumor	Microenvironment	Protein	DSP	Yes	ns	0.73	0.03	<0.001	0.031	0.02	0.045	0.005
CD45 Stroma	Microenvironment	Protein	DSP	Yes	ns	0.80	ns	0.003	0.007	0.003	ns	ns
CD3 Tumor	Microenvironment	Protein	DSP	Yes	ns	0.73	0.007	<0.001	ns	0.015	ns	0.015
CD3 Stroma	Microenvironment	Protein	DSP	Yes	ns	0.80	ns	<0.001	ns	0.003	ns	ns
CD8 Stroma	Microenvironment	Protein	DSP	Yes	ns	0.70	ns	0.005	0.032	0.003	ns	ns
GZMA Stroma	Microenvironment	Protein	DSP	Yes	ns	0.91	ns	0.001	ns	ns	ns	ns
PD-L1 Tumor	Microenvironment	Protein	DSP	Yes	0.73	0.78	0.022	0.035	ns	ns	ns	ns
ZAP-70	Microenvironment	Protein	RPPA	Yes	0.77	0.80	0.038	0.004	0.015	0.002	ns	0.021
TNBC Subtype	Tumor Intrinsic	RNA	RNAseq	N/A	N/A	0.9	0.006	0.002	0.001	0	0.04	0.005
mTNBC3s	Tumor Intrinsic	RNA	RNAseq	N/A	0.94	N/A	<0.001	0.002	ns	0.0001	0.046	0
THREE-PROTEIN RESPONSE PREDICTOR	Microenvironment	Protein	DSP	N/A	N/A	N/A	ns	<0.001	0.048	0.0001	ns	0.006

Extended Data Table 1d

Feature	Category	Omic	Assay	Boruta Selected?	Pre- to On- Paired T-Test p-val, NR	Pre- to On- Paired T-Test p-val, R	Change from Pre- to On- T-Test p-val
Effector Memory T Cells (TEM)	Microenvironment	RNA	RNAseq	Yes	ns	ns	0.0016
Activated CD8$^+$ T Cells	Microenvironment	RNA	RNAseq	Yes	ns	ns	0.00086
Immune Cells	Microenvironment	Protein	mIHC	Yes	0.036	ns	0.0018
CD3 Tumor	Microenvironment	Protein	DSP	Yes	ns	0.037	0.002
CD4 Tumor	Microenvironment	Protein	DSP	Yes	ns	0.0056	<0.001
GZMA Tumor	Microenvironment	Protein	DSP	Yes	0.00027	ns	<0.001
HLA-DR Stroma	Microenvironment	Protein	DSP	Yes	ns	0.0056	<0.001
HLA-DR Tumor	Microenvironment	Protein	DSP	Yes	ns	0.015	<0.001
PD-L1 Tumor	Microenvironment	Protein	DSP	Yes	0.027	ns	ns
p-GSK3B	Tumor Intrinsic	Protein	DSP	Yes	ns	ns	0.023
B7-H3	Microenvironment	Protein	RPPA	Yes	ns	ns	<0.001
COUP-TFII	Tumor Intrinsic	Protein	RPPA	Yes	ns	ns	0.004
GLI1	Tumor Intrinsic	Protein	RPPA	Yes	ns	0.00057	<0.001
HSP70	Tumor Intrinsic	Protein	RPPA	Yes	0.034	ns	0.002
MMP14	Tumor Intrinsic	Protein	RPPA	Yes	ns	0.046	<0.001
MTCO1	Tumor Intrinsic	Protein	RPPA	Yes	ns	0.0006	<0.001
PAR	Tumor Intrinsic	Protein	RPPA	Yes	0.00091	ns	0.011
SNAIL	Tumor Intrinsic	Protein	RPPA	Yes	ns	ns	<0.001

Extended Data Figure 1

a

b

Extended Data Figure 2

a

Participant S,
Pre-Treatment: Deficient

Participant K,
On-Olaparib: Intermediate

Participant X,
Pre-Treatment: Competent

b HRC/HRD Ratio

c γH2AX+ Ratio

d

Extended Data Figure 3 – Microenvironment Signatures

Extended Data Figure 3 – Microenvironment Signatures (cont.)

Extended Data Figure 3 – Microenvironment Signatures (cont.)

Extended Data Figure 3 – Microenvironment Signatures (cont.)

m

Overall Survival – Pre-Treatment
RNA – T Helper Cells

Strata — High — Low

p-value=0.12 ns

HR 2.21 (95% CI 0.73 - 6.68)
CoxPH p-value=0.16
Log rank p-value=0.15

Number at risk

Strata											
High	9	8	6	5	3	3	1	1	1	0	0
Low	10	7	5	4	0	0	0	0	0	0	0
	0	5	10	15	20	25	30	35	40	45	50

Overall Survival – On-Olaparib
RNA – T Helper Cells

Strata — High — Low

p-value=0.004 **

HR 3.97 (95% CI 1.16 - 13.61)
CoxPH p-value=0.028
Log rank p-value=0.02

Number at risk

Strata											
High	9	9	7	6	3	3	2	2	1	0	0
Low	11	6	4	2	0	0	0	0	0	0	0
	0	5	10	15	20	25	30	35	40	45	50

Progression Free Survival – Pre-Treatment
RNA – T Helper Cells

Strata — High — Low

HR 1.79 (95% CI 0.7 - 4.62)
CoxPH p-value=0.23
Log rank p-value=0.22

Number at risk

Strata					
High	9	6	3	2	0
Low	10	4	1	1	0
	0	5	10	15	20

Progression Free Survival – On-Olaparib
RNA – T Helper Cells

Strata — High — Low

HR 2.92 (95% CI 1.1 - 7.78)
CoxPH p-value=0.032
Log rank p-value=0.026

Number at risk

Strata					
High	9	8	2	2	0
Low	11	2	1	0	0
	0	5	10	15	20

n

RNA – Regulatory T Cells

o

RNA – Central Memory T Cells (TCM)

t

Overall Survival – Pre-Treatment
RNA – Gamma Delta T Cells

Strata — High — Low

HR 2.86 (95% CI 0.74 - 11.1)
CoxPH p-value=0.13
Log rank p-value=0.11

p-value=0.19 ns

Number at risk

Strata											
High	7	5	4	4	3	3	1	1	1	0	0
Low	12	10	7	5	0	0	0	0	0	0	0
	0	5	10	15	20	25	30	35	40	45	50

Overall Survival – On-Olaparib
RNA – Gamma Delta T Cells

Strata — High — Low

HR 3.71 (95% CI 1.07 - 12.84)
CoxPH p-value=0.039
Log rank p-value=0.029

p-value=0.004 **

Number at risk

Strata											
High	9	8	7	6	3	3	2	2	1	0	0
Low	11	7	4	2	0	0	0	0	0	0	0
	0	5	10	15	20	25	30	35	40	45	50

Progression Free Survival – Pre-Treatment
RNA – Gamma Delta T Cells

Strata — High — Low

HR 1.79 (95% CI 0.66 - 4.83)
CoxPH p-value=0.25
Log rank p-value=0.25

Number at risk

Strata					
High	7	5	2	2	0
Low	12	5	2	1	0
	0	5	10	15	20

Progression Free Survival – On-Olaparib
RNA – Gamma Delta T Cells

Strata — High — Low

HR 2.88 (95% CI 1.08 - 7.69)
CoxPH p-value=0.035
Log rank p-value=0.029

Number at risk

Strata					
High	9	8	2	2	0
Low	11	2	1	0	0
	0	5	10	15	20

u   RNA – B Cells

Pre-Treatment   p=0.0015

On-Olaparib   p=0.0015

Extended Data Figure 3 – Microenvironment Signatures (cont.)

v

w

x

Extended Data Figure 3 – Microenvironment Signatures (cont.)

Extended Data Figure 3 – Microenvironment Signatures (cont.)

ee

Extended Data Figure 3 – Microenvironment Signatures (cont.)

Extended Data Figure 3 – Microenvironment Signatures (cont.)

Extended Data Figure 4 – Olaparib-Induced Response Markers and Signatures

Extended Data Figure 5 – TNBC Subtypes

Extended Data Figure 5 – TNBC Subtypes (cont.)

c

d

Overall Survival – Pre-Treatment mTNBC3s

Overall Survival – On-Olaparib mTNBC3s