Routine Testing for *Chlamydia trachomatis* and *Neisseria gonorrhoeae* Infections within an

HIV Pre-Exposure Prophylaxis Program in Hanoi, Vietnam: Implications for Low- and Middle-

Income Countries

Paul C. Adamson[1], Hao T. M. Bui [2], Loc Q Pham[2], Le Minh Giang[2], Jeffrey D. Klausner[3]

Author Affiliations: [1] Division of Infectious Diseases, David Geffen School of Medicine at UCLA,

Los Angeles, CA; [2] Center for Training and Research on Substance Abuse and HIV, Hanoi Medical

University, Vietnam; [3] Department of Population and Public Health Sciences, Keck School of

Medicine, University of Southern California, Los Angeles, USA

Corresponding Author: Paul Adamson MD MPH, Division of Infectious Diseases, School of

Medicine, University of California, Los Angeles, 911 Broxton Ave, Suite 301, Los Angeles, CA

90024, Phone: (310) 794-5865; Email: PAdamson@mednet.ucla.edu

Summary: Our study found a high prevalence of *Neisseria gonorrhoeae* and *Chlamydia*

trachomatis, particularly pharyngeal and rectal infections, within an HIV PrEP program in Hanoi,

Vietnam. Our findings highlight the need for evidence-based screening guidelines in PrEP

programs in low-resource settings.

Abstract

Background: Data on *Neisseria gonorrhoeae* (NG) and *Chlamydia trachomatis* (CT) infections

within HIV pre-exposure prophylaxis (PrEP) programs in low- and middle-income countries

(LMICs) are limited. Our study reports the prevalence, anatomical distribution, and correlates of

NG and CT infections within an HIV PrEP program in Hanoi, Vietnam.

Methods: From January-December 2022, HIV PrEP program clients who were male at birth, ≥16

years old, reported ≥1 male sex partner in the prior 12 months, were enrolled. A questionnaire

collected sociodemographics, sexual behaviors, and clinical data. CT/NG testing was performed

on self-collected urine, rectal, and pharyngeal specimens. Multivariate logistic regression was

used to identify factors associated with CT and NG infections.

Results: There were 529 participants enrolled, the median age was 25.1 years. The overall

prevalence of CT or NG was 28.9% (153/529). The prevalence of NG was 14.3% and highest for

pharyngeal infections (11.7%), while for CT, the prevalence was 20.4% and highest for rectal

infections (14.0%). Symptoms in the prior week were reported by 45.8% (70/153) of those with

CT or NG infections. Condomless anal sex (aOR= 1.98; 95% CI: 1.27, 3.08) and sexualized drug

use in the prior 6 months (aOR= 1.68; 95% CI: 1.07, 2.65) were associated with CT/NG

infections.

Conclusions: Our study found a high prevalence of NG and CT infections, including pharyngeal

and rectal infections, within an HIV PrEP program in Hanoi, Vietnam. The findings underscore

the need for further research on CT/NG prevention and the development of evidence-based

guidelines for CT/NG screening in HIV PrEP programs in LMIC settings.

Introduction

Sexually transmitted infections (STIs) are a significant public health challenge globally.

Neisseria gonorrhoeae and *Chlamydia trachomatis* are the two most common bacterial STIs.

Men who have sex with men (MSM) are disproportionately affected by STIs. (2, 3) MSM on HIV

pre-exposure prophylaxis (PrEP) have high frequencies of bacterial STIs. [1–3] Thus, major HIV

PrEP guideline groups recommend routine screening for *C. trachomatis* and *N. gonorrhoeae*

among MSM on HIV PrEP, including testing at multiple anatomic sites of infection.

In low- and middle-income coutries (LMICs), HIV PrEP programs are expanding, yet data

on STIs among MSM in HIV PrEP programs are limited.[3] Understanding the epidemiology of

STIs in these setttings is important for planning and implementation of HIV PrEP programs.

Morever, in many LMIC settings, the availability of molecular *C. trachomatis* and *N.*

gonorrhoeae testing is limited and the costs associated testing remain significant barriers for

testing.[4,5] Identifying clinical and behavioral risk factors associated with those infections can

help optimize diagnosis and prevention of STIs, particularly in LMIC settings where resources

might be limited and there is ongoing rapid scale-up of HIV PrEP services.

Vietnam is a LMIC where HIV PrEP became available in 2018. Prior to that, baseline data

from the Hanoi-MSM (HIM) study found a high prevalence of *C. trachomatis* (22%) and *N.*

gonorrhoeae (12%) among HIV-uninfected MSM, who subsequently were enrolled into the pilot

HIV PrEP project.[6] The aims of this study are to fill gaps in research on STIs among MSM on

HIV PrEP in Vietnam, since the rollout of HIV PrEP. The study objectives were to determine the

prevalence of urethral, rectal, and pharyngeal *N. gonorrhoeae* and *C. trachomatis* infections

within an HIV PrEP program in Hanoi, Vietnam, as well as to determine behavioral and clinical

factors associated with *N. gonorrhoeae* and *C. trachomatis* infections.

Methods:

Study design and population

This was a observational, cross-sectional study from January 2022 to December 2022. The study was conducted within the HIV PrEP program at the Sexual Health and Promotion (SHP) Clinic at Hanoi Medical University. Study participants were eligible if they were male sex at birth, aged 16 years or older, reported having sex with men or transgender women in prior 12 months, and were enrolled in, or presenting for enrollment into, the HIV PrEP program; exclusion criteria were if they had *C. trachomatis* or *N. gonorrhoeae* testing done in the prior 3 months, unless they reported acute STI symptoms on the day of enrollment. Eligible participants were approached in the clinic and the study's objectives and procedures were explained by study research staff.

Data collection

Demographic, behavioral, and clinical characteristics were collected through the use of a tablet self-administered survey. Sexual behaviors were self-reported and included: number of sex partners in the prior month as the combined number of male, female, and transgender sex partners, anal sex position(s), condomless anal intercourse in the prior month, group sex (defined as having more than one sex partner at the same encounter), or having sex with partners met via mobile apps in the prior 6 months. Sexualized drug use was considered if the participant reported using a substance (heroine, ketamine, ecstasy, methamphetamine, cannabis, gamma hydroxybutyrate or gamma butyrolactone, poppers, or prostaglandin E medications) in the prior 6 months to enhance sexual pleasure. History of STIs in the prior three months was obtained by self-report.

Participants were asked about any genitourinary, rectal, or pharyngeal symptoms in the prior

one week. Genitourinary symptoms were classified as any of the following: pain with urination,

discharge, bleeding, pruritis, testicular pain, lymphadenopathy, or ulcers. Rectal symptoms were

classified as any of the following: tenesmus, dyschezia, pruritis, bleeding, discharge, ulcers, or

diarrhea. Pharyngeal symptoms included pain or itching in the throat.

Sample collection, testing, and treatment

Study participants received instructions on providing specimens for testing by study staff and

with visual aids. Clinician-collected samples were obtained if participants preferred or were

unable to perform self-collection. Study specimens were collected using either the Alinity m

multi-Collect Specimen Collection Kits (Abbott Molecular, USA) or cobas PCR Urine or Swab

Sample Kits (Roche Diagnostics, Branchburg, NJ, USA). Specimens were stored in the clinic

and transported daily to the laboraty. Testing was performed using either the Alinity m STI

Assay (Abbott Molecular, USA) or the cobas 4800 CT/NG assay (Roche Diagnostics,

Branchburg, NJ, USA) according to manufacturer's instructions. Test results for *C. trachomatis*

and *N. gonorrhoeae* were either positive, negative, or inconclusive, in the event of failed internal

controls or the presence of inhibitors. Participants were informed of their test results. For those

found to have infections, free treatment was offered. For *C. trachomatis* infections, doxycycline

100mg by mouth twice daily for 7 days was provided. For *N. gonorrhoeae* infections, ceftriaxone

500mg intramuscular injection once was the preferred treatment offered; cefixime 800mg by

mouth once was provided as an alternative treatment.

Data analysis

The primary outcome was the prevalence of *C. trachomatis* or *N. gonorrhoeae* infections at

different anatomic sites. Descriptive statistics were performed for demographic, behavioral and

clinical data; percentages for categorical variables, or median and interquartile range for

continuous variables, were reported. Categorical variables between those with and without C.

trachomatis or *N. gonorrhoeae* infections were compared using Pearson's Chi-squared test and

Fisher's exact test, and nonparametric continuous variables were compared using Wilcoxon

rank-sum tests.

Logistic regression modeling was used to evaluate factors associated with *N. gonorrhoeae* and

C. trachomatis infections separately as well as the outcome of having either infection. Variables

with p-values < 0.2 in univariate comparisons were included in the multivariable logistic

regression models; except for age, which was included in the final models. The number of sex

partners in the prior month was dichotomized to above or below the median (0-1 and ≥2) to

improve interpretation and model fit. Variables with missing data were excluded from the logistic

regression models. P-values ≤ 0.05 was defined as statistical significance. All data analyses

were performed using STATA version 18 (StataCorp LLC, College Station, TX, USA).

Ethics

The study was approved by the Institutional Review Boards (IRB) of Hanoi Medical University

(HMUIRB580), the University of California, Los Angeles, and the University of Southern

California. All study participants provided written informed consent.

132
Results

Participant Characteristics

From January to December 2022, there were 529 participants enrolled into the study

among 775 approached and 538 screened. The primary reasons for refusal of those who were

approached, but who did not undergo screening, was "too busy/not enough time" or "will

consider next time." Among the study participants, 81.6% (432/529) identified as a man, 1.7%

(9/529) identified as transgender woman, and 16.6% (88/529) identified as gender non-

conforming. The median age among all participants was 25.1 years (IQR: 21.7-29.5). There were

29.1% (154/529) currently undergoing university or post-secondary training, while 61.1%

(323/529) had completed university or post-secondary training.

In the prior six months, 95.8% (507/529) reported only male sex partners; the median

number of sex partners in the prior month was 1 (IQR: 1-2). Anal sex in the prior 6 months was

reported by 88.4% (396/529), of which 51.2% reported condomless anal sex. In the prior 6

months, group sex was reported by 12.3% (65/529) and sexualized substance use was reported

by 45.4% (240/529). A history of a STI in the prior 3 months was reported by 5.5% (29/529),

with 69.0% (20/29) of those being syphilis. Additional sexual bevavior data are shown in Table

1.

C. trachomatis and N. gonorrhoeae infections

In total, 28.9 % (153/529) participants had *C. trachomatis* or *N. gonorrhoeae* infections

at any anatomical site. The prevalence of *C. trachomatis* was 20.4% (108/529) and 14.3%

(76/529) for *N. gonorrhoeae*; 5.9% (31/529) of participants had *C. trachomatis* and *N.*

gonorrhoeae co-infections (Table 1). Among the 153 participants with either *C. trachomatis* or

N. gonorrhoeae infections, the median age was 25.1 years (IQR: 21.5 – 28.3). The median

number of sex partners in the prior month was 2 (IQR: 1-3), 14.2% (22/155) reported group sex

in the prior month, and 60.1% (94/155) reported sexualized substance use. Of the 153 with C.

trachomatis or *N. gonorrhoeae* infections, 45.8% (70/153) reported any symptoms in the prior

week; 33 (21.6%) reported pharyngeal symptoms, 29 (19.0%) reported rectal symptoms, and 26

(17.0%) reported urethral symptoms. More participants with *C. trachomatis* or *N. gonorrhoeae*

infections reported having symptoms on the day of enrollment, compared to those without

infections (9.2% vs. 3.0%, p=0.001).

C. trachomatis and *N. gonorrhoeae* infections by anatomic site are shown in Table 2. The

prevalence of *C. trachomatis* was highest in the rectum (14.0%; 74/529) and 3.4% (18/529) had

infections in multiple anatomic sites. For *N. gonorrhoeae*, pharyngeal site infections were most

prevalent (11.7%; 62/529) and 6.0% (32/529) had infections at multiple anatomic sites. Among

participants with either *C. trachomatis* or *N. gonorrhoeae*, the prevalence of rectal infections

was 17.8% (94/529) and 9.3% (49/529) had infections at multiple anatomic sites.

Among those reporting rectal symptoms in the prior week, the prevalence of having a

rectal infection with *C. trachomatis* or *N. gonorrhoeae* was 27.2% (22/81), followed by 25.0%

(18/72) for urethral symptoms and urethral infections, and 18.3% (23/126) for pharyngeal

symptoms and pharyngeal infections. Among those who did not report symptoms in the prior

week, the prevalence of urethral *C.trachomatis* was 3.1% (14/457) and 0.9% (4/457) for *N.*

gonorrhoeae. For both infections, the prevalence of both rectal and urethral infections were

higher among those reporting symptoms in the prior week; there were no differences in the

prevalence of pharyngeal infections by symptom status for either infection. The prevalence of

C. trachomatis or *N. gonorrhoeae* infections by anatomic site, stratified by reported symptoms

in the prior week are shown in Figure 1.

Factors associated with C*. trachomatis and* N. *gonorrhoeae infections*

In the multivariate logistic regression models, having either *C. trachomatis* or *N.*

gonorrhoeae infection was associated with condomless anal intercourse (aOR: 1.98; 95% CI:

1.27-3.08) and sexualized drug use (aOR: 1.68; 95% CI: 0.73, 1.95). Rectal and urethral

symptoms in the prior week were associated with infections in univariate analyses, but not in

the multivariate model (aOR: 1.19; 95% CI: 1.07, 2.65). When examining each infection

separately, meeting sex partners in mobile apps was an independent risk factor for *N.*

gonorrhoeae infections (aOR: 2.33; 95% CI: 1.19, 4.54), but not *C. trachomatis* (aOR: 0.83; 95%

CI: 0.48, 1.43). Both condomless anal sex (aOR: 2.09; 95% CI: 1.27-3.44) and sexualized drug

use (aOR: 2.18; 95% CI: 1.30-3.64) were associated with *C. trachomatis*, but not *N.*

gonorrhoeae. (Table 3).

Discussion

In this study including 529 primarily MSM in an HIV PrEP program in Hanoi, Vietnam,

testing for *C. trachomatis* and *N. gonorrhoeae* at multiple anatomic sites identified a high

prevalence (29%) of infections. Rectal infections were most common for *C. trachomatis* (14%),

while pharyngeal infections were most common for *N. gonorrhoeae* (11%), demonstrating the

utility of testing at multiple anatomic sites for the detection of infections outside of the genital

tract. While the prevalence of urethral and rectal *C. trachomatis* and *N. gonorrhoeae* infections

were higher among those who reported symptoms at those sites in the prior week, a

substantial proportion of overall infections (44%) were detected among those without

symptoms, underscoring the limitations of relying on syndromic management as an infection

control strategy in this setting. These findings highlight the opportunity to enhance STI

diagnosis and treatment within HIV PrEP programs in LMICs.

Our study provides important data on routine testing for *C. trachomatis* and *N.*

gonorrhoeae within a large HIV PrEP program in a LMIC setting, where prevalence data are

limited. A recent meta-analysis of STIs among people in HIV PrEP programs estimated a pooled

prevalence of *C. trachomatis* or *N. gonorrhoeae* was 24%, suggesting a slightly higher burden

among participants in our study population.[3] However, that study highlighted the scarcity of

data from LMICs, which makes comparisons difficult, but the study did not find a significant

difference in prevalence of chlamydia or gonorrhea by country income level. There is a paucity

of published prevalence estimates of *C. trachomatis* and *N. gonorrhoeae* among MSM on HIV

PrEP from other nearby countries in Asia, which could provide further context. One study done

among adolescent MSM and transgender women in Thailand found a *C. trachomatis* prevalence

of 15% and *N. gonorrhoeae* prevalence of 4.5%, estimates that are much lower than our study,

although pharyngeal testing was not performed in that study which would likely account for

some of that difference.[7] The prevalence of infections observed in our study align with

existing evidence from other, largely high-income, settings indicating a high burden of STIs

among MSM on HIV PrEP.[4,8]

Among participants in our study, we observed those reporting condomless anal sex

were about twice as likely to have *C. trachomatis* or *N. gonorrhoeae* infection and those having

sexualized drug use in the prior 6 months were 70% more likely to have *C. trachomatis* or *N.*

gonorrhoeae infection. Identifying factors associated with infections could help to focus testing

among those at highest risk for infections. For example, of the total 153 *C. trachomatis* or *N.*

gonorrhoeae infections in our study, testing those reporting either condomless anal sex or

sexualized drug use in the prior 6 months would have detected 75% of those infections.

Reporting rectal or urethral symptoms in the prior week was not associated with infections in

the multivariate analysis, highlighting the difficulty of using symptoms to identify infections,

given a high proportion of these types of infections are asymptomatic.[9] Despite this, the

prevalence of rectal and urethral *C. trachomatis* and *N. gonorrhoeae* was significantly higher

among those who reported symptoms in the prior week, suggesting symptoms are still

correlated with a higher frequency of infections at those sites. Our findings suggest that testing

for urethral infections among asymptomatic individuals is of limited value, as the prevalence in

this group was very low, 0.9% for *N. gonorrhoeae* and 3.1% for *C. trachomatis*. Meeting sex

partners on mobile apps is increasingly common globally and this behavior was associated with

twice the odds of *N. gonorrhoeae*, but not *C. trachomatis*, infections in our study, a finding that

was also observed in our previous work in a similar study population.[6] Possible explanations

for this observation might be there is a higher prevalence of *N. gonorrhoeae* among certain

sexual networks of online mobile app users or there is more oral sex within that network, since

the majority of these infections were pharyngeal. These findings indicate that focused testing

for *N. gonorrhoeae* among those meeting sex partners on apps could be a strategy to improve

gonorrhea control.

Our study further highlights the high prevalence of *C. trachomatis* and *N. gonorrhoeae*

infections among people in HIV PrEP programs and the need to improve evidence-based

guidelines for testing, particularly in LMICs. Given the high prevalence and incidence of STIs in

HIV PrEP programs, many guidelines recommend quarterly screening for *C. trachomatis* and *N.*

gonorrhoeae among MSM on HIV PrEP.[10,11] Yet while some modeling studies support

quarterly screening to reduce the population prevalence, the clinical and public health evidence

supporting the frequency of testing is lacking and the populations coverage needed to achieve a

population-benefit of decreased prevalence is much higher than observed in practice.[12–16]

The recent Gonoscreen Study, a randomized-controlled trial evaluating quarterly screening for

C. trachomatis and *N. gonorrhoeae* compared to no screening among MSM in Belgium found

quarterly screening was associated with fewer *C. trachomatis* infections and complications, but

observed no difference among *N. gonorrhoeae* infections.[17] Beyond the frequency of

screening, there is also a larger debate about the utility of screening for asymptomatic STIs in

populations of MSM, particularly those on HIV PrEP.[18,19] While *C. trachomatis* and *N.*

gonorrhoeae infections increase the risk for HIV transmission and provide a strong rationale for

using these as an entry point for HIV PrEP,[20] the increased risk for HIV acquisition caused by

bacterial STIs is largely mitigated among those on HIV PrEP.[21,22] Moreover, antimicrobial

resistance is one potential consequence of increased screening and treatment and is a

significant concern, particularly for *N. gonorrhoeae* and *Mycoplasma genitalium*; increased

antibiotic consumption was observed among those randomized to quarterly screening in the

Gonoscreen study.[17] Antimicrobial resistance in *N. gonorhoeae* is particularly concerning in

Vietnam, where the prevalence of ceftriaxone resistance, the first-line agent for treatment, is

more than 20% in some settings.[23]

Our study findings must be interpreted in light of the following limitations. First, the

study population might have been subject to slection bias. Efforts were made to expand

inclusivity by simplifying the study eligibility crtieria to reflect that of the HIV PrEP program.

Still, it was not possible to recruit all PrEP program clients and 70% of those who were recruited

agreed to be screened; thus, it is possible those who chose not to participate were different

than those who participated. Second, our study outcome was the prevalence of *C. trachomatis*

and *N. gonorrhoeae* infections at each visit, so while participants were allowed to participate

more than once, few did, and we were unable to estimate the incidence of *C. trachomatis* or *N.*

gonorrhoeae infections. Third, our study population of primarily MSM were largely well-

educated, employed, and living in Hanoi, the second most populated city in Vietnam, and might

not be generalizable to other populations or settings. Our study took place within the largest

HIV PrEP program in Hanoi, which is a strength of our study.

Our findings provide further evidence of the high prevalence of *C. trachomatis* and *N.*

gonorrhoeae infections among people on HIV PrEP, including the high proportions of rectal,

pharyngeal, and asymptomatic infections. Despite the high prevalence of infections, the high

cost of molecular STI testing is a major barrier to implementing routine testing in HIV PrEP

programs in LMICs.[4] Testing for urethral *C. trachomatis* and *N. gonorrhoeae* infections among

those without urethral symptoms is likely to be of limited value, based on our findings. Further

research is needed to establish evidence-based recommendations on the role and frequency of

testing for asymptomatic infections, including the cost-effectiveness of different approaches.

Biomedical interventions like doxycycline prophylaxis and vaccinations are promising tools for

STI prevention and further research is needed on the effectiveness and implementation in LMIC

settings, where key factors such as antimicrobial resistance and differences in user preferences

will influence their impact.

Acknowledgements: The research was supported by a grant of no charge materials from Roche

Molecular Systems and Abbott Molecular. PCA, HTB, GML, and JDK designed the study. PCA,

GML, JDK obtained funding. HTB and LQP performed data management and data analyses. PCA

wrote the first version of the manuscript. All authors reviewed, provided critical review, and

approved the manuscript.

Conflicts of Interest and Funding: This work was supported by the US National Institute of

Allergy and Infectious Diseases (R21 AI157817 to G. M. L and J. D. K.) and the Fogarty

International Center (K01TW012170 to P. C. A). The funders had no role in the data collection,

analysis, manuscript preparation, or decision to publish. JDK reports consulting fees from

Abbott in the prior 12 months. All other authors report no potential conflicts of interest.

Table 1. Baseline demographic, behavioral, and clinical characteristics of 529 participants enrolled from an

HIV PrEP program clinic in Hanoi, Vietnam.

	Overall (N = 529)	Positive for *C. trachomatis* (N=108)		Positive for *N. gonorrhoeae* (N=76)		Positive for *C. trachomatis* or *N. gonorrhoeae* (N=153)	
	n (%)	n (%)	p-value	n (%)	p-value	n (%)	p-value
Median Age (IQR)	25.1 (21.7-29.5)	25.3 (21.1 – 29.7)	0.869[†]	24.8 (22.1 – 27.2)	0.281[†]	25.1 (21.5 – 28.3)	0.336[†]
Age group			0.644		0.243		0.834
16- < 25 years old	261 (49.3)	51 (47.2)		38 (50.0)		75 (49.0)	
25- < 35 years old	221 (41.8)	45 (41.7)		35 (46.1)		66 (43.1)	
35 - 63 years old	47 (8.9)	12 (11.1)		3 (4.0)		12 (7.8)	
Gender self-identification			0.126[‡]		0.316[‡]		0.232[‡]
Man	432 (81.7)	85 (78.7)		58 (76.3)		121 (79.1)	
Transgender woman	9 (1.7)	0		1 (1.3)		1 (0.7)	
Other/Unsure*	88 (16.6)	23 (21.3)		17 (22.4)		31 (20.3)	
Education			0.687		0.942		0.797
High school or below	52 (9.8)	13 (12.0)		8 (10.5)		17 (11.1)	
In post-secondary training	154 (29.1)	31 (28.7)		21 (27.6)		45 (29.4)	
Completed post-secondary training or higher	323 (61.1)	64 (59.3)		47 (61.8)		91 (59.5)	
Monthly income in million VND (median, IQR)	9.0 (5 – 15)	8.5 (5-14)	0.813[†]	8 (4.5 – 14.5)	0.566[†]	8 (5 – 14.5)	0.493[†]
Employed	419 (79.2)	84 (77.8)	0.682	63 (82.9)	0.392	123 (80.4)	0.668
Gender of sex partners prior 6 months							
Men only	507 (95.8)	105 (97.2)	0.420	75 (98.7)	0.180	150 (98.0)	0.106
Women only	46 (8.7)	11 (10.2)	0.538	3 (4.0)	0.112	11 (7.2)	0.433
Men and women	1 (0.2)	0	1.000[‡]	0	1.000[‡]	0	1.000[‡]
No sex partners	9 (1.7)	1 (0.9)	0.694[‡]	0	0.371[‡]	1 (0.7)	0.458[‡]
Number of sex partners in prior 1 month[#]	1 (1 – 2)	2 (1 – 3)	<0.001[†]	2 (1 – 2)	0.150[†]	2 (1 – 3)	<0.001[†]
Anal sex with male partners in prior 1 month[#]	396 (88.4)	95 (94.1)	0.043	64 (92.8)	0.219	131 (92.9)	0.043
Anal sex position with male partner[#]			0.633		0.109		0.385
Insertive anal sex always	137 (34.6)	29 (30.5)		22 (34.4)		44 (33.6)	
Receptive anal sex always	149 (37.6)	38 (40.0)		18 (29.1)		45 (34.4)	
Both insertive and receptive	110 (27.8)	28 (29.5)		24 (37.5)		42 (32.1)	
Condomless anal sex in prior 1 month[#]	203 (51.3)	63 (66.3)	0.001	39 (60.9)	0.091	83 (63.4)	0.001
Vaginal sex with female partners in prior month[#]	38 (7.2)	8 (7.4)	0.064	2 (2.6)	0.814	8 (5.2)	0.064
Deep kissing in prior week[#]	260 (86.1)	61 (84.7)	0.700	41 (89.1)	0.518	85 (86.7)	0.823
Any group sex, prior 6 months	65 (12.3)	13 (12.0)	0.929	14 (18.4)	0.078	22 (14.4)	0.350
Any sexualized drug use, prior 6 months	240 (45.4)	68 (63.0)	<0.001	40 (52.6)	0.169	88 (57.5)	<0.001

	Overall (N = 529)	Positive for *C. trachomatis* (N=108)		Positive for *N. gonorrhoeae* (N=76)		Positive for *C. trachomatis* or *N. gonorrhoeae* (N=153)	
	n (%)	n (%)	p-value	n (%)	p-value	n (%)	p-value
Met sex partners via mobile apps	297 (56.1)	66 (61.1)	0.244	55 (72.4)	0.002	97 (63.4)	0.032
Pharyngeal symptoms in prior week	126 (23.8)	21 (19.4)	0.232	20 (26.3)	0.581	33 (21.6)	0.438
Rectal symptoms in prior week	87 (16.5)	25 (23.2)	0.035	15 (19.7)	0.403	29 (19.0)	0.321
Genitourinary symptoms in prior week	72 (13.6)	17 (15.7)	0.469	15 (19.7)	0.092	26 (17.0)	0.148
Any genitourinary symptoms at the day of enrollment	14 (2.7)	7 (6.5)	0.012‡	8 (10.5)	<0.001‡	12 (7.8)	<0.001‡
Any symptoms in prior week	217 (41.0)	49 (45.4)	0.303	41 (54.0)	0.013	70 (45.8)	0.158
Self-reported STI diagnosis in the prior 3 months	29 (5.5)	6 (5.6)	0.970	6 (7.9)	0.318	9 (5.9)	0.796
Used antiseptic mouthwash in the previous month	286 (54.1)	65 (60.2)	0.152	38 (50.0)	0.442	84 (54.9)	0.805

*Participants who were gender nonconforming or gender incongruent, or unsure of their gender identity.
†Wilcoxon rank-sum (Mann-Whitney) test was performed
‡Fisher's exact test was performed
§Variables that contains missing values because "Don't know" or "Don't remember" choice were treated as missing.
#Sample size was lower than 529 due to skip pattern (e.g., sample size for "Condomless anal sex" was among people
who reported engaging in anal sex)

Table 2. Prevalence of *Chlamydia trachomatis* and *Neisseria gonorrhoeae* infections by anatomic site

among 529 participants enrolled from an HIV PrEP program clinic in Hanoi, Vietnam.

Anatomic Site	C. trachomatis (n/N*, %)	N. gonorrhoeae (n/N*, %)	*C. trachomatis* or *N. gonorrhoeae* (n/N*, %)	*C. trachomatis* and *N. gonorrhoeae* (n/N*, %)
Total Infections (Any Site)	108 (20.4)	76 (14.4)	153 (28.9)	31 (5.9)
Single Site Infections				
Pharyngeal only	19/528 (3.6)	32/528 (6.1)	41/528 (7.7)	3/528 (0.6)
Rectal only	57/493 (11.6)	10/494 (2.0)	52/493 (10.6)	4/493 (0.8)
Urethral only	14/529 (2.7)	2/529 (0.4)	11/529 (2.1)	0/529 (0)
Multisite Infections				
Pharyngeal and rectal	9/493 (1.8)	18/494 (3.6)	26/494 (5.3)	13/493 (2.6)
Pharyngeal and urethral	1/528 (0.2)	5/528 (1.0)	7/528 (1.3)	3/528 (0.6)
Urethral and rectal	7/493 (1.4)	2/494 (0.4)	7/493 (1.4)	1/493 (0.2)
Pharyngeal, rectal, and urethral	1/493 (0.2)	7/494 (1.4)	9/494 (1.8)	7/493 (1.4)

313   **Figure 1.** Prevalence of *C. trachomatis* (Panel A) and *N. gonorrhoeae* (Panel B) infections at pharyngeal,

314   rectal, and urethral sites, stratified by reported symptoms at that site in the prior week.

316
317
318   ***p-value ≤0.05; **p-value ≤0.01**

319 **Table 3.** Multivariable logistic regression with factors associated with *C. trachomatis* only, *N.*

320 *gonorrhoeae* only, and *C. trachomatis* or *N. gonorrhoeae* infections.

Characteristics Age group	*C. trachomatis*		*N. gonorrhoeae*		*C. trachomatis* or *N. gonorrhoeae*	
	aOR (95% CI)	p-value	aOR (95% CI)	p-value	aOR (95% CI)	p-value
16 to < 25 years old	1.00		1.00		1.00	
25 to < 35 years old	0.75 (0.44, 1.26)	0.278	1.03 (0.58, 1.85)	0.908	0.81 (0.51, 1.29)	0.382
≥ 35 years old	1.27 (0.58, 2.79)	0.553	0.39 (0.11, 1.38)	0.144	0.76 (0.36, 1.64)	0.490
Two or more sex partners in prior 1 month	1.68 (0.98, 2.88)	0.061	1.16 (0.63, 2.15)	0.630	1.53 (0.95, 2.47)	0.080
Condomless anal sex in prior 1 month	**2.09 (1.27, 3.44)**	**0.004**	1.52 (0.86, 2.68)	0.152	**1.98 (1.27, 3.08)**	**0.003**
Sexualized drug use in prior 6 months	**2.18 (1.30, 3.64)**	**0.003**	1.07 (0.60, 1.91)	0.819	**1.68 (1.07, 2.65)**	**0.025**
Met sex partners via mobile apps	0.83 (0.48, 1.43)	0.505	**2.33 (1.19, 4.54)**	**0.013**	1.21 (0.74, 1.96)	0.449
Any anal or urethral symptoms in prior week	1.44 (0.84, 2.45)	0.182	1.30 (0.72, 2.36)	0.385	1.19 (0.73, 1.95)	0.487
Used an antiseptic mouthwash in the prior 1 month	1.14 (0.70, 1.86)	0.592	0.72 (0.41, 1.25)	0.240	0.85 (0.55, 1.32)	0.462
Model information						
Sample size included	396		396		396	
Hosmer-Lemeshow p-value	0.724		0.434		0.3621	
McFadden's R2	0.069		0.053		0.053	
Log likelihood	-203.127		-165.818		-238.097	

References

323
1. Kojima N, Davey DJ, Klausner JD. Pre-exposure prophylaxis for HIV infection and new
sexually transmitted infections among men who have sex with men. AIDS **2016**; 30:2251–
2252.

2. Traeger MW, Schroeder SE, Wright EJ, et al. Effects of Pre-exposure Prophylaxis for the
Prevention of Human Immunodeficiency Virus Infection on Sexual Risk Behavior in Men
Who Have Sex With Men: A Systematic Review and Meta-analysis. Clin Infect Dis **2018**;
67:676–686.

3. Ong JJ, Baggaley RC, Wi TE, et al. Global Epidemiologic Characteristics of Sexually
Transmitted Infections Among Individuals Using Preexposure Prophylaxis for the
Prevention of HIV Infection: A Systematic Review and Meta-analysis. JAMA Netw Open
2019; 2:e1917134.

4. Ong JJ, Fu H, Baggaley RC, et al. Missed opportunities for sexually transmitted infections
testing for HIV pre-exposure prophylaxis users: a systematic review. J Int AIDS Soc **2021**;
24:e25673.

5. Wi TE, Ndowa FJ, Ferreyra C, et al. Diagnosing sexually transmitted infections in resource-
constrained settings: challenges and ways forward. J Int AIDS Soc **2019**; 22 Suppl
6:e25343.

6. Adamson PC, Bhatia R, Tran KDC, et al. Prevalence, Anatomic Distribution, and Correlates
of Chlamydia trachomatis and Neisseria gonorrhoeae Infections Among a Cohort of Men
Who Have Sex With Men in Hanoi, Vietnam. Sex Transm Dis **2022**; 49:504–510.

7. Songtaweesin WN, Pornpaisalsakul K, Kawichai S, et al. Sexually transmitted infections
incidence in young Thai men who have sex with men and transgender women using HIV
pre-exposure prophylaxis. Int J STD AIDS **2022**; 33:447–455.

8. World Health Organization. WHO implementation tool for pre-exposure prophylaxis (PrEP)
of HIV infection. Module 13. Integrating STI services. Geneva, Switzerland.: 2022.
Available at: https://www.who.int/publications/i/item/9789240057425. Accessed 11 June
2024.

9. Kent CK, Chaw JK, Wong W, et al. Prevalence of Rectal, Urethral, and Pharyngeal
Chlamydia and Gonorrhea Detected in 2 Clinical Settings among Men Who Have Sex with
Men: San Francisco, California, 2003. Clinical Infectious Diseases **2005**; 41:67–74.
Available at: https://doi.org/10.1086/430704.

10. Centers for Disease Control and Prevention. Preexposure prophylaxis for the prevention of
HIV infection in the United States—2021 Update: a clinical practice guideline. 2021.
Available at: https://www.cdc.gov/hiv/pdf/risk/prep/cdc-hiv-prep-guidelines-2021.pdf.
Accessed 12 June 2024.

11. British HIV Association, British Association of Sexual Health and HIV. BHIVA/BASHH guidelines on the use of HIV pre-exposure prophylaxis (PrEP), 2018. 2018. Available at: https://www.bhiva.org/PrEP-guidelines. Accessed 12 June 2024.

12. Jenness SM, Weiss KM, Goodreau SM, et al. Incidence of Gonorrhea and Chlamydia Following Human Immunodeficiency Virus Preexposure Prophylaxis Among Men Who Have Sex With Men: A Modeling Study. Clinical Infectious Diseases **2017**; 65:712–718. Available at: https://doi.org/10.1093/cid/cix439.

13. Reitsema M, Hoek AJ van, van der Loeff MS, et al. Preexposure prophylaxis for men who have sex with men in the Netherlands: impact on HIV and Neisseria gonorrhoeae transmission and cost-effectiveness. AIDS **2020**; 34:621. Available at: https://journals.lww.com/aidsonline/fulltext/2020/03150/preexposure_prophylaxis_for_men _who_have_sex_with.14.aspx.

14. Hocking JS, Temple-Smith M, Guy R, et al. Population effectiveness of opportunistic chlamydia testing in primary care in Australia: a cluster-randomised controlled trial. The Lancet **2018**; 392:1413–1422. Available at: https://www.sciencedirect.com/science/article/pii/S0140673618318166.

15. Tsoumanis A, Hens N, Kenyon CR. Is Screening for Chlamydia and Gonorrhea in Men Who Have Sex With Men Associated With Reduction of the Prevalence of these Infections? A Systematic Review of Observational Studies. Sexually Transmitted Diseases **2018**; 45:615. Available at: https://journals.lww.com/stdjournal/fulltext/2018/09000/is_screening_for_chlamydia_and_ gonorrhea_in_men.7.aspx.

16. Tao G, Patel CG, He L, Workowski K. STI/HIV testing, STIs, and HIV PrEP use among men who have sex with men (MSM) and men who have sex with men and women (MSMW) in United States, 2019-2022. Clin Infect Dis **2024**; :ciae314.

17. Vanbaelen T, Tsoumanis A, Florence E, et al. Effect of screening for *Neisseria gonorrhoeae* and *Chlamydia trachomatis* on incidence of these infections in men who have sex with men and transgender women taking HIV pre-exposure prophylaxis (the Gonoscreen study): results from a randomised, multicentre, controlled trial. The Lancet HIV **2024**; 11:e233–e244. Available at: https://www.sciencedirect.com/science/article/pii/S2352301823002990.

18. Williams E, Williamson DA, Hocking JS. Frequent screening for asymptomatic chlamydia and gonorrhoea infections in men who have sex with men: time to re-evaluate? The Lancet Infectious Diseases **2023**; 23:e558–e566. Available at: https://www.thelancet.com/journals/laninf/article/PIIS1473-3099(23)00356-0/abstract.

19. Ridpath AD, Chesson H, Marcus JL, et al. Screening Peter to Save Paul: The Population-Level Effects of Screening Men Who Have Sex With Men for Gonorrhea and Chlamydia. Sexually Transmitted Diseases **2018**; 45:623. Available at: https://journals.lww.com/stdjournal/citation/2018/09000/screening_peter_to_save_paul__th e_population_level.8.aspx.

20. Kasaie P, Schumacher CM, Jennings JM, et al. Gonorrhoea and chlamydia diagnosis as an
entry point for HIV pre-exposure prophylaxis: a modelling study. BMJ Open **2019**;
9:e023453. Available at: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6429744/.

21. Volk JE, Marcus JL, Phengrasamy T, et al. No New HIV Infections With Increasing Use of
HIV Preexposure Prophylaxis in a Clinical Practice Setting. Clin Infect Dis **2015**; 61:1601–
1603.

22. Hoornenborg E, Coyer L, Achterbergh RCA, et al. Sexual behaviour and incidence of HIV
and sexually transmitted infections among men who have sex with men using daily and
event-driven pre-exposure prophylaxis in AMPrEP: 2 year results from a demonstration
study. The Lancet HIV **2019**; 6:e447–e455. Available at:
https://www.sciencedirect.com/science/article/pii/S2352301819301365.

23. Adamson PC, Hieu VN, Nhung PH, Whiley DM, Chau TM. Ceftriaxone resistance in
Neisseria gonorrhoeae associated with the penA-60.001 allele in Hanoi, Viet Nam. The
Lancet Infectious Diseases **2024**; 0. Available at:
https://www.thelancet.com/journals/laninf/article/PIIS1473-3099(24)00230-5/fulltext.