Original article

Efficacy and safety of neoadjuvant stereotactic body radiation therapy plus dalpiciclib and exemestane for hormone receptor-positive, HER2-negative breast cancer: A prospective pilot study

Authors: Yu Zhang[1,2,3,4†], Shuo Cao[1†], Nan Niu[1,2,3†], Huilian Shan[1†], Jinqi Xue[1,2,3†], Guanglei Chen[1,2,3], Yongqing Xu[1], Jianqiao Yin[1], Chao Liu[1,2,3], Lisha Sun[1,2,3], Xiaofan Jiang[1,2,3], Meiyue Tang[1], Qianshi Xu[1,2], Mingxuan Jia[1], Xu Zhang[1], Zhenyong Zhang[1], Qingfu Zhang[5], Jianfei Wang[6], Ailin Li[1], Yongliang Yang[7*], Caigang Liu[1,2,3*]

Author Affiliations:

[1] Department of Oncology, Shengjing Hospital of China Medical University, Shenyang, China;

[2] Cancer Stem Cell and Translational Medicine Laboratory, Shengjing Hospital of China Medical University, Shenyang, China;

[3] Innovative Cancer Drug Research and Development Engineering Center of Liaoning Province, Shenyang, China.;

[4] Department of Gastrointestinal Surgery, Yantai Affiliated Hospital of Binzhou Medical University, Yantai, China;

[5] Department of Pathology, the First Affiliated Hospital and College of Basic Medical Sciences of China Medical University, Shenyang, China;

[6] Jiangsu Hengrui Pharmaceuticals, Shanghai, China;

[7] School of Bioengineering, Dalian University of Technology, Dalian, China.

Authors' contributions

Y. Zhang, Data curation, formal analysis, methodology. S. Cao, Data curation, formal analysis, methodology, writing–original draft. N. Niu, Validation, data curation, formal analysis, methodology, writing–review and editing. H. Shan, Data curation, methodology. J. Xue, Data curation, methodology. G. Chen, Data curation, methodology. Y. Xu, Methodology. J. Yin, Methodology. C. Liu, Formal analysis, methodology. L. Sun, Methodology. X. Jiang, Writing-original draft. M. Tang, Data curation. Q. Xu, Formal analysis. M. Jia, Methodology. X. Zhang, Methodology. Z. Zhang, Methodology. Q. Zhang, Formal analysis. J. Wang, Methodology. A. Li, Conceptualization, methodology. Y. Yang, Methodology, writing–review and editing. Caigang Liu, Conceptualization, supervision, project administration, writing–review and editing.

Yu Zhang, Shuo Cao, Nan Niu, Huilian Shan and Jinqi Xue contributed equally to this work.

Corresponding author:

Caigang Liu, MD, PhD

Department of Oncology, Shengjing Hospital of China Medical University, Shenyang 110004, China

Tel.: 86-18940254967

E-mail: liucg@sj-hospital.org

Abstract

Background: Both neoadjuvant chemotherapy and endocrine therapy only result in trivial pathological complete response rates and moderate objective response rates (ORR) in hormone receptor (HR)-positive, human epidermal growth factor receptor-2 (HER2)-negative breast cancer, more promising alternatives are urgently needed. With proven synergistic effect of cyclin dependent kinase 4/6 (CDK4/6) inhibitor and radiotherapy in preclinical studies, this pilot study aimed to explore the efficacy and safety of neoadjuvant stereotactic body radiation therapy (SBRT) followed by dalpiciclib and exemestane in HR-positive, HER2-negative breast cancer.

Methods: This was a single-arm, non-controlled prospective pilot study. Treatment-naïve patients with unilateral HR-positive, HER2-negative breast cancer received neoadjuvant radiotherapy (24Gy/3F) followed by dalpiciclib and exemestane for 6 cycles. The primary endpoint was the proportion of patients with residual cancer burden (RCB) score of 0-I. Key secondary endpoints included ORR, breast-conservation rate, biomarker analysis, and safety.

Results: All 12 enrolled patients completed the study treatment and surgery. Two (16.7%) of them achieved the RCB 0-I with the ORR of 91.7% (11/12). Analyses of tumor specimens showed significant increase of infiltrating T cells rather than alteration of PD-L1 positive immune cells. The most common grade 3 adverse events (AEs) were neutropenia (66.7%) and leukopenia (25.0%), but no grade 4-5 AE or death occurred.

Conclusions: Our results suggested neoadjuvant SBRT followed by dalpiciclib and

exemestane are effective and tolerable, and provides novel insights for the neoadjuvant

treatment of HR+/HER2- breast cancer, which may be considered as a feasible option

for patients with HR-positive, HER2-negative breast cancer.

Funding: None

Clinical trial number: NCT05132790

Impact statement: Neoadjuvant SBRT with dalpiciclib and exemestane resulted

in a 16.7% RCB 0-I rate and a 91.7% ORR, with acceptable toxicities in HR-

positive, HER2-negative breast cancer patients.

Introduction

Hormone receptor (HR)-positive, human epidermal growth factor receptor 2 (HER2)-negative breast cancer is the most prevalent subtype of breast cancer, which accounts for approximately 70% of all breast cancer (Parker et al., 2009). Neoadjuvant chemotherapy (NCT) is recommended as the first-choice for patients with HR-positive, HER2-negative breast cancer, who are candidates for preoperative therapy, according to National Comprehensive Cancer Network (NCCN) guidelines(Gradishar et al., 2022). Neoadjuvant endocrine therapy (NET) is an acceptable, less toxic alternative of NCT. However, less than 10% of patients with HR-positive, HER2-negative breast cancer can achieve pathological complete response (pCR), and only 60% of patients respond to NCT or NET (Sella et al., 2021). Hence, more promising alternatives are urgently needed.

Cyclin dependent kinase 4/6 (CDK4/6) inhibitors are breakthroughs in the treatment of HR-positive, HER2-negative breast cancer, and have a remarkable efficacy in the advanced setting when combined with endocrine therapy (ET) (Finn et al., 2015; Goetz et al., 2017; Hortobagyi et al., 2016). Presently, abemaciclib has been approved for high-risk, early HR-positive breast cancer plus ET as adjuvant treatment and its application is being explored in neoadjuvant setting (neoMONARCH and NCT04293393). However, therapies with combination of CDK4/6 inhibitors and NET only achieve a pCR rate of less than 5% (Hurvitz et al., 2020; Johnston et al., 2019).In the PALLET study, objective response rate (ORR) was not statistically significantly

different between the combinational therapy and monotherapy (54.3% vs 49.5%, P = 0.20) (Johnston et al., 2019). Likewise, in the FELINE study, there was no statistical difference in clinical, mammographic, ultrasound or magnetic resonance imaging (MRI) response between letrozole alone and combination of letrozole with a CDK4/6 inhibitor(Khan et al., 2020). Thus, it is reasonable to explore a novel neoadjuvant therapy for HR-positive, HER2-negative breast cancer.

Radiotherapy is one of the most important local methods to control malignancy by breaking DNA in tumor cells (Cox & Swanson, 2013). Stereotactic body radiation therapy (SBRT) is a novel and relatively safe radiotherapy technique and delivers high doses of radiation in a small number of fractions to the tumor lesion while minimizing radiation exposure to the surrounding tissues(Chmura et al., 2021). Currently, neoadjuvant radiotherapy can be applicable for inoperable breast cancer to allow an effective and sometimes more conservative surgery(Cardoso et al., 2018). Preclinical studies have indicated that CDK4/6 inhibitors can sensitize breast cancer cells to radiotherapy, possibly by inhibition of DNA damage repair response, enhancement of apoptosis and blockage of cell cycle progression, and inhibition of intra-tumor cellular metabolism and angiogenesis (Han et al., 2013; A. M. Pesch et al., 2020; G. Petroni et al., 2021). A preclinical study has explored different schedules of combination of radiotherapy and CDK4/6 inhibitors in human and mouse model and demonstrated that radiotherapy followed by CDK4/6 inhibitors can enhance antineoplastic effects, compared with their monotherapy alone or CDK4/6 inhibitors concurrent with or

followed by radiotherapy (G. Petroni et al., 2021). Moreover, the combination of CDK4/6 inhibitors and radiotherapy can lead to a satisfying overall disease control and safety based on several retrospective studies in metastatic breast cancer (Bosacki et al., 2021). Nevertheless, the efficacy and safety of this regimen in neoadjuvant treatment of early or locally advanced HR-positive, HER2-negative breast cancer are unknown. We have conducted a prospective pilot study to determine the efficacy and safety of neoadjuvant radiotherapy followed by dalpiciclib and exemestane in patients with newly-diagnosed early or locally advanced HR-positive, HER2-negative breast cancer. Dalpiciclib (SHR6390) is a potent selective inhibitor of CDK4/6 and exerts synergistic anti-tumor effect when combined with endocrine therapy for advanced luminal breast cancer in the DAWNA-1 trials (Xu et al., 2021).

Methods

Study design

This single-arm, non-controlled prospective pilot study in women with HR-positive, HER2-negative early or locally advanced breast cancer was registered with ClinicalTrials.gov (NCT05132790).

The study was approved by the Institutional Review Board and Ethics Committee of Shengjing Hospital of China Medical University, according to the Declaration of Helsinki and Good Clinical Practice guidelines. Written informed consent was obtained from each patient.

Participants

Eligible patients included pre- and post-menopausal women aged between 18 and 75

years, with histologically confirmed HR-positive, HER2-negative (0 or 1+ by

immunohistochemistry [IHC], or 2+ by IHC with negative results of fluorescence in

situ hybridization), early or locally advanced invasive breast cancer with the lesion of

≥2cm by MRI. Other main inclusion criteria were Eastern Cooperative Oncology

Group performance status 0-1, adequate marrow, hepatic and renal function. The key

exclusion criteria included stage IV tumors, inflammatory breast cancer, prior systemic

treatment of the current cancer, other malignancies, cardiac disease or history of cardiac

dysfunction, pregnancy, lactation, and refusal to use contraception.

Procedures

SBRT was performed every other day in breast target lesion with a total radiation dose

of 24Gy separated by 3 fractions. Within 2-4 weeks after the completion of radiotherapy,

individual patients received 6 cycles of combination of oral 150 mg Dalpiciclib daily

on days 1-21 of each 4-week cycle and 25 mg exemestane with a total duration of 24

weeks. Patients without menopause were also injected subcutaneously with 3.6 mg

goserelin every 4 weeks. All patients received study treatment until completion of all

prescribed protocol therapy, disease progression, unacceptable toxicity, or withdrawal

of consent. If disease progression, the patient was either proceed to surgery or received

alternative neoadjuvant therapy. Surgery was performed 3-5 weeks after the last dose of dalpiciclib to allow adverse event (AE) recovery. Recommended surgery and adjuvant therapy followed per local guidelines or institutional standards. The necessity of additional chemotherapy for patients with pCR was judged by investigators. A 5-year follow-up is conducted every 3 months during the first 2 years, and every 6 months for the subsequent 3 years. Additionally, safety data are collected within 90 days after surgery for subjects who discontinue study treatment.

Outcomes

The primary endpoint was the rate of residual cancer burden (RCB) 0-I. Secondary endpoints included ORR (defined as the proportion of patients with complete or partial response according to the Response Evaluation Criteria In Solid Tumors [RECIST] version 1.1 by MRI), pCR rate in the breast and axillary lymph nodes (tpCR; ypT0/is ypN0) and in the breast (bpCR; ypT0/is ypNx), breast-conservation rate, preoperative endocrine prognostic index (PEPI) score for breast cancer-specific survival (BCSS) (PEPI 0), and safety, according to the Common Terminology Criteria for Adverse Events (CTCAE) version 5.0.

Patient-Reported Outcome (PRO) assessments

In the study, the patient-reported outcome (PRO) assessments was measured using the European Organization for Research and Treatment of Cancer (EORTC) Quality of

Life Questionnaire Core-30 (QLQ-C30; version 3) (Cull, 1998) at baseline, at the end

of cycles 1 to 6, and before surgery. The PRO data were scored, according to the

EORTC scoring manual. A raw score was calculated as the average of items

contributing to a scale and standardized to a range of 0 to 100 points. On global health

status and functioning scales, a higher score indicated better status. Higher scores on

symptom scales indicated worse symptoms.

Tumor-infiltrating lymphocytes (TILs)

Tumor samples were collected when performing needle biopsy and surgery. The

specimens were fixed in 10% formalin and paraffin-embedded. The tissue sections (3

µm) were stained with hematoxylin and eosin (H&E) and the frequency of TILs were

quantified manually by a pathologist in a blinded manner. The TILs scores were

determined as the percentage of TILs in tumor area (tumor cells and stroma). The

percentage of stromal TILs (sTILs) was calculated in accordance with the

recommendations by an International TILs Working Group 2014 (Salgado et al., 2015).

Programmed cell death-ligand 1 (PD-L1) expression

The expression of programmed cell death-ligand 1 (PD-L1) was characterized by

immunohistochemistry (IHC) using anti-PD-L1 (clone SP142, Spring Bioscience, USA)

and the percentages of PD-L1 positive tumor cells (TC) or infiltrating inflammatory

cells (IC) were quantified.

Statistical analysis

This exploratory study involves 12 patients, with the sample size determined based on

clinical considerations, not statistical factors.

All statistical analyses were conducted using SAS 9.4 (North Carolina, USA).

Continuous data are presented as mean and standard deviation (SD) or mean and 95%

confidence interval (CI). Categorical data are expressed as frequency and percentage.

The 95% CIs of pathological complete response rate, proportion of patients with RCB-

0 or RCB-I, and ORR were estimated using the Clopper-Pearson method.

Results

Patient characteristics

Between November 2021 and January 2022, 12 HR-positive, HER2-negative patients

were screened and recruited in the trial. Their baseline demographic and clinical

characteristics are shown in **Table 1**. They had a median age of 48.5 years (range, 36-

56) and the median tumor size of 33.5mm (range, 23-90) assessed by breast MRI.

Patient outcomes

After the completion of study treatment, 2 (16.7%, 95% CI 3.5%-46.0%) out of 12

patients achieved RCB 0–I tumor response, tpCR and PEPI 0 (**Table 2**). There were 3

(25.0%) patients reached bpCR. Overall, this study led to an ORR (as assessed by breast

MRI) of 91.7% (95% CI 62.5%-100%, **Fig. 1**). Rate of conversion from mastectomy at

baseline to breast conservation at surgery was 25.0% (95% CI 8.3%-53.9%). The

median Ki67 of surgical specimens for patients without pCR was 7% (range, 5-70).

Toxicity

All patients completed the study treatment and AEs are summarized in **Table 3**. The

most frequent grade 1-2 AEs were leukopenia (58.3%), hot flushes (50.0%) and

lymphopenia (50.0%). There were some patients with Grade 3 AEs, including

neutropenia (66.7%) and leukopenia (25.0%), but not grade 4 AEs. There was no

serious AEs, no therapy-related death and no patients with dose interruption and

reduction or treatment discontinuation in this population. Furthermore, there was no

patient with radiation-related dermatitis and skin hyperpigmentation in this population.

Patient-reported outcome

All patients completed the EORTC QLQ-C30 and had available results of PRO

assessment (**Fig. 2**). There was no clinically significant deterioration in global health

status, physical, role, emotional, cognitive and social functioning in this population.

Alteration in tumor immune microenvironment

To examine the impact of this new neoadjuvant therapy on tumor environment, we

performed H&E staining of tissue sections in 10 evaluable patients (**Fig. 3**). Compared

with those in pre-treatment biopsied tissues, treatment significantly increased the

percentages of tumor-infiltrating lymphocytes (TILs) in the tumor environment (22.9%

vs.10.7%, P=0.043). IHC staining of PD-L1 revealed that there appeared no significant

difference in the percent of PD-L1 positive immune cells and tumor cells between the

pre-treatment and post-treatment specimens.

Discussion

This pioneering study reported the efficacy and safety of sequential neoadjuvant

radiotherapy and combination of dalpiciclib and exemestane in patients with HR-

positive, HER2-negative breast cancer. This new therapeutic regimen achieved

considerable anti-tumor activity with the RCB 0-I rate of 16.7% and ORR rate of 91.7%,

respectively. More importantly, this combinatory therapeutic strategy was well

tolerated in this population.

CDK4/6 inhibitors have recently been preclinically identified as radiosensitizers, acting

mainly through hindering DNA damage repair response after radiation exposure, and

enhancing cell apoptosis and blocking cell cycle, which provide a conceptual basis of

combining radiotherapy with a CDK4/6 inhibitor (Khasraw et al., 2017; Xie et al.,

2019). Moreover, both radiotherapy and CDK4/6 inhibitors exert immunomodulatory

effects and may synergize anticancer immune responses (Andrea M Pesch et al., 2020).

SBRT with hypofractionation and higher radiation doses is more effective than

conventional fractionation in triggering immune responses (Gandhi et al., 2015).

Consistent with the basic research (Giulia Petroni et al., 2021), we used the sequential

treatment with SBRT and combination of dalpiciclib and exemestane.

Unlike HER2-positive and triple-negative breast cancer, NCT or NET for HR-positive,

HER2-negative breast cancer only achieves a pCR rate of $\leqslant 10\%$ (Cortazar et al., 2014).

Our study indicated that this novel therapeutic regiment achieved a pCR and RCB 0-I

rate of 16.7%，while neoadjuvant therapies with combination of a CDK4/6 inhibitor

and letrozole led to a pCR rate of 0-3.8% and RCB 0-I rate of 6.1-7.7% and treatment

with NCT resulted in a pCR rate of 5.8-5.9% and RCB 0-I rate of 11.8-15.7% in HR-

positive, HER2-negative breast cancer patients (C. X. Ma et al., 2017; Prat et al., 2020).

It is notable that pCR rate (commonly defined as ypT0/isN0) is not significantly

associated with the prognosis of HR-positive, HER2-negative breast cancer (Cynthia X

Ma et al., 2017). In addition, clinical trials to test different therapies for patients with

HR-positive, HER2-negative breast cancer have focused on clinical response rates as

important clinical endpoints (Hurvitz et al., 2020; Johnston et al., 2019; Khan et al.,

2020; Khasraw et al., 2017; C. X. Ma et al., 2017). In our study, treatment with the new

therapeutic regiment led to eleven (91.7%) patients achieving objective responses by

MRI, while previous studies have revealed that NET achieves an objective response

rate of approximately 60% in HR-positive, HER2-negative breast cancer patients (Sella

et al., 2021). One of the principal objectives of neoadjuvant therapy is to facilitate

downstaging and enhance the rate of breast conservation. With the addition of SBRT,

a slightly higher rate of patients changed from mastectomy to breast conservation

compared with PALLET study (25.0% vs 14.1%). (Johnston et al., 2019). Consequently,

our new neoadjuvant therapeutic regiment appeared to improve the likelihood of pathological downstaging and achieve a margin-free resection, particularly for those with locally advanced and high-risk breast cancer.

Considering that treatment with radiotherapy or a CDK4/6 inhibitor can enhance anti-tumor immunity in vitro, We analyzed the impact of new neoadjuvant therapeutic regiment on alternation in biomarker expression in the numbers of TILs before and after the treatment., Our study showed a modest but statistically significant increase in the tumor tissues with the addition of SBRT. Apparently, sequential neoadjuvant therapies with radiotherapy, a CDK 4/6 inhibitor and exemestane turned non-responsive 'cold' tumors into responsive 'hot' ones. Therefore, our findings may provide significant insights into designing effective therapeutic strategies against HR-positive, HER2-negative breast cancer.

Patients with HR-positive, HER2-negative breast cancer were well-tolerated to the combinational regimen. During the observation period, those patients only developed grade 1-3 of AEs, including grade 3 neutropenia (66.7%), similar to that of combination of dalpiciclib and fulvestrant in the DAWNA-1 trial (84.2%) (Xu et al., 2021). There was no new AEs following therapies with radiotherapy and dalpiciclib in this population.

We recognized that our study had limitations, including a one-arm preliminary exploratory trial with small sample size. The small sample size limited the comparison of our data with historical data in the literature due to the potential bias. Consequently,

cross-trial comparisons like this hold limited significance. Thus, further prospective

randomized clinical trials with a larger population for longer treatment duration are

warranted to validate the findings.

In conclusion, our study provides novel insights into the neoadjuvant treatment of

HR+/HER2- breast cancer. The sequential radiotherapy and a CDK4/6 inhibitor plus

ET were effective and well-tolerated, which could serve as a feasible option for

neoadjuvant therapy for HR-positive, HER2 -negative breast cancers. Further

validation of these findings is warranted in a large-scale study.

Acknowledgments

Jiangsu Hengrui Pharmaceuticals provided the study drug dalpiciclib free of charge for

patients enrolled in the study. We thank the patients, their families involved in this study.

Conflict of interest

Jianfei Wang is the employee of Jiangsu Hengrui Pharmaceuticals Co., Ltd. Caigang

Liu is a senior editior of eLife. Yongliang Yang is a reviewing editor of eLife. The other

authors declare that no competing interests exist.

Funding

This research did not receive any specific grant from funding agencies in the public,

commercial, or not-for-profit sectors.

Author ORCIDs：Caigang Liu https://orcid.org/0000-0003-2083-235X

Data availability statement

The raw clinical and imaging data are protected due to patient privacy laws. The

datasets generated and/or analyzed during the current study are available from the

corresponding author Caigang Liu on request for 10 years; de-identified clinical data

and experimental data are available on request sharing, which will need the approval of

the Institutional Ethical Committees. De-identified data will then be transferred to the

inquiring investigator over secure file transfer.

References

Bosacki, C., Bouleftour, W., Sotton, S., Vallard, A., Daguenet, E., Ouaz, H., Cojoracu, I., Moslemi, D., Molekzadehmoghani, M., & Magné, N. (2021). CDK 4/6 inhibitors combined with radiotherapy: A review of literature. *Clin Transl Radiat Oncol*, *26*, 79-85. https://doi.org/10.1016/j.ctro.2020.11.010

Cardoso, F., Senkus, E., Costa, A., Papadopoulos, E., Aapro, M., André, F., Harbeck, N., Aguilar Lopez, B., Barrios, C. H., Bergh, J., Biganzoli, L., Boers-Doets, C. B., Cardoso, M. J., Carey, L. A., Cortés, J., Curigliano, G., Diéras, V., El Saghir, N. S., Eniu, A., . . . Winer, E. P. (2018). 4th ESO-ESMO International Consensus Guidelines for Advanced Breast Cancer (ABC 4)†. *Ann Oncol*, *29*(8), 1634-1657. https://doi.org/10.1093/annonc/mdy192

Chmura, S., Winter, K. A., Robinson, C., Pisansky, T. M., Borges, V., Al-Hallaq, H., Matuszak, M., Park, S. S., Yi, S., Hasan, Y., Bazan, J., Wong, P., Yoon, H. A., Horton, J., Gan, G., Milano, M. T., Sigurdson, E. R., Moughan, J., Salama, J. K., & White, J. (2021). Evaluation of Safety of Stereotactic Body Radiotherapy for the Treatment of Patients With Multiple Metastases: Findings From the NRG-BR001 Phase 1 Trial. *JAMA Oncol*, *7*(6), 845-852. https://doi.org/10.1001/jamaoncol.2021.0687

Cortazar, P., Zhang, L., Untch, M., Mehta, K., Costantino, J. P., Wolmark, N., Bonnefoi, H., Cameron, D., Gianni, L., Valagussa, P., Swain, S. M., Prowell, T., Loibl, S., Wickerham, D. L., Bogaerts, J., Baselga, J., Perou, C., Blumenthal, G., Blohmer, J., . . . von Minckwitz, G. (2014). Pathological complete response and long-term clinical benefit in breast cancer: the CTNeoBC pooled analysis. *Lancet*, *384*(9938), 164-172. https://doi.org/10.1016/s0140-6736(13)62422-8

Cox, J. A., & Swanson, T. A. (2013). Current modalities of accelerated partial breast irradiation. *Nat Rev Clin Oncol*, *10*(6), 344-356. https://doi.org/10.1038/nrclinonc.2013.65

Cull, A. (1998). EORTC QLQ-30. A breach of copyright. *Int J Pancreatol*, *24*(1), 64-65. https://doi.org/10.1007/bf02787535

Finn, R. S., Crown, J. P., Lang, I., Boer, K., Bondarenko, I. M., Kulyk, S. O., Ettl, J., Patel, R., Pinter, T., Schmidt, M., Shparyk, Y., Thummala, A. R., Voytko, N. L., Fowst, C., Huang, X., Kim, S. T., Randolph, S., & Slamon, D. J. (2015). The cyclin-dependent kinase 4/6 inhibitor palbociclib in combination with letrozole versus letrozole alone as first-line treatment of oestrogen receptor-positive, HER2-negative, advanced breast cancer (PALOMA-1/TRIO-18): a randomised phase 2 study. *Lancet Oncol*, *16*(1), 25-35. https://doi.org/10.1016/s1470-2045(14)71159-3

Gandhi, S. J., Minn, A. J., Vonderheide, R. H., Wherry, E. J., Hahn, S. M., & Maity, A. (2015). Awakening the immune system with radiation: optimal dose and fractionation. *Cancer*

398   *letters*, *368*(2), 185-190. https://doi: 10.1016/j.canlet.2015.03.024.

400   Goetz, M. P., Toi, M., Campone, M., Sohn, J., Paluch-Shimon, S., Huober, J., Park, I. H., Trédan, O.,
401   Chen, S. C., Manso, L., Freedman, O. C., Garnica Jaliffe, G., Forrester, T., Frenzel, M., Barriga,
402   S., Smith, I. C., Bourayou, N., & Di Leo, A. (2017). MONARCH 3: Abemaciclib As Initial
403   Therapy for Advanced Breast Cancer. *J Clin Oncol*, *35*(32), 3638-3646.
404   https://doi.org/10.1200/jco.2017.75.6155

406   Gradishar, W. J., Moran, M. S., Abraham, J., Aft, R., Agnese, D., Allison, K. H., Anderson, B., Burstein,
407   H. J., Chew, H., Dang, C., Elias, A. D., Giordano, S. H., Goetz, M. P., Goldstein, L. J., Hurvitz,
408   S. A., Isakoff, S. J., Jankowitz, R. C., Javid, S. H., Krishnamurthy, J., . . . Kumar, R. (2022).
409   Breast Cancer, Version 3.2022, NCCN Clinical Practice Guidelines in Oncology. *J Natl*
410   *Compr Canc Netw*, *20*(6), 691-722. https://doi.org/10.6004/jnccn.2022.0030

412   Han, Y. K., Lee, J. H., Park, G. Y., Chun, S. H., Han, J. Y., Kim, S. D., Lee, J., Lee, C. W., Yang, K., & Lee,
413   C. G. (2013). A possible usage of a CDK4 inhibitor for breast cancer stem cell-targeted
414   therapy. *Biochem Biophys Res Commun*, *430*(4), 1329-1333.
415   https://doi.org/10.1016/j.bbrc.2012.10.119

417   Hortobagyi, G. N., Stemmer, S. M., Burris, H. A., Yap, Y. S., Sonke, G. S., Paluch-Shimon, S.,
418   Campone, M., Blackwell, K. L., André, F., Winer, E. P., Janni, W., Verma, S., Conte, P., Arteaga,
419   C. L., Cameron, D. A., Petrakova, K., Hart, L. L., Villanueva, C., Chan, A., . . . O'Shaughnessy,
420   J. (2016). Ribociclib as First-Line Therapy for HR-Positive, Advanced Breast Cancer. *N Engl*
421   *J Med*, *375*(18), 1738-1748. https://doi.org/10.1056/NEJMoa1609709

423   Hurvitz, S. A., Martin, M., Press, M. F., Chan, D., Fernandez-Abad, M., Petru, E., Rostorfer, R.,
424   Guarneri, V., Huang, C. S., Barriga, S., Wijayawardana, S., Brahmachary, M., Ebert, P. J.,
425   Hossain, A., Liu, J., Abel, A., Aggarwal, A., Jansen, V. M., & Slamon, D. J. (2020). Potent
426   Cell-Cycle Inhibition and Upregulation of Immune Response with Abemaciclib and
427   Anastrozole in neoMONARCH, Phase II Neoadjuvant Study in HR(+)/HER2(-) Breast
428   Cancer. *Clin Cancer Res*, *26*(3), 566-580. https://doi.org/10.1158/1078-0432.CCR-19-
429   1425

431   Johnston, S., Puhalla, S., Wheatley, D., Ring, A., Barry, P., Holcombe, C., Boileau, J. F., Provencher,
432   L., Robidoux, A., Rimawi, M., McIntosh, S. A., Shalaby, I., Stein, R. C., Thirlwell, M., Dolling,
433   D., Morden, J., Snowdon, C., Perry, S., Cornman, C., . . . Jacobs, S. A. (2019). Randomized
434   Phase II Study Evaluating Palbociclib in Addition to Letrozole as Neoadjuvant Therapy in
435   Estrogen Receptor-Positive Early Breast Cancer: PALLET Trial. *J Clin Oncol*, *37*(3), 178-189.
436   https://doi.org/10.1200/jco.18.01624

438   Khan, Q. J., O'Dea, A., Bardia, A., Kalinsky, K., Wisinski, K. B., O'Regan, R., Yuan, Y., Ma, C. X.,
439   Jahanzeb, M., Trivedi, M. S., Spring, L., Makhoul, I., Wagner, J. L., Winblad, O., Amin, A. L.,

440   Blau, S., Crane, G. J., Elia, M., Hard, M., & Sharma, P. (2020). Letrozole + ribociclib versus
441   letrozole + placebo as neoadjuvant therapy for ER+ breast cancer (FELINE trial). *Journal*
442   *of Clinical Oncology*, *38*(15_suppl), 505-505.
443   https://doi.org/10.1200/JCO.2020.38.15_suppl.505

445   Khasraw, M., Chung, S., Day, B., Johns, T., Joshi, S., Madani, D., McDonald, K., & Whittaker, S. (2017).
446   Combination of palbociclib and radiotherapy for glioblastoma. Cell Death Discov
447   2017;3:17033. https://doi: 10.1038/cddiscovery.2017.33.

449   Ma, C. X., Gao, F., Luo, J., Northfelt, D. W., Goetz, M., Forero, A., Hoog, J., Naughton, M., Ademuyiwa,
450   F., Suresh, R., Anderson, K. S., Margenthaler, J., Aft, R., Hobday, T., Moynihan, T., Gillanders,
451   W., Cyr, A., Eberlein, T. J., Hieken, T., . . . Ellis, M. J. (2017). NeoPalAna: Neoadjuvant
452   Palbociclib, a Cyclin-Dependent Kinase 4/6 Inhibitor, and Anastrozole for Clinical Stage 2
453   or 3 Estrogen Receptor-Positive Breast Cancer. *Clin Cancer Res*, *23*(15), 4055-4065.
454   https://doi.org/10.1158/1078-0432.Ccr-16-3206

456   Parker, J. S., Mullins, M., Cheang, M. C., Leung, S., Voduc, D., Vickery, T., Davies, S., Fauron, C., He,
457   X., Hu, Z., Quackenbush, J. F., Stijleman, I. J., Palazzo, J., Marron, J. S., Nobel, A. B., Mardis,
458   E., Nielsen, T. O., Ellis, M. J., Perou, C. M., & Bernard, P. S. (2009). Supervised risk predictor
459   of breast cancer based on intrinsic subtypes. *J Clin Oncol*, *27*(8), 1160-1167.
460   https://doi.org/10.1200/jco.2008.18.1370

462   Pesch, A. M., Hirsh, N. H., Chandler, B. C., Michmerhuizen, A. R., Ritter, C. L., Androsiglio, M. P.,
463   Wilder-Romans, K., Liu, M., Gersch, C. L., Larios, J. M., Pierce, L. J., Rae, J. M., & Speers, C.
464   W. (2020). Short-term CDK4/6 Inhibition Radiosensitizes Estrogen Receptor-Positive
465   Breast Cancers. *Clin Cancer Res*, *26*(24), 6568-6580. https://doi.org/10.1158/1078-
466   0432.Ccr-20-2269

468   Petroni, G., Buqué, A., Yamazaki, T., Bloy, N., Liberto, M. D., Chen-Kiang, S., Formenti, S. C., &
469   Galluzzi, L. (2021). Radiotherapy Delivered before CDK4/6 Inhibitors Mediates Superior
470   Therapeutic Effects in ER(+) Breast Cancer. *Clin Cancer Res*, *27*(7), 1855-1863.
471   https://doi.org/10.1158/1078-0432.Ccr-20-3871

473   Prat, A., Saura, C., Pascual, T., Hernando, C., Muñoz, M., Paré, L., Farré, B. G., Fernández, P. L.,
474   Galván, P., & Chic, N. (2020). Ribociclib plus letrozole versus chemotherapy for
475   postmenopausal women with hormone receptor-positive, HER2-negative, luminal B
476   breast cancer (CORALLEEN): an open-label, multicentre, randomised, phase 2 trial. *The*
477   *lancet oncology*, *21*(1), 33-43. https://doi: 10.1016/s1470-2045(19)30786-7.

479   Salgado, R., Denkert, C., Demaria, S., Sirtaine, N., Klauschen, F., Pruneri, G., Wienert, S., Van den
480   Eynden, G., Baehner, F. L., Penault-Llorca, F., Perez, E. A., Thompson, E. A., Symmans, W.
481   F., Richardson, A. L., Brock, J., Criscitiello, C., Bailey, H., Ignatiadis, M., Floris, G., . . . Loi, S.

482  (2015). The evaluation of tumor-infiltrating lymphocytes (TILs) in breast cancer:
483    recommendations by an International TILs Working Group 2014. *Ann Oncol*, *26*(2), 259-
484    271. https://doi.org/10.1093/annonc/mdu450

486 Sella, T., Weiss, A., Mittendorf, E. A., King, T. A., Pilewskie, M., Giuliano, A. E., & Metzger-Filho, O.
487    (2021). Neoadjuvant Endocrine Therapy in Clinical Practice: A Review. *JAMA Oncol*, *7*(11),
488    1700-1708. https://doi.org/10.1001/jamaoncol.2021.2132

490 Xie, X., Zheng, W., Chen, T., Lin, W., Liao, Z., Liu, J., & Ding, Y. (2019). CDK4/6 inhibitor palbociclib
491    amplifies the radiosensitivity to nasopharyngeal carcinoma cells via mediating apoptosis
492    and suppressing DNA damage repair. *OncoTargets and therapy*, 11107-11117.
493    https://doi: 10.2147/ott.S234221.

495 Xu, B., Zhang, Q., Zhang, P., Hu, X., Li, W., Tong, Z., Sun, T., Teng, Y., Wu, X., Ouyang, Q., Yan, X.,
496    Cheng, J., Liu, Q., Feng, J., Wang, X., Yin, Y., Shi, Y., Pan, Y., Wang, Y., . . . Zou, J. (2021).
497    Dalpiciclib or placebo plus fulvestrant in hormone receptor-positive and HER2-negative
498    advanced breast cancer: a randomized, phase 3 trial. *Nat Med*, *27*(11), 1904-1909.
499    https://doi.org/10.1038/s41591-021-01562-9

501 **Table 1: Baseline characteristics of the patients.**

	Patients (n=12)
Median age (range), years	48.5 (36-56)
⩽ 50	6 (50.0%)
> 50	6 (50.0%)
Menopausal status	
Premenopausal	6 (50.0%)
Postmenopausal	6 (50.0%)
Tumor size	
T2	11 (91.7%)
T3	1 (8.3%)
Lymph node status	
N0	5 (41.7%)
N1	2 (16.7%)
N2	5 (41.7%)
Clinical stage	
IIA	4 (33.3%)
IIB	3 (25.0%)
IIIA	5 (41.7%)
Tumor grade	
II	11 (91.7%)
III	1 (8.3%)
ER expression	
> 10%	12 (100.0%)
PgR expression	
< 20%	3 (25.0%)
⩾ 20%	9 (75.0%)
HER2 expression	
0	7 (58.3%)
1+	1 (8.3%)
2+, FISH	4 (33.3%)
Median baseline Ki67 (range), %	35 (15-70)
⩽ 14%	3 (25.0%)
> 14%	9 (75.0%)

Data are n (%) or median (range).

ER: estrogen receptor; PgR: progesterone receptor; HER2: human epidermal growth factor receptor 2; FISH: fluorescence in situ hybridization.

Table 2: Pathological and clinical response (n=12).

	n (%)
Total pathological complete response	2 (16.7%)
Breast pathological complete response	3 (25.0%)
Residual cancer burden score	
0	2 (16.7%)
I	0
II	6 (50.0%)
III	4 (33.3%)
Radiological response	
Complete response	2 (16.7%)
Partial response	9 (75.0%)
Stable disease	1 (8.3%)
Objective response rate	11 (91.7%)
Preoperative endocrine prognostic index	
0	2 (16.7%)
1-3	0
≥4	10 (83.3%)

Data are presented as n (%)

Table 3: Treatment-emergent adverse events (n=12).

	n (%)		
	Grade 1 or 2	Grade 3	Grade 4
Neutropenia	3 (25.0%)	8 (66.7%)	0
Leukopenia	7 (58.3%)	3 (25.0%)	0
Hot flushes	6 (50.0%)	0	0
Lymphopenia	6 (50.0%)	0	0
Insomnia	4 (33.3%)	0	0
Alopecia	4 (33.3%)	0	0
Anemia	4 (33.3%)	0	0
Hyperuricemia	4 (33.3%)	0	0
Blood urea increased	4 (33.3%)	0	0
Rash	3 (25.0%)	0	0
Thrombocytopenia	3 (25.0%)	0	0
Hyperglycemia	3 (25.0%)	0	0
Creatinine increased	3 (25.0%)	0	0
Fatigue	3 (25.0%)	0	0
Arthralgia	2 (16.7%)	0	0
γ-glutamyl transferase increased	2 (16.7%)	0	0
Hypocalcemia	2 (16.7%)	0	0
Abnormal T wave of electrocardiogram	2 (16.7%)	0	0
Hyponatremia	2 (16.7%)	0	0
Lactate dehydrogenase increased	2 (16.7%)	0	0
Diarrhea	1 (8.3%)	0	0
Headache	1 (8.3%)	0	0
Mucosal inflammation Constipation	1 (8.3%)	0	0
Proteinuria	1 (8.3%)	0	0
Hypertriglyceridemia	1 (8.3%)	0	0
Alanine aminotransferase increased	1 (8.3%)	0	0

Data are presented as n (%).

Figure legends

Figure 1: Waterfall plot of best reduction in tumor measurement from basline.

Figure 2: Mean change from baseline in QLQ-C30 over time. (A) Global health

status. (B-F) Physical, role, emotional, cognitive and social functioning.

Figure 3: TILs and PD-L1 changes before and after treatment. (A) Study schema. (B)

Representative TILs. Scale bar, 100 μM. (C-E) Percentage of TILs, PD-L1 expression

in immune cell and tumor cell in paired specimens (n=10). Bars, boxes, and whiskers

represent median, interquartile range, and range, respectively.