Interplay Between the Gut Microbiome and Typhoid Fever: Insights from Endemic Countries and a Controlled Human Infection Model

Philip M. Ashton[1,2], Leonardos Mageiros[3$], James E. Meiring[1,2,4$], Angeziwa Chunga Chirambo[1,5$], Farhana Khanam[6$], Sabina Dongol[9,10$], Happy Banda[1], Abhilasha Karkey[9,10], Lorena Preciado Llanes[12], Helena Thomaides-Brears[12], Malick Gibani[12,13], Nazmul Hasan Rajib[6], Nazia Rahman[6], Prasanta Kumar Biswas[6], Md Amirul Islam Bhuiyan[6], Sally Kay[7], Kate Auger[7], Olivier Seret[7], Nicholas R. Thomson[7,8], Andrew J Pollard[12,14], Stephen Baker[15], Buddha Basnyat[9,17,], John D. Clemens[6], Christiane Dolecek[17], Sarah J. Dunstan[18], Gordon Dougan[15,16], Robert S. Heyderman[19], Virginia E. Pitzer[20], Firdausi Qadri[6], Melita A. Gordon[1,2], Kathryn E. Holt[3,8*], Thomas C. Darton[4*], STRATAA Study Group[21]

1. Malawi-Liverpool-Wellcome Programme, Blantyre, Malawi
2. Institute of Infection, Veterinary and Ecological Sciences, University of Liverpool, Liverpool, UK
3. Department of Infectious Diseases, School of Translational Medicine, Monash University, Melbourne, Victoria, Australia
4. Clinical Infection Research Group, Division of Clinical Medicine, School of Medicine & Population Health, and The Florey Institute for Infection, University of Sheffield, UK
5. Department of Medical Laboratory Sciences, School of Life Sciences and Allied Health Professions, Kamuzu University of Health Sciences, Malawi
6. International Centre for Diarrhoeal Diseases Research, Bangladesh, Dhaka, Bangladesh
7. Wellcome Sanger Institute, Hinxton, UK
8. Department of Infection Biology, London School of Hygiene & Tropical Medicine, London, UK
9. Oxford University Clinical Research Unit, Patan Academy of Health Sciences, Kathmandu, Nepal
10. Patan Academy of Health Sciences, Patan Hospital, Lalitpur, Nepal
11. MRC Human Immunology Unit, MRC Weatherall Institute of Molecular Medicine, University of Oxford, Oxford, OX3 9DS, UK.
12. Oxford Vaccine Group, Department of Paediatrics, University of Oxford, Oxford, UK
13. Department of Infectious Disease, Imperial College London, London, UK
14. NIHR Oxford Biomedical Research Centre, Oxford, UK
15. Department of Medicine, University of Cambridge, Cambridge, UK
16. Centre for Translational Stem Cell Biology, Hong Kong
17. Nuffield Department of Medicine, Centre for Tropical Medicine and Global Health, University of Oxford, Oxford, UK
18. Department of Infectious Diseases, The University of Melbourne at The Peter Doherty Institute for Infection and Immunity, Melbourne, Australia
19. Research Department of Infection, Division of Infection and Immunity, University College London, London, UK

20. Department of Epidemiology of Microbial Diseases, Yale School of Public Health, Yale University, New Haven, CT, USA

21. Strategic Typhoid Alliance Across Asia and Africa Consortium

$ contributed equally
* contributed equally

Working title: Gut microbiome in relation to typhoid fever

Abstract

Typhoid fever is a systemic infection caused by *Salmonella enterica* serovar Typhi (*S.* Typhi) invasion from the gut lumen. Transmission between people occurs through ingestion of contaminated food and water, particularly in settings with poor water and sanitation infrastructure, resulting in over 10 million illnesses annually. As the pathogen invades via the gastrointestinal tract, it is plausible that the gut microbiome may influence the outcome of *S.* Typhi exposure. There is some evidence that bacteria producing short-chain fatty acids (SCFAs) may create an environment unfavourable to invasive *Salmonella,* but data from humans is limited.

To investigate the association between gut microbiome and typhoid fever, we analysed samples collected from three all-age cohorts enrolled in a prospective surveillance study conducted across three settings where typhoid fever is endemic (Dhaka, Bangladesh; Blantyre, Malawi; and Kathmandu, Nepal). Cohorts consisted of acute typhoid fever patients (n=92), asymptomatic household contacts of typhoid fever patients (representing individuals who were likely exposed to *S.* Typhi but did not develop disease, n=97), and asymptomatic serosurvey participants with high Vi antibody titres (representing individuals who were exposed to *S.* Typhi and may be carriers, n=69). The stool microbiomes of each cohort were characterised using shotgun metagenomics, and bacterial diversity, composition, and function were compared.

We identified 4 bacterial species that were significantly lower in abundance in typhoid fever patients compared with household contacts (i.e. probably exposed), in two of the three participant populations (Bangladesh and Malawi). These bacteria may represent taxa that provide protection against development of clinical infection upon exposure to *S.* Typhi, and include the inflammation-associated species *Prevotella copri* clade A and *Haemophilus parainfluenzae*. Our functional analysis identified 28 specific metabolic gene clusters (MGCs) negatively associated with typhoid fever in Bangladesh and Malawi, including seven MGCs involved in SCFA metabolism. The putative protection provided by microbiome SCFA metabolism was supported by data from a controlled human infection model conducted in a UK population, in which participants who did not develop typhoid fever following ingestion of *S.* Typhi had higher abundance of a putative SCFA-metabolising MGC (q-value = 0.22).

79 This study identified the same protective associations between taxonomic and functional
80 microbiota characteristics and non-susceptibility to typhoid fever across multiple human
81 populations. Future research should explore the potential functional role of SCFAs and
82 inflammation-associated bacteria in resistance to *S.* Typhi and other enteric infections.

Introduction

Typhoid fever, caused by invasive *Salmonella enterica* serovar Typhi (*S.* Typhi) infection, causes an estimated 10.9 million illnesses and 116,800 deaths per year disproportionately affecting people in South and Southeast Asia and sub-Saharan Africa (Stanaway et al. 2019; Meiring et al. 2023). The introduction of typhoid conjugate vaccines is expected to decrease the number of typhoid fever cases in these settings, along with improvements in water, sanitation and hygiene (WASH). However, the large burden of disease, the role of asymptomatic gall bladder carriage as a source of infection, and the fact that typhoid fever often impacts the most impoverished and marginalised communities, means that eradicating this infection will be difficult (Meiring et al. 2021; Patel et al. 2021; Steele et al. 2016).

The human gut microbiota, a complex ecosystem comprising trillions of microbial cells, plays an indispensable role in shaping our overall health and susceptibility to diseases (Shreiner, Kao, and Young 2015). This densely populated microbial environment is not only pivotal in the processes of digestion and nutrient absorption but also intimately connected to our immune system, determining its responses to various stimuli (Belkaid and Hand 2014). As the nexus between the external environment and our internal physiology, the gut microbiome is a critical determinant of the outcome of gastrointestinal infectious disease exposures, via a phenomenon termed 'colonization resistance' (Stecher et al. 2007; Winter et al. 2010; Lopez et al. 2016; Libertucci and Young 2019). This protective effect arises from multifaceted interactions: direct competition for nutrients, production of antimicrobial compounds, and modulation of the host's innate and adaptive immune responses (Leshem, Liwinski, and Elinav 2020). Some of the earliest work on colonisation resistance showed that suppressing the gut microbiome of mice with antibiotics dramatically reduced the dose of *Salmonella* required to establish infection (Bohnhoff, Drake, and Miller 1954). Recent work has made great strides in understanding mechanisms by which *Salmonella* and the microbiome interact (Rogers, Tsolis, and Bäumler 2021). Short chain fatty acids (SCFAs) are important in interaction between the microbiota and *Salmonella*, reducing the availability of oxygen in the gut (Byndloss et al. 2017) and acidifying the cytosol of *Salmonella,* inhibiting growth (Jacobson et al. 2018). Despite well-established knowledge about the gut microbiome's function in preventing *Salmonella* infections and the established faecal-oral transmission pathway of *S.* Typhi, the precise influence of the gut microbiome on the outcome of *S.* Typhi exposure remains underexplored, complicated by the human host-specificity of *S.* Typhi.

While insights from animal studies of invasive infection with other *S. enterica* serovars can be

informative, detailed studies of the interaction between the human gut microbiome and *S.* Typhi

specifically are lacking. Controlled human infection model (CHIM) studies provide an

experimental system in which to interrogate infections in humans, complementing observational

studies of natural infections (Waddington et al. 2014; Gibani et al. 2020). Previous data

generated from this model revealed an association between typhoid fever susceptibility and the

presence of *Methanobrevibacter* in the gut microbiome (Zhang et al. 2018). Analysing natural

infections, Haak et al. determined that typhoid fever patients in Bangladesh had reduced

microbial diversity and fewer SCFA producers than healthy counterparts (Haak et al. 2020).

Better understanding of the causal relationships between the gut microbiome and typhoid fever

could lead to novel preventative mechanisms and diagnostics, as well as an improved

understanding of colonisation resistance to bacteria causing serious invasive diseases in high

burden settings.

To fill this knowledge gap, we sequenced the stool microbiome of 258 participants enrolled in

the STRATAA study, conducted in three diverse high typhoid fever burden settings in Asia and

Africa (Meiring et al. 2021). Signals that replicated in more than one study population were

further investigated in a UK-based CHIM cohort.

Materials & Methods

Participants and sample collection

Detailed methods for the STRATAA study have been published previously (Darton et al. 2017; Meiring et al. 2021). In brief, approximately 100,000 individuals were enumerated in a demographic census in three communities: Ndirande in Blantyre, Malawi; Lalitpur in Kathmandu, Nepal; and Mirpur in Dhaka, Bangladesh. Febrile patients, of any age, within the study populations were recruited via passive surveillance, and "acute typhoid fever" patients were defined as those with positive blood cultures yielding *S.* Typhi. When participants presented with fever, stool samples were collected where possible, preferably prior to antimicrobial use. Stool specimens were transferred to -80°C within 6 hours of collection. Data on antimicrobial use prior to their enrolment was recorded (Darton et al. 2017). Blood-culture confirmed typhoid fever patients were then followed up and stool samples were collected from their household contacts; asymptomatic stool culture-negative individuals were included in microbiome sequencing. High Vi-titre individuals were identified from a community-based serological survey of up to 8,500 age-stratified participants per site from the original demographic census. From the serological survey, samples were analysed for anti-Vi IgG antibody (Meiring et al. 2021). The participants at each site with the highest Vi responses were followed up and stool sample collected for microbiome analysis (Khanam et al. 2021). Stool cultures were *S.* Typhi-negative in all but one participant (Khanam et al. 2021), however it is known that carriers shed only intermittently and high-Vi individuals is an acceptable predictor for *S.* Typhi carriage (Robbins and Robbins 1984; Qureshi et al. 2024). For the microbiome sub-study, we randomly selected for sequencing stool samples from up to 40 participants from each site, from each of three groups: acute typhoid fever cases, household contacts, and serosurvey participants with high-Vi titres.

Research ethics committee approval for a joint study protocol across all three surveillance sites was obtained within each country and from the Oxford Tropical Research Ethics Committee (University of Oxford, Oxford, UK).

DNA extraction and sequencing

DNA extraction from stool was done using the Fast-Prep 24 FastDNA Spin Kit (MP Biomedicals, CA, USA) according to manufacturer's instructions. DNA was sent to the Wellcome Sanger Institute (UK) for metagenomic sequencing using Illumina HiSeq 2500 or HiSeq 4000 to generate 150 bp paired end reads, yielding an average 14.8 million read pairs (standard deviation 2.3 million) per sample.

Bioinformatics

Raw sequencing data was quality trimmed and adapters were removed using bbduk v38.96 with the parameters `ktrim=r k=23 mink=11 hdist=1 tbo tpe qtrim=r trimq=20 minlength=50`. Sequence deriving from human DNA was removed by mapping to a human reference genome (GCF_009914755.1) and the hostile tool (Constantinides, Hunt, and Crook 2023). Taxonomic profiling was carried out with metaphlan v4.0.6 with database version mpa_vOct22_CHOCOPhlAnSGB_202212 (Blanco-Míguez et al. 2023).

We used the BiG-MAP program (Victória Pascal Andreu et al. 2021) (https://github.com/medema-group/BiG-MAP, commit e7b8042) to compare metagenomic readsets against a database of non-redundant metabolic gene clusters (MGCs) identified using gutSMASH (Victòria Pascal Andreu et al. 2023) from a collection of reference genomes from the Culturable Genome Reference, the Human Microbiome Project and other Clostridia genomes (https://zenodo.org/records/7252625#.ZFVTrexBz0r). Each MGC is assigned to a species which is the species from which the representative sequence representing the cluster was obtained. BiG-MAP calculated the number of reads per kilobase per million reads (RPKM) for each metabolic gene cluster.

A sub-sample of 8.8 million reads was taken from each readset using the `sample` command of the seqtk tool (v1.3-r106). Sub-sampled reads were then analysed with Resistance Gene Identifier (RGI) v6.0.2 bwt command against the CARD v3.2.7 reference database (Alcock et al. 2023). Due to uncertainty regarding some CARD database classifications (e.g. OXA-1 was classified as a carbapenemase, a curation error which has subsequently been resolved), we used the drug class information for each gene from the NCBI AMR Reference Gene Database (refgene) v2023-08-08.2 (https://www.ncbi.nlm.nih.gov/pathogens/refgene). Only "Core" genes

from NCBI refgene were analysed; these genes are almost entirely mobilisable AMR
determinants and do not include mutational resistance. Extended-spectrum beta-lactamases
(ESBLs) were identified based on the gene product descriptions in NCBI refgene. Only genes
with an Average Percent Coverage of 100% in the RGI analysis were included in further
analysis. The number of reads mapped to each gene (RGI Completely Mapped Reads) was
normalised by the length of the gene (RGI Reference Length) to generate a "Reads per kilobase
of AMR gene" metric. There was no need to normalise for the number of reads, as all readsets
had been sub-sampled to the same number of reads. RGI results were parsed and analysed
using the scripts described in the Data Availability section of this manuscript. AMR genes
identified in *S.* Typhi were obtained from (Argimón et al. 2021).
204

205 Statistical analysis

Statistical analyses were done in R (v4.1.0). To quantify alpha diversity, Shannon's index was
calculated using the alpha.div function in the rbiom package (v1.0.3), and ANOVA analysis
carried out with the aov function (stats package, v4.1.0). The Bray-Curtis metric of beta diversity
was calculated with the vegdist function of the R package vegan (v2.6.4). Principal co-ordinates
analysis (PCoA) was carried out with the cmdscale function (package stats, v4.1.0), and
statistics comparing beta diversity between participant groups (i.e. acute typhoid, household
contacts, presumptive carrier) was carried out using a PERMANOVA approach implemented by
the adonis2 function of the vegan package (v2.6.4). Associations between taxa and phenotypes
of interest (e.g. participant group) were explored using multivariable linear modelling
implemented in the maaslin2 function of the maaslin2 R package (v1.6.0) (Mallick et al. 2021).
Depending on the analysis, co-variates such as age, sex, antibiotic exposure, and sequencing
run were included, as described. For maaslin2 analyses, Benjamini-Hochberg correction was
applied for multiple testing. For analyses of data from endemic cohorts a q-value threshold of
0.05 was used to identify significant associations, while for CHIM data analyses, due to small
sample size, the default maaslin2 q-value threshold of 0.25 was used.

221 Controlled human infection model analysis

We used data from a *Salmonella* Typhi and Paratyphi A CHIM cohort to validate the
associations identified in the endemic countries. Briefly, healthy adults aged 18-60 were
recruited to be challenged with a single oral dose of either *S.* Typhi or *S.* Paratyphi A (Gibani et
al. 2020). The primary endpoint for the model was a diagnosis of typhoid or paratyphoid fever,

defined as a temperature of ≥38°C persisting for ≥12 hours and/or *S.* Typhi/Paratyphi A

bacteraemia in a sample collected ≥72 hours after oral challenge. Only data from participants

who were being challenged for the first time were analysed here. Stool samples were taken

from participants prior to challenge and stored at -80°C. DNA extraction, sequencing and

analyses was undertaken using the methods described above. The primary maaslin2 analyses

for this cohort included only species or MGC classes that were significantly associated with

participant group in at least two endemic country cohorts; a secondary analysis was carried out

without this restriction. Analyses were done separately for participants challenged with either *S.*

Typhi or *S.* Paratyphi A, in addition to a combined analysis including both pathogens.

Results

Description of population

Stool microbiome sequences were successfully generated for 258 participants from Bangladesh (n = 80), Malawi (n = 102), and Nepal (n = 76). Across the three populations, there were three participant groups; 92 patients with acute typhoid fever, 97 household contacts of acute typhoid cases, and 69 people with high anti-Vi titres. The baseline characteristics of participants are described in Table 1. Overall, acute typhoid fever patients were significantly younger than household contacts, while in Bangladesh and Malawi high-Vi titre participants were significantly older than household contacts (Supplementary Figure 1, Table 1). Overall, in Malawi, Bangladesh, and Nepal, 62%, 53% and 54% of participants were female, respectively (Table 1, Supplementary Figure 2). In Bangladesh and Malawi, sex distribution was similar between the participant groups, whereas in Nepal, 82% of household contacts and 67% of carriers were female, while only 24% of typhoid fever cases were female (Table 1, Supplementary Figure 2). Of the acute typhoid fever cases, 37.5%, 73.9% and 44.8% from Bangladesh, Malawi, and Nepal, respectively, reported antimicrobial use in the 2 weeks prior to stool sample collection.

252 *Table 1: Characteristics of participants in the microbiome sub-study of STRATAA surveillance*

		Bangladesh	Malawi	Nepal
Acute typhoid	N	40	23	29
	Median age, years (range)	6 (1-60)	10 (3-40)	17 (5-38)
	Female (%)	48%	61%	24%
	Antibiotics in last 2 weeks (%)	38%	74%	45%
Carrier	N	NA*	39	30
	Median age	NA*	32	43.9
	Female (%)	NA*	56%	67%
	Antibiotics in last 2 weeks (%)	NA*	NA†	NA†
Control	N	40	40	17
	Median age, years (range)	29 (4-65)	24 (18-41)	35 (15-48)
	Female (%)	65%	65%	82%
	Antibiotics in last 2 weeks (%)	NA†	NA†	NA†

253 * Carriers from Bangladesh were not included in the analysis due to sample protocol differences

254 † Antibiotic usage in the two weeks previous was an exclusion criterion for carriers and

255 household contacts.

Acute typhoid fever patients compared with household contacts

First, we compared stool metagenomic profiles of acute typhoid patients with those of

household contacts, within each of the three study populations, to explore microbiome

signatures associated with typhoid fever. Participant group was not significantly associated with

differences in alpha diversity in any population (Supplementary Figure 3, Supplementary Table

1). Beta diversity varied significantly between typhoid fever patients and household contacts

across all sites (Figure 1, Supplementary Table 2). The proportion of beta diversity variance

explained by participant group was highest in Malawi (R^2 = 0.25, FDR = 0.01), followed by

Bangladesh (R^2 = 0.07, FDR = 0.01), Nepal (R^2 = 0.06, FDR = 0.01 and the combined analysis

(R^2 = 0.04, FDR = 0.01) (Figure 1, Supplementary Table 2, Supplementary Table 3,

Supplementary Table 4, Supplementary Table 5). Participant group explained the greatest beta

diversity variance of the factors investigated (which also included sex, age, and reported

antibiotic use) in a combined analysis of all three sites and for each site individually. In Malawi,

prior antibiotic usage and the interaction of prior antibiotic usage with age and sex were also

significantly associated with beta diversity, although these variables explained a lower

proportion of variance than participant group (R^2 0.02-0.05, Supplementary Table 4). In Nepal,

sex was significantly associated with beta diversity and explained almost as much variance as

participant group (R^2 = 0.04, Supplementary Table 5), which was itself associated with sex at

this site. Firmicutes was the dominant phylum in typhoid fever patients and household contacts

at all three sites, followed by Bacteroidetes in Malawi and Nepal and Actinobacteria in

Bangladesh (Supplementary Figure 4, Supplementary Table 6).

Figure 1: Principal co-ordinate analysis of the beta diversity between household contacts and typhoid fever
participants across all three sites, and for each site individually. The proportion of variance explained (R^2) and the
Bonferroni corrected FDR from a PERMANOVA analysis including participant group, sex, and antibiotic usage are
displayed on each plot. In the combined analysis, country of sampling was also included in the PERMANOVA
analysis.

Maaslin2 analysis of taxonomic profiles identified 92, 23 and 0 species significantly associated

with household contact vs typhoid fever participant groups in Malawi, Bangladesh, and Nepal,

respectively (Figure 2, Supplementary Table 7, Supplementary Table 8). No taxa showed

significant associations across all three sites; however, four showed consistent associations at

two sites (Bangladesh and Malawi), all of which were negatively associated with typhoid fever

(Figure 2). These four species were *Prevotella copri* clade A, *Haemophilus parainfluenzae,*

Clostridium SGB6179 and a *Veillonellaceae* spp represented by a metagenome assembled

genome, GGB4266_SGB5809. Although there were differences in age between cases and

household contacts at all sites, age was included as a covariate in maaslin2, and the

associations with typhoid status were also evident within age groups where data from that age

group was available (Supplementary Figures 5-8). Of these four species, none were significantly

associated with typhoid fever in Nepal or the CHIM; most samples from the CHIM did not

contain any reads assigned to these species (Supplementary Table 29).

Species	Cohort	N	Coefficient	Standard Error	Q-value	Median abundance typhoid	Median abundance control
s__Clostridium_SGB6179	Bangladesh	80	8.79	1.34	0.000043	0	0.166
s__Clostridium_SGB6179	Malawi	63	5.52	1.6	0.049	0	0.149
s__Clostridium_SGB6179	Nepal	46	-1.01	1.34	1	0.0362	0
s__Clostridium_SGB6179	CHIM	26	-0.0825	1.17	0.94	0	0
s__Prevotella_copri_clade_A	Bangladesh	80	4.54	0.978	0.0097	0.0488	0.331
s__Prevotella_copri_clade_A	Malawi	63	10.1	1.98	0.00099	0.0991	18.2
s__Prevotella_copri_clade_A	Nepal	46	-1.54	1.47	0.98	5.72	0.903
s__Prevotella_copri_clade_A	CHIM	26	-0.33	1.78	0.94	0	0
s__GGB4266_SGB5809	Bangladesh	80	4.58	1.09	0.015	0.00924	0.213
s__GGB4266_SGB5809	Malawi	63	8.73	1.29	0.0000076	0	1.89
s__GGB4266_SGB5809	Nepal	46	2.96	1.77	0.9	0	0.0155
s__GGB4266_SGB5809	CHIM	26	0.424	0.287	0.93	0	0
s__Haemophilus_parainfluenzae	Bangladesh	80	3.64	0.979	0.03	0	0.0147
s__Haemophilus_parainfluenzae	Malawi	63	6.85	1.11	0.00005	0	0.824
s__Haemophilus_parainfluenzae	Nepal	46	-0.44	1.49	1	0.00797	0.00514
s__Haemophilus_parainfluenzae	CHIM	26	0.62	1.34	0.94	0	0

Figure 2: Forest plot of species that significantly differed between household contacts and typhoid fever participants
in at least two endemic countries. Results from all three endemic countries and the CHIM cohort are shown for
context. CHIM maaslin2 analysis only included species that were significantly associated in at least two of the
endemic country sites. Species are labelled with their GTDB identifier, note that 's__ GGB4266_SGB5809' refers to a
proposed novel genus/species of Veillonellaceae.

Some species were associated with typhoid fever at a single endemic country site. In

Bangladesh, there were 19 species for which the relative abundance significantly differed

between typhoid fever participants and household contacts, that didn't differ at any other site, or

in the CHIM (Supplementary Table 7). Three species were negatively associated with typhoid

fever (*Prevotella copri* clade C, *Romboutsia timonensis*, and *Ligilactobacillus* ruminis), while 16

species including 7 species of Actinomyces were positively associated with typhoid fever

(Supplementary Table 7). In Malawi, 70 species were negatively associated with typhoid fever,

including *Ruminococcus gnavus*, *Roseburia intestinalis*, *Roseburia inulinivorans,* and

Faecalibacterium prausnitzii (Supplementary Table 8), while 18 species were positively

associated with typhoid fever including 11 (61%) that are only in the Metaphlan4 database as

metagenome assembled genomes (Supplementary Table 8).

To complement our taxonomic analysis, we also carried out functional gene analysis. We

explored two dimensions of variation in metabolic gene clusters (MGCs). First, we examined the

distribution of specific MGCs, investigating participant group differences at a granular, sub-type

level (e.g. a specific RNF complex from a strain of *Bacteroides ovatus*). Second, we compared

the abundance of different MGC types between participant groups (e.g. all RNF complex

MGCs). This dual approach enables us to capture both detailed and broad patterns of functional

gene distribution within the microbiome.

We identified 264 specific MGCs that significantly differed between typhoid fever patients and

household contacts in the Malawi cohort, 126 in Bangladesh, and 2 in Nepal (Supplementary

Table 24, Supplementary Table 25, Supplementary Table 26). Neither of the specific MGC

associations identified in the Nepal data were replicated in the Malawi or Bangladesh cohorts.

There were 28 specific MGCs significantly associated with typhoid fever in both Bangladesh and

Malawi (summarised in Table 2, full information in Supplementary Table 9), all were negatively

associated with typhoid fever in both settings. Six of these specific MGCs were linked to SCFA

metabolism ('pyruvate2acetate.formate') and five to anaerobic metabolism ('Rnf complex'),

associated with *Prevotella* and *Haemophilus* (see Figure 2, Supplementary Table 9). None of

these 28 specific MGCs showed negative associations with acute typhoid fever in Nepal.

Table 2: Summary of the specific MGCs negatively associated with typhoid fever in both Bangladesh and Malawi, and the species of each MGC reference sequence. Full information in Supplementary Table 9.

MGC_class	n	Species
Pyruvate2acetate.formate	6	Haemophilus aegyptius, Haemophilus haemolyticus, Haemophilus parainfluenzae, Prevotella copri, Prevotella sp., Prevotella sp.
Rnf_complex	5	Haemophilus parainfluenzae, Prevotella copri, Prevotella sp., Prevotella sp., Prevotella sp.
TPP_AA_metabolism	4	Clostridium celatum, Haemophilus pittmaniae, Haemophilus sputorum, Prevotella sp.
Respiratory_glycerol	2	Aggregatibacter sp., Haemophilus parainfluenzae
Arginine2putrescine.Putrescine2spermidine	1	Romboutsia sp.
Formate_dehydrogenase	1	Haemophilus sp.
Fumarate2succinate	1	Haemophilus sp.
OD_eut_pdu_related.PFOR_II_pathway	1	Paeniclostridium sordellii
OD_fatty_acids	1	Dialister succinatiphilus
Others_HGD_unassigned	1	Prevotella copri
PFOR_II_pathway	1	Prevotella sp.
TPP_AA_metabolism.Arginine2putrescine	1	Haemophilus sp.
TPP_fatty_acids	1	Prevotella copri
porA	1	Prevotella copri
succinate2propionate	1	Dialister invisus

There were 31 MGC types that significantly differed between typhoid fever patients and
household contacts in the Malawi cohort, 8 in Bangladesh, and 1 in Nepal (Figure 3,
Supplementary Table 30). There was one MGC type that was negatively associated with
typhoid fever in both Malawi and Bangladesh, "TPP_AA_metabolism.Arginine2putrescine".
Other MGC types that were negatively associated with typhoid fever in Malawi only included
"acetate2butyrate.TPP_fatty_acids", "TPP_fatty_acids.aminobutyrate2Butyrate",
"acrylate2propionate" and "Rnf_complex.succinate2propionate", while in Bangladesh,
"Pyruvate2acetate" was negatively associated with typhoid fever. In Bangladesh,
"Others_HGD_unassigned.Nitrate_reductase" was positively associated with typhoid fever,
while in Malawi, two MGC types were positively associated with typhoid fever –
"Molybdopterin_dependent_oxidoreductase" and "pdu". In Nepal,
"Fumarate2succinate.fatty_acids" was significantly negatively associated with typhoid fever.

MGC	Cohort	N	Coefficient	Standard Error	Q-value	Effect Size
acetate2butyrate.TPP_fatty_acids	Malawi	46	7.57	1.02	0.0000035	
acetate2butyrate.TPP_fatty_acids	Bangladesh	80	-0.573	0.944	0.88	
acetate2butyrate.TPP_fatty_acids	Nepal	45	-1.08	0.427	0.61	
acetate2butyrate.TPP_fatty_acids	CHIM	26	0.0946	0.627	1	
TPP_AA_metabolism.Arginine2putrescine	Malawi	46	8.26	1.16	0.0000047	
TPP_AA_metabolism.Arginine2putrescine	Bangladesh	80	3.93	0.951	0.012	
TPP_AA_metabolism.Arginine2putrescine	Nepal	45	-0.624	1.33	1	
TPP_AA_metabolism.Arginine2putrescine	CHIM		NA	NA	NA	
TPP_fatty_acids.aminobutyrate2Butyrate	Malawi	46	5.9	1.15	0.00093	
TPP_fatty_acids.aminobutyrate2Butyrate	Bangladesh	80	0.148	0.849	0.97	
TPP_fatty_acids.aminobutyrate2Butyrate	Nepal	45	-0.575	0.884	1	
TPP_fatty_acids.aminobutyrate2Butyrate	CHIM		NA	NA	NA	
acrylate2propionate	Malawi	46	4.19	0.948	0.0052	
acrylate2propionate	Bangladesh	80	-2.63	0.963	0.18	
acrylate2propionate	Nepal	45	0.768	0.976	1	
acrylate2propionate	CHIM		NA	NA	NA	
pdu	Malawi	46	-2.81	0.664	0.0069	
pdu	Bangladesh	80	-1.24	0.633	0.35	
pdu	Nepal	45	2.07	0.478	0.054	
pdu	CHIM		NA	NA	NA	
Molybdopterin_dependent_oxidoreductase	Malawi	46	-2.27	0.616	0.021	
Molybdopterin_dependent_oxidoreductase	Bangladesh	80	-0.944	0.476	0.35	
Molybdopterin_dependent_oxidoreductase	Nepal	45	2.02	0.641	0.39	
Molybdopterin_dependent_oxidoreductase	CHIM	26	-0.302	0.41	0.93	
Rnf_complex.succinate2propionate	Malawi	46	2.73	0.804	0.039	
Rnf_complex.succinate2propionate	Bangladesh	80	-0.782	0.486	0.49	
Rnf_complex.succinate2propionate	Nepal	45	0.105	0.337	1	
Rnf_complex.succinate2propionate	CHIM	26	-0.314	0.259	0.93	
Pyruvate2acetate	Malawi	46	0.435	0.137	0.056	
Pyruvate2acetate	Bangladesh	80	0.677	0.174	0.017	
Pyruvate2acetate	Nepal	45	-0.173	0.152	0.99	
Pyruvate2acetate	CHIM	26	0.155	0.24	0.93	
Others_HGD_unassigned.Nitrate_reductase	Malawi	46	-2.61	1.13	0.2	
Others_HGD_unassigned.Nitrate_reductase	Bangladesh	80	-3.06	0.6	0.00076	
Others_HGD_unassigned.Nitrate_reductase	Nepal	45	3.54	0.87	0.067	
Others_HGD_unassigned.Nitrate_reductase	CHIM	26	-1.8	0.647	0.74	
Fumarate2succinate.fatty_acids	Malawi	46	-1.7	0.934	0.34	
Fumarate2succinate.fatty_acids	Bangladesh	80	2.03	0.776	0.18	
Fumarate2succinate.fatty_acids	Nepal	45	3.66	0.681	0.0039	
Fumarate2succinate.fatty_acids	CHIM		NA	NA	NA	

Effect Size: -6 -4 -2 0 2 4 6 8 10 12

Figure 3: Forest plot of selected MGC types that differed significantly between household contacts and typhoid fever participants in at least one cohort. If an entry is NA, then there was insufficient of that MGC type identified in that cohort for maaslin analysis.

We hypothesised the taxa and MGCs that were found in higher abundance in asymptomatic

household contacts of typhoid fever patients may be protective against developing disease upon

exposure to *S.* Typhi. We sought to explore this more directly by assessing their abundance in

CHIM participants who did or did not develop enteric fever upon challenge with typhoidal

Salmonella, in a UK-based study (Gibani et al. 2020). We sequenced the microbiomes of 13

participants who were subsequently challenged with *S.* Typhi and 13 with *S.* Paratyphi A, of

which 7 and 6 were diagnosed with enteric (typhoid or paratyphoid) fever, respectively

(Supplementary Table 18). There was no significant difference (FDR < 0.05) in alpha or beta

diversity at baseline among CHIM participants by age at challenge, sex, or subsequent enteric

fever diagnosis (Supplementary Table 19, Supplementary Table 20, Supplementary Figure 15).

In an analysis including only the 28 MGC classes that were identified as negatively associated

with typhoid in both Malawi and Bangladesh, two were also negatively associated with typhoid

fever diagnosis in CHIM participants: an "Rnf_complex.Glycine_cleavage.succinate2propionate"

MGC (coefficient = -3.4, q-value = 0.22) and an "Rnf_complex" MGC (coefficient = -6.2, q-value

= 0.22) (Supplementary Figure 16, full results available in Supplementary Table 23). None of

the 28 MGC classes tested were associated with paratyphoid fever or a combination of both

typhoid and paratyphoid fever. We investigated whether any of the species associated with

typhoid in both Malawi and Bangladesh were associated with susceptibility to typhoid and/or

paratyphoid fever in the CHIM cohort; none of them were (see Figure 2). All four species that

were associated with typhoid fever in the endemic cohorts were not detected in the majority of

samples from the CHIM (Supplementary Table 29).

374 Analysis of high-Vi participants

High-Vi participants may represent asymptomatic gall-bladder carriers, or individuals with recent

exposure or sub-clinical infection, either of whom may be a source in transmission. Detection of

carriers is important for typhoid fever control, and will remain so even as disease incidence

reduces due to vaccination and improvements in WASH, therefore we sought to explore the

microbiome signature of this group. Samples from Bangladesh were not included in this

analysis, as laboratory processing of the high Vi-titre participant samples from Bangladesh was

not consistent with the other sites or participant groups. Using PERMANOVA, participant group

was significantly associated with beta diversity in both countries (age and sex were not

significant, Supplementary Table 11, Supplementary Table 12). Participant group explained

greater variance in the Malawi cohort (R^2 0.25, FDR=0.01) than Nepal (R^2 0.07, FDR=0.01,

Supplementary Table 11, Supplementary Table 12,

Supplementary Figure 10). The three participant groups have distinct beta-diversity signatures,

with high-Vi individuals closer to typhoid patients than to household contacts (Figure 4). In the

Malawi cohort, 125 bacterial species significantly differed between household contacts and high

Vi-titre individuals (Supplementary Table 13), and 41 of these also significantly differed between

household contacts and typhoid fever (Supplementary Table 21). These 41 species include *H.*

parainfluenzae, Blautia obeum, and *Ruminococcus gnavus* (all less prevalent in acute typhoid

and high-Vi individuals than household contacts, Supplementary Figure 11). Amongst Nepal

samples, only one species significantly differed between household controls and high-Vi

individuals (the Firmicutes SGB GGB9790 SGB15413). There were no species associated with

household contacts compared with typhoid patients from Nepal, and SGB15413 did not differ

between these groups (Supplementary Figure 12).

The combined relative abundance of species only described as species genome bins (SGBs)

was significantly lower in household contacts in Malawi (13.9%), compared with both typhoid

fever patients (34.8%, P=0.012) and high Vi-titre participants (47.1%, P=2.2x10^{-16}) in this setting

(Supplementary Figure 13). The 50 SGBs with the highest summed abundance across all

participant types in Malawi belonged to Firmicutes (n = 41), Actinobacteria (n = 4),

Bacteroidetes (n = 3), and Proteobacteria (n = 2) (Supplementary Table 22). The most common

families of the 50 most abundant SGBs were Oscillospiraceae (n = 13), Lachnospiraceae (n =

7), and Clostridiaeceae (n = 4) (Supplementary Table 22). In Nepal, high Vi-titre individuals had

lower SGB relative abundance than acute typhoid cases (16.8% vs 29.9%, p = 0.03,

Supplementary Figure 13).

AMR genes

We recently reported AMR genotypes and phenotypes for *S.* Typhi isolated from acute typhoid fever cases in Bangladesh, Malawi, and Nepal (Dyson et al. 2024). Quinolone resistance was common in Bangladesh and Nepal but rare in Malawi; here we saw a similar pattern reflected in the gut microbiomes, with 2% quinolone resistance in Malawi, 30% in Bangladesh and 15% in Nepal (only mobile genetic element encoded quinolone determinants were analysed in the metagenomic data, Supplementary Table 14). Macrolide resistance was identified in one *S.* Typhi isolate from Bangladesh (0.6%) but not in Nepal or Malawi, here we found higher rates of macrolide resistance in microbiomes from individuals in Bangladesh (38%) and Nepal (35%) and lower in Malawi (11%). For the older drugs, *S.* Typhi isolates from Malawi had near-universal resistance to older sulfonamides and tetracyclines, and this was mirrored in high rates of resistance genes to these drugs in the microbiome samples (92% sulfonamides, 52% tetracycline). ESBLs were not detected in any *S.* Typhi isolates but were common in the microbiomes in Nepal (17%) and Bangladesh (9%); less so in Malawi (3%).

The proportion of samples with macrolide and trimethoprim resistance genes that have been observed in *S.* Typhi was significantly higher in participants with acute typhoid fever than in household contacts or high Vi-titre participants (macrolides = 42%, 14%, 22% respectively, Chi-square test p-value =$4.4x10^{-5}$; trimethoprim = 32%, 12%, 20% respectively, Chi-square test p-value = 0.005). There was no significant difference between participant types in the proportion of samples with resistance genes to any other drug classes. There was no statistically significant association between prior reported antibiotic usage and the number of AMR genes (Wilcoxon-rank sum P-value = 0.52) or the number of AMR gene classes (Wilcoxon Rank Sum P-value = 0.56) in typhoid fever patients across all sites (Supplementary Figure 14).

Discussion

In this study, we examined the relationship between the stool microbiome and typhoid fever in three settings with endemic disease. Our analyses identified four microbial species (Figure 2), linked to anaerobic fermentation and SCFA metabolism (Table 2), that differentiated typhoid fever patients from household contacts in Bangladesh and Malawi. While these taxa were not found in the UK-based CHIM participants, a different MGC with a potential link to SCFA metabolism was higher in participants who were not susceptible to typhoid fever. Looking at the abundance of different MGC types (as opposed to specific MGCs) we found four types of MGC involved in SCFA metabolism that were negatively associated with typhoid fever in Malawi, and one in Bangladesh. Overall, our data support a protective role for SCFA-producing microbes, although with our study design it is not possible to deconvolute the impact of the microbiome on typhoid from the impact of typhoid on the microbiome.

The strengths of our study include a larger sample size than previous investigations of the interaction between typhoid fever and the microbiome, and a multi-site study design to increase generalisability. In contrast to previous studies of the microbiome and typhoid fever, we used shotgun metagenomics to characterise the gut flora, enabling in depth functional analysis and investigation of AMR genes. Notably, using the same methodology to analyse participants from different countries, most of the taxa we associated with typhoid fever differed between locations. This suggests that susceptibility to disease in different human populations may be modified by different species, which could hamper the generalisability of study findings between populations. It is notable that the most consistent microbial signature for disease susceptibility, identified across multiple study populations with small sample sizes, was a functional signature (specifically gene clusters associated with SCFA metabolism) rather than any specific taxa that perform this function.

One of the key limitations of our study is the lack of age and sex matching between the typhoid fever and household contact groups. The age and sex composition of the household controls reflects the occupants of the household during the daytime visits of the study workers. This discrepancy may introduce bias, as age and sex are known to influence the composition and function of the microbiome (Yatsunenko et al. 2012). Although these variables were included in the statistical analyses, the lack of matching weakens our findings. Our use of a case-control study design in the endemic countries generated numerous hypotheses for subsequent

research, but it does not prove causal relationships. Additionally, we did not collect dietary

information from participants. Diet is a major determinant of microbiome composition and

function, and variations in dietary habits could confound the associations we observed

(Bourdeau-Julien et al. 2023). Lastly, accounting for the effect of antibiotics on the microbiome

in these cohorts presents a complex challenge, as it can significantly alter microbiome

composition and potentially mask or mimic associations with typhoid fever. These challenges

could largely be addressed using CHIM studies. Unfortunately, the only available CHIM data

was from a different study population (in the UK) and with a small sample size (n=13 exposed to

S. Typhi). Use of larger high-income country CHIMs and development of CHIMs in populations

from settings with endemic disease would be helpful to better elucidate specific microbial

signatures of protection.

Our use of shotgun metagenomics enabled us to also investigate the prevalence of AMR genes

associated with resistance in *S.* Typhi across the three sites and participant groups. Macrolide

and trimethoprim resistance genes were more common in acute typhoid cases compared with

the household controls and high Vi-titre participants, highlighting that typhoid and associated

antimicrobial usage imposes a selective pressure on the gut microbiome. This supports the idea

that reducing disease via immunisation could reduce AMR beyond just *S.* Typhi. Higher levels of

resistance to sulphonamides in Malawi likely reflects the common use of trimethoprim-

sulfamethoxazole in HIV/AIDS programs (MacPherson et al. 2022), while higher microbiome

prevalence of acquired ESBL and fluoroquinolone-resistance genes in Bangladesh than Malawi

reflects the local epidemiology (e.g. recent studies of blood stream infections show that in

Bangladesh, 72% and 75% were resistant to ciprofloxacin and third-generation cephalosporins

respectively, compared with 31% and 30% in Malawi) (Musicha et al. 2017; Ahmed et al. 2017).

The gut microbiome prevalence of resistance genes followed a similar pattern to the prevalence

of resistance genes in *S.* Typhi observed from these sites (Dyson et al. 2024) for tetracycline,

sulphonamides and quinolones, which likely reflects a shared evolutionary pressure on the

isolates and the microbiome. The fact that non-ESBL beta-lactamases were identified in fewer

metagenomes than ESBL beta-lactamases is because we only counted beta-lactamase

encoding genes that are commonly found in *S.* Typhi.

Zhang et al. investigated 16S community profile and metatranscriptome features associated

with susceptibility to typhoid fever in a CHIM (Zhang et al. 2018). One of their primary findings

was that the archaeal genus *Methanobrevibacter* was enriched in people who were challenged

with *S.* Typhi but did not develop disease. In contrast, we only identified one association with a

Methanobrevibacter species (*M. smithii*); a positive association with typhoid fever in Malawi.

Furthermore, Zhang et al., identified that *Prevotella* was higher in people who developed typhoid

fever, in contrast to our findings that it was negatively associated with typhoid fever in natural

infections in endemic settings. One source of these discrepancies could be that Zhang et al.

employed 16S, while we used shotgun metagenomics. Haak et al. observed more SCFA

producing bacteria in, in both healthy control and febrile but non-typhoid fever participants in

their study in Bangladesh compared with typhoid fever patients, and higher faecal load of

SCFAs in controls than any febrile group (Haak et al. 2020), which corresponds well with our

findings. It has also been shown that higher SCFAs in an acidic environment, and a strict

anaerobic environment, can inhibit the growth of Enterobacteriaceae such as *Salmonella* via

both direct and indirect mechanisms (Byndloss et al. 2017; Jacobson et al. 2018; Rogers,

Tsolis, and Bäumler 2021). Recently, higher relative abundance of butyrate producing bacteria

was associated with a reduced risk of hospitalisation for infections in a prospective study carried

out in high income European countries (Kullberg et al. 2024). Two MGC types positively

associated with typhoid fever in Malawi were molybdopterin dependent oxidoreductase and

propanediol utilisation microcompartments, both of which enable growth in the inflamed gut

(Faber et al. 2017; Zhu et al. 2018), suggesting that the microbiota of typhoid fever participants

in Malawi may be in a state of dysbiosis. Arginine to putrescine was the only MGC type

associated with health in both Bangaldesh and Malawi, polyamines such as putrescine promote

gut barrier integrity (Rao, Xiao, and Wang 2020), and can act as immunomodulators (Proietti et

al. 2020).

Our findings from the STRATAA cohorts can be interpreted from two perspectives: i) the

microbiota protecting against typhoid and, ii) typhoid causing microbiota changes. In contrast,

the CHIM study design enables us to specifically assess whether microbiota present prior to

pathogen exposure are associated with the outcome of that exposure (i.e. developing disease

or not). Unfortunately, none of the taxa negatively associated with typhoid fever in the endemic

settings were present in most of the CHIM participants, making it impossible to validate the

association. It is well established that microbiome species composition differs between human

populations, and particularly between those in high-income countries and low-and-middle-

income countries (Gupta, Paul, and Dutta 2017). However, the metabolic gene cluster findings

identified a recurring finding across the endemic settings and the CHIM; an association between

species encoding SCFA metabolising genes and non-susceptibility to typhoid fever. While it

should be noted that the CHIM sample size was small, the fact that we identified a statistically

significant difference supports the idea that the effect is quite strong. While our CHIM analyses

lacked power, which is reflected in the high q-values obtained, the q-value for the SCFA MGC

association was below the default maaslin2 threshold for significant findings.

Both *P. copri* and *H. parainfluenzae,* which were significantly lower in typhoid fever patients

compared with household contacts in both Bangladesh and Malawi, are associated with

increased gut inflammation (Sohn et al. 2023; Larsen 2017). *P. copri* is thought to increase

Th17 inflammation (Godinez et al. 2011) while *H. parainfluenzae* stimulates intestinal (IFN-γ)+

CD4+ T cells (Sohn et al. 2023), two mechanisms which play a key role in the response to

Salmonella infections (Bao et al. 2000; Raffatellu et al. 2008). Increased inflammation could

prime the host to respond more rapidly to pathogenic exposures, leading to enhanced control of

infections. It is highly plausible that the gut microbiome plays a role in shaping the response of

the immune system to pathogen challenge (Durack and Lynch 2019). This intriguing association

needs further investigation, as the role of inflammation in enabling *Salmonella* to overcome

colonisation resistance in mice is well established (Winter et al. 2010; Rogers, Tsolis, and

Bäumler 2021). Among the species negatively associated with typhoid fever in Malawi only was

Ruminococcus gnavus, which could influence susceptibility to enteric infection via IgA

stimulation (Bunker et al. 2019), protects against enteropathogenic *E. coli* (McGrath et al. 2022),

stimulates host tryptophan catabolism (Hoffmann et al. 2016) (*S.* Typhi requires tryptophan to

grow in macrophages (Blohmke et al. 2016)), and produces secondary bile acids including

chenodeoxycholic acid and iso-LCA that have anti-virulence effects on *Salmonella* (Lee et al.

2013; Yang, Stein, and Hang 2023).

Direction of causality cannot be determined from our study design, and a murine model of

Salmonella Typhimurium infection recently reported shifts in gut microbiome composition,

including a reduction in *Ruminococcaceae* taxa associated with acetate and butyrate

production, following infection (Rogers et al. 2024). It is therefore plausible that the microbiome

is modified in people suffering from typhoid fever. Gut disruption is known to alter the gut

microbiome, for example, colorectal cancer patients from Morocco and Kenya had reduced *P.*

copri in their gut, (Allali et al. 2018; Obuya et al. 2022). Potential triggers for microbiome

alterations in typhoid fever could include antibiotic exposure, dietary changes or anorexia due to

sickness, *S.* Typhi-induced malabsorption in the small intestine resulting in alterations to the

nutrient composition within the large intestine, and the direct impact of fever on microbiome

composition as the increase in temperature could favour the growth of particular bacterial

species (Chuttani, Jain, and Misra 1971; Huus and Ley 2021).

We cannot be certain about the exact cause of higher levels of anti-Vi antibodies in the high Vi-

titre participants. Potential sources of immune stimulation include recent asymptomatic *S.* Typhi

exposure/infection, exposure/infection with *Citrobacter freundii* which can be Vi-antigen positive,

or chronic *S.* Typhi carriage. It is striking that there was a clear divergence in microbiota profile

between these participants and acute cases and household contacts in Malawi, as this suggests

that there is an interaction between the cause of the high Vi-titre and the gut microbiota. The 41

species that were significantly lower in both typhoid fever cases and high Vi-titre participants

included the potentially immune-modulatory *H. parainfluenzae* (Sohn et al. 2023), the bile acid

modifying *B. obeum* (formerly *Ruminococcus obeum*), and *R. gnavus*. Notably, it has been

demonstrated that bile salt hydrolases encoded by *B. obeum* can inhibit *Vibrio cholerae*

virulence gene activation and colonization (Alavi et al. 2020), and it is intriguing to hypothesise a

similar mechanism might protect against *S.* Typhi gallbladder carriage and/or systemic infection.

The influence of these species on susceptibility to *S.* Typhi infection, gallbladder carriage, and

immune-reactivity should be investigated further.

589 Conclusions

We found that typhoid fever patients in Malawi and Bangladesh display distinct microbiome

signatures compared to household contacts, marked by lower abundance of SCFA producers

and inflammation-related species. The negative association between SCFA producing genes

and typhoid fever susceptibility was also replicated to some extent in our UK-based challenge

study. The differences in species composition in the microbiota of these distinct cohorts make

direct comparison and therefore validation difficult, highlighting the importance of establishing

challenge studies in endemic settings to directly address important mechanistic questions.

Data availability

The code used in the analysis of these data is available on github -
https://github.com/flashton2003/STRATAA_metagenomics. Sequencing data for this project are
available at the European Nucleotide Archive, project accessions PRJEB14050 and
PRJEB22175.

Acknowledgements

We thank all the STRATAA participants for their willingness to take part in this research. We
also acknowledge the sequencing and pathogen informatics teams at the Wellcome Sanger
Institute for sequencing and data processing, and Zoe Dyson for assisting with sequence data
transfer. We would like to acknowledge Ndaru Jambo, Andreas Bäumler, and Renee Tsolis for
helpful discussions. KA, SK, NRT, OS were funded by the Wellcome Trust (220540/Z/20/A). For
the purpose of open access, the authors have applied a CC-BY public copyright licence to any
author-accepted manuscript version arising from this submission. This research was funded by
the Wellcome Trust (STRATAA grant number 106158/Z/14/Z, and core funding to the Wellcome
Sanger Institute, grant number 206194) and the Bill & Melinda Gates Foundation (grant number
OPP1141321). MAG, ACC, and PMA were supported by a research professorship
(NIHR300039) from the National Institute for Health and Care Research, UK Department of
Health and Social Care. GD is supported by the Health@InnoHK, Innovation Technology
Commission Funding. MMG is supported by the NIHR Imperial Biomedical Research Centre
and the Wellcome Trust (224029/Z/21/Z).

Members of the STRATAA study group are Christoph Blohmke, Yama Farooq, Jennifer Hill,
Nhu Tran Hoang, Tikhala Makhaza Jere, Harrison Msuku, Tran Vu Thieu Nga, Rose Nkhata,
Sadia Isfat Ara Rahman, Nazia Rahman, Neil J Saad, Trinh Van Tan, Deus Thindwa, Merryn
Voysey, and Richard Wachepa. Their affiliations are: Malawi-Liverpool-Wellcome Programme,
Blantyre, Malawi (Tikhala Makhaza Jere, Harrison Msuku, Rose Nkhata, Deus Thindwa,
Richard Wachepa); International Centre for Diarrhoeal Disease Research, Dhaka, Bangladesh
(Sadia Isfat Ara Rahman, Nazia Rahman); Oxford Vaccine Group, Department of Paediatrics,
University of Oxford, and the NIHR Oxford Biomedical Research Centre, Oxford, UK (Christoph
Blohmke, Yama Farooq, Jennifer Hill, Merryn Voysey); The Hospital for Tropical Diseases,
Wellcome Trust Major Overseas Programme, Oxford University Clinical Research Unit, Ho Chi

628 Minh City, Viet Nam (Nhu Tran Hoang, Tran Vu Thieu Nga, Trinh Van Tan); Department of
629 Epidemiology of Microbial Diseases and the Public Health Modeling Unit, Yale School of Public
630 Health, Yale University, New Haven, CT, USA (Neil J Saad)
631
632
633

634 Author contributions

Philip M. Ashton – conceptualisation, data curation, formal analysis, investigation, methodology,
resources, software, visualisation, writing – original draft preparation.
Leonardos Mageiros - conceptualisation, data curation, formal analysis, investigation,
methodology, software, writing – review and editing.
James Meiring - data curation, investigation, methodology, writing – review and editing.
Angeziwa Chunga Chirambo – data curation, investigation, writing – review and editing.
Farhana Khanam – data curation, investigation, writing – review and editing.
Happy Banda – investigation, writing – review and editing.
Abhilasha Karkey – investigation, supervision, writing – review and editing.
Sabina Dongol – data curation, investigation, writing – review and editing.
Lorena Preciado Llanes – conceptualisation, data curation, investigation, methodology, writing –
review and editing.
Helena Thomaides-Brears - investigation, methodology, writing – review and editing.
Malick Gibani – data curation, investigation, methodology, writing – review and editing.
Nazmul Hasan Rajib – investigation, writing – review and editing.
Nazia Rahman – investigation, writing – review and editing.
Prasanta Kumar Biswas – investigation, writing – review and editing.
Md Amirul Islam Bhuiyan – investigation, writing – review and editing.
Andrew Pollard – conceptualisation, funding acquisition, supervision, writing – review and
editing.
Stephen Baker – funding acquisition, writing – review and editing.
Buddha Basnyat – funding acquisition, supervision, resources, writing – review and editing.
John D Clemens – funding acquisition, supervision, writing – review and editing.
Christiane Dolecek – funding acquisition, writing – review and editing.
Sarah J Dunstan – funding acquisition, writing – review and editing.

Gordon Dougan – conceptualisation, funding acquisition, methods, resources, writing – review
and editing.

Robert S. Heyderman – funding acquisition, writing – review and editing.

Virginia E. Pitzer – funding acquisition, writing – review and editing.

Firdausi Qadri – funding acquisition, supervision, resources, writing – review and editing.

Melita A. Gordon – conceptualisation, funding acquisition, resources, supervision, writing –
review and editing.

Thomas C. Darton – conceptualisation, methodology, writing – review and editing.

Kathryn E. Holt – conceptualisation, funding acquisition, resources, methodology, supervision,
writing – original draft preparation, writing – review and editing.

671 References

Ahmed, Dilruba, Md Ausrafuggaman Nahid, Abdullah Bashar Sami, Farhana Halim, Nasrin
Akter, Tuhin Sadique, Md Sohel Rana, Md Shahriar Bin Elahi, and Md Mahbubur
Rahman. 2017. 'Bacterial Etiology of Bloodstream Infections and Antimicrobial
Resistance in Dhaka, Bangladesh, 2005–2014'. *Antimicrobial Resistance & Infection
Control* 6 (1): 2. https://doi.org/10.1186/s13756-016-0162-z.
Alavi, Salma, Jonathan D. Mitchell, Jennifer Y. Cho, Rui Liu, John C. Macbeth, and Ansel Hsiao.
2020. 'Interpersonal Gut Microbiome Variation Drives Susceptibility and Resistance to
Cholera Infection'. *Cell* 181 (7): 1533-1546.e13.
https://doi.org/10.1016/j.cell.2020.05.036.
Alcock, Brian P., William Huynh, Romeo Chalil, Keaton W. Smith, Amogelang R. Raphenya,
Mateusz A. Wlodarski, Arman Edalatmand, et al. 2023. 'CARD 2023: Expanded
Curation, Support for Machine Learning, and Resistome Prediction at the
Comprehensive Antibiotic Resistance Database'. *Nucleic Acids Research* 51 (D1):
D690–99. https://doi.org/10.1093/nar/gkac920.
Allali, Imane, Noureddine Boukhatem, Leila Bouguenouch, Hanaa Hardi, H. Abir Boudouaya, M.
Belen Cadenas, Karim Ouldim, Saaïd Amzazi, M. Andrea Azcarate-Peril, and Hassan
Ghazal. 2018. 'Gut Microbiome of Moroccan Colorectal Cancer Patients'. *Medical
Microbiology and Immunology* 207 (3): 211–25. https://doi.org/10.1007/s00430-018-
0542-5.
Argimón, Silvia, Corin A. Yeats, Richard J. Goater, Khalil Abudahab, Benjamin Taylor, Anthony
Underwood, Leonor Sánchez-Busó, et al. 2021. 'A Global Resource for Genomic
Predictions of Antimicrobial Resistance and Surveillance of Salmonella Typhi at
Pathogenwatch'. *Nature Communications* 12 (1): 2879. https://doi.org/10.1038/s41467-
021-23091-2.
Bao, S, K W Beagley, M P France, J Shen, and A J Husband. 2000. 'Interferon-γ Plays a Critical
Role in Intestinal Immunity against Salmonella Typhimurium Infection'. *Immunology* 99
(3): 464–72. https://doi.org/10.1046/j.1365-2567.2000.00955.x.
Belkaid, Yasmine, and Timothy Hand. 2014. 'Role of the Microbiota in Immunity and
Inflammation'. *Cell* 157 (1): 121–41. https://doi.org/10.1016/j.cell.2014.03.011.

Blanco-Míguez, Aitor, Francesco Beghini, Fabio Cumbo, Lauren J. McIver, Kelsey N. Thompson, Moreno Zolfo, Paolo Manghi, et al. 2023. 'Extending and Improving Metagenomic Taxonomic Profiling with Uncharacterized Species Using MetaPhlAn 4'. *Nature Biotechnology* 41 (11): 1633–44. https://doi.org/10.1038/s41587-023-01688-w.

Blohmke, Christoph J., Thomas C. Darton, Claire Jones, Nicolas M. Suarez, Claire S. Waddington, Brian Angus, Liqing Zhou, et al. 2016. 'Interferon-Driven Alterations of the Host's Amino Acid Metabolism in the Pathogenesis of Typhoid Fever'. *Journal of Experimental Medicine* 213 (6): 1061–77. https://doi.org/10.1084/jem.20151025.

Bohnhoff, Marjorie, Barbara L. Drake, and C. Phillip Miller. 1954. 'Effect of Streptomycin on Susceptibility of Intestinal Tract to Experimental Salmonella Infection.' *Proceedings of the Society for Experimental Biology and Medicine* 86 (1): 132–37. https://doi.org/10.3181/00379727-86-21030.

Bourdeau-Julien, Isabelle, Sophie Castonguay-Paradis, Gabrielle Rochefort, Julie Perron, Benoît Lamarche, Nicolas Flamand, Vincenzo Di Marzo, Alain Veilleux, and Frédéric Raymond. 2023. 'The Diet Rapidly and Differentially Affects the Gut Microbiota and Host Lipid Mediators in a Healthy Population'. *Microbiome* 11 (1): 26. https://doi.org/10.1186/s40168-023-01469-2.

Bunker, Jeffrey J., Christoph Drees, Andrea R. Watson, Catherine H. Plunkett, Cathryn R. Nagler, Olaf Schneewind, A. Murat Eren, and Albert Bendelac. 2019. 'B Cell Superantigens in the Human Intestinal Microbiota'. *Science Translational Medicine* 11 (507): eaau9356. https://doi.org/10.1126/scitranslmed.aau9356.

Byndloss, Mariana X., Erin E. Olsan, Fabian Rivera-Chávez, Connor R. Tiffany, Stephanie A. Cevallos, Kristen L. Lokken, Teresa P. Torres, et al. 2017. 'Microbiota-Activated PPAR-γ Signaling Inhibits Dysbiotic Enterobacteriaceae Expansion'. *Science* 357 (6351): 570–75. https://doi.org/10.1126/science.aam9949.

Chuttani, H. K., K. Jain, and R. C. Misra. 1971. 'Small Bowel in Typhoid Fever'. *Gut* 12 (9): 709–12.

Constantinides, Bede, Martin Hunt, and Derrick W Crook. 2023. 'Hostile: Accurate Decontamination of Microbial Host Sequences'. *Bioinformatics* 39 (12): btad728. https://doi.org/10.1093/bioinformatics/btad728.

Darton, Thomas C, James E Meiring, Susan Tonks, Md Arifuzzaman Khan, Farhana Khanam, Mila Shakya, Deus Thindwa, et al. 2017. 'The STRATAA Study Protocol: A Programme to Assess the Burden of Enteric Fever in Bangladesh, Malawi and Nepal Using Prospective Population Census, Passive Surveillance, Serological Studies and Healthcare Utilisation Surveys'. *BMJ Open* 7 (6): e016283. https://doi.org/10.1136/bmjopen-2017-016283.

Durack, Juliana, and Susan V. Lynch. 2019. 'The Gut Microbiome: Relationships with Disease and Opportunities for Therapy'. *The Journal of Experimental Medicine* 216 (1): 20–40. https://doi.org/10.1084/jem.20180448.

Dyson, Zoe A., Philip M. Ashton, Farhana Khanam, Angeziwa Chunga Chirambo, Mila Shakya, James E. Meiring, Susan Tonks, et al. 2024. 'Pathogen Diversity and Antimicrobial Resistance Transmission of Salmonella Enterica Serovars Typhi and Paratyphi A in Bangladesh, Nepal, and Malawi: A Genomic Epidemiological Study'. *The Lancet Microbe* 0 (0). https://doi.org/10.1016/S2666-5247(24)00047-8.

Faber, Franziska, Parameth Thiennimitr, Luisella Spiga, Mariana X. Byndloss, Yael Litvak, Sara Lawhon, Helene L. Andrews-Polymenis, Sebastian E. Winter, and Andreas J. Bäumler. 2017. 'Respiration of Microbiota-Derived 1,2-Propanediol Drives Salmonella Expansion during Colitis'. *PLoS Pathogens* 13 (1): e1006129. https://doi.org/10.1371/journal.ppat.1006129.

Gibani, Malick M., Celina Jin, Sonu Shrestha, Maria Moore, Lily Norman, Merryn Voysey, Elizabeth Jones, et al. 2020. 'Homologous and Heterologous Re-Challenge with

Salmonella Typhi and Salmonella Paratyphi A in a Randomised Controlled Human Infection Model'. *PLOS Neglected Tropical Diseases* 14 (10): e0008783. https://doi.org/10.1371/journal.pntd.0008783.

Godinez, Ivan, A. Marijke Keestra, Alanna Spees, and Andreas J. Bäumler. 2011. 'The IL-23 Axis in Salmonella Gastroenteritis'. *Cellular Microbiology* 13 (11): 1639–47. https://doi.org/10.1111/j.1462-5822.2011.01637.x.

Gupta, Vinod K., Sandip Paul, and Chitra Dutta. 2017. 'Geography, Ethnicity or Subsistence-Specific Variations in Human Microbiome Composition and Diversity'. *Frontiers in Microbiology* 8 (June):1162. https://doi.org/10.3389/fmicb.2017.01162.

Haak, Bastiaan W, Hanna K de Jong, Sarantos Kostidis, Martin Giera, Rapeephan R Maude, Rasheda Samad, Lalith Wijedoru, et al. 2020. 'Altered Patterns of Compositional and Functional Disruption of the Gut Microbiota in Typhoid Fever and Nontyphoidal Febrile Illness'. *Open Forum Infectious Diseases* 7 (7). https://doi.org/10.1093/ofid/ofaa251.

Hoffmann, Thomas W., Hang-Phuong Pham, Chantal Bridonneau, Camille Aubry, Bruno Lamas, Camille Martin-Gallausiaux, Marco Moroldo, et al. 2016. 'Microorganisms Linked to Inflammatory Bowel Disease-Associated Dysbiosis Differentially Impact Host Physiology in Gnotobiotic Mice'. *The ISME Journal* 10 (2): 460–77. https://doi.org/10.1038/ismej.2015.127.

Huus, Kelsey E., and Ruth E. Ley. 2021. 'Blowing Hot and Cold: Body Temperature and the Microbiome'. *mSystems* 6 (5): 10.1128/msystems.00707-21. https://doi.org/10.1128/msystems.00707-21.

Jacobson, Amanda, Lilian Lam, Manohary Rajendram, Fiona Tamburini, Jared Honeycutt, Trung Pham, Will Van Treuren, et al. 2018. 'A Gut Commensal-Produced Metabolite Mediates Colonization Resistance to Salmonella Infection'. *Cell Host & Microbe* 24 (2): 296-307.e7. https://doi.org/10.1016/j.chom.2018.07.002.

Khanam, Farhana, Thomas C. Darton, James E. Meiring, Protup Kumer Sarker, Prasanta Kumar Biswas, Md Amirul Islam Bhuiyan, Nazmul Hasan Rajib, et al. 2021. 'Salmonella Typhi Stool Shedding by Patients With Enteric Fever and Asymptomatic Chronic Carriers in an Endemic Urban Setting'. *The Journal of Infectious Diseases* 224 (12 Suppl 2): S759–63. https://doi.org/10.1093/infdis/jiab476.

Kullberg, Robert F J, Irina Wikki, Bastiaan W Haak, Anni Kauko, Henrike Galenkamp, Hessel Peters-Sengers, Joe M Butler, et al. 2024. 'Association between Butyrate-Producing Gut Bacteria and the Risk of Infectious Disease Hospitalisation: Results from Two Observational, Population-Based Microbiome Studies'. *The Lancet Microbe* 5 (9): 100864. https://doi.org/10.1016/S2666-5247(24)00079-X.

Larsen, Jeppe Madura. 2017. 'The Immune Response to Prevotella Bacteria in Chronic Inflammatory Disease'. *Immunology* 151 (4): 363–74. https://doi.org/10.1111/imm.12760.

Lee, Ja-Young, Hisashi Arai, Yusuke Nakamura, Satoru Fukiya, Masaru Wada, and Atsushi Yokota. 2013. 'Contribution of the 7β-Hydroxysteroid Dehydrogenase from Ruminococcus Gnavus N53 to Ursodeoxycholic Acid Formation in the Human Colon'. *Journal of Lipid Research* 54 (11): 3062–69. https://doi.org/10.1194/jlr.M039834.

Leshem, Avner, Timur Liwinski, and Eran Elinav. 2020. 'Immune-Microbiota Interplay and Colonization Resistance in Infection'. *Molecular Cell* 78 (4): 597–613. https://doi.org/10.1016/j.molcel.2020.03.001.

Libertucci, Josie, and Vincent B. Young. 2019. 'The Role of the Microbiota in Infectious Diseases'. *Nature Microbiology* 4 (1): 35–45. https://doi.org/10.1038/s41564-018-0278-4.

Lopez, Christopher A., Brittany M. Miller, Fabian Rivera-Chávez, Eric M. Velazquez, Mariana X. Byndloss, Alfredo Chávez-Arroyo, Kristen L. Lokken, Renée M. Tsolis, Sebastian E. Winter, and Andreas J. Bäumler. 2016. 'Virulence Factors Enhance Citrobacter Rodentium Expansion through Aerobic Respiration'. *Science (New York, N.Y.)* 353 (6305): 1249–53. https://doi.org/10.1126/science.aag3042.

MacPherson, Eleanor E., Joanna Reynolds, Esnart Sanudi, Alexander Nkaombe, John
Mankhomwa, Justin Dixon, and Clare I. R. Chandler. 2022. 'Understanding Antimicrobial
Use in Subsistence Farmers in Chikwawa District Malawi, Implications for Public
Awareness Campaigns'. *PLOS Global Public Health* 2 (6): e0000314.
https://doi.org/10.1371/journal.pgph.0000314.
Mallick, Himel, Ali Rahnavard, Lauren J. McIver, Siyuan Ma, Yancong Zhang, Long H. Nguyen,
Timothy L. Tickle, et al. 2021. 'Multivariable Association Discovery in Population-Scale
Meta-Omics Studies'. *PLOS Computational Biology* 17 (11): e1009442.
https://doi.org/10.1371/journal.pcbi.1009442.
McGrath, Conor J., Edgaras Laveckis, Andrew Bell, Emmanuelle Crost, Nathalie Juge, and
Stephanie Schüller. 2022. 'Development of a Novel Human Intestinal Model to Elucidate
the Effect of Anaerobic Commensals on Escherichia Coli Infection'. *Disease Models &*
Mechanisms 15 (4): dmm049365. https://doi.org/10.1242/dmm.049365.
Meiring, James E., Farhana Khanam, Buddha Basnyat, Richelle C. Charles, John A. Crump,
Frederic Debellut, Kathryn E. Holt, et al. 2023. 'Typhoid Fever'. *Nature Reviews Disease*
Primers 9 (1): 1–18. https://doi.org/10.1038/s41572-023-00480-z.
Meiring, James E., Mila Shakya, Farhana Khanam, Merryn Voysey, Maile T. Phillips, Susan
Tonks, Deus Thindwa, et al. 2021. 'Burden of Enteric Fever at Three Urban Sites in
Africa and Asia: A Multicentre Population-Based Study'. *The Lancet Global Health* 9
(12): e1688–96. https://doi.org/10.1016/S2214-109X(21)00370-3.
Musicha, Patrick, Jennifer E Cornick, Naor Bar-Zeev, Neil French, Clemens Masesa, Brigitte
Denis, Neil Kennedy, et al. 2017. 'Trends in Antimicrobial Resistance in Bloodstream
Infection Isolates at a Large Urban Hospital in Malawi (1998–2016): A Surveillance
Study'. *The Lancet Infectious Diseases* 17 (10): 1042–52. https://doi.org/10.1016/S1473-
3099(17)30394-8.
Obuya, Sarah, Amr Elkholy, Nagavardhini Avuthu, Michael Behring, Prachi Bajpai, Sumit
Agarwal, Hyung-Gyoon Kim, et al. 2022. 'A Signature of Prevotella Copri and
Faecalibacterium Prausnitzii Depletion, and a Link with Bacterial Glutamate Degradation
in the Kenyan Colorectal Cancer Patients'. *Journal of Gastrointestinal Oncology* 13 (5):
2282–92. https://doi.org/10.21037/jgo-22-116.
Pascal Andreu, Victòria, Hannah E. Augustijn, Koen van den Berg, Justin J. J. van der Hooft,
Michael A. Fischbach, and Marnix H. Medema. 2021. 'BiG-MAP: An Automated Pipeline
To Profile Metabolic Gene Cluster Abundance and Expression in Microbiomes'.
mSystems 6 (5): e00937-21. https://doi.org/10.1128/mSystems.00937-21.
Pascal Andreu, Victòria, Hannah E. Augustijn, Lianmin Chen, Alexandra Zhernakova, Jingyuan
Fu, Michael A. Fischbach, Dylan Dodd, and Marnix H. Medema. 2023. 'gutSMASH
Predicts Specialized Primary Metabolic Pathways from the Human Gut Microbiota'.
Nature Biotechnology 41 (10): 1416–23. https://doi.org/10.1038/s41587-023-01675-1.
Patel, Priyanka D., Pratiksha Patel, Yuanyuan Liang, James E. Meiring, Theresa Misiri, Felistas
Mwakiseghile, J. Kathleen Tracy, et al. 2021. 'Safety and Efficacy of a Typhoid
Conjugate Vaccine in Malawian Children'. *New England Journal of Medicine* 385 (12):
1104–15. https://doi.org/10.1056/NEJMoa2035916.
Proietti, Elisa, Sofia Rossini, Ursula Grohmann, and Giada Mondanelli. 2020. 'Polyamines and
Kynurenines at the Intersection of Immune Modulation'. *Trends in Immunology* 41 (11):
1037–50. https://doi.org/10.1016/j.it.2020.09.007.
Qureshi, Sonia, Noshi Maria, Tabish Chawla, Junaid Iqbal, Abdul Momin Kazi, Mehreen Adnan,
Aneeta Hotwani, et al. 2024. 'The Frequency and Associated Factors of Typhoid
Carriage in Patients Undergoing Cholecystectomy for Gallbladder Disease in Pakistan: A
Cross-Sectional Study'. *PLoS Neglected Tropical Diseases* 18 (6): e0011775.
https://doi.org/10.1371/journal.pntd.0011775.

Raffatellu, Manuela, Renato L. Santos, David E. Verhoeven, Michael D. George, R. Paul
Wilson, Sebastian E. Winter, Ivan Godinez, et al. 2008. 'Simian Immunodeficiency
Virus–Induced Mucosal Interleukin-17 Deficiency Promotes Salmonella Dissemination
from the Gut'. *Nature Medicine* 14 (4): 421–28. https://doi.org/10.1038/nm1743.
Rao, Jaladanki N., Lan Xiao, and Jian-Ying Wang. 2020. 'Polyamines in Gut Epithelial Renewal
and Barrier Function'. *Physiology* 35 (5): 328–37.
https://doi.org/10.1152/physiol.00011.2020.
Robbins, J. D., and J. B. Robbins. 1984. 'Reexamination of the Protective Role of the Capsular
Polysaccharide (Vi Antigen) of Salmonella Typhi'. *The Journal of Infectious Diseases*
150 (3): 436–49. https://doi.org/10.1093/infdis/150.3.436.
Rogers, Andrew W. L., Lauren C. Radlinski, Henry Nguyen, Connor R. Tiffany, Thaynara
Parente Carvalho, Hugo L. P. Masson, Michael L. Goodson, et al. 2024. '*Salmonella* Re-
Engineers the Intestinal Environment to Break Colonization Resistance in the Presence
of a Compositionally Intact Microbiota'. *Cell Host & Microbe* 32 (10): 1774-1786.e9.
https://doi.org/10.1016/j.chom.2024.07.025.
Rogers, Andrew W. L., Renée M. Tsolis, and Andreas J. Bäumler. 2021. 'Salmonella versus the
Microbiome'. *Microbiology and Molecular Biology Reviews: MMBR* 85 (1): e00027-19.
https://doi.org/10.1128/MMBR.00027-19.
Shreiner, Andrew B., John Y. Kao, and Vincent B. Young. 2015. 'The Gut Microbiome in Health
and in Disease'. *Current Opinion in Gastroenterology* 31 (1): 69–75.
https://doi.org/10.1097/MOG.0000000000000139.
Sohn, Jiho, Lu Li, Lixia Zhang, Robert J. Genco, Karen L. Falkner, Hervé Tettelin, Aryn M.
Rowsam, et al. 2023. 'Periodontal Disease Is Associated with Increased Gut
Colonization of Pathogenic Haemophilusparainfluenzae in Patients with Crohn's
Disease'. *Cell Reports* 42 (2): 112120. https://doi.org/10.1016/j.celrep.2023.112120.
Stanaway, Jeffrey D., Robert C. Reiner, Brigette F. Blacker, Ellen M. Goldberg, Ibrahim A.
Khalil, Christopher E. Troeger, Jason R. Andrews, et al. 2019. 'The Global Burden of
Typhoid and Paratyphoid Fevers: A Systematic Analysis for the Global Burden of
Disease Study 2017'. *The Lancet Infectious Diseases* 19 (4): 369–81.
https://doi.org/10.1016/S1473-3099(18)30685-6.
Stecher, Bärbel, Riccardo Robbiani, Alan W. Walker, Astrid M. Westendorf, Manja Barthel,
Marcus Kremer, Samuel Chaffron, et al. 2007. 'Salmonella Enterica Serovar
Typhimurium Exploits Inflammation to Compete with the Intestinal Microbiota'. *PLOS*
Biology 5 (10): e244. https://doi.org/10.1371/journal.pbio.0050244.
Steele, A. Duncan, Deborah C. Hay Burgess, Zoey Diaz, Megan E. Carey, and Anita K. M.
Zaidi. 2016. 'Challenges and Opportunities for Typhoid Fever Control: A Call for
Coordinated Action'. *Clinical Infectious Diseases: An Official Publication of the Infectious*
Diseases Society of America 62 (Suppl 1): S4–8. https://doi.org/10.1093/cid/civ976.
Waddington, Claire S., Thomas C. Darton, Claire Jones, Kathryn Haworth, Anna Peters, Tessa
John, Ben A. V. Thompson, et al. 2014. 'An Outpatient, Ambulant-Design, Controlled
Human Infection Model Using Escalating Doses of Salmonella Typhi Challenge
Delivered in Sodium Bicarbonate Solution'. *Clinical Infectious Diseases* 58 (9): 1230–40.
https://doi.org/10.1093/cid/ciu078.
Winter, Sebastian E., Parameth Thiennimitr, Maria G. Winter, Brian P. Butler, Douglas L.
Huseby, Robert W. Crawford, Joseph M. Russell, et al. 2010. 'Gut Inflammation
Provides a Respiratory Electron Acceptor for Salmonella'. *Nature* 467 (7314): 426–29.
https://doi.org/10.1038/nature09415.
Yang, Xinglin, Kathryn R. Stein, and Howard C. Hang. 2023. 'Anti-Infective Bile Acids Bind and
Inactivate a Salmonella Virulence Regulator'. *Nature Chemical Biology* 19 (1): 91–100.
https://doi.org/10.1038/s41589-022-01122-3.

Yatsunenko, Tanya, Federico E. Rey, Mark J. Manary, Indi Trehan, Maria Gloria Dominguez-Bello, Monica Contreras, Magda Magris, et al. 2012. 'Human Gut Microbiome Viewed across Age and Geography'. *Nature* 486 (7402): 222–27. https://doi.org/10.1038/nature11053.

Zhang, Yan, Arthur Brady, Cheron Jones, Yang Song, Thomas C. Darton, Claire Jones, Christoph J. Blohmke, et al. 2018. 'Compositional and Functional Differences in the Human Gut Microbiome Correlate with Clinical Outcome Following Infection with Wild-Type Salmonella Enterica Serovar Typhi'. *mBio* 9 (3): e00686-18. https://doi.org/10.1128/mBio.00686-18.

Zhu, Wenhan, Maria G. Winter, Mariana X. Byndloss, Luisella Spiga, Breck A. Duerkop, Elizabeth R. Hughes, Lisa Büttner, et al. 2018. 'Precision Editing of the Gut Microbiota Ameliorates Colitis'. *Nature* 553 (7687): 208–11. https://doi.org/10.1038/nature25172.

Supplementary Figures

Supplementary Figure 1: Age distribution of participants in the healthy control, acute typhoid, and presumptive carrier groups, broken down by study site. Statistical annotations are the p-values resulting from Wilcoxon rank sum tests.

921
922   *Supplementary Figure 2: Sex split of the different participant groups across the different sites.*

923
924 *Supplementary Figure 3: Alpha diversity for acute typhoid fever patients and healthy household contacts for each*
925 *endemic country study site.*

929 *Supplementary Figure 4: Bar chart showing the prevalence of eight common phyla in acute typhoid patients and*
930 *healthy household contacts, for each study site.*

931
932 *Supplementary Figure 5: Prevalence of Prevotella copri clade A across age brackets in Malawi, Bangladesh, and*
933 *Nepal.*

936
937 *Supplementary Figure 6: Prevalence of Clostridium SGB6179 across age brackets in Malawi, Bangladesh, and Nepal*

938
939 *Supplementary Figure 7: Prevalence of GGB4266_SGB5809 across age brackets in Malawi, Bangladesh, and Nepal*

940
941 *Supplementary Figure 8: Prevalence of Haemophilus parainfluenzae across age brackets in Malawi, Bangladesh,*
942 *and Nepal*

943
944 *Supplementary Figure 9: Alpha diversity of acute typhoid patients, high Vi-titre participants, and household contacts*
945 *from Malawi and Nepal. Statistical annotations are the p-values resulting from Wilcoxon rank sum tests. A single*
946 *asterisk indicates a p-value < 0.01, triple asterisk indicates a p-value < 0.0001.*

948   *Supplementary Figure 10: Beta diversity of acute typhoid fever, high Vi-titre participants, and household contacts*
949   *from Malawi and Nepal*

950
951   *Supplementary Figure 11: Prevalence of B. obeum, H. parainfluenzae, and R. gnavus across three participant groups*
952   *(household contacts, typhoid cases, and high Vi-titre) in Malawi.*

953
954
955 *Supplementary Figure 12: Heatmap of top 400 taxa from Nepalese participants – household contacts, acute typhoid*
956 *cases and high-Vi titre participants.*

958
959 *Supplementary Figure 13: The abundance of species only described as species genome bins in the metaphlan*
960 *database from Malawian and Nepalese household contacts, acute typhoid, and high-Vi titre participants.*

962
963   *Supplementary Figure 14: Comparison of A) the number of AMR gene and B) the number of AMR gene classes*
964   *identified in participants from all three sites who did and did not report antibiotic usage prior to sampling*

965
966 *Supplementary Figure 15: Beta-diversity plot of 26 CHIM participants challenged with S. Typhi or S. Paratyphi,*
967 *coloured by whether they were diagnosed with typhoid/paratyphoid fever following pathogen exposure.*

968
969 *Supplementary Figure 16: Comparison of prevalence of A) succinate2priopionate and B) rnf MGCs between people*
970 *who were susceptible (disease) and non-susceptible (no_disease) to challenge with S. Typhi.*

971
972

Supplementary Table captions

Supplementary Table 1: ANOVA analysis of alpha diversity from healthy household contacts and typhoid fever patients from all three sites

Supplementary Table 2: PERMANOVA analysis of beta diversity from healthy household contacts and typhoid fever patients from all three sites

Supplementary Table 3: PERMANOVA analysis of beta diversity from healthy household contacts and typhoid fever patients from all Bangladesh

Supplementary Table 4: PERMANOVA analysis of beta diversity from healthy household contacts and typhoid fever patients from Malawi

Supplementary Table 5: PERMANOVA analysis of beta diversity from healthy household contacts and typhoid fever patients from Nepal

Supplementary Table 6: Median proportion of reads assigned to different phyla from healthy household contacts and typhoid fever patients from all three sites

Supplementary Table 7: Species associated with health or disease in Bangladesh only. Positive co-efficient (coef) means the species is associated with health, negative co-efficient means it's associated with typhoid fever.

Supplementary Table 8: Species associated with health or disease in Malawi only. Positive co-efficient (coef) means the species is associated with health, negative co-efficient means it's associated with typhoid fever.

Supplementary Table 9: Full information about Metabolic Gene Clusters associated with health in both Bangladesh and Malawi.

Supplementary Table 10: ANOVA analysis of alpha diversity from healthy household contacts, typhoid fever patients, and high Vi-titre participants from Malawi and Nepal

Supplementary Table 11: PERMANOVA analysis of beta diversity from healthy household contacts, typhoid fever patients, and high Vi-titre participants from Malawi

Supplementary Table 12: PERMANOVA analysis of beta diversity from healthy household contacts, typhoid fever patients, and high Vi-titre participants from Nepal

Supplementary Table 13: Species associated with high-Vi titre compared with household controls in Malawi only. Positive coefficient (coef) is associated with health, negative co-efficient is associated with high-Vi titre.

Supplementary Table 14: Metagenome identification of AMR genes commonly identified in Salmonella Typhi per country

Supplementary Table 15: Metagenome identification of AMR genes commonly identified in Salmonella Typhi per participant group

Supplementary Table 16: The average weighted importance of species to random forest classification of samples as being from control or presumptive carrier participants.

Supplementary Table 17: The correlation between the proportion of S. Typhi isolates and the proportion of microbiome samples from each site with AMR genes to quinolones, sulphonamides and tetracycline.

Supplementary Table 18: Demographic information on CHIM participants

Supplementary Table 19: Alpha diversity ANOVA results for CHIM data

Supplementary Table 20: PERMANOVA analysis of beta-diversity amongst CHIM participants challenged with S. Typhi and S. Paratyphi

Supplementary Table 21: Species associated with household controls vs both typhoid fever and high Vi titre from Malawi.