Brain White Matter Microstructure Associations with Blood Markers of the GSH Redox cycle in Schizophrenia

Tommaso Pavan[1], Pascal Steullet[2], Yasser Alemán-Gómez[1], Raoul Jenni[2], Zoé Schilliger[2,6], Martine Cleusix[2], Luis Alameda[3,4,5], Kim Q. Do[2], Philippe Conus[3], Patric Hagmann[1], Daniella Dwir[2], Paul Klauser[2,6], and Ileana Jelescu[1]

1. Department of Radiology, Lausanne University Hospital (CHUV) and University of Lausanne (UNIL), Lausanne, Switzerland

2. Center for Psychiatric Neuroscience, Department of Psychiatry, Lausanne University Hospital and the University of Lausanne, Lausanne, Switzerland

3. Service of General Psychiatry, Treatment and Early Intervention in Psychosis Program. Lausanne University Hospital (CHUV), Lausanne, Switzerland

4. Department of Psychosis Studies, Institute of Psychiatry, Psychology and Neuroscience. King's College of London, London, UK.

5. Centro Investigacion Biomedica en Red de Salud Mental (CIBERSAM); Instituto de Biomedicina de Sevilla (IBIS), Hospital Universitario Virgen del Rocio, Departamento de Psiquiatria, Universidad de Sevilla, Sevilla, Spain.

6. Service of Child and Adolescent Psychiatry, Department of Psychiatry, Lausanne University Hospital and the University of Lausanne, Lausanne, Switzerland

Corresponding author: Tommaso Pavan; address: *Centre de Recherche en Radiologie*, PET3, CHUV, Rue du Bugnon 46, 1011, Lausanne, Switzerland ; telephone:+41213146020; fax:+41213146020; email: tommaso.pavan@chuv.ch

Abstract

In groups of patients suffering from schizophrenia (SZ), redox dysregulation was reported in both peripheral fluids and brain. It has been hypothesized that such dysregulation, including alterations of the glutathione (GSH) cycle could participate in the brain white matter (WM) abnormalities in SZ due to the oligodendrocytes' susceptibility to oxidative stress.

In this study we aim to assess the differences between 82 schizophrenia patients (PT) and 86 healthy controls (HC) in GSH-redox peripheral blood markers: GSH peroxidase (GPx), reductase (GR) enzymatic activities and their ratio (GPx/GR-ratio), evaluating the hypotheses that alterations in the homeostasis of the systemic GSH cycle may be associated with pathological mechanisms in the brain WM in PT. To do so, we employ the advanced diffusion MRI methods: Diffusion Kurtosis Imaging (DKI) and White Matter Tract Integrity-Watson (WMTI-W), which provide excellent sensitivity to demyelination and neuroinflammation.

We show that GPx levels are higher (p=0.00041) in female control participants and decrease with aging (p=0.026). We find differences between PT and HC in the association of GR and mean kurtosis (MK, p<0.0001). Namely, lower MK was associated with higher blood GR activity in HC, but not in PT, suggesting that high GR activity (a hallmark of reductive stress) in HC was linked to changes in myelin integrity. However, GSH-redox peripheral blood markers did not explain the WM anomalies detected in PT, or the design of the present study could not detect subtle phenomenon, if present.

Introduction

Psychosis and schizophrenia (SZ) present significant challenges for patients, their families, and society (Wittchen et al., 2011). Only partially understood, its etiology is attributed to a multifaceted interplay of genetic, environmental and developmental factors that affect a variety of biological processes, including the redox balance (McCutcheon et al., 2020; Perkins et al., 2020). In SZ, a redox dysregulation was reported in peripheral fluids, suggested by a diminished antioxidant capacity, oxidative damage of lipids, DNA, and RNA (Jorgensen et al., 2022; Rambaud et al., 2022; Tsugawa et al., 2019), while in the brain, lower glutathione (GSH), and lower NAD+/NADH ratio were found (Das et al., 2019; Kim et al., 2017). The causes of such redox dysregulation and the resulting oxidative stress are multifactorial, including, chronic subclinical inflammation and weakened antioxidant capacities. Anomalies in the GSH redox system, a key antioxidant system for cell protection and detoxification, have been widely observed in SZ. Meta-analyses showed lower mean levels of GSH in blood (Tsugawa et al., 2019), and brain (Das et al., 2019; Gawryluk et al., 2011; Yao et al., 2006; Zhang et al., 2018) in patients vs. controls, as well as higher variability of brain GSH levels in SZ patients than in healthy subjects, suggesting the presence of subgroups of patients with different regulation of the GSH system (Palaniyappan et al., 2021).

The redox regulation and antioxidant function of GSH acts largely via the GSH redox cycle that involves the GSH peroxidase (GPx) and GSH reductase (GR). GPx neutralizes peroxides by oxidizing GSH into GSH-disulfide (GSSG). In turn, GR reduces GSSG back to GSH. The ratio between GSSG and GSH determines the redox potential of the GSH system that plays a central role in regulating redox-sensitive signaling pathways and mitochondrial oxidative phosphorylation. Low redox potentials (caused by too low GSH/GSSG ratio) and too high redox potentials (caused by too high GSH/GSSG ratio) are both detrimental to proper cellular function by provoking respectively oxidative stress and reductive stress. Reductive stress associated with too negative redox potential of the GSH/GSSG couple can indeed trigger mitochondrial oxidation and reduce cell viability (Huali Zhang et al., 2012). A proper coordination of GPx and GR activities is therefore essential to neutralize reactive oxygen species and maintain a proper redox potential of the GSH/GSSG couple.

Hence, GPx and GR activities and the GPx/GR-ratio could inform about the homeostasis of the GSH redox cycle. Two out of three postmortem studies have found lower brain GPx in SZ patients vs. controls (Gawryluk et al., 2011; Yao et al., 2006; Zhang et al., 2018). Similarly, meta-analyses suggest that SZ patients tend to also display lower blood GPx than controls (Tsugawa et al., 2019). However, the decrease in peripheral GPx activity seems to be mostly found in chronic patients (Flatow et al., 2013). In contrast to GPx, there is no evidence of consistent alterations of GR in the brain (Yao et al., 2006) or the blood (Tsugawa et al., 2019).

A possible reason for the discrepancies between studies is the influence of confounding factors such as age (Jones et al., 2002; Martínez de Toda et al., 2019), sex (Schilliger et al., 2024) and possibly medication (Raffa et al., 2009). This highlights the importance of accounting for the confounding effects of these demographic variables on the GSH-redox measures in relation to disease.

It has been hypothesized that a GSH deficit could contribute to the brain white matter (WM) abnormalities in SZ (Monin et al., 2016, 2015), due to the susceptibility of oligodendrocytes to oxidative stress (Juurlink et al., 1998) as

a consequence of their elevated metabolic activity and iron content required for myelin production and
maintenance (Bradl and Lassmann, 2010; Thorburne and Juurlink, 1996).
The GSH redox cycle supports the WM integrity by promoting the survival, proliferation and differentiation of the
oligodendrocyte precursor cells (Back et al., 1998; Monin et al., 2015), protecting the oligodendrocytes from
oxidative stress thanks to the GPx expression (Baud et al., 2004; Ravera et al., 2015). Beyond oligodendrocytes, the
GSH system dysregulation can also impair the energy metabolism of axons (Nave et al., 2023), the mitochondrial
functioning (Smith et al., 2019), and lead to subclinical activation of microglia (Dwir et al., 2020), affecting the WM.
Post-mortem studies of SZ patients have reported reduced oligodendrocyte densities in the cortex and subcortical
regions (Bernstein et al., 2009; Hof et al., 2003), swollen or dystrophic oligodendroglia (Uranova et al., 2018, 2011;
Williams, 2021), adjacent microglia (Uranova et al., 2020), and decompaction and splitting of the myelin sheath,
with vacuole inclusions between the myelin sheath (Uranova et al., 2018, 2011; Williams, 2021).
Abnormal white matter integrity in SZ is also widely reported using *in-vivo* diffusion magnetic resonance imaging
(dMRI). Most studies investigating schizophrenia show widespread increases in mean diffusivity (MD) and/or
reductions in fractional anisotropy (FA) (Barth et al., 2023; Cetin-Karayumak et al., 2020; Kelly et al., 2018) with
medium effect sizes (Kelly et al., 2018) (*Cohen's d*=0.42), consistent with anomalies observed in the *post-mortem*
studies.
However, direct evidence linking white matter abnormalities to the GSH system in schizophrenia remains limited.
GSH levels in frontal brain regions have been found to correlate positively with generalized FA in the cingulum of
both early psychosis patients and controls (Monin et al., 2015), while it was shown in earlier work that the
precursor of GSH and antioxidant, N-acetylcysteine, improves the integrity of the fornix (i.e., increases generalized
FA) (Klauser et al., 2018) and increases brain GSH (Conus et al., 2018) in early psychosis patients.
Here, we aim to assess whether the activity of GPx, GR, and the GPx/GR ratio measured in blood cells (which
inform about the systemic homeostasis of the GSH redox cycle) are associated with the brain WM microstructure
estimated using dMRI, in patients with a diagnosis from the schizophrenia spectrum. For this purpose, we utilized
advanced dMRI methods: Diffusion Kurtosis Imaging (DKI) (Jensen et al., 2005; Jensen and Helpern, 2010), an
extension of diffusion tensor imaging (DTI), and the biophysical model White Matter Tract Integrity-Watson
(Fieremans et al., 2011; Jespersen et al., 2018) (WMTI-W). In addition to classical DTI measures such as MD and
FA, DKI enables the estimation of the mean kurtosis (MK), a measure that provides complementary information
about tissue heterogeneity. The WMTI-W model allows the estimation of specific microstructural cellular features
such as the axonal density (f) in each voxel with fewer *ad hoc* simplifying assumptions and fit constraints (Novikov
et al., 2018; Hui Zhang et al., 2012), and increased validity (Novikov et al., 2018) as compared to other biophysical
methods such as free-water imaging (Pasternak et al., 2009) (FWI) and Neurite Orientation Dispersion and Density
Imaging (NODDI) (Hui Zhang et al., 2012). We chose WMTI-W over FWI, which is commonly found in studies on
inflammation in schizophrenia, due to reserves about interpreting the free-water fraction as a direct marker of
inflammation. Notably, diffusivities tend to decrease during acute gliosis, challenging the specificity of free-water to
inflammation (Guglielmetti et al., 2016). The FWI model also separates signal contributions into free isotropic
water and an anisotropic tissue compartment, which conflates intra- and extracellular components. In contrast,
our use of the Standard Model of white matter (like WMTI-W) provides greater specificity in distinguishing
inflammation from cell loss or myelin damage (Jelescu and Fieremans, 2023).
We focused our analysis on these four metrics (MD, FA, MK, f) because of several factors. Firstly, we wanted to limit
the number of multiple comparisons to retain good statistical power. Furthermore, MD, FA and MK are
complementary and highly sensitive to changes in the tissue microstructure, with similar or higher effect size as
compared to RD and RK in early psychosis and schizophrenia (Pavan et al., 2025). As for WMTI-W metrics, f was
shown to be sensitive to both demyelination and changes in axonal densities at earlier stages than the extra-axonal
perpendicular diffusivity (Falangola et al., 2014; Jelescu et al., 2016), and showed similar sensitivity to the extra-
axonal perpendicular diffusivity in discriminating groups (Pavan et al., 2025). Together, this set of metrics has the
ability to detect abnormal myelination or axonal densities (⬆MD, ⬇FA, ⬇MK, ⬇f) (Falangola et al., 2014;
Guglielmetti et al., 2016; Jelescu et al., 2016; Wang et al., 2011), but also neuroinflammation which has the effect of
reducing diffusivities (⬇MD) and increasing kurtosis and apparent axonal water fraction (⬆MK, ⬆f) due to higher
cellular crowding associated with microgliosis and astrocytosis (Guglielmetti et al., 2016; Jelescu et al., 2016;
Jelescu and Fieremans, 2023; Wijtenburg and Rowland, 2023).
In our previous work using this same cohort, which focused on WM alterations in EP and SZ vs healthy controls
(HC) (Pavan et al., 2025). We showed that WM MD was higher, while FA, MK, and f were lower in early psychosis
and SZ (⬆MD, ⬇FA, ⬇MK, ⬇f) than in HC, suggesting the predominance of myelin integrity abnormalities rather
than active inflammatory processes, which is also consistent with the cohort treatment history.
Here we hypothesize that alterations in the homeostasis of the systemic GSH redox cycle may be associated with
pathological mechanisms in the brain (i.e., redox dysregulation and microglia dysfunction, altered oligodendrocyte
density and myelination) that affect WM. Therefore, we expect that peripheral markers related to the GSH redox

cycle may correlate with WM microstructure metrics sensitive to axonal density, neuroinflammation and
myelination.
More specifically, we hypothesize that individuals with a high GPx/GR-ratio would exhibit a more oxidized state of
the GSH/GSSG couple compared to those with a low GPx/GR-ratio. An elevated GPx/GR ratio and/or increased GPx
activity in the blood may indicate systemic oxidative stress, whereas a low ratio and/or high GR activity may
suggest systemic reductive stress. Notably, patients who benefited from N-acetylcysteine treatment were primarily
those with high blood GPx activity (Conus et al., 2018). Conversely, reductive stress can disrupt redox-sensitive
processes and paradoxically elevate mitochondrial reactive oxygen species production, ultimately impairing
mitochondrial function and causing oxidative damage (Korge et al., 2015). Based on this, we stratified participants
into high and low GPx/GR-ratio groups to investigate potential differences in brain microstructure related to
distinct oxidative or reductive states. A comparable approach was used in a study assessing baseline GPx activity
(Conus et al., 2018).
Finally, due to the known effect of age (Martínez de Toda et al., 2019; Stadtman, 2002) and sex (Martínez de Toda
et al., 2023; Ruszkiewicz et al., 2019) on the expressions of the GSH redox cycle markers, we explored the role that
these confounding variables could have on the link between our peripheral GSH-redox measurements and WM
microstructure.

Methods

Participants

Data were collected from 168 individuals (**Table 1**) divided into two groups: 86 healthy controls (HC) and 82
patients (PT) with a diagnosis from the schizophrenia spectrum disorder (**Supplementary Table S1**). Patients
were recruited from the Lausanne University Hospital. PT with psychosis related to intoxication or organic brain
disease, IQ<70, reporting alcoholism, history of drug abuse (nicotine excluded), major somatic disease, documented
anamnestic or current organic brain damage were excluded. HC were recruited from the same sociodemographic
area as the patients. HC were excluded if they or a first-degree family member reported to have suffered from
neurological, traumatic, major mood disorder, psychosis, prodromal symptoms, and current or past antipsychotic
treatments. Symptoms in PT were assessed via the Positive and Negative Syndrome Scale (PANSS) (Kay et al., 1987),
and global functioning was estimated for the whole sample via the Global Assessment of Function scale (GAF)
(American Psychiatric Association, 2000) by a trained psychologist (not blinded). The study was approved by the
local Ethics Committee of the Canton of Vaud (Switzerland) under authorization numbers CER-VD 382/11 and
2018-01731.

MRI acquisition

172 MRI scanning sessions were performed on two different 3-Tesla systems (Magnetom TrioTim and PRISMA, Siemens
173 Healthineers, Erlangen, Germany), each equipped with a 32-channel head coil. A 1-mm isotropic T1-weighted
174 image was acquired for anatomical reference. Whole-brain diffusion-weighted images (DWI) were acquired using
175 diffusion spectrum imaging (DSI) scheme across 15 b-values, ranging from 0 to 8000 s/mm², spatial resolution of
176 2.2 x 2.2 x 3 mm³ - See Supplementary Material for further acquisition details.

Image preprocessing

178 The MPRAGE image was bias field corrected (Tustison et al., 2010) and skull-stripped via nonlinear registration to
179 the MNI-152 template (Grabner et al., 2006) using Advanced Normalization Tools (ANTs) (Avants et al., 2008). The
180 diffusion preprocessing pipeline included MP-PCA denoising, Gibbs ringing-, EPI-, eddy current and motion
181 corrections, following most recent guidelines (Ades-Aron et al., 2018) - see Supplementary Material for
182 preprocessing details.

Microstructure estimation

For Diffusion Kurtosis Imaging and White Matter Tract Integrity - Watson estimation, the diffusion dataset was
truncated (Jensen et al., 2005) at b≤2500 s/mm² (refer to Supplementary Methods for details). DKI was fit voxel-
wise in the entire brain (Veraart et al., 2013) using a weighted linear-least squares algorithm in Matlab (Veraart et
al., 2013), from which seven scalar maps were derived. However, only three parameter maps were included in the
statistical analysis: MD, FA, and MK. WMTI-W parameters were estimated voxel-wise from the seven DKI scalars,
using an in-house Python script, yielding five parameter maps. However, only axonal density f was included in the
statistical analysis. We focused our analysis on these four metrics (MD, FA, MK, f) because of their ability at
detecting abnormal myelination or axonal densities (▯MD, ▯FA, ▯MK, ▯f) (Falangola et al., 2014; Guglielmetti et al.,

2016; Jelescu et al., 2016; Wang et al., 2011), and neuroinflammation (⊠MD, ⊠MK, ⊠*f*) (Guglielmetti et al., 2016;
Jelescu et al., 2016; Jelescu and Fieremans, 2023; Wijtenburg and Rowland, 2023).
195
196
Blood analyses

Given that diurnal fluctuations are known for both GPx and GR (Budkowska et al., 2022), blood sampling was
tipically done early in the morning (8-10am) under fasting conditions. GR and GPx activity were assessed in
hemolyzed blood cells incubated in phosphate buffer solution (100mM, pH 7.5) containing EDTA (0.6mM), and
non-limiting levels of either oxidized glutathione (GSSG, 2.5mM) and NADPH (0.25mM) (for GR), or GSH (2.5mM),
NADPH (0.25mM), GR (0.84U/ml; Fluka) and tert-butyl hydroperoxide (TBHP, 0.8mM, Fluka) (for GPx). We
expressed GPx and GR activities in nmoles NADPH used/min/g hemoglobin. These enzymatic activities were
measured with two different plate readers. Batch effects between plate readers were corrected via ComBat for
biological data (Johnson et al., 2007) (see additional details on the blood analyses and harmonization procedure in
the Supplementary Methods).

Statistical analysis

Blood markers analysis

Statistical analyses were run in R, group comparisons were computed via Welch's t-test. Robust linear regression
(Maechler et al., 2023) was used for all the regression and contrast analyses, always including a correction by sex
and quadratic age (see Age Correction section in Supplementary Methods for additional details).

Association between blood markers and WM microstructure

Participants with a delay between MRI scan and blood samples larger than 60 days were excluded whenever an
association between blood biomarkers and WM microstructure metrics was tested (86% of the original cohort,
N_{HC}=80, N_{PT}=65, delay$_{HC}$=10.0±11.7, delay$_{PT}$=17.4±16.0, p=0.0012). The 60-days cutoff was chosen to retain the
highest number of subjects while reducing the group-level delay and controlling the effect of seasonal or longer
timescale fluctuations on the blood markers (see **Fig. S4** for details). All analyses included a correction for
quadratic age, sex, and delay. Associations were tested at voxel level using FSL's Tract-Based Spatial Statistics
(TBSS) (Smith et al., 2006). First, individual FA maps were used to build a study-specific FA template using ANTs
(Avants et al., 2008) and the estimated warps were used to spatially normalize each individual FA, MD, MK and *f*
map. Before any statistical analysis, all the microstructure parameter skeletons were harmonized for scanner type
at voxel level in an unbiased way (mean normalized towards the center of both sites, without reference batch) via
ComBat harmonization (Fortin et al., 2017). ComBat was previously proven efficient at correcting scanner effects
in the ENIGMA-consortium (Radua et al., 2020), in the current cohort (Alemán-Gómez et al., 2020; Pavan et al.,
2025), and validated in travelling participants (Richter et al., 2022). After permutation testing (FSL randomize
(Winkler et al., 2014), 5000 permutations), the resulting statistical maps were false discovery rate and Threshold-
Free Cluster Enhancement corrected.

Low and High GPx/GR ratio analysis

To study if there could be differences between participants showing different levels of GPx and GR activity, we
repeated the analysis described above after re-classifying the participants into two groups according to their
GPx/GR-ratio. Low-ratio and high-ratio were defined as having a GPx/GR-ratio respectively lower and higher than
the median GPx/GR ratio of the HC group (**Fig.1A,** Median$_{HC}$=6.64).

Results

Demographics

Summary demographics of the cohort can be found in **Table 1**. PT (29±9 y/o) and HC (28±8 y/o) did not differ in
age (p=0.21). As expected, differences were found between PT and HC in functional levels as measured by the GAF
scale (p<0.0001). Differences between groups were also found in the delay between MRI acquisition and blood
collection (p<0.0001), leading to the inclusion of the delay as a covariate.

GSH-redox cycle system

Overall, GPx (p=0.69) and GR (p=0.24) activities or the GPx/GR-ratio (p=0.32) did not differ between PT and HC
groups. HC female participants showed higher GPx activity than males (p=0.00041) but the same trend was not
found in PT. GPx and GR activities correlated significantly for the whole cohort (r=0.38, p<0.0001, **Fig. 1A**), but also
for PT (r=0.38, p=0.00038) and HC (r=0.40, p=0.00013) when considered independently. The slope contrast
between PT and HC did not show any significant difference (GPx ⊠ group interaction: p=0.80, **Fig. 1B**). GPx and GR
activities correlated also in males and females separately (r_{male}=0.33, p=0.00019 and r_{female}=0.46, p=0.00066) with
no significant sex difference for the slopes either (sex ⊠ group interaction: p=0.19, **Fig. 1C**). Age was only found
significantly associated with a decrease in GPx activity when pooling HC and PT together (**Fig. 1D**, p=0.026). No
significant association was found with medication. No significant association was found between psychiatric
functioning/symptoms scores and redox markers after multiple comparison correction (**Table S2**).
253
254
Associations between GSH-redox cycle and WM microstructure

TBSS analysis showed significant associations only between GR and MK in HC. No other association for PT or HC
was found between GPx, GR or GPx/GR-ratio with either MD, FA, MK or f. In HC, MK was significantly and
negatively associated with GR in multiple WM clusters (**Fig. 2A**, p=0.00018). Furthermore, the slope (GR–MK) in
HC differed significantly from the regression slope of PT (**Fig. 2B**, p<0.0001). The significant contrast was localized
bilaterally in the anterior and superior corona radiata, genu and body of the corpus callosum.
261
262

Associations between GSH-redox cycle and dMRI WM microstructure in the low and high GPx/GR-ratio
groups

We then assessed how low and high GPx/GR-ratio reflected at the WM microstructure level. First, we established
how the low- vs high-ratio is driven by both GPx and GR levels. Considering both PT and HC together, participants
with low ratio (p<0.0001) had significantly lower GPx (mean GPx=13.09) and higher GR (mean GR=2.61) as
compared to subjects with high ratio (mean GPx=19.86, mean GR=2.18). GPx and GR significantly correlated within
both low- (r=0.75, p<0.0001) and high-ratio (r=0.67, p<0.0001) groups.

TBSS voxel-wise analysis of the 3-way interaction model of GPx or GR interacting with low/high-ratio, and clinical
status, showed widespread significant differences in the slopes of MK vs GR between low ratio HC and PT (**Fig. 2C**,
p=0.00031) but not for any other comparison, indicating the low-ratio group drives the difference between HC and
PT found in the previous section. Identified areas for the GR-MK association were the bilateral anterior and
superior corona radiata, genu, splenium and body of the corpus callosum, right posterior limb, and right
retrolenticular part of the internal capsule. Notably, many voxels of the peripheral WM were also significant.
276
277

Discussion

In this work, we quantified peripheral biomarkers of the GSH redox cycle (GPx, GR activities and their ratio) in
patients with schizophrenia and control subjects, and we explored the relationships of these redox markers with
the WM microstructure estimated with advanced dMRI methods. We did not find significant differences in GPx, GR
or GPx/GR-ratio between groups. However, female participants showed higher GPx activity than males,
independently of their clinical status. GPx was also the only biomarker whose activity decreased with age,
suggesting a decrease in antioxidant defense. Finally, GR activity was negatively associated with MK in HC, but not
in PT, mostly driven by the subgroup of individuals with a low GPx/GR-ratio where the differences between HC and
PT were enhanced and more widespread across the WM.
287
288 *Peripheral GSH redox cycle*
289
Many studies have investigated blood GPx and GR activities in cohorts of patients suffering from schizophrenia
(Flatow et al., 2013; Koga et al., 2016; Tsugawa et al., 2019; Zhang et al., 2018), sometimes with mixed results.
However, meta-analyses (Flatow et al., 2013; Tsugawa et al., 2019) show overall significantly lower GPx, but not GR,
in PT as compared to HC. The analysis of Flatow et al. (2013) indicates that a decrease in GPx activity was prevalent
in patients with chronic SZ or in PT with acute relapses but not with first psychosis episode.

Some of the discrepancies in findings related to GSH/GPx/GR levels among studies on PT with schizophrenia could
be explained not only by the stage of the disease, but also by effects of age (Jones et al., 2002; Martínez de Toda et
al., 2019; Stadtman, 2002), sex (Schilliger et al., 2024), and medication (Raffa et al., 2009) on the peripheral
markers. We observed higher GPx activity in females than males in HC, and decreased GPx activity with age,
independently of the clinical status. However, in our cohort, which includes PT with early psychosis and chronic SZ,
we did not find any difference between PT and HC regarding GPx, GR and their ratio, even after correction for age
and sex.
Other sources of variability include methodological differences in sample collection, processing, storage, and
quantification (see Supplementary material in Perkins et al. (2020)).
Our findings of sex differences in HC and age effects on blood GPx activity regardless of the clinical status,
corroborate results from numerous studies (Abdalla et al., 1986; Jones et al., 2002; Martínez de Toda et al., 2019;
Schilliger et al., 2024; Stadtman, 2002). Higher levels of GPx, but not GR, have been widely reported in the brain of
females as compared to males in both animal and human studies (Dukhande et al., 2009; Martínez de Toda et al.,
2023; Ruszkiewicz et al., 2019). A higher GPx activity in females is consistent with more efficient antioxidant
systems and less basal inflammation in females than in males (Martínez de Toda et al., 2023). Our observation of a
diminished GPx activity with age is also in line with the literature (Martínez De Toda et al., 2019) and with the
notion that the antioxidant defenses weaken over the years, contributing to the accumulation of oxidative stress
and its harmful effects (Stadtman, 2002).
In our work, we found that GPx and GR activities correlated positively within both PT and HC and within both
males and females, suggesting a generally quite well-coordinated GSH redox cycle. Likewise, the GPx/GR ratio,
which was not yet examined before to our knowledge, did not differ between PT and HC groups. Previous work
showed that the activities of both enzymes were correlated in female but not male adolescents from the general
population (Schilliger et al., 2024), while, EP patients with a history of childhood trauma exhibited impaired
coordination between GPx and GR activity (Alameda et al., 2018). This suggests that the relationship between GPx
and GR activities may differ in a sex-dependent manner during adolescence but also among subgroups of
individuals according to environmental and other yet unidentified factors.
321
322 *Associations between peripheral GSH redox cycle and WM microstructure*
323
Due to the diversity in the GSH redox cycle status across individuals and the complex effects of sex and age, it was
therefore essential to tackle the relationships between the peripheral redox markers and WM microstructure by
carefully considering age, sex and their interactions, and by investigating the links between the GSH redox cycle and
WM separately in subgroups of individuals characterized by different peripheral redox profiles (oxidative vs
reductive stress).
The strongest differences between PT and HC were found in the association of GR with the WM microstructure (**Fig.**
2B, C), particularly with mean kurtosis. MK is considered a measure of tissue complexity (Jensen et al., 2005;
Jensen and Helpern, 2010), which has demonstrated sensitivity at detecting abnormal myelination (Falangola et al.,
2014; Guglielmetti et al., 2016; Jelescu et al., 2016; Wang et al., 2011), neurodegeneration and neuroinflammation
(Jelescu and Fieremans, 2023). Our previous work on the same data focusing on JHU region-of-interest (ROI)
analysis demonstrated that MK was significantly lower in early psychosis and chronic SZ as compared to age-range
matched controls (Pavan et al., 2025). Our current voxel-wise analysis on the white matter skeleton revealed that
lower MK was associated with higher blood GR activity in HC, but not in patients. This suggests that high GR activity
was linked with changes in myelin and tissue heterogeneity in HC. As high GR activity may be a sign of reductive
stress, we looked at the relationship between GR and MK in individuals with low GPx/GR-ratio, which proved to be
much stronger in this sub-group. This confirmed that the difference in the GR-MK association between PT and HC
was indeed mostly driven by individuals displaying a low GPx/GR-ratio.
Furthermore, it suggests that reductive stress in blood is linked to a low MK in the white matter of healthy
volunteers. Under the assumption that reductive stress in blood is accompanied by similar reductive stress in the
brain, our data would indicate that white matter is susceptible to reductive stress in the form of reduced
complexity, possibly from demyelination and cell density reduction. The latter could be a consequence of impaired
metabolism regulation, inflammation and cytotoxicity that are known to be associated with reductive stress (Huali
Zhang et al., 2012). However, high GR is not associated with low MK in the white matter of patients. The lack of a
direct association between MK and GR in patients suggests that other competing mechanisms contribute more to
the WM integrity impairment in patients.
Prior research has shown that N-acetylcysteine treatment can improve brain GSH levels after six months (Conus et
al., 2018) and WM integrity as measured by generalized FA in PT (Klauser et al., 2018). Notably, improvements in
cognitive abilities and in symptoms (Conus et al., 2018), particularly in individuals with high GPx levels, have also
been reported, highlighting the interplay between blood markers, WM integrity and cognitive function. Here,
however, we found no association of GPx or high GPx/GR-ratio with WM microstructure in PT, contrary to our

expectation. Instead, our results showed that PT and HC had a similar redox profile with no significant difference in
blood markers, indicating that patients do not exhibit a redox imbalance at the group level, which could explain the
lack of correlations between blood markers and WM alterations. No patient sub-group or cluster that would show a
significant difference in redox balance to the HC could be identified either.
Furthermore, Lesh et al. (2021) correlated average free-water fraction of the WM and GM with magnetic resonance
spectroscopy measures of GSH/creatine ratio in the dorsolateral prefrontal cortex. The authors found a negative
association between GSH/creatine and the free-water fraction in the GM but not WM, possibly suggesting that
changes due to GSH, and GPx/GR, may be more evident in the GM than the WM.
Interestingly, the spatial localization of the strongest GR-MK significant contrasts between HC and patients (**Fig.**
2D) coincides with the FA WM region of interests (anterior and superior corona radiata, genu and body of the
corpus callosum) showing the strongest effect size between HC and patients found in the large meta-analysis of the
ENIGMA study group (Kelly et al., 2018) and similarly reported in our previous study (Pavan et al., 2025).
Future research on the WM microstructure alterations in SZ should explore their relationship with brain
metabolism more directly. One potential approach is to examine the direct association between GSH-redox levels in
the brain, measured through MR spectroscopy, and WM microstructure. Furthermore, exploring the involvement of
other brain metabolites could also provide new insights into WM alterations.

Limitations

Our study has a few limitations that need to be considered. First, we used harmonization procedures to remove
both the effects of MRI scanner and of plate readers used to quantify the redox markers. In addition, the delay
between MRI acquisition and blood collection could be another source of bias (Perkins et al., 2020) despite being
capped and controlled during the statistical analyses. Second, the heterogeneity among the patient's cohort (that
includes early psychosis patients and subjects with chronic SZ) may significantly reduce the statistical power due to
potential conflicting trends that may arise at WM microstructure or GSH-system levels across the different stages
and types of psychosis. Third, we did not apply additional corrections across all modalities and contrasts for the
cluster findings (e.g., a second false discovery rate pass). This decision stems from the large number of voxels
resulting from pooling output images from multiple TBSS runs, and the inability of a second correction to account
for correlations within the blood and microstructure measures. No finding will survive such conservative
correction. Nevertheless, our clusters' results, already corrected with the classical correction (Threshold-Free
Cluster Enhancement + false discovery rate) found in TBSS, are large and spatially consistent, suggesting they are
unlikely to be false positives. Fourth, lifestyle habits, such as smoking (Mons et al., 2016) and diet (Gould and
Pazdro, 2019), could influence GSH, GPx and GR activities, having a confounding role on our results. Accounting for
these factors in a reliable way is challenging as they are often self-reported. Lastly, GPx and GR activities and their
ratio in blood cells inform us about the systemic redox homeostasis which does not necessarily reflect the
conditions found in the brain. However, several studies have found associations of peripheral markers of the GSH-
redox cycle with brain GSH levels (Xin et al., 2016), brain atrophy and enlargement of the lateral ventricles
(Buckman et al., 1987), or EEG sensory evoked responses (Geiser et al., 2017).

Conclusions

Altogether, our association analyses indicate that there is no direct correlation between the peripheral GSH-redox
cycle and WM anomalies detected in PT, although a link between reductive stress and WM microstructure was
found in HC. Our results do not therefore exclude the involvement of the GSH redox cycle during the
development/maturation of WM. The dysregulation of this system during brain development could still contribute
to WM anomalies in psychosis and schizophrenia.

Acknowledgments

This work was supported by the Swiss National Science Foundation (PCEFP2_194260, to I.J.; 320030_197787 to PH
and YA), the National Center of Competence in Research (NCCR) "SYNAPSY - The Synaptic Bases of Mental
Diseases" from the Swiss National Science Foundation (n° 51NF40 – 185897 to KQD & PC) and the Foundation
Alamaya. Dr. Alameda is supported by Carigest fellowship and by Frutiger Adrian et Simone fellowship. Dr. Dwir
and P. Klauser are supported by Frutiger Adrian & Simone fellowship. We are grateful to Gloria Reuteler, Adeline
Cottier and Morgane Baumgartner for expert technical assistance.

Competing Interests

The authors have no competing interests or other interests that might be perceived to influence the results and/or
discussion reported in this paper.
414
415
416 **Data Availability**

417 Data will not be publicly available due to absence of consent from the participants. The code is available here:
418 *github.com/Mic-map/WMTI-Watson_Python*
419
420
421 **Author Contributions**

422 The authors confirm contribution to the paper as follows: study conceptualization: T.P, P.S., I.J, P.K., D.D.;
423 investigation and formal analysis: T.P.;
424 interpretation of results: T.P, P.S., I.J, P.K., D.D., R.J., Z.S.; writing — original draft: T.P., P.S, I.J.; MR data collection:
425 M.C.; writing — review and editing: L.A., K.D., P.C., and P.H.
426
427
428 **Declaration of generative AI and AI-assisted technologies in the writing process**

429 During the preparation of this work the author(s) used ChatGPT to improve language and readability. After using
430 this tool/service, the author(s) reviewed and edited the content as needed and take(s) full responsibility for the
431 content of the publication.

433 References

Abdalla, D.S., Monteiro, H.P., Oliveira, J.A., Bechara, E.J., 1986. Activities of superoxide dismutase and glutathione

peroxidase in schizophrenic and manic-depressive patients. Clin. Chem. 32, 805–807.

https://doi.org/10.1093/clinchem/32.5.805

Ades-Aron, B., Veraart, J., Kochunov, P., McGuire, S., Sherman, P., Kellner, E., Novikov, D.S., Fieremans, E., 2018.

Evaluation of the accuracy and precision of the diffusion parameter EStImation with Gibbs and NoisE removal

pipeline. NeuroImage 183, 532–543. https://doi.org/10.1016/j.neuroimage.2018.07.066

Alameda, L., Fournier, M., Khadimallah, I., Griffa, A., Cleusix, M., Jenni, R., Ferrari, C., Klauser, P., Baumann, P.S.,

Cuenod, M., Hagmann, P., Conus, P., Do, K.Q., 2018. Redox dysregulation as a link between childhood trauma and

psychopathological and neurocognitive profile in patients with early psychosis. Proc. Natl. Acad. Sci. U. S. A. 115,

12495–12500. https://doi.org/10.1073/pnas.1812821115

Alemán-Gómez, Y., Najdenovska, E., Roine, T., Fartaria, M.J., Canales-Rodríguez, E.J., Rovó, Z., Hagmann, P., Conus, P.,

Do, K.Q., Klauser, P., Steullet, P., Baumann, P.S., Bach Cuadra, M., 2020. Partial-volume modeling reveals reduced gray

matter in specific thalamic nuclei early in the time course of psychosis and chronic schizophrenia. Hum. Brain

Mapp. 41, 4041–4061. https://doi.org/10.1002/hbm.25108

American Psychiatric Association, 2000. Diagnostic and Statistical Manual of Mental Disorders (4th ed., text rev.).

Avants, B.B., Epstein, C.L., Grossman, M., Gee, J.C., 2008. Symmetric diffeomorphic image registration with cross-

correlation: evaluating automated labeling of elderly and neurodegenerative brain. Med. Image Anal. 12, 26–41.

https://doi.org/10.1016/j.media.2007.06.004

Back, S.A., Gan, X., Li, Y., Rosenberg, P.A., Volpe, J.J., 1998. Maturation-Dependent Vulnerability of Oligodendrocytes

to Oxidative Stress-Induced Death Caused by Glutathione Depletion. J. Neurosci. 18, 6241–6253.

https://doi.org/10.1523/JNEUROSCI.18-16-06241.1998

Barth, C., Kelly, S., Nerland, S., Jahanshad, N., Alloza, C., Ambrogi, S., Andreassen, O.A., Andreou, D., Arango, C., Baeza,

I., Banaj, N., Bearden, C.E., Berk, M., Bohman, H., Castro-Fornieles, J., Chye, Y., Crespo-Facorro, B., de la Serna, E.,

Díaz-Caneja, C.M., Gurholt, T.P., Hegarty, C.E., James, A., Janssen, J., Johannessen, C., Jönsson, E.G., Karlsgodt, K.H.,

Kochunov, P., Lois, N.G., Lundberg, M., Myhre, A.M., Pascual-Diaz, S., Piras, F., Smelror, R.E., Spalletta, G., Stokkan, T.S.,

Sugranyes, G., Suo, C., Thomopoulos, S.I., Tordesillas-Gutiérrez, D., Vecchio, D., Wedervang-Resell, K., Wortinger, L.A.,

Thompson, P.M., Agartz, I., 2023. In vivo white matter microstructure in adolescents with early-onset psychosis: a

multi-site mega-analysis. Mol. Psychiatry 28, 1159–1169. https://doi.org/10.1038/s41380-022-01901-3

Baud, O., Greene, A.E., Li, J., Wang, H., Volpe, J.J., Rosenberg, P.A., 2004. Glutathione peroxidase-catalase

cooperativity is required for resistance to hydrogen peroxide by mature rat oligodendrocytes. J. Neurosci. Off. J. Soc.

Neurosci. 24, 1531–1540. https://doi.org/10.1523/JNEUROSCI.3989-03.2004

Beaulieu, C., 2002. The basis of anisotropic water diffusion in the nervous system - a technical review. NMR

Biomed. 15, 435–455. https://doi.org/10.1002/nbm.782

Bernstein, H.-G., Steiner, J., Bogerts, B., 2009. Glial cells in schizophrenia: pathophysiological significance and

possible consequences for therapy. Expert Rev. Neurother. 9, 1059–1071. https://doi.org/10.1586/ern.09.59

Bradl, M., Lassmann, H., 2010. Oligodendrocytes: biology and pathology. Acta Neuropathol. (Berl.) 119, 37–53.

https://doi.org/10.1007/s00401-009-0601-5

Buckman, T.D., Kling, A.S., Eiduson, S., Sutphin, M.S., Steinberg, A., 1987. Glutathione peroxidase and CT scan

abnormalities in schizophrenia. Biol. Psychiatry 22, 1349–1356. https://doi.org/10.1016/0006-3223(87)90069-2

Budkowska, M., Cecerska-Heryć, E., Marcinowska, Z., Siennicka, A., Dołęgowska, B., 2022. The Influence of Circadian

Rhythm on the Activity of Oxidative Stress Enzymes. Int. J. Mol. Sci. 23, 14275.

https://doi.org/10.3390/ijms232214275

Cetin-Karayumak, S., Di Biase, M.A., Chunga, N., Reid, B., Somes, N., Lyall, A.E., Kelly, S., Solgun, B., Pasternak, O.,

Vangel, M., Pearlson, G., Tamminga, C., Sweeney, J.A., Clementz, B., Schretlen, D., Viher, P.V., Stegmayer, K., Walther, S.,

Lee, J., Crow, T., James, A., Voineskos, A., Buchanan, R.W., Szeszko, P.R., Malhotra, A.K., Hegde, R., McCarley, R.,

Keshavan, M., Shenton, M., Rathi, Y., Kubicki, M., 2020. White matter abnormalities across the lifespan of

schizophrenia: a harmonized multi-site diffusion MRI study. Mol. Psychiatry 25, 3208–3219.
https://doi.org/10.1038/s41380-019-0509-y

Conus, P., Seidman, L.J., Fournier, M., Xin, L., Cleusix, M., Baumann, P.S., Ferrari, C., Cousins, A., Alameda, L., Gholam-
Rezaee, M., Golay, P., Jenni, R., Woo, T.-U.W., Keshavan, M.S., Eap, C.B., Wojcik, J., Cuenod, M., Buclin, T., Gruetter, R.,
Do, K.Q., 2018. N-acetylcysteine in a Double-Blind Randomized Placebo-Controlled Trial: Toward Biomarker-
Guided Treatment in Early Psychosis. Schizophr. Bull. 44, 317–327. https://doi.org/10.1093/schbul/sbx093

Das, T.K., Javadzadeh, A., Dey, A., Sabesan, P., Théberge, J., Radua, J., Palaniyappan, L., 2019. Antioxidant defense in
schizophrenia and bipolar disorder: A meta-analysis of MRS studies of anterior cingulate glutathione. Prog.
Neuropsychopharmacol. Biol. Psychiatry 91, 94–102. https://doi.org/10.1016/j.pnpbp.2018.08.006

Dukhande, V.V., Isaac, A.O., Chatterji, T., Lai, J.C.K., 2009. Reduced glutathione regenerating enzymes undergo
developmental decline and sexual dimorphism in the rat cerebral cortex. Brain Res. 1286, 19–24.
https://doi.org/10.1016/j.brainres.2009.05.029

Dwir, D., Giangreco, B., Xin, L., Tenenbaum, L., Cabungcal, J.-H., Steullet, P., Goupil, A., Cleusix, M., Jenni, R., Chtarto, A.,
Baumann, P.S., Klauser, P., Conus, P., Tirouvanziam, R., Cuenod, M., Do, K.Q., 2020. MMP9/RAGE pathway
overactivation mediates redox dysregulation and neuroinflammation, leading to inhibitory/excitatory imbalance: a
reverse translation study in schizophrenia patients. Mol. Psychiatry 25, 2889–2904.
https://doi.org/10.1038/s41380-019-0393-5

Falangola, M.F., Guilfoyle, D.N., Tabesh, A., Hui, E.S., Nie, X., Jensen, J.H., Gerum, S.V., Hu, C., LaFrancois, J., Collins, H.R.,
Helpern, J.A., 2014. Histological correlation of diffusional kurtosis and white matter modeling metrics in cuprizone-
induced corpus callosum demyelination. NMR Biomed. 27, 948–957. https://doi.org/10.1002/nbm.3140

Fieremans, E., Jensen, J.H., Helpern, J.A., 2011. White matter characterization with diffusional kurtosis imaging.
NeuroImage 58, 177–188. https://doi.org/10.1016/j.neuroimage.2011.06.006

Flatow, J., Buckley, P., Miller, B.J., 2013. Meta-Analysis of Oxidative Stress in Schizophrenia. Biol. Psychiatry 74, 400–
409. https://doi.org/10.1016/j.biopsych.2013.03.018

Fortin, J.-P., Parker, D., Tunç, B., Watanabe, T., Elliott, M.A., Ruparel, K., Roalf, D.R., Satterthwaite, T.D., Gur, R.C., Gur,
R.E., Schultz, R.T., Verma, R., Shinohara, R.T., 2017. Harmonization of multi-site diffusion tensor imaging data.
NeuroImage 161, 149–170. https://doi.org/10.1016/j.neuroimage.2017.08.047

Fournier, M., Scolamiero, M., Gholam-Rezaee, M.M., Cleusix, M., Jenni, R., Ferrari, C., Golay, P., Baumann, P.S., Cuenod,
M., Conus, P., Do, K.Q., Hess, K., 2021. Topology predicts long-term functional outcome in early psychosis. Mol.
Psychiatry 26, 5335–5346. https://doi.org/10.1038/s41380-020-0826-1

Gawryluk, J.W., Wang, J.-F., Andreazza, A.C., Shao, L., Young, L.T., 2011. Decreased levels of glutathione, the major
brain antioxidant, in post-mortem prefrontal cortex from patients with psychiatric disorders. Int. J.
Neuropsychopharmacol. 14, 123–130. https://doi.org/10.1017/S1461145710000805

Geiser, E., Retsa, C., Knebel, J.-F., Ferrari, C., Jenni, R., Fournier, M., Alameda, L., Baumann, P.S., Clarke, S., Conus, P., Do,
K.Q., Murray, M.M., 2017. The coupling of low-level auditory dysfunction and oxidative stress in psychosis patients.
Schizophr. Res. 190, 52–59. https://doi.org/10.1016/j.schres.2017.02.002

Gould, R.L., Pazdro, R., 2019. Impact of Supplementary Amino Acids, Micronutrients, and Overall Diet on
Glutathione Homeostasis. Nutrients 11, 1056. https://doi.org/10.3390/nu11051056

Grabner, G., Janke, A.L., Budge, M.M., Smith, D., Pruessner, J., Collins, D.L., 2006. Symmetric Atlasing and Model
Based Segmentation: An Application to the Hippocampus in Older Adults, in: Larsen, R., Nielsen, M., Sporring, J.
(Eds.), Medical Image Computing and Computer-Assisted Intervention – MICCAI 2006, Lecture Notes in Computer
Science. Springer, Berlin, Heidelberg, pp. 58–66. https://doi.org/10.1007/11866763_8

Gravina, P., Spoletini, I., Masini, S., Valentini, A., Vanni, D., Paladini, E., Bossù, P., Caltagirone, C., Federici, G., Spalletta,
G., Bernardini, S., 2011. Genetic polymorphisms of glutathione S-transferases GSTM1, GSTT1, GSTP1 and GSTA1 as
risk factors for schizophrenia. Psychiatry Res. 187, 454–456. https://doi.org/10.1016/j.psychres.2010.10.008

Guglielmetti, C., Veraart, J., Roelant, E., Mai, Z., Daans, J., Van Audekerke, J., Naeyaert, M., Vanhoutte, G., Delgado y
Palacios, R., Praet, J., Fieremans, E., Ponsaerts, P., Sijbers, J., Van der Linden, A., Verhoye, M., 2016. Diffusion kurtosis
imaging probes cortical alterations and white matter pathology following cuprizone induced demyelination and
spontaneous remyelination. NeuroImage 125, 363–377. https://doi.org/10.1016/j.neuroimage.2015.10.052

Gysin, R., Kraftsik, R., Sandell, J., Bovet, P., Chappuis, C., Conus, P., Deppen, P., Preisig, M., Ruiz, V., Steullet, P., Tosic, M.,
Werge, T., Cuénod, M., Do, K.Q., 2007. Impaired glutathione synthesis in schizophrenia: Convergent genetic and
functional evidence. Proc. Natl. Acad. Sci. 104, 16621–16626. https://doi.org/10.1073/pnas.0706778104

Hof, P.R., Haroutunian, V., Friedrich, V.L., Byne, W., Buitron, C., Perl, D.P., Davis, K.L., 2003. Loss and altered spatial
distribution of oligodendrocytes in the superior frontal gyrus in schizophrenia. Biol. Psychiatry 53, 1075–1085.
https://doi.org/10.1016/S0006-3223(03)00237-3

Jelescu, I.O., Fieremans, E., 2023. Chapter 2 - Sensitivity and specificity of diffusion MRI to neuroinflammatory
processes, in: Laule, C., Port, J.D. (Eds.), Advances in Magnetic Resonance Technology and Applications, Imaging
Neuroinflammation. Academic Press, pp. 31–50. https://doi.org/10.1016/B978-0-323-91771-1.00010-1

Jelescu, I.O., Zurek, M., Winters, K.V., Veraart, J., Rajaratnam, A., Kim, N.S., Babb, J.S., Shepherd, T.M., Novikov, D.S.,
Kim, S.G., Fieremans, E., 2016. In vivo quantification of demyelination and recovery using compartment-specific

diffusion MRI metrics validated by electron microscopy. NeuroImage 132, 104–114.
https://doi.org/10.1016/j.neuroimage.2016.02.004

Jensen, J.H., Helpern, J.A., 2010. MRI quantification of non-Gaussian water diffusion by kurtosis analysis. NMR
Biomed. 23, 698–710. https://doi.org/10.1002/nbm.1518

Jensen, J.H., Helpern, J.A., Ramani, A., Lu, H., Kaczynski, K., 2005. Diffusional kurtosis imaging: The quantification of
non-gaussian water diffusion by means of magnetic resonance imaging. Magn. Reson. Med. 53, 1432–1440.
https://doi.org/10.1002/mrm.20508

Jespersen, S.N., Olesen, J.L., Hansen, B., Shemesh, N., 2018. Diffusion time dependence of microstructural
parameters in fixed spinal cord. NeuroImage 182, 329–342. https://doi.org/10.1016/j.neuroimage.2017.08.039

Johnson, W.E., Li, C., Rabinovic, A., 2007. Adjusting batch effects in microarray expression data using empirical
Bayes methods. Biostat. Oxf. Engl. 8, 118–127. https://doi.org/10.1093/biostatistics/kxj037

Jones, D.P., Mody, V.C., Carlson, J.L., Lynn, M.J., Sternberg, P., 2002. Redox analysis of human plasma allows
separation of pro-oxidant events of aging from decline in antioxidant defenses. Free Radic. Biol. Med. 33, 1290–
1300. https://doi.org/10.1016/S0891-5849(02)01040-7

Jorgensen, A., Baago, I.B., Rygner, Z., Jorgensen, M.B., Andersen, P.K., Kessing, L.V., Poulsen, H.E., 2022. Association of
Oxidative Stress-Induced Nucleic Acid Damage With Psychiatric Disorders in Adults: A Systematic Review and
Meta-analysis. JAMA Psychiatry 79, 920–931. https://doi.org/10.1001/jamapsychiatry.2022.2066

Juurlink, B.H., Thorburne, S.K., Hertz, L., 1998. Peroxide-scavenging deficit underlies oligodendrocyte susceptibility
to oxidative stress. Glia 22, 371–378. https://doi.org/10.1002/(sici)1098-1136(199804)22:4<371::aid-
glia6>3.0.co;2-6

Kay, S.R., Fiszbein, A., Opler, L.A., 1987. The Positive and Negative Syndrome Scale (PANSS) for Schizophrenia.
Schizophr. Bull. 13, 261–276. https://doi.org/10.1093/schbul/13.2.261

Kelly, S., Jahanshad, N., Zalesky, A., Kochunov, P., Agartz, I., Alloza, C., Andreassen, O.A., Arango, C., Banaj, N., Bouix, S.,
Bousman, C.A., Brouwer, R.M., Bruggemann, J., Bustillo, J., Cahn, W., Calhoun, V., Cannon, D., Carr, V., Catts, S., Chen, J.,
Chen, J.-X., Chen, X., Chiapponi, C., Cho, K.K., Ciullo, V., Corvin, A.S., Crespo-Facorro, B., Cropley, V., De Rossi, P., Diaz-
Caneja, C.M., Dickie, E.W., Ehrlich, S., Fan, F.-M., Faskowitz, J., Fatouros-Bergman, H., Flyckt, L., Ford, J.M., Fouche, J.-
P., Fukunaga, M., Gill, M., Glahn, D.C., Gollub, R., Goudzwaard, E.D., Guo, H., Gur, R.E., Gur, R.C., Gurholt, T.P.,
Hashimoto, R., Hatton, S.N., Henskens, F.A., Hibar, D.P., Hickie, I.B., Hong, L.E., Horacek, J., Howells, F.M., Hulshoff Pol,
H.E., Hyde, C.L., Isaev, D., Jablensky, A., Jansen, P.R., Janssen, J., Jönsson, E.G., Jung, L.A., Kahn, R.S., Kikinis, Z., Liu, K.,
Klauser, P., Knöchel, C., Kubicki, M., Lagopoulos, J., Langen, C., Lawrie, S., Lenroot, R.K., Lim, K.O., Lopez-Jaramillo, C.,
Lyall, A., Magnotta, V., Mandl, R.C.W., Mathalon, D.H., McCarley, R.W., McCarthy-Jones, S., McDonald, C., McEwen, S.,

McIntosh, A., Melicher, T., Mesholam-Gately, R.I., Michie, P.T., Mowry, B., Mueller, B.A., Newell, D.T., O'Donnell, P., Oertel-Knöchel, V., Oestreich, L., Paciga, S.A., Pantelis, C., Pasternak, O., Pearlson, G., Pellicano, G.R., Pereira, A., Pineda Zapata, J., Piras, F., Potkin, S.G., Preda, A., Rasser, P.E., Roalf, D.R., Roiz, R., Roos, A., Rotenberg, D., Satterthwaite, T.D., Savadjiev, P., Schall, U., Scott, R.J., Seal, M.L., Seidman, L.J., Shannon Weickert, C., Whelan, C.D., Shenton, M.E., Kwon, J.S., Spalletta, G., Spaniel, F., Sprooten, E., Stäblein, M., Stein, D.J., Sundram, S., Tan, Y., Tan, S., Tang, S., Temmingh, H.S., Westlye, L.T., Tønnesen, S., Tordesillas-Gutierrez, D., Doan, N.T., Vaidya, J., van Haren, N.E.M., Vargas, C.D., Vecchio, D., Velakoulis, D., Voineskos, A., Voyvodic, J.Q., Wang, Z., Wan, P., Wei, D., Weickert, T.W., Whalley, H., White, T., Whitford, T.J., Wojcik, J.D., Xiang, H., Xie, Z., Yamamori, H., Yang, F., Yao, N., Zhang, G., Zhao, J., van Erp, T.G.M., Turner, J., Thompson, P.M., Donohoe, G., 2018. Widespread white matter microstructural differences in schizophrenia across 4322 individuals: results from the ENIGMA Schizophrenia DTI Working Group. Mol. Psychiatry 23, 1261–1269. https://doi.org/10.1038/mp.2017.170

Kim, S.-Y., Cohen, B.M., Chen, X., Lukas, S.E., Shinn, A.K., Yuksel, A.C., Li, T., Du, F., Öngür, D., 2017. Redox Dysregulation in Schizophrenia Revealed by in vivo NAD+/NADH Measurement. Schizophr. Bull. 43, 197–204. https://doi.org/10.1093/schbul/sbw129

Klauser, P., Xin, L., Fournier, M., Griffa, A., Cleusix, M., Jenni, R., Cuenod, M., Gruetter, R., Hagmann, P., Conus, P., Baumann, P.S., Do, K.Q., 2018. N-acetylcysteine add-on treatment leads to an improvement of fornix white matter integrity in early psychosis: a double-blind randomized placebo-controlled trial. Transl. Psychiatry 8, 1–8. https://doi.org/10.1038/s41398-018-0266-8

Koga, M., Serritella, A.V., Sawa, A., Sedlak, T.W., 2016. Implications for reactive oxygen species in schizophrenia pathogenesis. Schizophr. Res. 176, 52–71. https://doi.org/10.1016/j.schres.2015.06.022

Korge, P., Calmettes, G., Weiss, J.N., 2015. Increased reactive oxygen species production during reductive stress: The roles of mitochondrial glutathione and thioredoxin reductases. Biochim. Biophys. Acta BBA - Bioenerg. 1847, 514–525. https://doi.org/10.1016/j.bbabio.2015.02.012

Lesh, T.A., Maddock, R.J., Howell, A., Wang, H., Tanase, C., Daniel Ragland, J., Niendam, T.A., Carter, C.S., 2021. Extracellular free water and glutathione in first-episode psychosis-a multimodal investigation of an inflammatory model for psychosis. Mol. Psychiatry 26, 761–771. https://doi.org/10.1038/s41380-019-0428-y

Lloyd, A.F., Davies, C.L., Miron, V.E., 2017. Microglia: origins, homeostasis, and roles in myelin repair. Curr. Opin. Neurobiol. 47, 113–120. https://doi.org/10.1016/j.conb.2017.10.001

Ma, J., Li, D.-M., Zhang, R., Yang, X.-D., Gao, C.-G., Lu, S.-M., Wu, S.-N., Wang, L., 2010. Genetic analysis of glutamate cysteine ligase modifier (GCLM) gene and schizophrenia in Han Chinese. Schizophr. Res. 119, 273–274. https://doi.org/10.1016/j.schres.2009.12.017

Maechler, M., Rousseeuw, P., Croux, C., Todorov, V., Ruckstuhl, A., Salibian-Barrera, M., Verbeke, T., Koller, M.,

Conceicao, E.L.T., Palma, M.A. di, 2023. robustbase: Basic Robust Statistics.

Martínez de Toda, I., González-Sánchez, M., Díaz-Del Cerro, E., Valera, G., Carracedo, J., Guerra-Pérez, N., 2023. Sex

differences in markers of oxidation and inflammation. Implications for ageing. Mech. Ageing Dev. 211, 111797.

https://doi.org/10.1016/j.mad.2023.111797

Martínez De Toda, I., Vida, C., Garrido, A., De La Fuente, M., 2019. Redox Parameters as Markers of the Rate of Aging

and Predictors of Life Span. J. Gerontol. Ser. A. https://doi.org/10.1093/gerona/glz033

Martínez de Toda, I., Vida, C., Sanz San Miguel, L., De la Fuente, M., 2019. Function, Oxidative, and Inflammatory

Stress Parameters in Immune Cells as Predictive Markers of Lifespan throughout Aging. Oxid. Med. Cell. Longev.

2019, 4574276. https://doi.org/10.1155/2019/4574276

McCutcheon, R.A., Reis Marques, T., Howes, O.D., 2020. Schizophrenia—An Overview. JAMA Psychiatry 77, 201–210.

https://doi.org/10.1001/jamapsychiatry.2019.3360

Monin, A., Baumann, P.S., Griffa, A., Xin, L., Mekle, R., Fournier, M., Butticaz, C., Klaey, M., Cabungcal, J.H., Steullet, P.,

Ferrari, C., Cuenod, M., Gruetter, R., Thiran, J.P., Hagmann, P., Conus, P., Do, K.Q., 2015. Glutathione deficit impairs

myelin maturation: relevance for white matter integrity in schizophrenia patients. Mol. Psychiatry 20, 827–838.

https://doi.org/10.1038/mp.2014.88

Monin, A., Fournier, M., Baumann, P.S., Cuénod, M., Do, K.Q., 2016. Role of Redox Dysregulation in White Matter

Anomalies Associated with Schizophrenia, in: Handbook of Behavioral Neuroscience. Elsevier, pp. 481–500.

https://doi.org/10.1016/B978-0-12-800981-9.00028-6

Mons, U., Muscat, J.E., Modesto, J., Richie, J.P., Brenner, H., 2016. Effect of smoking reduction and cessation on the

plasma levels of the oxidative stress biomarker glutathione – Post-hoc analysis of data from a smoking cessation

trial. Free Radic. Biol. Med. 91, 172–177. https://doi.org/10.1016/j.freeradbiomed.2015.12.018

Nave, K.-A., Asadollahi, E., Sasmita, A., 2023. Expanding the function of oligodendrocytes to brain energy

metabolism. Curr. Opin. Neurobiol. 83, 102782. https://doi.org/10.1016/j.conb.2023.102782

Novikov, D.S., Kiselev, V.G., Jespersen, S.N., 2018. On modeling. Magn. Reson. Med. 79, 3172–3193.

https://doi.org/10.1002/mrm.27101

Palaniyappan, L., Sabesan, P., Li, X., Luo, Q., 2021. Schizophrenia Increases Variability of the Central Antioxidant

System: A Meta-Analysis of Variance From MRS Studies of Glutathione. Front. Psychiatry 12, 796466.

https://doi.org/10.3389/fpsyt.2021.796466

Pasternak, O., Sochen, N., Gur, Y., Intrator, N., Assaf, Y., 2009. Free water elimination and mapping from diffusion

MRI. Magn. Reson. Med. 62, 717–730. https://doi.org/10.1002/mrm.22055

Pavan, T., Alemán-Gómez, Y., Jenni, R., Steullet, P., Schilliger, Z., Dwir, D., Cleusix, M., Alameda, L., Do, K.Q., Conus, P.,

Hagmann, P., Klauser, P., Jelescu, I., 2025. White matter microstructure alterations in early psychosis and

schizophrenia. Transl. Psychiatry 15, 1–10. https://doi.org/10.1038/s41398-025-03397-1

Perkins, D.O., Jeffries, C.D., Do, K.Q., 2020. Potential Roles of Redox Dysregulation in the Development of

Schizophrenia. Biol. Psychiatry 88, 326–336. https://doi.org/10.1016/j.biopsych.2020.03.016

Radua, J., Vieta, E., Shinohara, R., Kochunov, P., Quidé, Y., Green, M.J., Weickert, C.S., Weickert, T., Bruggemann, J.,

Kircher, T., Nenadić, I., Cairns, M.J., Seal, M., Schall, U., Henskens, F., Fullerton, J.M., Mowry, B., Pantelis, C., Lenroot, R.,

Cropley, V., Loughland, C., Scott, R., Wolf, D., Satterthwaite, T.D., Tan, Y., Sim, K., Piras, Fabrizio, Spalletta, G., Banaj, N.,

Pomarol-Clotet, E., Solanes, A., Albajes-Eizagirre, A., Canales-Rodríguez, E.J., Sarro, S., Di Giorgio, A., Bertolino, A.,

Stäblein, M., Oertel, V., Knöchel, C., Borgwardt, S., du Plessis, S., Yun, J.-Y., Kwon, J.S., Dannlowski, U., Hahn, T.,

Grotegerd, D., Alloza, C., Arango, C., Janssen, J., Díaz-Caneja, C., Jiang, W., Calhoun, V., Ehrlich, S., Yang, K., Cascella,

N.G., Takayanagi, Y., Sawa, A., Tomyshev, A., Lebedeva, I., Kaleda, V., Kirschner, M., Hoschl, C., Tomecek, D., Skoch, A.,

van Amelsvoort, T., Bakker, G., James, A., Preda, A., Weideman, A., Stein, D.J., Howells, F., Uhlmann, A., Temmingh, H.,

López-Jaramillo, C., Díaz-Zuluaga, A., Fortea, L., Martinez-Heras, E., Solana, E., Llufriu, S., Jahanshad, N., Thompson,

P., Turner, J., van Erp, T., Glahn, D., Pearlson, G., Hong, E., Krug, A., Carr, V., Tooney, P., Cooper, G., Rasser, P., Michie, P.,

Catts, S., Gur, Raquel, Gur, Ruben, Yang, F., Fan, F., Chen, J., Guo, H., Tan, S., Wang, Z., Xiang, H., Piras, Federica,

Assogna, F., Salvador, R., McKenna, P., Bonvino, A., King, M., Kaiser, S., Nguyen, D., Pineda-Zapata, J., 2020. Increased

power by harmonizing structural MRI site differences with the ComBat batch adjustment method in ENIGMA.

NeuroImage 218, 116956. https://doi.org/10.1016/j.neuroimage.2020.116956

Raffa, M., Mechri, A., Othman, L.B., Fendri, C., Gaha, L., Kerkeni, A., 2009. Decreased glutathione levels and

antioxidant enzyme activities in untreated and treated schizophrenic patients. Prog. Neuropsychopharmacol. Biol.

Psychiatry 33, 1178–1183. https://doi.org/10.1016/j.pnpbp.2009.06.018

Rambaud, V., Marzo, A., Chaumette, B., 2022. Oxidative Stress and Emergence of Psychosis. Antioxid. Basel Switz.

11, 1870. https://doi.org/10.3390/antiox11101870

Ravera, S., Bartolucci, M., Cuccarolo, P., Litamè, E., Illarcio, M., Calzia, D., Degan, P., Morelli, A., Panfoli, I., 2015.

Oxidative stress in myelin sheath: The other face of the extramitochondrial oxidative phosphorylation ability. Free

Radic. Res. 49, 1156–1164. https://doi.org/10.3109/10715762.2015.1050962

Richter, S., Winzeck, S., Correia, M.M., Kornaropoulos, E.N., Manktelow, A., Outtrim, J., Chatfield, D., Posti, J.P.,

Tenovuo, O., Williams, G.B., Menon, D.K., Newcombe, V.F.J., 2022. Validation of cross-sectional and longitudinal

662 ComBat harmonization methods for magnetic resonance imaging data on a travelling subject cohort. Neuroimage

663 Rep. 2, 100136. https://doi.org/10.1016/j.ynirp.2022.100136

664 Rodríguez-Santiago, B., Brunet, A., Sobrino, B., Serra-Juhé, C., Flores, R., Armengol, L., Vilella, E., Gabau, E., Guitart,

665 M., Guillamat, R., Martorell, L., Valero, J., Gutiérrez-Zotes, A., Labad, A., Carracedo, A., Estivill, X., Pérez-Jurado, L.A.,

666 2010. Association of common copy number variants at the glutathione S-transferase genes and rare novel genomic

667 changes with schizophrenia. Mol. Psychiatry 15, 1023–1033. https://doi.org/10.1038/mp.2009.53

668 Ruszkiewicz, J.A., Miranda-Vizuete, A., Tinkov, A.A., Skalnaya, M.G., Skalny, A.V., Tsatsakis, A., Aschner, M., 2019. Sex-

669 Specific Differences in Redox Homeostasis in Brain Norm and Disease. J. Mol. Neurosci. 67, 312–342.

670 https://doi.org/10.1007/s12031-018-1241-9

671 Schilliger, Z., Alemán-Gómez, Y., Magnus Smith, M., Celen, Z., Meuleman, B., Binz, P.-A., Steullet, P., Do, K.Q., Conus, P.,

672 Merglen, A., Piguet, C., Dwir, D., Klauser, P., 2024. Sex-specific interactions between stress axis and redox balance

673 are associated with internalizing symptoms and brain white matter microstructure in adolescents. Transl.

674 Psychiatry 14, 1–11. https://doi.org/10.1038/s41398-023-02728-4

675 Smith, G.A., Lin, T.-H., Sheehan, A.E., Van der Goes van Naters, W., Neukomm, L.J., Graves, H.K., Bis-Brewer, D.M.,

676 Züchner, S., Freeman, M.R., 2019. Glutathione S-Transferase Regulates Mitochondrial Populations in Axons through

677 Increased Glutathione Oxidation. Neuron 103, 52-65.e6. https://doi.org/10.1016/j.neuron.2019.04.017

678 Smith, S.M., Jenkinson, M., Johansen-Berg, H., Rueckert, D., Nichols, T.E., Mackay, C.E., Watkins, K.E., Ciccarelli, O.,

679 Cader, M.Z., Matthews, P.M., Behrens, T.E.J., 2006. Tract-based spatial statistics: voxelwise analysis of multi-subject

680 diffusion data. NeuroImage 31, 1487–1505. https://doi.org/10.1016/j.neuroimage.2006.02.024

681 Stadtman, E.R., 2002. Importance of individuality in oxidative stress and aging. Free Radic. Biol. Med. 33, 597–604.

682 https://doi.org/10.1016/s0891-5849(02)00904-8

683 Thorburne, S.K., Juurlink, B.H., 1996. Low glutathione and high iron govern the susceptibility of oligodendroglial

684 precursors to oxidative stress. J. Neurochem. 67, 1014–1022. https://doi.org/10.1046/j.1471-

685 4159.1996.67031014.x

686 Tosic, M., Ott, J., Barral, S., Bovet, P., Deppen, P., Gheorghita, F., Matthey, M.-L., Parnas, J., Preisig, M., Saraga, M., Solida,

687 A., Timm, S., Wang, A.G., Werge, T., Cuénod, M., Quang Do, K., 2006. Schizophrenia and Oxidative Stress: Glutamate

688 Cysteine Ligase Modifier as a Susceptibility Gene. Am. J. Hum. Genet. 79, 586–592.

689 https://doi.org/10.1086/507566

690 Tsugawa, S., Noda, Y., Tarumi, R., Mimura, Y., Yoshida, K., Iwata, Y., Elsalhy, M., Kuromiya, M., Kurose, S., Masuda, F.,

691 Morita, S., Ogyu, K., Plitman, E., Wada, M., Miyazaki, T., Graff-Guerrero, A., Mimura, M., Nakajima, S., 2019.

692 Glutathione levels and activities of glutathione metabolism enzymes in patients with schizophrenia: A systematic

review and meta-analysis. J. Psychopharmacol. (Oxf.) 33, 1199–1214.
https://doi.org/10.1177/0269881119845820

Tustison, N.J., Avants, B.B., Cook, P.A., Zheng, Y., Egan, A., Yushkevich, P.A., Gee, J.C., 2010. N4ITK: improved N3 bias
correction. IEEE Trans. Med. Imaging 29, 1310–1320. https://doi.org/10.1109/TMI.2010.2046908

Uranova, N.A., Vikhreva, O.V., Rachmanova, V.I., Orlovskaya, D.D., 2011. Ultrastructural Alterations of Myelinated
Fibers and Oligodendrocytes in the Prefrontal Cortex in Schizophrenia: A Postmortem Morphometric Study.
Schizophr. Res. Treat. 2011, e325789. https://doi.org/10.1155/2011/325789

Uranova, N.A., Vikhreva, O.V., Rakhmanova, V.I., Orlovskaya, D.D., 2020. Dystrophy of Oligodendrocytes and Adjacent
Microglia in Prefrontal Gray Matter in Schizophrenia. Front. Psychiatry 11.

Uranova, N.A., Vikhreva, O.V., Rakhmanova, V.I., Orlovskaya, D.D., 2018. Ultrastructural pathology of
oligodendrocytes adjacent to microglia in prefrontal white matter in schizophrenia. Npj Schizophr. 4, 1–10.
https://doi.org/10.1038/s41537-018-0068-2

Veraart, J., Sijbers, J., Sunaert, S., Leemans, A., Jeurissen, B., 2013. Weighted linear least squares estimation of
diffusion MRI parameters: Strengths, limitations, and pitfalls. NeuroImage 81, 335–346.
https://doi.org/10.1016/j.neuroimage.2013.05.028

Wang, Y., Wang, Q., Haldar, J.P., Yeh, F.-C., Xie, M., Sun, P., Tu, T.-W., Trinkaus, K., Klein, R.S., Cross, A.H., Song, S.-K.,
2011. Quantification of increased cellularity during inflammatory demyelination. Brain 134, 3590–3601.
https://doi.org/10.1093/brain/awr307

Wijtenburg, S.A., Rowland, L.M., 2023. Chapter 19 - Schizophrenia spectrum disorders, in: Laule, C., Port, J.D. (Eds.),
Advances in Magnetic Resonance Technology and Applications, Imaging Neuroinflammation. Academic Press, pp.
469–487. https://doi.org/10.1016/B978-0-323-91771-1.00008-3

Williams, M. (Ed.), 2021. The Neuropathology of Schizophrenia. Springer International Publishing, Cham.
https://doi.org/10.1007/978-3-030-68308-5

Winkler, A.M., Ridgway, G.R., Webster, M.A., Smith, S.M., Nichols, T.E., 2014. Permutation inference for the general
linear model. NeuroImage 92, 381–397. https://doi.org/10.1016/j.neuroimage.2014.01.060

Wittchen, H.U., Jacobi, F., Rehm, J., Gustavsson, A., Svensson, M., Jönsson, B., Olesen, J., Allgulander, C., Alonso, J.,
Faravelli, C., Fratiglioni, L., Jennum, P., Lieb, R., Maercker, A., van Os, J., Preisig, M., Salvador-Carulla, L., Simon, R.,
Steinhausen, H.-C., 2011. The size and burden of mental disorders and other disorders of the brain in Europe 2010.
Eur. Neuropsychopharmacol. J. Eur. Coll. Neuropsychopharmacol. 21, 655–679.
https://doi.org/10.1016/j.euroneuro.2011.07.018

Xin, L., Mekle, R., Fournier, M., Baumann, P.S., Ferrari, C., Alameda, L., Jenni, R., Lu, H., Schaller, B., Cuenod, M., Conus,
P., Gruetter, R., Do, K.Q., 2016. Genetic Polymorphism Associated Prefrontal Glutathione and Its Coupling With
Brain Glutamate and Peripheral Redox Status in Early Psychosis. Schizophr. Bull. 42, 1185–1196.
https://doi.org/10.1093/schbul/sbw038

Yan, C., Duan, L., Fu, C., Tian, C., Zhang, B., Shao, X., Zhu, G., 2020. Association Between Glutathione S-Transferase
(GST) Polymorphisms and Schizophrenia in a Chinese Han Population. Neuropsychiatr. Dis. Treat. 16, 479–487.
https://doi.org/10.2147/NDT.S235043

Yao, J.K., Leonard, S., Reddy, R., 2006. Altered Glutathione Redox State in Schizophrenia. Dis. Markers 22, 83–93.
https://doi.org/10.1155/2006/248387

Zhang, Huali, Limphong, P., Pieper, J., Liu, Q., Rodesch, C.K., Christians, E., Benjamin, I.J., 2012.
Glutathione-dependent reductive stress triggers mitochondrial oxidation and cytotoxicity. FASEB J. 26, 1442–1451.
https://doi.org/10.1096/fj.11-199869

Zhang, Hui, Schneider, T., Wheeler-Kingshott, C.A., Alexander, D.C., 2012. NODDI: practical in vivo neurite
orientation dispersion and density imaging of the human brain. NeuroImage 61, 1000–1016.
https://doi.org/10.1016/j.neuroimage.2012.03.072

Zhang, Y., Catts, V.S., Shannon Weickert, C., 2018. Lower antioxidant capacity in the prefrontal cortex of individuals
with schizophrenia. Aust. N. Z. J. Psychiatry 52, 690–698. https://doi.org/10.1177/0004867417728805

Tables

<table>
<tr><td></td><td></td><th colspan="2">HC (N=86)</th><th colspan="2">PT (N=82)</th><th></th><th></th></tr>
<tr><td></td><td></td><th>Mean</th><th>Std.</th><th>Men</th><th>Std.</th><th>Stat.</th><th>p-value</th></tr>
<tr><td>Age (years)</td><td></td><td>28.1</td><td>8.6</td><td>29.8</td><td>9.6</td><td>W=3130</td><td>p=2.1e-01ns</td></tr>
<tr><td>GPx</td><td></td><td>16.7</td><td>6.6</td><td>16.3</td><td>6.2</td><td></td><td></td></tr>
<tr><td>GR</td><td></td><td>2.5</td><td>0.7</td><td>2.3</td><td>0.7</td><td></td><td></td></tr>
<tr><td>GPx/GR-ratio</td><td></td><td>6.9</td><td>2.7</td><td>7.4</td><td>3.3</td><td></td><td></td></tr>
<tr><td>Delay MRI-Biomarker (days)†</td><td></td><td>17.9</td><td>31.7</td><td>36.7†</td><td>44.5†</td><td>W=2249</td><td>p=4.4e-05****</td></tr>
<tr><td>Age at Psychosis Onset (years)</td><td></td><td></td><td></td><td>23.4</td><td>7.7</td><td></td><td></td></tr>
<tr><td>Duration of illness (months)</td><td></td><td></td><td></td><td>68.7</td><td>86.7</td><td></td><td></td></tr>
<tr><td>CPZ-equivalent dose (mg/day)</td><td></td><td></td><td></td><td>357.1</td><td>288.9</td><td></td><td></td></tr>
<tr><td>PANSS: total</td><td></td><td></td><td></td><td>63.5</td><td>14.9</td><td></td><td></td></tr>
<tr><td>PANSS: positive symptoms</td><td></td><td></td><td></td><td>13.6</td><td>4.4</td><td></td><td></td></tr>
<tr><td>PANSS: negative symptoms</td><td></td><td></td><td></td><td>17.2</td><td>5.7</td><td></td><td></td></tr>
<tr><td>PANSS: general psychopathology</td><td></td><td></td><td></td><td>33.0</td><td>8.0</td><td></td><td></td></tr>
<tr><td>GAF</td><td></td><td>84.9</td><td>4.1</td><td>54.9</td><td>11.8</td><td>W=6608</td><td>p=3.9e-28****</td></tr>
<tr><td></td><td></td><th>N</th><th>Pct.</th><th>N</th><th>Pct.</th><td></td><td></td></tr>
<tr><td>Scanner</td><td>Prisma</td><td>68</td><td>79.1</td><td>60</td><td>73.2</td><td>χ^2:0.81</td><td>p=0.37</td></tr>
<tr><td></td><td>Trio</td><td>18</td><td>20.9</td><td>22</td><td>26.8</td><td></td><td></td></tr>
<tr><td>Sex</td><td>Female</td><td>32</td><td>37.2</td><td>18</td><td>22.0</td><td>χ^2:4.68</td><td>p=0.031*</td></tr>
<tr><td></td><td>Male</td><td>54</td><td>62.8</td><td>64</td><td>78.0</td><td></td><td></td></tr>
<tr><td>Reader</td><td>Infinite</td><td>66</td><td>76.7</td><td>75</td><td>91.5</td><td>χ^2:6.742</td><td>p=0.009*</td></tr>
<tr><td></td><td>SPARK</td><td>20</td><td>23.3</td><td>7</td><td>8.5</td><td></td><td></td></tr>
<tr><td>GPx/GR-Ratio</td><td>High</td><td>44</td><td>51.2</td><td>40</td><td>48.8</td><td></td><td></td></tr>
<tr><td></td><td>Low</td><td>42</td><td>48.8</td><td>42</td><td>51.2</td><td></td><td></td></tr>
</table>

Table 1 Cohort demographics. P-values refer to Wilcox's tests between patients and healthy control. χ2 test is computed for the scanner, sex, and reader contingency tables. GPx and GR activities are expressed in nmoles NADPH used/min/g hemoglobin. HC: healthy controls, PT: patients, GPx: glutathione peroxidase, GR: glutathione reductase, GAF: Global Assessment of Functioning, PANSS: Positive and Negative Syndrome Scale. †: the delay MRI-biomarker statistics are reported excluding one PT participant with an extreme delay of 271 days.

Figures Captions

Figure 1

Biomarkers of the blood GSH-redox cycle and the effects of sex, age and clinical status. Correlations between GPx and GR activity in the whole cohort, colored by the quartiles of the ratio between GPx and GR (A), clinical status (B), and sex (C). Association between GPx and age (D) for both the entire cohort (shaded gray area) and by clinical status. GPx and GR activities are expressed in nmoles NADPH used/min/g hemoglobin. GPx: GSH peroxidase; GR: GSH reductase; HC: healthy controls; PT: patients; M: males; F: females.

Figure 2

Significant associations between biomarkers of the blood GSH-redox cycle and the dMRI microstructure estimates. (A, B): Brain plots show the significant association clusters between MK and GR activity ($p_{fdr} \leq 0.05$). GR was found significantly associated with MK in HC (A), with a slope that differed from the PT slope (B). (C): Brain plots show the significant clusters where the association of MK with GR in individuals with low GPx/GR-ratio yielded a different slope between HC and PT ($p_{fdr} \leq 0.05$). The regression plots show the association average values within significant clusters. Clusters were identified in the bilateral anterior and superior corona radiata, as well as in the genu and body of the corpus callosum and sections of the internal capsule. Results are corrected for quadratic age, sex, and delay. GR and $GR_{Low\text{-}ratio}$ are centered. Excluding the HC participant with MK < 0.80 does not change the results. Green: WM skeleton; HC: healthy controls; PT: patients; GR: glutathione reductase; MK: mean kurtosis; β_{HC}: slope of HC; β_{PT}: slope of the PT; Low-ratio: participants with GPx/GR-ratio < median ratio of the HC group (6.64).