Traumatic Amputations - A Nationwide Epidemiological Analysis of a developing country over 16 years

Authors: Marcella Moura Ceratti (MD)[1], Carolina Carvalho Jansen Sorbello (MD)[1], Isabela Roskamp Sunye (MD)[1], Felipe Soares Portela (MD)[1], Marcelo Fiorelli Alexandrino da Silva (MD)[1], Marcelo Passos Teivelis (MD)[1], Miguel Cendoroglo Neto (PhD)[123], Nelson Wolosker (PhD)[123].

Affiliations:

1: Hospital Israelita Albert Einstein, São Paulo, SP, Brasil.

2: Universidade de São Paulo - USP, Faculdade de Medicina, Departamento de Cirurgia Vascular e Endovascular, São Paulo, SP, Brasil.

3. Faculdade Israelita de Ciências da Saúde Albert Einstein, São Paulo, SP, Brasil.

ABSTRACT

Background:
Defining the impact of amputation is essential for developing cost-effective preventive health policies. Trauma is one of the most common causes of limb loss, affecting mainly the young working population. To date, few studies have investigated the epidemiology of patients undergoing trauma-related amputations and their public health implications in developing countries. The aim of this study was to analyze all limb amputations due to traumatic injuries performed in the Brazilian public health system over a 16-year period, studying their incidence, demographics, hospitalization and costs.

Objective:
To analyze the epidemiologic data on traumatic amputations within Brazil's public health system.

Methods:

This study was a cross-sectional and retrospective population-based analysis of traumatic amputations performed in the Brazilian public health system from 2008 to 2023. DATASUS, a public database of the Brazilian public health system, was used to select trauma cases, which were filtered by the traumatic amputation code. The dataset included the number of procedures, regional distribution, patient demographics, length of hospital stay, ICU stay, lethality and financial reimbursement.

Results:
There were 202,940 traumatic amputations in Brazil between 2008 and 2023. This condition was most common in males (78.7%), with an average age of 44.77 years, mainly involving fingers (62.7%). Cases involving lower limbs led to longer hospital stays and more ICU admissions. The mean length of hospital stay was 4.57 days, culminating in a lethality of 3.15%, which remained stable during the follow-up period. The estimated total cost to the Brazilian public health system was US$ 54,870,097.79, equivalent to approximately US$ 34,29381 spent per year on traumatic amputations, with the lower limb representing twice as much per patient.

Conclusion:
Traumatic amputation is still common in all regions of Brazil and has remained stable over the past 16 years, as has its lethality. Lesions leading to lower limb amputation account for the majority of hospital stays, intensive care unit admissions, and higher lethality and costs.

INTRODUCTION

Traumatic amputations, characterized by the abrupt loss of a limb due to external forces, represent a significant medical challenge with varied incidence rates across different regions and contexts. A study conducted at a Level I trauma center in North India reported a 2.5% incidence among trauma cases (1). In contrast, in the United States, the incidence of major lower extremity amputations resulting from trauma was 0.18% of all trauma admissions (2).

In the United States, trauma is the second most common cause of limb loss, following vascular disease (3). The incidence and prevalence of traumatic amputations rose by 16.4% and 49.2%, respectively, from 1990 to 2019. However, age-standardized incidence and prevalence rates have declined over the years (4). In Brazil, a study on lower limb amputations showed that trauma is decreasing as a cause of amputation in the country, decreasing from 12% of amputations in 2008 to 7% in 2020. At the same time, diabetes became the leading cause of amputations (27%), followed by gangrene (25%) and peripheral arterial disease (23%) (5).

The causes of traumatic amputations vary depending on the location and time period. During wars, for example, the primary causes involve improvised explosive devices, as seen in conflicts in Afghanistan and Iraq. However, in the civilian population on a global scale, the leading cause

of traumatic amputation before the age of 60 is exposure to mechanical forces. In contrast, after 60, the main cause is falls (4).

The lethality for traumatic amputations ranges from 6% to 13% in the literature (1,2). However, another significant problem with amputations in general is their morbidity and loss of functionality. Despite advances in adaptation techniques for individuals with amputated limbs, this type of injury continues to have a substantial impact on quality of life. The main implications of living with an amputated limb include complications of the surgical wound (such as infection and necrosis (6); pain (such as phantom limb pain in 10.9% to 50.9% of cases) (7); fractures due to osteoporosis and muscle atrophy (8); psychosocial distress and dependence on daily activities (9).

Although there are numerous studies analyzing the prevalence of amputations worldwide, no national epidemiological study has evaluated traumatic amputations in Brazil. Brazil is the largest country in Latin America with a population of 203,080,756 inhabitants (10), 71% of whom depend entirely on the Brazilian public health system (SUS - 'Sistema Único de Saúde') for healthcare services (11). All the information on SUS hospitalizations is available on a publicly accessible platform on the DATASUS (Departamento de Informação e Informática do SUS) website (12).

This study aimed to present the main epidemiological data on traumatic amputations in Brazil. Data from DATASUS, referring to the entire national territory, contain information on the number of traumatic amputations, body segment amputated, age, biological sex, lethality, hospital stay, need for an intensive care unit (ICU) and costs for the public health system.

METHODS

Study Design and Ethical Approval:

This is a cross-sectional and retrospective population-based study. The institution's Research Ethics Committee approved it (CAAAE 35826320.2.0000.0071). Since DATASUS is a public repository of anonymized data, informed consent was not required (12).

Data Source, Extraction Process, and Period:

Data were retrieved from DATASUS, the online database of the Brazilian Health System (SUS), which compiles information on government-funded hospitalizations nationwide. The focus was on trauma amputation procedures from 2008 to 2023. The extraction of data was automated using a Python-based protocol (v. 2.7.13; Beaverton, OR, USA) developed by our IT department, running under Windows 10. Selenium WebDriver (v. 3.1.8; Selenium HQ) and Pandas (v. 2.7.13;

Lambda Foundry, Inc. and PyData Development Team, NY, USA) tools were used for data segregation and refinement.

Study Population:

The average Brazilian population from 2008 to 2022 was 196,918,278 (10). Considering that 71% of the population uses exclusively the Brazilian public health service, this study's population sample was approximately 140,000,000 patients (11).

Selection Criteria and Extracted Data:

Data related to trauma cases were identified using ICD-10 codes (S00-S99 and T00-T98). From these, cases involving "disarticulation of the shoulder joint (scapulohumeral joint)", "amputation/disarticulation of the hand and wrist", "amputation/disarticulation of the upper limbs", "reimplantation from the shoulder to the mid-third of the forearm", "reimplantation from the distal third of the forearm to the metacarpals", "reimplantation or revascularization at the level of the hand and other fingers (except the thumb)", "reimplantation or revascularization of the thumb", "hip disarticulation (coxofemoral disarticulation)", "amputation/disarticulation of the lower limbs", "amputation/disarticulation of the foot and tarsus", "reimplantation from the thigh to the proximal third of the leg", "reimplantation from the mid-third of the leg to the foot", and "amputation/disarticulation of a finger". The corresponding DATASUS codes are, respectively: 0408010070, 0408020016, 0408020024,0408020253,040802026, 04080202701, 0408020288, 0408040106, 0408050012, 0408050020, 0408050306, 0408050314, 0408060042 . The dataset included annual procedure counts, regional distributions, patient demographics (age and biological sex), lengths of hospital stays, ICU durations, lethality and financial reimbursements. Financial data in Brazilian Reais (R$) was converted to U.S. dollars (U$), based on the median exchange rate from 2008 to 2023, amounting to U$ 1 = R$ 3,2018 (13).

Data Preparation:

Data compilation and tabulation were performed in .csv format using Microsoft Office Excel 2019 (Redmond, WA, USA). Population data by age group were obtained from the Brazilian Institute of Geography and Statistics (IBGE) (14). Statistical analyses were based on the population relying solely on the Brazilian public health system, which is approximately 140,000,000 patients (11).

Statistical Analysis:

Statistical analyses were conducted using SPSS version 20.0 for Windows (IBM Corp, Armonk, NY). Linear regression was employed to analyze trends in traumatic amputations across various age groups. Chi-square and likelihood ratio tests were used to evaluate differences in trauma

rates and lethality across the country's macro-regions. A significance level of $P \leq 0.05$ was set for all statistical tests.

RESULTS

From 2008 to 2023, there were 202,940 traumatic amputations. Among these, the majority (62.7%) were finger amputations, 29.6% lower limb amputations, and 7.7% upper limb amputations. The frequency of amputations remained practically stable over the years (**Figure 1**).

Figure 1: Number of traumatic amputations from 2008 to 2023.

Figure 2 shows the distribution of traumatic amputations across the five regions of Brazil. Most amputations occurred in the Northeast and Southeast regions. Adjusting for population size (10), the region with the highest incidence of traumatic amputations over 16 years was the south (122 amputations per 100,000 inhabitants), followed by the northeast (119 per 100,000 inhabitants), the center-west (116 per 100,000 inhabitants), the north (101 per 100,000 inhabitants) and finally the southeast (77 per 100,000 inhabitants). The proportion of finger amputations, lower limb amputations and upper limb amputations was nearly the same across all the country.

Figure 2: Amputations per region of Brazil.

Traumatic amputations were more common in males (78.7%). For finger amputations, males accounted for 82.6%; for upper limb amputations, 84.1%; and for lower limb amputations, 69%. The distribution of traumatic amputations by biological sex can be visualized in **Figure 3**.

Figure 3: Distribution of traumatic amputations per biological sex.

Traumatic amputations were more common between the ages of 20 and 59 (**Figure 4**). The mean age was 44,77 years. Cases of lower limb amputations peaked around the ages of 65-69 and again in patients aged 80 or over. In the pediatric age group, the vast majority of amputations were due to finger amputations.

Figure 4: Traumatic amputations per age group.

Figure 5 shows the number of hospitalization days for traumatic amputations of lower limbs, upper limbs, and fingers. Most patients stayed in the hospital for one day. Interestingly, patients with lower limb amputations stayed in the hospital for a longer duration, with most being hospitalized for 3 days. The mean duration of hospitalization was 4,57 days.

Days in hospital of Traumatic Amputation cases

Figura 5: Days in hospital of patients with traumatic amputations.

Most of the patients who required hospitalization in the Intensive Care Unit (ICU) were victims of traumatic lower limb amputations (**Figure 6**). However, the vast majority of patients (94%) in the study did not require intensive care. For those who needed ICU care, the duration of stay in the ICU was 5,85 days.

Figure 6: Days in the Intensive Care Unit (ICU) of patients with traumatic amputations.

The overall lethality of traumatic amputations was 3.15%.The highest lethality was in the lower limb amputation group (9.32%), followed by upper limbs (2.59%) and fingers (0.30%). The region with the highest lethality was the Southeast, followed by the Northeast (**Figure 7**). Lethality remained relatively stable from 2008 to 2023, as can be observed in **Figure 8**.

Figure 7: Lethality of traumatic amputations per region.

Figure 8: Lethality of traumatic amputations per year.

The Brazilian public health system spent a total of $54,870,097.79 on traumatic amputations from 2008 to 2023. Most of the expenses occurred in the Southeast and Northeast regions, which is consistent with the number of cases in these areas. Additionally, the treatment of traumatic lower limb amputations was more costly than other types of amputations. **Table 1** provides more details.

Topography of amputation	Region of Brazil					
	Midwest	Northeast	North	Southeast	South	Brazil
Upper Limb	$307.081,06	$947.674,84	$347.324,63	$1.289.719,95	$671.213,18	$3.563.013,65
Lower Limb	$2.310.901,69	$10.974.837,34	$1.917.519,35	$11.192.910,46	$6.243.049,15	$32.639.217,99
Fingers	$1.954.574,99	$5.253.346,35	$1.663.599,75	$6.303.489,39	$3.492.855,68	$18.667.866,15
Total	**$4.572.557,74**	**$17.175.858,52**	**$3.928.443,73**	**$18.786.119,79**	**$10.407.118,01**	**$54.870.097,79**

Table 1: Cost of Treatment for Traumatic Amputations by Region in Brazil

The overall cost per patient for traumatic amputations was $270.38. For lower limb amputations, this cost increased to $542.81. The graphic distribution of cost per patient is presented in **Figure 9.**

Figure 9: Cost per patient with traumatic amputations.

DISCUSSION

Limb loss is a major problem both at individual and public health levels, resulting in decreased quality of life, working years and dependance. Traumatic amputation is one of its most common causes and an especially relevant one in developing countries, such as Brazil, where almost 75% of the population rely uniquely on governmental health care (11,14).

Traumatic amputations are generally caused by serious trauma, such as high-energy traffic accidents, explosions, and machine injuries (15). In these cases, limb salvage is often not possible. A study using data from the German Trauma Society showed that 13.6% of patients with traumatic amputations were able to undergo replantation, while 86.4% suffered limb loss (16). The decision whether or not to replant a limb depends on multiple factors, including the

severity and extent of the injury, the presence of vascular lesions, the patient's haemodynamic status and the resources available in the care service (17–19).

Between 2008 and 2023, 202,940 amputations were performed in the Brazilian public health system as a consequence of trauma. In terms of yearly incidence rate, an average of 9.058 per 100 000 inhabitants/year was observed compared to a total of 13.23 million globally found in 2019 (4). The figures remained stable during the observed period despite the previously reported tendency of a significant increase in overall lower limb amputations in the country (5).

For a continental-sized country such as Brazil, the incidence of traumatic amputations in the last 16 years was surprisingly even among most regions. While geographically and economically distinct, the south, the northeast, and the center-west demonstrated similar rates, varying from 122 per 100,000 inhabitants in the first to 116 in the latter. Perhaps even more interestingly, the southeast, home to São Paulo and Rio de Janeiro, the country's two biggest cities (14), showed the lowest rate by far, with 77 amputations per 100,000 inhabitants. More studies will be necessary to clarify its reasons, but it may be explained by the more accessible health system in this region. The proportion of incidence of fingers or limb amputations also did not vary significantly across the country.

Regarding biological sex and age, the affected population had 4 times more men than women. Most were between 20 and 59 years, which aligns with previous reports (3–5,7,22). A possible explanation is partly occupational, with men often doing most of the high-risk physical and machinery work, in which accidents are more frequent. Young men also account for the majority of amputations caused by road traffic collisions. Together, those causes comprehend two of the most frequent etiologies leading to traumatic amputations (4).

These epidemiological results corroborate the impact of trauma on a country's economic structure. Apart from the direct costs of hospitalization and rehab, the injury, which predominantly affects the most economically active population, culminates in absenteeism from work and loss in productive capacity.

Regarding the site, amputation was found to be most common in fingers, as shown in prior studies (5,7,22). This is probably due to the vulnerability of the hand to violent wounds and the lack of muscle protection. Finger wounds were particularly relevant in the pediatric age group, where they accounted for the vast majority of amputations. Traumatic amputation of the upper limbs occurred more rarely (7.7 percent), with an incidence similar to that described in developed countries (23).

Lower limb amputations accounted for a third of the sample, were increasingly frequent in the geriatric population, and differed less between the sexes than other types of amputations. A similar epidemiology was observed in a global, multicentre study, apparently because falls are

the main mechanism of trauma in older adults (4). In addition, the presence of osteoporosis in this age group, which particularly affects women, can also potentiate the injury and lead to a more frequent need for amputation (4). Moreover, higher energy traumas are more likely to result in vascular injuries, carrying a worse prognosis (3,5,22). In the context of a rapidly aging country, the need for preventive measures adapted to different populations is evident.

The morbidity associated with leg lesions may help to explain why, in our study, victims of lower limb amputation required more intensive care admissions, longer hospital stays, and a higher lethality than the other patients. Regardless of advances in medical treatment, the overall lethality of 3,15, shown in our figures, remained relatively stable over the period studied. Nevertheless, the mortality rate is still lower than the 6% to 13% found in the literature (1,2).

As a universal public health system, the total cost of $54,870,097.79 spent in the period studied was covered in its totality by SUS, representing an expense of $34,29381 per year in traumatic amputations. Due to its severity, a patient with lower limb amputation was found to cost almost twice as much as the overall cost per patient ($542.81 to $270.38). It is also important to highlight that the sum does not include indirect hospital costs, rehab, or loss of working life years, which means that the total economic impact of this event for the country is probably higher.

This national epidemiological study, the first of its kind in Latin America, aims to present the main data on traumatic amputations in Brazil, including their impact on the national public health system. Similar to other types of trauma, identifying the patterns of these injuries is essential for their prevention. Therefore, our results can help guide strategies to reduce the significant impact and morbidity of traumatic amputations in our population.

Limitations of this study include those of population-based retrospective studies. In addition, as DATASUS is mainly based on epidemiological data from ICD-10 codes, the true number of traumatic amputations is likely to be underestimated due to coding errors. Finally, our analysis does not include private and supplementary healthcare systems, thus evaluating approximately three-fourths of the country's population (11).

CONCLUSION

Traumatic amputation in Brazil from 2008 to 2023 was most common among young men, affecting particularly the fingers but accounting for longer hospital stays and more ICU admissions when involving lower limbs. The mean duration of hospitalization was 4,57 days, culminating at a 3.15% lethality rate, which remained stable during the overlooked period. The total cost to the Brazilian public health system was $54,870,097.79, representing $34,29381 spent per year in traumatic amputations, with lower limb accounting for a figure twice as high per patient.

REFERENCES

1. Dhillon MS, Saini UC, Rana A, Aggarwal S, Srivastava A, Hooda A. The burden of post-traumatic amputations in a developing country - An epidemiological study from a level I trauma centre. Injury. 2022 Apr;53(4):1416–21.
2. Low EE, Inkellis E, Morshed S. Complications and revision amputation following trauma-related lower limb loss. Injury. 2017 Feb;48(2):364–70.
3. Sadoma BR, Sheets NW, Plurad DS, Dubina ED. Traumatic Amputations Treated in US Emergency Departments: A Review of the NEISS Database. Am Surg. 2023 Oct;89(10):4123–8.
4. Yuan B, Hu D, Gu S, Xiao S, Song F. The global burden of traumatic amputation in 204 countries and territories. Front Public Health. 2023;11:1258853.
5. Portela FSO, Louzada ACS, da Silva MFA, Teivelis MP, Kuzniec S, Wolosker N. Editor's Choice - Analysis of Lower Limb Amputations in Brazil's Public Health System over 13 Years. Eur J Vasc Endovasc Surg Off J Eur Soc Vasc Surg. 2024 Jul;68(1):91–8.
6. Potts JR, Wendelken JR, Elkins RC, Peyton MD. Lower extremity amputation: review of 110 cases. Am J Surg. 1979 Dec;138(6):924–8.
7. Takala J, Hämäläinen P, Sauni R, Nygård CH, Gagliardi D, Neupane S. Global-, regional- and country-level estimates of the work-related burden of diseases and accidents in 2019. Scand J Work Environ Health. 2024 Mar 1;50(2):73–82.
8. Finco MG, Kim S, Ngo W, Menegaz RA. A review of musculoskeletal adaptations in individuals following major lower-limb amputation. J Musculoskelet Neuronal Interact. 2022 Jun 1;22(2):269–83.
9. Hogan WB, Anderson G, Kovoor M, Alsoof D, McDonald CL, Zhang AS, et al. Phantom limb syndrome: Assessment of psychiatric and medical comorbidities associated with Phantom pain in 44,028 below knee amputees. Injury. 2022 Nov;53(11):3697–701.
10. Panorama do Censo 2022 [Internet]. [cited 2024 Jul 25]. Panorama do Censo 2022. Available from: https://censo2022.ibge.gov.br/panorama/
11. Agência de Notícias - IBGE [Internet]. 2020 [cited 2024 Aug 8]. PNS 2019: sete em cada dez pessoas que procuram o mesmo serviço de saúde vão à rede pública | Agência de Notícias. Available from: https://agenciadenoticias.ibge.gov.br/agencia-sala-de-imprensa/2013-agencia-de-noticias/releases/28793-pns-2019-sete-em-cada-dez-pessoas-que-procuram-o-mesmo-servico-de-saude-vao-a-rede-publica
12. DATASUS – Ministério da Saúde [Internet]. [cited 2024 Aug 12]. Available from: https://datasus.saude.gov.br/
13. Receita Federal [Internet]. [cited 2024 Aug 8]. Conversão de dólares para reais. Available from: https://www.gov.br/receitafederal/pt-br/assuntos/meu-imposto-de-renda/tabelas/conversao
14. IBGE | Portal do IBGE | IBGE [Internet]. [cited 2024 Aug 19]. Available from: https://www.ibge.gov.br/
15. Rankin IA, Nguyen TT, McMenemy L, Clasper JC, Masouros SD. The Injury Mechanism of Traumatic Amputation. Front Bioeng Biotechnol. 2021;9:665248.
16. Delhey P, Huber S, Hanschen M, Häberle S, Trentzsch H, Deiler S, et al. Significance of Traumatic Macroamputation in Severely Injured Patients: An Analysis of the Traumaregister

DGU®. Shock Augusta Ga. 2015 Mar;43(3):233–7.

17. Wolosker N, Guadêncio A, Kuzniec S, Rosoky RA, Kalume C, Neves CA, et al. Surgical treatment of noniatrogenic trauma of the femoral arteries. Sao Paulo Med J Rev Paul Med. 1996;114(1):1079–82.

18. Wolosker N, Nakano L, Anacleto MMM, Puech-Leão P. Primary utilization of stents in angioplasty of superficial femoral artery. Vasc Endovascular Surg. 2003;37(4):271–7.

19. Puech-Leão P, Kauffman P, Wolosker N, Anacleto AM. Endovascular grafting of a popliteal aneurysm using the saphenous vein. J Endovasc Surg Off J Int Soc Endovasc Surg. 1998 Feb;5(1):64–70.

20. Guimarães RA, de Sena KG, de Morais Neto OL, Malta DC. Magnitude and factors associated with motor road traffic injuries in Brazil: Results from the National Health Survey, 2019. Injury. 2023 Mar 11;S0020-1383(23)00244-9.

21. Murray J, Cerqueira DR de C, Kahn T. Crime and violence in Brazil: Systematic review of time trends, prevalence rates and risk factors. Aggress Violent Behav. 2013 Sep;18(5):471–83.

22. Moini M, Rasouli MR, Khaji A, Farshidfar F, Heidari P. Patterns of extremity traumas leading to amputation in Iran: results of Iranian National Trauma Project. Chin J Traumatol Zhonghua Chuang Shang Za Zhi. 2009 Apr;12(2):77–80.

23. Pomares G, Coudane H, Dap F, Dautel G. Epidemiology of traumatic upper limb amputations. Orthop Traumatol Surg Res OTSR. 2018 Apr;104(2):273–6.

Attachment

	Upper Limb Traumatic Amputations	Lower Limb Traumatic Amputations	Finger Traumatic Amputations	Total Traumatic Amputations
Total	15.598	60.130	127.212	202.940
Year				
2008	757	3.585	7.056	11.398
2009	909	3.627	6.968	11.504
2010	914	3.899	6.981	11.794
2011	1.020	4.439	7.098	12.557
2012	1.066	4.320	7.614	13.000
2013	1.127	3.913	8.128	13.168
2014	1.124	4.186	8.593	13.903
2015	976	3.615	8.194	12.785
2016	1.008	3.430	8.104	12.542

2017	1.003	3.527	7.824	12.354
2018	1.012	3.790	8.188	12.990
2019	1.043	3.610	8.668	13.321
2020	880	3.382	8.322	12.584
2021	922	3.315	8.159	12.396
2022	921	3.675	8.462	13.058
2023	916	3.817	8.853	13.586
Region				
Midwest	1.612	3.984	13.350	18.946
Northeast	4.858	23.833	36.277	64.968
North	1.905	4.214	11.478	17.597
Southeast	4.633	18.038	42.294	64.965
South	2.590	10.061	23.813	36.464
Sex				
Female	2.485	18.667	22.132	43.284
Male	13.113	41.463	105.080	159.656
Age group (years)				
00 - 04	447	320	5.296	6.063
05 - 09	370	326	3.782	4.478
10 - 14	477	409	3.416	4.302
15 - 19	1.110	1.828	7.892	10.830
20 - 24	1.589	2.959	10.640	15.188
25 - 29	1.545	2.899	10.237	14.681
30 - 34	1.567	2.715	10.281	14.563
35 - 39	1.481	2.753	10.377	14.611
40 - 44	1.356	3.012	10.490	14.858
45 - 49	1.353	3.394	10.650	15.397

50 - 54	1.182	4.204	10.232	15.618
55 - 59	1.013	5.195	9.537	15.745
60 - 64	769	5.945	7.951	14.665
65 - 69	502	6.150	6.079	12.731
70 - 74	370	5.789	4.378	10.537
75 - 79	209	4.824	2.861	7.894
≥ 80	258	7.408	3.113	10.779
Overall Lethality				
	2,59%	**9,32%**	**0,30%**	**3,15%**
Lethality per region				
Midwest	2,05%	8,86%	0,17%	2,16%
Northeast	3,19%	9,39%	0,44%	3,93%
North	1,63%	5,36%	0,22%	1,60%
Southeast	2,81%	10,30%	0,30%	3,25%
South	2,12%	9,20%	0,21%	2,82%
Lethality per year				
2008	3,30%	8,51%	0,24%	3,04%
2009	1,98%	8,88%	0,24%	3,10%
2010	1,53%	9,10%	0,39%	3,36%
2011	2,55%	9,91%	0,25%	3,85%
2012	2,44%	8,96%	0,28%	3,34%
2013	2,66%	9,38%	0,34%	3,23%
2014	2,31%	10,30%	0,29%	3,47%
2015	2,46%	9,18%	0,34%	3,00%

2016	2,38%	9,45%	0,31%	2,97%
2017	2,69%	9,95%	0,31%	3,25%
2018	2,27%	10,21%	0,31%	3,35%
2019	3,07%	9,17%	0,33%	2,94%
2020	3,30%	9,25%	0,30%	2,92%
2021	2,82%	9,59%	0,31%	2,98%
2022	3,04%	9,20%	0,32%	3,01%
2023	2,84%	7,91%	0,25%	2,58%