MEDICAL RISK CLASSIFICATION FOR SEVERE COVID-19 BASED ON CHRONIC MEDICAL CONDITIONS: A COMPARATIVE ANALYSIS

5083 words

Ilse Westerhof[1*], Annemarijn de Boer[1], Angela Lupattelli[2], Isabel Slurink[1], Nicolaas P.A. Zuithoff[1], Otilia Boldea[3], Hedvig Marie Egeland Nordeng[2,4], Jizzo R. Bosdriesz[5,6,7,8], Frank Pijpers[9,10], Maarten Schim van der Loeff[6,7,8], Mirjam Knol[5], Janneke van de Wijgert[1,5], Patricia Bruijning-Verhagen[1], and Ganna Rozhnova[1,11,12,13]

[1]Julius Centre for Health Sciences and Primary Care, University Medical Centre Utrecht, Utrecht University, Utrecht, The Netherlands
[2] Pharmacoepidemiology and Drug Safety Research group, Department of Pharmacy, University of Oslo, Oslo, Norway
[3]Department of Econometrics and OR, Tilburg School of Economics and Management, Tilburg University, Tilburg, The Netherlands
[4] Department of Child Health and Development, National Institute of Public Health, Oslo, Norway
[5]Department of Infectious Disease Control, National Institute for Diseases, Public Health and the Environment, Bilthoven
[6]Amsterdam UMC location University of Amsterdam, Infectious Diseases, Meibergdreef 9, Amsterdam, the Netherlands
[7]Amsterdam Institute for Immunology & Infectious Diseases, Amsterdam, the Netherlands
[8]Department of Infectious Diseases, Public Health Service (GGD) of Amsterdam, Amsterdam, Netherlands.
[9]Statistics Netherlands, The Hague, Netherlands
[10]Korteweg-de Vries Institute for Mathematics, University of Amsterdam, Amsterdam, The Netherlands
[11]Center for Complex Systems Studies (CCSS), Utrecht University, Utrecht, The Netherlands
[12]BioISI—Biosystems & Integrative Sciences Institute, Faculdade de Ciências, University of Lisbon, Lisbon, Portugal
[13]Faculdade de Ciências, University of Lisbon, Lisbon, Portugal
[*] Corresponding author: Ilse Westerhof; i.westerhof-4@umcutrecht.nl

Abstract (250 words)

Background: The European Centre for Disease Prevention and Control lists medical

conditions conferring high- or moderate-risk for severe COVID-19. Individual European

countries also developed risk classifications to guide COVID-19 vaccination

recommendations.

Objectives: To examine discordance between national and European risk classifications for

severe COVID-19 in the Netherlands and Norway across general, hospitalized, and deceased

populations, and investigate related factors.

Methods: This multi-country data-linkage study included 17.4 million inhabitants in the

Netherlands and 5.6 million in Norway in 2020. Medical conditions were defined using ICD-

10 codes in the European and Dutch classifications, and ICD-10 and ICPC-2 codes in the

Norwegian classification. Individuals were classified as high-, moderate-, or low-risk per

each classification. Performance was evaluated using sensitivity, specificity, positive and

negative predictive values (PPV, NPV), Brier score, and area under the curve (AUC).

Results: Discordance between classifications was 12–14% in the general population and 36-

47% among hospitalized or deceased individuals. The European classification assigned more

individuals to high-risk, while national classifications assigned more as moderate-risk.

National classifications demonstrated higher sensitivity, particularly in Norway, while the

European classification showed higher specificity. PPVs were ≤0.9% and NPVs near 100%

across all classifications. Norway's national classification had better discrimination and

calibration, while in the Netherlands, the European classification better predicted mortality.

Discordances frequently involved subjects with cardiovascular disease, lung disease, and

diabetes.

Conclusion: The European classification defines a substantially broader high-risk population

compared to national classifications. No single classification is universally optimal. Further

research should refine which chronic conditions best predict severe COVID-19.

Funding: ZonMw, EU, FCT, Norwegian Research Council's COVID-19 Emergency Call,

iAPOGEE.

Key words: Data linkage study | Severe COVID-19 | COVID-19 vaccination | Chronic
medical conditions | Medical risk classification

Introduction

The impact of public health measures, including COVID-19 vaccination, depends on the accurate identification of populations at elevated risk for severe outcomes, including hospitalization and death [1]. The World Health Organization recommends targeting vaccination programs to population groups at the highest risk of severe COVID-19 [1], but disparities exist in the risk classifications developed by different health authorities. While COVID-19 currently presents as common cold- or flu-like illness, older individuals and those with certain chronic medical conditions still face higher hospitalization and death rates compared to the general population [2,3]. Most risk classifications group individuals into high-, moderate-, and low-risk for severe COVID-19, based on factors like medical conditions and age. High-risk individuals have chronic conditions that lead to a much higher likelihood of severe outcomes, while those in the moderate-risk group face an elevated but lower risk.

Informed by influenza risk classifications and studies on risk factors for severe COVID-19, national (e.g., the Health Council of the Netherlands[4] and the Public Health Institute in Norway [5]) and international (e.g., the European Centre for Disease Prevention and Control [6]) public health authorities assigned high- or moderate-risk of developing severe COVID-19 to specific pre-existing medical conditions. These medical risk classifications have remained unchanged [6,7] or undergone minor alterations [4,5,8,9] since their development at the beginning of the pandemic. They have continued to guide epidemiological research and policy decisions on eligibility criteria and prioritization for vaccination programs aimed at reducing the incidence of severe COVID-19. While these international and national

classifications can be applied within the same country, it is unclear to what extent they differ in assigning medical risk status to individuals.

To inform evidence-based decision-making for severe COVID-19, we conducted a comparative analysis of risk status based on a national versus the European classification. This observational national registry-based cohort study used Dutch and Norwegian health and administrative data from 1 January 2020, linked at the individual level, including 17.4 million inhabitants of the Netherlands and 5.6 million inhabitants of Norway. We aimed to examine discordance between the respective national and European medical risk classifications for severe COVID-19 for the Netherlands and Norway across general, hospitalized, and deceased populations, and investigate related factors.

Methods

At the start of the COVID-19 pandemic, the Netherlands and Norway adopted national vaccination strategies, with the shared objective of minimizing severe COVID-19 outcomes such as hospitalization and death, reducing pressure on the health system, and preventing social disruption [10]. Individuals with chronic medical conditions associated with increased risk of severe COVID-19 were prioritized early in the national vaccination campaigns, which began in January 2021. In the Netherlands, high-risk individuals were identified through the International Statistical Classification of Diseases and Related Health Problems (10th revision; ICD-10) diagnostic codes [11]. The National Institute of Health and the Environment (RIVM) and Health Council defined the medical risk groups recommended for vaccination, including people with cardiovascular disease, diabetes, chronic lung or kidney disease, immunosuppression, cancer, and severe obesity [9,12]. The Norwegian Institute of Public Health (FHI) used both ICD-10 and International Classification of Primary Care (ICPC-2) [13] codes in the national health registries to define and flag individuals with moderate- or high-risk conditions for prioritized vaccination [5].

Data Collection

Data collection – The Netherlands

The Dutch dataset included hospital discharge registry (DHD) and the National Cause of Death Registry (NCDR) data for all inhabitants of the Netherlands as per the Dutch Population Registry on 1 January 2020, as well as individuals hospitalized and deceased due to COVID-19 in 2020. The three registries were linked using unique personal identification numbers. The Population Registry was used to extract age, immigration, and emigration dates. The DHD provided information on chronic medical conditions diagnosed in hospitals

since 1 January 1995, including the date of diagnosis. All diagnoses were registered according to the ICD-10 codes. From the NCDR, the date of death was extracted.

Data collection – Norway

The Norwegian dataset included data from multiple Norwegian registries for all inhabitants of Norway as per the National Population Registry on 1 January 2020, as well as individuals hospitalized and deceased due to COVID-19 in 2020. The registries included the Statistics Norway (SSB) registry, the Norwegian Patient Registry (NPR), and Norway Control and Payment of Health Reimbursements (KUHR) registry, and the Norwegian Cause of Death Registry (DAR). These registries were linked using unique personal identification numbers. The NPR and SSB registries were used to extract age, immigration, and emigration dates for all inhabitants of Norway. The KUHR registry provided information on medical diagnoses in primary and secondary outpatient care using ICPC-2 and ICD-10 codes, respectively. The NPR provided information on medical diagnoses in specialist outpatient and inpatient care, using ICD-10 coding. Both KUHR and NPR provided medical diagnosis data since 1 January 2010. DAR provided information on mortality.

Definitions of severe COVID-19

COVID-19 Hospitalization was defined as hospital admission for at least 24 hours and i) a primary diagnosis with ICD-10 code U071, U072, or U109; or ii) a secondary diagnosis with ICD-10 code U071, U072, or U109 combined with a primary code of respiratory illness (defined as any of the ICD-10 J-codes). This definition was based on prior research in Norway [14].

Death due to COVID-19 was defined as individuals with either i) COVID-19 listed as the primary cause of death (ICD-10 codes U071, U072, or U109 or ii) with a COVID-19 hospitalization in the 7 days preceding date of death. In the Dutch data registry, only the second definition was used, because cause of death was not available in death registry.

Risk classifications for severe COVID-19

The Dutch risk classification was obtained from the 'Adviesnota Vaststelling volwassen medische risicogroepen COVID-19-vaccinatiecampagne' ['Recommendation on establishing adult medical risk groups for the COVID-19 vaccination campaign'] in February 2023 [4]. This classification contains a list of chronic medical condition categories defined as either high- or moderate-risk for severe COVID-19. Each medical condition was independently coded using ICD-10 by two authors (AvB and IW), with no discrepancies found.

The Norwegian risk classification was obtained from the Norwegian Health Institute in July 2023 [5]. This scheme includes chronic medical condition categories with corresponding ICPC-2 and ICD-10 codes classified as either high-risk or increased-risk, which we renamed as high- and moderate-risk for consistency with other classifications.

The European risk classification was obtained from the 'Core protocol for ECDC studies of COVID-19 vaccine effectiveness against hospitalization with Severe Acute Respiratory Infection (SARI), laboratory-confirmed with SARS-CoV-2 or with seasonal influenza – Version 3.0' dated February 2024 [6]. This protocol provides ICD-10 codes for medical condition categories that are mandatory or optional for reporting in the ECDC studies. We classified the former medical condition categories as high-risk and the latter as moderate-risk.

Individuals in the Dutch or Norwegian datasets were assigned to have moderate- or high-risk chronic medical conditions if at least one relevant ICD-10 (both datasets) or ICPC-2 code (Norwegian cohort only) was recorded for that individual within the six months prior to 1 January 2020. For cancer diagnoses, records within five years prior to 1 January 2020 were used. These timeframes were defined according to the criteria specified in the classifications. Different medical risk classifications may use different codes for the same chronic medical condition, as each classification was developed with its own coding system. The ICD-10 and ICPC-2 codes used for this analysis are detailed in **Supplementary Tables 1-3**.

Individuals with one or more high-risk chronic medical condition categories, regardless of whether they also had one or more moderate-risk conditions, were defined as being at high-risk for severe COVID-19. Individuals with one or more moderate-risk chronic medical condition categories and no high-risk chronic medical condition were defined as being at moderate-risk for severe COVID-19. Individuals without any listed chronic medical condition categories were defined as low-risk. Pregnancy was not considered a criterion for categorizing individuals into the high- or moderate-risk groups in all classifications.

Statistical Analysis

Data for the Netherlands and Norway were analyzed separately. Descriptive analyses were performed to compare the respective national with the ECDC classification. The prevalence of high- and moderate-risk medical condition categories, and the most frequent combinations of medical condition categories, were shown in upset plots which visualize co-occurance of chronic medical condition categories in the population. Due to privacy reasons, numbers below 10 for the Dutch data and below 5 for the Norwegian data were not shown.

We reported both absolute numbers and proportions of individuals by 10-year age groups, classified into high-, moderate-, and low-risk groups for severe COVID-19. The discordance between the European and each national classification was calculated as the proportion of the population with a different European versus national risk status for the full national population, COVID-19 hospitalized population, and COVID-19 deceased population, and visualized with matrix plots. Different medical risk classifications may use different codes for the same chronic medical condition because they were formulated as such by the classifications. The 95% confidence intervals (CI) of discordance and concordance proportions were calculated using the Clopper-Pearson method [15]. Moreover, cumulative incidence of COVID-19 hospitalizations and deaths within the full national populations were presented.

Logistic regression models were applied for each classification to predict hospitalization and death from COVID-19 in each country. Individuals were dichotomized into high- or moderate-risk versus low-risk groups, and classification performance was evaluated using sensitivity, specificity, positive predictive value (PPV), and negative predictive value (NPV). Additionally, model performance was compared with Brier scores (i.e., mean squared errors) of the national and European classifications. Differences between these scores were tested using the Wilcoxon signed-rank test. Moreover, the Area Under the Curve (AUC) values were computed and statistically compared using DeLong's test to determine if there were significant differences.

To understand the factors related to discordance, we visualized the co-occurance of chronic medical condition categories in the discordant general, hospitalized, and deceased populations with upset plots. Also, to understand how coding differences affect overall discordance, we reviewed the most prevalent diseases in hospitalized and deceased

populations one by one by assessing the differences in population sizes between classifications. Since the Netherlands and Norway consider individuals above a certain age as high-risk for severe COVID-19, we also assessed the discordance in the general population in three scenarios: 1) the entire population regardless of age, 2) individuals younger than 60 years of age, aligning with the Dutch schemes [16], and 3) individuals younger than 65 years of age, aligning with the Norwegian schemes [5].

A sensitivity analysis was conducted to assess the importance of the reference date by evaluating the discordance in the general population for 1 January 2018, 1 January 2019, and 1 January 2021, instead of 1 January 2020.

Code Availability

The codes reproducing the results of this study are publicly available at https://github.com/IWesterhof/COVID-19-risk-classifications.

Data Availability

The aggregated tables of counts generated and analyzed in this study are publicly available at https://github.com/IWesterhof/COVID-19-risk-classifications.
Results are based on calculations by the authors using non-public microdata from Statistics Netherlands. Under certain conditions, these microdata are accessible for statistical and scientific research. For further information: microdata@cbs.nl.

Ethics

The analyses conducted with data from the Netherlands did not fall under the scope of the Dutch Medical Research Involving Human Subjects Act (WMO) and, therefore, did not require approval from an accredited ethics committee in the Netherlands. For the analyses conducted with data from Norway, the Regional Committee for Medical and Health Ethics of South/East Norway (no. 285687) approved the study. The Norwegian Data Protection Services for research and the University of Oslo approved the Data Protection Impact Assessment – DPIA (no. 341884).

Role of the funding source

The funders of the study had no role in study design, data collection, data analysis, data interpretation, or writing of the report. The corresponding author had full access to all the data in the study and had final responsibility for the decision to submit for publication.

Results

The Dutch cohort contained records of 17.4 million individuals (median age 42, interquartile range (IQR) 22-60), while the Norwegian cohort contained records of 5.6 million individuals (median age 39, IQR 21-57).

Comparison of risk classifications for severe COVID-19

Chronic medical condition categories included in the European, Dutch, and Norwegian risk classifications and their respective risk groups for severe COVID-19 are shown in **Table 1**.

Several chronic medical condition categories were consistently considered high-risk (e.g., cancer, immunodeficiency, and organ transplants) and moderate-risk (e.g., dementia) in the European and national classifications. There was some variability in the risk classifications of other conditions. Obesity was considered high-risk in the European classification but moderate-risk in both national classifications; lung disease, cardiovascular disease, and diabetes mellitus were considered high-risk in the European classification but moderate-risk in both national classifications. Hypertension was only listed in the European classification. Down syndrome and neurological disorders with impaired breathing were not included in the European classification but were considered high-risk in both national classifications.

Table 1. Comparison of European and national risk classifications for severe COVID-19. Chronic medical condition categories based on ICD-10 and ICPC-2 codes (see **Supplementary Tables 1-3**) and their respective risk groups for severe COVID-19 are shown. Different medical risk classifications may use different codes for the same chronic medical condition.

Chronic medical condition categories	Classification		
	European	Dutch	Norwegian
Cancer (solid organ and haematological)	orange	orange	orange
Immunodeficiency	orange	orange	orange
Organ transplant	orange	orange	orange
Overweight and obesity	orange	orange	blue
Lung disease	orange	blue	blue
Cardiovascular disease	orange	blue	blue
Diabetes mellitus	orange	blue	blue
Hypertension	orange	(blank)	(blank)
Kidney disease dialysis required	blue	orange	orange
Anaemia	blue	orange	(blank)
Kidney disease	blue	blue	orange
Liver disease/cirrhosis	blue	blue	orange
Neurological/muscular disorder (without impaired breathing)	blue	blue	orange
Dementia	blue	blue	blue
Tuberculosis	blue	blue	(blank)
Stroke	blue	(blank)	blue
Asplenia	blue	(blank)	(blank)
Rheumatologic disease	blue	(blank)	(blank)
Sickle cell disease	(blank)	orange	blue
Down syndrome	(blank)	orange	orange
Neurological disorder with impaired breathing	(blank)	orange	orange
Reduced immunity	(blank)	blue	blue
HIV infection	(blank)	blue	(blank)

Cochlear implants		

Risk due to chronic medical conditions   ▮ = High   ▮ = Moderate   ▢ = low

Classification of the general population by risk for severe COVID-19

Based on the national classifications, the most prevalent chronic medical condition categories in the Netherlands were lung disease, followed by cardiovascular disease, reduced immunity, and neurological/-muscular disorder; all classified as moderate-risk (**Figure 1 A**). In Norway, the most prevalent conditions were cardiovascular disease, lung disease, and diabetes mellitus; these were also classified as moderate-risk (**Figure 1 B**). Based on the European classification, the most prevalent chronic medical condition categories in the Netherlands were cardiovascular disease, hypertension, and cancer, and in Norway cardiovascular disease, diabetes mellitus, and cancer; all classified as high-risk (**Figure 1 C-D**). Differences in the prevalence between classifications arise from variations in the diagnosis codes to define a specific medical risk condition. For example, in the Netherlands, 0.54% of the population was classified as having cancer using the national classification, compared to 2.0% using the European classification because this classification included seven additional ICD-10 codes (**Supplementary Tables 1-2**). In Norway, 2.3% of the population was classified as having cancer using the national classification compared to 2.1% using the European classification due to differences in included codes.

In both countries and across all classifications, most individuals under 80 years of age were assigned low-risk for developing severe COVID-19 (**Figure 2**). Among children and adolescents (under 20 years old), 92.8% were classified as low-risk, 6.9% as moderate-risk, and 0.3% as high-risk, according to national schemes, while this was 98.5%, 0.4%, and 1.2%, respectively, according to the European classification. In Norway, the national classification

identified 98.5% as low-risk, 0.4% as moderate-risk, and 1.2% as high-risk, while this was

96.4%, 0.5%, and 3.1%, respectively, according to the European classification. In both

countries and across all classifications, the percentage of individuals in moderate- and high-

risk groups increased with age, reaching 41.1% and 53.6% for those over 80 years in the

national classifications, and 41.8% and 37.7% in the European classifications in the

Netherlands and Norway, respectively.

In the Netherlands and Norway, risk classification differed for 12.0% and 13.8% of the

general population, respectively, between the national and the European classification

schemes (**Figure 3 A-B**). The European classification assigned more people as high-risk (9%

in the Netherlands, 9.2% in Norway) compared to national classifications (1.5% and 3.0%,

respectively). Conversely, the national classifications identified more people as moderate-risk

(11.3% and 12.7%, respectively) than the European classification (1.0% and 1.3%,

respectively). According to the national classifications, 87.2% of the population in the

Netherlands and 84.4% in Norway were classified as low-risk. These percentages were

slightly higher using the European classification (90.1% and 89.5%, respectively).

Discordance for population with severe COVID-19 in 2020

The Netherlands had 39,063 hospitalizations (median age 70 years [IQR 58-78]) and 7,031

deaths (median age 78 years [IQR 71-83]). In Norway, there were 2,228 hospitalizations

(median age 64 years [IQR: 51-78]) and 514 deaths (median age 84 years [IQR: 75-90]).

The cumulative incidence of COVID-19 hospitalizations and deaths increased with higher

risk levels for both the national and European classifications (**Table 2**). For example, the

cumulative incidence of hospitalizations was 0.15% for the low-risk group, 0.68% for the

moderate-risk group, and 0.14% for the high-risk group for the national classification

compared to 0.15%, 0.52%, and 0.94%, respectively, for the European classification.

Similarly, the cumulative incidence of COVID-19 deaths was 0.02% for low-risk, 0.17% for

moderate-risk, and 0.29% for high-risk groups for the national classification compared to

0.02%, 0.13%, and 0.24%, respectively, for the European classification.

Cardiovascular disease, classified as high-risk in the European classification and moderate-

risk in the national classifications, is the most prevalent chronic condition among hospitalized

and deceased individuals (**Supplementary Figure 1-2**). Another important disease that is

prevalent and differently classified is Diabetes Mellitus.

For all ages and both countries, and similar to the general population, the European

classification assigns more individuals to high-risk, while national classifications define more

individuals as moderate-risk (**Supplementary Figures 3-4**). Discordance was even higher

among hospitalized (the Netherlands 36.5% and Norway 38.2%), and deceased (the

Netherlands 46.6% and Norway 45.1%) populations, compared to the general population

(discordance in the Netherlands 12.0% and Norway 13.8%), as shown in **Figure 3C-F.**

Moreover, a large part of the hospitalized (51.5% in the Netherlands and 49.5% in Norway)

and deceased (40.2% and 36.2%, respectively) populations was defined as low-risk in both

classifications, which means these individuals had none of the chronic medical conditions

listed in classifications.

National classifications showed consistently higher sensitivity than European classifications

(**Table 3**). For example, in Norway, sensitivity for hospitalization was 45.6% for the national

classification versus 32.9% for the European classification; for COVID-19 deaths, sensitivity

Table 2: Cumulative COVID-19 hospitalizations and deaths in 2020. Risk groups were defined on January 1, 2020. Individuals born after January 1, 2020, are not included.

	The Netherlands					Norway				
	General population	Hospitalization		Death		General population	Hospitalization		Death	
Classification	n	n	Cumulative Incidence	n	Cumulative Incidence	n	n	Cumulative Incidence	n	Cumulative Incidence
National classification										
- *Low-risk*	15,171,220	22230	0.15%	3004	0.02%	4,708,944	1170	0.03%	199	0.00%
- *Moderate-risk*	1,973,422	13444	0.68%	3285	0.17%	706,399	786	0.11%	225	0.03%
- *High-risk*	257,637	3068	1.19%	742	0.29%	165,361	272	0.16%	90	0.05%
European classification										
- *Low-risk*	15,671,292	23174	0.15%	3069	0.02%	4,995,283	1433	0.03%	258	0.01%
- *Moderate-risk*	165,325	865	0.52%	208	0.13%	72,122	71	0.10%	24	0.03%
- *High-risk*	1,565,662	14703	0.94%	3754	0.24%	513,299	724	0.14%	232	0.05%

Table 3: Performance COVID-19 classifications for severe COVID-19. Best performing risk-classification in bold.

	National classification				European classification				Brier score/MSE (SD)			AUC (95% CI)		
	Sens	Spec	PPV	NPV	Sens	Spec	PPV	NPV	National class.	European class.	P-value	National class.	European class.	P-value
The Netherlands														
Hospitalizations	**42.2%**	87.2%	0.70%	99.9%	39.7%	**90.1%**	**0.90%**	99.9%	0.002132 (0.04585)	**0.002131 (0.04582)**	0.38	0.649	**0.650**	0.13
Deaths	**57.3%**	87.2%	0.18%	>99.9%	56.4%	**90.1%**	**0.23%**	>99.9%	0.000404 (0.02005)	**0.000403 (0.02004)**	<0.001	0.725	**0.733**	<0.001
Norway														
Hospitalizations	**45.6%**	84.4%	0.12%	>99.9%	32.9%	**89.5%**	0.12%	>99.9%	**0.00039395 (0.01982)**	0.00039398 (0.01983)	<0.001	**0.651**	0.612	<0.001
Deaths	**63.2%**	84.4%	0.04%	>99.9%	47.1%	**89.5%**	0.04%	>99.9%	**0.0000920796 (0.00959)**	0.0000920824 (0.00959)	<0.001	**0.741**	0.684	<0.001
Abbreviations: Sens: Sensitivity, Spec: Specificity, PPV: Positive Predicted Value, NPV: Negative Predicted value, MSE: Mean Squared Error, AUC: Area Under the Curve														

was 63.2% versus 47.1%, respectively. Specificity was always higher for the European

classification. PPV and NPV were comparable across classifications, with very low PPVs

(≤0.9%) and NPVs near 100%. The Brier scores for all classifications were below 0.01,

consistent with the low incidence of outcomes and the correspondingly low predicted risks.

Discrimination and calibration metrics (Brier score and AUC) favored the national

classification in Norway (both $p<0.001$), while in the Netherlands, no significant differences

were seen for hospitalizations ($p\geq0.13$), and the European classification performed better for

COVID-19 mortality ($p<0.001$).

Comparison between the Norwegian and Dutch classifications for the hospitalized and

deceased populations in Norway shows a discordance of 17.1% and 21.8%, respectively

(**Supplementary Figure 5**). This is mainly caused by differences in selecting diseases (ICD-

19 and ICPC-2 codes) within the disease categories lung disease, cardiovascular disease, and

dementia.

Discordance between risk classifications and the related factors

Discordance was higher in elderly because the number of individuals with one or more

medical risk conditions was higher than in younger age groups (**Figure 3-4; Supplementary**

Figure 6). **Figure 4** shows the chronic medical condition categories in the general population

who were classified differently by the European and respective national classification

schemes (the discordant population). In the discordant population in the Netherlands, 23.5%

had neurological/-muscular disorders, 23.8% reduced immunity, and 13.6% kidney disease

according to the national classification (**Figure 4A**). In the discordant population in Norway,

6.7% had reduced immunity, and 2.3% neurological/muscular disorders, classified as

moderate- and high-risk, respectively (**Figure 4B**). On the contrary, the European

classification included chronic medical condition categories that were not included in the

national classifications. In the Netherlands, examples were hypertension and kidney disease

with prevalences of 25.5% and 6.3% in the discordant population. In Norway, 8.2% of the

discordant population had hypertension and 3.6% had kidney disease according to European

schemes, but not included in national schemes. Unlike Norway, none of the discordant

individuals in the Netherlands were classified as overweight/obese or diabetic based on the

European classification. Similar was observed for the hospitalized and deceased discordant

populations (**Supplementary Figures 7-8**).

Some of the discordance in the most prevalent chronic medical condition categories (lung

disease, cardiovascular disease, diabetes mellitus, cancer, and kidney disease) among

hospitalized and deceased populations may stem from differences in ICD-10 and ICPC-2

codes. In the Netherlands, the national classification identified more individuals with lung

disease and diabetes, while the Norwegian system captured more cases across all disease

categories except kidney disease, indicating that it includes more ICD-10 codes

(**Supplementary Table 5**). The European classification identified more individuals with

cardiovascular disease and cancer in the Netherlands and more individuals with kidney

disease in Norway. Notably, in the Netherlands, cancer was better captured with the

European classification, identifying more cases in the general (2.0% versus 0.5%),

hospitalized (9.1% versus 4.1%), and deceased (10.8% versus 4.1%) populations compared to

the Dutch classification. Similarly, in Norway, the European classification identified more

individuals with kidney disease in the general (0.8% versus 0.2%), hospitalized (10.1%

versus 2.4%), and deceased (13.3% versus 2.3%) populations compared to the national

classification.

In the Netherlands, the most common chronic medical condition categories varied by age

group (**Supplementary Figures 9-10**). In those younger than 60 or 65, lung disease was the

most prevalent condition, followed by neurological/-muscular disorder and cardiovascular

disease. In Norway, the national classification showed no differences between all ages and

individuals younger than 65 and 60 years of age in most pronounced chronic conditions. In

the Netherlands, based on the European classification there was no change in the order of

most prominent conditions contributing to discordance in the population aged below 60 years

compared to all ages. In contrast, for those aged below 65 years, the most prevalent

conditions were cardiovascular disease, diabetes mellitus, and lung disease, but not

hypertension and cancer.

Sensitivity analyses

No meaningful differences in discordance percentages were observed when using different

reference dates to define the study populations (data not shown).

Discussion

In this multi-country, nationwide registry-linkage study, we compared the respective national

with the European medical risk classification for severe COVID-19, and classified the

populations of the Netherlands and Norway according to this risk, across different age

groups. For clarity, we reiterate that individuals with at least one high-risk chronic condition

were classified as high-risk, those with only moderate-risk conditions as moderate-risk, and

those without listed conditions as low-risk (see Methods for details). Our findings revealed

substantial differences between the national and European risk classifications in the general

population (discordance in the Netherlands 12.0% and Norway 13.8%), hospitalized patients

(the Netherlands 36.5% and Norway 38.2%), and deceased patients (the Netherlands 46.6%

and Norway 45.1%), with the European classification scheme assigning a broader range of

chronic medical condition categories to the high-risk category than the two national

classifications. Consequently, a larger proportion of the population is classified as high-risk

when using the European classification compared to the national classifications. Conversely,

a larger proportion of the population is classified as moderate-risk when using the respective

national classifications compared to the European classification. The Dutch national

classification may be underinclusive for cancer and cardiovascular disease, and the

Norwegian national classification for kidney disease, while those disease categories are

probably more clinically relevant as they are prevalent among hospitalized and deceased

populations. Interestingly, a large part of the hospitalized (51.5% in the Netherlands and

49.5% in Norway) and deceased (40.2% and 36.2%) populations were defined as low-risk in

both national and European classifications. This may be due to chronic medical conditions

not captured by either classification. This suggests that the classifications differ even more

when applied to severe COVID-19 outcomes and fail to identify a considerable share of

affected individuals. Moreover, our results indicate that the national classifications provide

better sensitivity, while the European classifications offer better specificity. Additionally, the

accuracy and discrimination of individuals at risk for severe COVID-19 outcomes is greater

for the national classification in Norway; however, this was not demonstrated for the national

classification in the Netherlands. Notably, none of the classifications showed significantly

better accuracy or discrimination for predicting hospitalizations. In contrast, the European

classification was significantly better in identifying individuals at risk of death due to

COVID-19.

These discrepancies may have implications for COVID-19 vaccination programs, which

prioritize specific risk populations. Although primarily designed for research focusing on

COVID-19 vaccine effectiveness and enhancing SARI surveillance in Europe, the European

protocols have influenced public health policy in the past, as demonstrated, for example, with

influenza vaccination strategies [17]. However, we recognize that this classification is

primarily surveillance-oriented and not an officially mandated European clinical standard.

While not a universal policy tool, this classification provides a structured framework that can

support public health decision-making alongside other clinical and epidemiological

considerations. Therefore, the European classification for severe COVID-19 could remain

relevant for broader public health decisions and policy. Our comparative analysis suggests

that using the European classification, 9.0% of the population in the Netherlands and 9.2% of

the population in Norway are high-risk due to chronic medical condition categories and

should be vaccinated, while these percentages are only 1.5% and 3.0%, respectively, when

using the national classifications. Vaccination programs that prioritize both moderate- and

high-risk groups, the European classification would result in 10.0% of the Dutch population

and 10.5% of the Norwegian population having to be vaccinated, compared to 12.8% and

13.7%, respecitively, if using the respective national classifications. Moreover, chronic

medical condition categories are defined based on different ICD-10 and ICPC-2 codes in the

three classifications, resulting in differences in prevalence of individual medical risk

categories across classifications. Therefore, the choice between the European or national

classifications may have important consequences for vaccination prioritization and coverage,

potentially influencing the impact of vaccination campaigns in preventing severe COVID-19.

In our analyses, the national classification performed better in Norway for both COVID-19

hospitalization and mortality, while in the Netherlands, the European classification performed

better for mortality, with no clear differences observed for hospitalization. Although the

statistical performance measures provide useful comparison, they highlight that no single

classification is universally optimal. Sensitivity was consistently higher for national

classifications, indicating better identification of individuals at high- or moderate-risk of

severe COVID-19 outcomes. In contrast, the European classification demonstrated higher

specificity, more accurately excluding low-risk individuals who did not experience severe

outcomes. Given the potential severity of COVID-19 and the high effectiveness of vaccines

alongside the rarity of serious adverse events [18], it may be preferable to prioritize

sensitivity when designing vaccination policies to ensure that individuals with chronic

medical conditions are not overlooked.

The dominant medical condition categories generating the discordance in risk status were

lung and cardiovascular disease in both Norway and the Netherlands, neurological disorders

in the Netherlands, and diabetes in Norway. Hypertension was one of the most pronounced

discordant chronic condition categories between classifications. According to the European

classification, but not to the national ones, hypertension was considered a high-risk disorder

for severe COVID-19, and its prevalence estimate was 3.5% in the Netherlands and 1.4% in

Norway. Several studies have indicated that hypertension was not an independent predictor of

severe COVID-19 [19,20]. Therefore, the high-risk population might be overestimated in the

European classification. Lung disease, cardiovascular disease, and diabetes mellitus were also

among the most pronounced medical condition categories leading to discordance, which were

diagnosed in 1.7-4.6% of the population according to the national and European

classifications. These conditions were classified as moderate-risk in the national

classifications but as high-risk in the European classification. Although multiple studies have

shown an increased risk of severe COVID-19 in individuals with these medical condition

categories [21–23], it is unknown whether they should be part of the high- or moderate-risk

group. Interestingly, the European classification does not include Down syndrome and

neurological/-muscular disorder as risk factors for severe COVID-19, while the national

classifications consider them as moderate- or high-risk chronic conditions. Several studies

have demonstrated an increased risk of COVID-19 hospitalization and death among

individuals with Down syndrome [24,25] and multiple sclerosis [26,27]. However, our

findings show that only a very small fraction of the discordant population has Down

syndrome (≤0.2% in both countries), while those with neurological/muscular disorders make

up a larger proportion—23.5% in the Netherlands and 2.3% in Norway. This supports the

need to assess the discordant conditions in relation to severe COVID-19 to ensure vaccination

prioritization and coverage of individuals at high-risk.

Our study has several limitations. First, the available data did not allow us to assess how well

the classifications predict severe COVID-19 outcomes after 2020, when vaccination became

available and reinfections were frequently observed. As individuals were prioritized for

vaccination based on the national guidelines which align with the data used in our study, this

would introduce bias. Second, due to limited testing capacity in 2020, we were unable to

clearly identify exposed individuals and analyze their risk of severe COVID-19 outcomes

upon infection. Therefore, we assessed the risk of severe COVID-19 for the entire national

population rather among those infected. Lastly, a limitation specific to the Norwegian data is

the absence of ICD-10 codes for asplenia in the registries. However, since asplenia is a very

rare condition, estimated at 0.44 cases of surgical asplenia per 100,000 person-years in

Norway, this limination is unlikely to affect our overall conclusions [28].

Future research should aim to establish a consensus on which chronic medical condition

categories are universally recognized as conferring moderate- and high-risk for severe

COVID-19, especially in the post-pandemic era, where the population has substantial

immunity due to vaccination and infections. We propose a matched multi-country cohort

study to estimate the risk of hospitalization and death due to COVID-19 in individuals with

and without specific chronic conditions, accounting for confounders like age, vaccination

status, and circulating variants. Additionally, the importance of chronic medical conditions in

the hospitalized and deceased populations that are not included in the national and European

classifications, and therefore classified as low-risk, should be further assessed. This will help

understand which additional conditions should be included in a risk classification for severe

COVID-19. The analyses should ideally be conducted at the diagnostic code level, as our

findings highlight substantial differences between classifications for specific disease

categories. This approach will also help determine whether certain ICD-10 and ICPC-2 codes

are overly broad or clinically unsubstantiated. This research will be crucial for deciding

which classification most accurately captures the population at highest risk. Moreover,

mathematical modeling studies can help to understand the impact of different risk

classifications and optimal vaccination strategies [29–32]. Incorporating risk stratification

into modeling studies has the potential to design more effective strategies, a consideration

often overlooked in current research efforts. Furthermore, exploring emerging data sources

and digital health technologies could improve the identification and monitoring of moderate-

and high-risk populations.

Conclusion

There are substantial differences between the European and the respective national medical

risk classifications for severe COVID-19 based on pre-existing chronic medical condition

categories, which may have important implications for vaccination strategies. The European

classification suggests that a larger proportion of the population is at high-risk for severe

COVID-19 and may require vaccination. Classification performance showed that no single

classification is universally optimal. Further research should identify which chronic

conditions are associated with risk of developing severe COVID-19 and should be included

in risk classifications.

Author Contributions

Conceptualization: I Westerhof, G Rozhnova. Data management: I Westerhof, A de Boer.

Methodology: I Westerhof, A de Boer, G Rozhnova. Input statistical methodology: N

Zuithoff. Investigation: I Westerhof, A de Boer. Formal analysis: I Westerhof, A de Boer.

Visualization: I Westerhof. Writing—original draft: I Westerhof, A de Boer, G Rozhnova.

Writing—review and editing: I Westerhof, A de Boer, A Lupattelli, I Slurink, N Zuithoff, O

Boldea, HME Nordeng, JR Bosdriesz, F Pijpers, M Schim van der Loeff, M Knol, JHHM van

de Wijgert, P Bruijning, and G Rozhnova.

Declaration of interests

N. Zuithoff was involved in studies financed by Sanofi, Regeneron, Eli Lilly, Leo Pharma,

Abbvie, and Pfizer; fees were paid directly to the institution.

Acknowledgments

We are grateful to all the participating families in the Netherlands and Norway who are part

of the health registries and made this research possible. GR, IW, AdB, and OB were

supported by the ZonMw project (10430362220002). GR was supported by Fundação para a

Ciência e a Tecnologia project 2022.01448.PTDC, DOI 10.54499/2022.01448.PTDC. GR,

IW, AL, and PB were supported by the VERDI project (101045989), funded by the European

Union. Views and opinions expressed are however those of the author(s) only and do not

necessarily reflect those of the European Union or the Health and Digital Executive Agency.

Neither the European Union nor the granting authority can be held responsible for them. IW

was supported by the scholarship program iAPOGEE "International Alliance for

PharmacoGenetic Epidemiology Excellence" at the University of Oslo, Norway. Data

307   acquisition of the Norwegian data in this study was funded by HN's EU-COVID-19 project,

308   funded by the Norwegian Research Council's COVID-19 Emergency Call (project no.

309   31270). The work on the Norwegian data was performed on the TSD (Tjeneste for Sensitive

310   Data) facilities, owned by the University of Oslo, operated and developed by the TSD service

311   group at the University of Oslo, IT-Department (USIT) (tsd-drift@usit.uio.no).

313 Declaration of generative AI in scientific writing
314 During the preparation of this work the author(s) used Chat GPT in order to improve

315 readability and language. After using this tool/service, the author(s) reviewed and edited the

316 content as needed and take(s) full responsibility for the content of the publication.

References

[1]   World Health Organization. Global Covid-19 Vaccination Strategy in a Changing World July 2022 update. 2022.

[2]   Antos A, Kwong ML, Balmorez T, Villanueva A, Murakami S. Unusually High Risks of COVID-19 Mortality with Age-Related Comorbidities: An Adjusted Meta-Analysis Method to Improve the Risk Assessment of Mortality Using the Comorbid Mortality Data. Infect Dis Rep 2021;13:700–11. https://doi.org/10.3390/idr13030065.

[3]   Zhang J, Dong X, Liu G, Gao Y. Risk and Protective Factors for COVID-19 Morbidity, Severity, and Mortality. Clin Rev Allergy Immunol 2022;64:90–107. https://doi.org/10.1007/s12016-022-08921-5.

[4]   Gezondheidsraad. Adviesnota Vaststelling volwassen medische risicogroepen COVID-19-vaccinatiecampagne. 2023.

[5]   Norwegian Institute of Public Health. Coronavirus vaccine 2023. https://www.fhi.no/en/id/corona/coronavirus-immunisation-programme/coronavirus-vaccine (accessed April 17, 2024).

[6]   European Centre for Disease Prevention and Control. Core protocol for ECDC studies of COVID-19 vaccine effectiveness against hospitalisation with Severe Acute Respiratory Infection, laboratory-confirmed with SARS-CoV-2 or with seasonal influenza - Version 3.0. 2024.

[7]   European Centre for Disease Prevention and Control. Core protocol for ECDC studies of COVID-19 vaccine effectiveness against hospitalisation with Severe Acute Respiratory Infection laboratory-confirmed with SARS-CoV-2, version 1.0. Stockholm: 2021.

[8]   Norwegian Institute of Public Health. Coronavirus vaccine 2020. https://www.fhi.no/en/id/corona/coronavirus-immunisation-programme/coronavirus-vaccine (accessed September 1, 2022).

[9]   Gezondheidsraad. Advies Strategieën voor COVID-19-vaccinatie. 2020.

[10]   Kamerbrief over strategie vaccinatie tegen COVID-19 (coronavirus) | Kamerstuk | Rijksoverheid.nl n.d. https://www.rijksoverheid.nl/documenten/kamerstukken/2020/11/20/kamerbrief-over-covid-19-vaccinatiestrategie (accessed June 10, 2025).

[11]   World Health Organization. International Statistical Classification of Diseases Related Health Problems 10th Revision Geneva. Https://IcdWhoInt/Browse10/2019/En 2019.

[12]   Rijksinstituut voor Volksgezondheid en Milieu. The National Immunisation Programme in the Netherlands. Surveillance and developments in 2021-2022. 2021.

[13]   World Health Organization. International Classification of primary care, 2nd edition (ICPC-2). Https://WwwWhoInt/Standards/Classifications/Other-Classifications/International-Classification-of-Primary-Care 2024.

[14]   Whittaker R, Toikkanen S, Dean K, Lyngstad TM, Buanes EA, Kløvstad H, et al. A comparison of two registry-based systems for the surveillance of persons hospitalised with COVID-19 in Norway, February 2020 to May 2022. Eurosurveillance 2023;28. https://doi.org/10.2807/1560-7917.ES.2023.28.33.2200888.

[15]   Clopper CJ, Pearson ES. The Use of Confidence or Fiducial Limits Illustrated in the Case of the Binomial. Biometrika 1934;26:404. https://doi.org/10.2307/2331986.

[16]   Rijksinstituut voor Volksgezondheid en Milieu. COVID-19 vaccination 2024. https://www.rivm.nl/en/coronavirus-covid-19/vaccination (accessed April 22, 2024).

[17]   Mereckiene J, Cotter S, Nicoll A, Lopalco P, Noori T, Weber JT, et al. Seasonal influenza immunisation in Europe. Overview of recommendations and vaccination

coverage for three seasons: pre-pandemic (2008/09), pandemic (2009/10) and post-
pandemic (2010/11). Eurosurveillance 2014;19. https://doi.org/10.2807/1560-
7917.ES2014.19.16.20780.

[18] Safety of COVID-19 vaccines | European Medicines Agency (EMA) n.d.
https://www.ema.europa.eu/en/human-regulatory-overview/public-health-
threats/coronavirus-disease-covid-19/covid-19-medicines/safety-covid-19-vaccines
(accessed June 10, 2025).

[19] D'Elia L, Giaquinto A, Zarrella AF, Rendina D, Iaccarino Idelson P, Strazzullo P, et
al. Hypertension and mortality in SARS-COV-2 infection: A meta-analysis of
observational studies after 2 years of pandemic. Eur J Intern Med 2023;108:28–36.
https://doi.org/10.1016/j.ejim.2022.11.018.

[20] Fresán U, Guevara M, Trobajo-Sanmartín C, Burgui C, Ezpeleta C, Castilla J.
Hypertension and Related Comorbidities as Potential Risk Factors for COVID-19
Hospitalization and Severity: A Prospective Population-Based Cohort Study. J Clin
Med 2021;10:1194. https://doi.org/10.3390/jcm10061194.

[21] Ge E, Li Y, Wu S, Candido E, Wei X. Association of pre-existing comorbidities with
mortality and disease severity among 167,500 individuals with COVID-19 in Canada:
A population-based cohort study. PLoS One 2021;16:e0258154.
https://doi.org/10.1371/journal.pone.0258154.

[22] Wee LE, Tan JYJ, Chiew CJ, Abisheganaden JA, Chotirmall SH, Lye DCB, et al. A
Nationwide Cohort Study of Delta and Omicron SARS-CoV-2 Outcomes in
Vaccinated Individuals With Chronic Lung Disease. Chest 2024.
https://doi.org/10.1016/j.chest.2024.05.017.

[23] Rainer L, Bachner F, Eglau K, Ostermann H, Siebert U, Zuba M. Comorbidities and
COVID-19 hospitalization, ICU admission and hospital mortality in Austria. Wien
Klin Wochenschr 2022;134:856–67. https://doi.org/10.1007/s00508-022-02036-9.

[24] Bedston S, Almaghrabi F, Patterson L, Agrawal U, Woolford L, Anand SN, et al. Risk
of severe COVID-19 outcomes after autumn 2022 COVID-19 booster vaccinations: a
pooled analysis of national prospective cohort studies involving 7.4 million adults in
England, Northern Ireland, Scotland and Wales. The Lancet Regional Health - Europe
2024;37:100816. https://doi.org/10.1016/j.lanepe.2023.100816.

[25] Semenzato L, Botton J, Drouin J, Cuenot F, Dray-Spira R, Weill A, et al. Chronic
diseases, health conditions and risk of COVID-19-related hospitalization and in-
hospital mortality during the first wave of the epidemic in France: a cohort study of 66
million people. The Lancet Regional Health - Europe 2021;8:100158.
https://doi.org/10.1016/j.lanepe.2021.100158.

[26] Prosperini L, Tortorella C, Haggiag S, Ruggieri S, Galgani S, Gasperini C. Increased
risk of death from COVID-19 in multiple sclerosis: a pooled analysis of observational
studies. J Neurol 2022;269:1114–20. https://doi.org/10.1007/s00415-021-10803-3.

[27] Moreno‐Torres I, Meca Lallana V, Costa‐Frossard L, Oreja‐Guevara C, Aguirre C,
Alba Suárez EM, et al. Risk and outcomes of COVID‐19 in patients with multiple
sclerosis. Eur J Neurol 2021;28:3712–21. https://doi.org/10.1111/ene.14990.

[28] Orangzeb S, Watle SV, Caugant DA. Adherence to vaccination guidelines of patients
with complete splenectomy in Norway, 2008–2020. Vaccine 2023;41:4579–85.
https://doi.org/10.1016/j.vaccine.2023.06.034.

[29] Béraud G. Mathematical models and vaccination strategies. Vaccine 2018;36:5366–72.
https://doi.org/10.1016/j.vaccine.2017.10.014.

[30] Bubar KM, Reinholt K, Kissler SM, Lipsitch M, Cobey S, Grad YH, et al. Model-
informed COVID-19 vaccine prioritization strategies by age and serostatus. Science
(1979) 2021;371:916–21. https://doi.org/10.1126/science.abe6959.

[31]   Viana J, van Dorp CH, Nunes A, Gomes MC, van Boven M, Kretzschmar ME, et al.
417   Controlling the pandemic during the SARS-CoV-2 vaccination rollout. Nat Commun
418   2021;12:3674. https://doi.org/10.1038/s41467-021-23938-8.
419   [32]   Matrajt L, Eaton J, Leung T, Brown ER. Vaccine optimization for COVID-19: Who to
420   vaccinate first? Sci Adv 2021;7. https://doi.org/10.1126/sciadv.abf1374.
421

Figure 1. Chronic conditions in the general population. A-B) National and **C-D)** European medical risk

classification for severe COVID-19. In all classifications, chronic medical condition categories were

assigned if registered in the past six months prior to 1 January 2020, except for cancer which was assigned

if registered in the prior five years. The left bars show the number and proportion of each chronic medical

condition category in the population. Orange bars indicate the high-risk chronic conditions according to the

classification, while blue bars indicate moderate-risk chronic conditions. The dots represent the presence of

one or more chronic medical condition categories in an individual, with the upper bars showing the

combinations that occur in at least 1% of the general population.

Figure 2. Classification of the general population by risk for severe COVID-19. Population shown as

absolute numbers (top) and proportions (bottom) classified by age and risk for severe COVID-19 due to

chronic medical condition categories in the **A-B)** national and **C-D)** European classifications.

Figure 3. Discordance in assigning risk status to individuals between the European and national

classifications in the A-B) general population, **C-D)** hospitalized population, and **E-F)** deceased

population. The discordance between the European and each national classification was calculated as the

proportion of the general populations with a different risk status (off-diagonal elements). Concordance is

the proportion of the population classified into the same risk group (diagonal elements). Since every

individual belongs to one of the risk groups, the elements of the matrix plots add up to 100%.

Figure 4. Chronic conditions in the discordant general population. A-B) National and **C-D)** European

medical risk classification for severe COVID-19. In all classifications, chronic medical condition

categories were those that had been registered in the past six months prior to 1 January 2020, except for

cancer which was included as a chronic medical condition if registered in the past five years preceding 1

January, 2020. The dots represent the presence of one or more chronic medical condition categories. The

left bars show the number and proportion of each chronic medical condition category in the population.

Orange bars indicate the high-risk chronic conditions according to the classification, while blue bars

indicate moderate-risk chronic conditions. The upper bars show the combinations of chronic conditions

that occur in at least 1% of the discordant general population. [6][4][5]

A The Netherlands; General population

	High	Moderate	Low	Total
High	1.3%	**6.4%**	**1.3%**	**9.0%**
Moderate	**0.0%**	0.9%	**0.0%**	**1.0%**
Low	**0.2%**	**4.1%**	85.8%	**90.1%**
Total	**1.5%**	**11.3%**	**87.2%**	100%

European classification (y-axis); National classification (x-axis)

Discordance = 12.0% (95%CI 12.0–12.1%)
Concordance = 88.0% (95%CI 87.9–88.0%)

B Norway; General population

	High	Moderate	Low	Total
High	2.5%	**6.1%**	**0.7%**	**9.2%**
Moderate	**0.1%**	0.6%	**0.6%**	**1.3%**
Low	**0.4%**	**6.0%**	83.1%	**89.5%**
Total	**3.0%**	**12.7%**	**84.4%**	100%

European classification (y-axis); National classification (x-axis)

Discordance = 13.8% (95%CI 13.8–13.9%)
Concordance = 86.2% (95%CI 86.1–86.2%)

C The Netherlands; Hospitalized population

	High	Moderate	Low	Total
High	10.0%	**29.4%**	**2.2%**	**41.7%**
Moderate	**0.1%**	2.0%	**0.1%**	**2.3%**
Low	**0.2%**	**4.4%**	51.5%	**56.1%**
Total	**10.3%**	**35.9%**	**53.8%**	100%

European classification (y-axis); National classification (x-axis)

Discordance = 36.5% (95%CI 36.0–37.0%)
Concordance = 63.5% (95%CI 63.0–64.0%)

D Norway; Hospitalized population

	High	Moderate	Low	Total
High	11.0%	**19.8%**	**1.7%**	**32.5%**
Moderate	**0.5%**	1.3%	**1.3%**	**3.2%**
Low	**0.8%**	**14.1%**	49.5%	**64.3%**
Total	**12.2%**	**35.3%**	**52.5%**	100%

European classification (y-axis); National classification (x-axis)

Discordance = 38.2% (95%CI 36.2–40.3%)
Concordance = 61.8% (95%CI 59.7–63.8%)

E The Netherlands; Deceased population

	High	Moderate	Low	Total
High	10.4%	**40.6%**	**2.4%**	**53.4%**
Moderate	**0.1%**	2.7%	**0.1%**	**2.9%**
Low	**0.1%**	**3.4%**	40.2%	**43.7%**
Total	**10.6%**	**46.7%**	**42.7%**	100%

European classification (y-axis); National classification (x-axis)

Discordance = 46.6% (95%CI 45.5–47.5%)
Concordance = 53.4% (95%CI 52.5–54.5%)

F Norway; Deceased population

	High	Moderate	Low	Total
High	16.0%	**28.0%**	**1.2%**	**45.1%**
Moderate	**0.6%**	2.7%	**1.4%**	**4.7%**
Low	**1.0%**	**13.0%**	36.2%	**50.2%**
Total	**17.5%**	**43.8%**	**38.7%**	100%

European classification (y-axis); National classification (x-axis)

Discordance = 45.1% (95%CI 40.8–49.6%)
Concordance = 54.9% (95%CI 50.4–59.2%)