Association between plausible genetic factors and weight loss from GLP1-RA and bariatric surgery: a multi-ancestry study in 10 960 individuals from 9 biobanks

Jakob German*[1,2], Mattia Cordioli*[1,3], Veronica Tozzo[4], Sarah Urbut[5,6,7], Kadri Arumäe[8], Roelof A.J. Smit[9,10,11,12], Jiwoo Lee[3,6], Josephine H. Li[6,13,14,15], Adrian Janucik[14,16], Yi Ding[4], Akintunde Akinkuolie[4], Henrike Heyne[17], Andrea Eoli[18,19], Chadi Saad[20], Yasser Al-Sarraj[20], Rania Abdel-latif[20], Shaban Mohammed[21], Moza Al Hail[21], Alexandra Barry[6,14], Zhe Wang[9], Estonian Biobank research team[38], Tatiana Cajuso[1,22], Andrea Corbetta[1,23,24], Pradeep Natarajan[6,14,25,26], Samuli Ripatti[1,14,27,40], Anthony Philippakis[2,3], Lukasz Szczerbinski[6,13,28,29,30], Bogdan Pasaniuc[4,31,32,33,34], Zoltan Kutalik[35,36,37], Hamdi Mbarek[20], Ruth J.F. Loos[9,10,11,12], Uku Vainik[8,38,39,+], Andrea Ganna[1,40,+]

* These authors contributed equally
+ These authors jointly supervised the project

1. Institute for Molecular Medicine Finland (FIMM), University of Helsinki, Helsinki, Finland;
2. Eric and Wendy Schmidt Center, Broad Institute of MIT and Harvard, Cambridge, MA, USA, 02142;
3. Broad Institute of MIT and Harvard, Cambridge, MA, USA;
4. Department of Computational Medicine, David Geffen School of Medicine at UCLA, Los Angeles, CA, USA;
5. Division of Cardiovascular Medicine, Massachusetts General Hospital, Boston, MA;
6. Center for Genomic Medicine Massachusetts General Hospital, Boston, MA;
7. Cardiovascular Research Center, Massachusetts General Hospital, Boston, MA;
8. Institute of Psychology, Faculty of Social Sciences, University of Tartu, Tartu, Estonia;
9. Charles Bronfman Institute for Personalized Medicine, Icahn School of Medicine at Mount Sinai, New York, NY, USA;
10. Department of Environmental Medicine and Climate Science, Icahn School of Medicine at Mount Sinai, New York, NY, USA;
11. Novo Nordisk Foundation Center for Basic Metabolic Research, Faculty of Health and Medical Sciences, University of Copenhagen, Copenhagen, Denmark;
12. Novo Nordisk Foundation Center for Genomic Mechanisms of Disease, Broad Institute of MIT and Harvard, Cambridge, MA, USA;
13. Diabetes Unit, Massachusetts General Hospital, Boston, MA, USA;
14. Program in Medical & Population Genetics, Broad Institute of Harvard and MIT, Cambridge, MA, USA;
15. Harvard Medical School, Boston, MA, USA;
16. Center for Digital Medicine, Medical University of Bialystok, Bialystok, Poland;
17. Hasso Plattner Institut, Potsdam, Germany;
18. Digital Engineering Faculty, University of Potsdam, Potsdam, Germany, Prof.-Dr.-Helmert-Str. 2-3, 14482;
19. Hasso Plattner Institute for Digital Health at Mount Sinai, Icahn School of Medicine at Mount Sinai, New York City, NY, USA;
20. Qatar Genome Program, Qatar Precision Health Institute, Qatar Foundation, Doha, Qatar;
21. Department of Pharmacy, Hamad Medical Corporation, Doha, Qatar;

22. Department of Pathology, Dana-Farber Cancer Institute and Harvard Medical School, Boston, MA, 02115, USA;

23. Health Data Science Centre, Human Technopole, Milan, Italy;

24. MOX - Laboratory for Modeling and Scientific Computing, Department of Mathematics, Politecnico di Milano, Milan,Italy;

25. Personalized Medicine, Mass General Brigham, Boston, MA, USA;

26. Department of Medicine, Harvard Medical School, Boston, MA, USA;

27. Department of Public Health, Clinicum, Faculty of Medicine, University of Helsinki, Helsinki, Finland;

28. Programs in Metabolism and Medical & Population Genetics, Broad Institute of MIT and Harvard, Cambridge, MA, USA;

29. Department of Endocrinology, Diabetology and Internal Medicine, Medical University of Bialystok, Bialystok, Poland;

30. Clinical Research Centre, Medical University of Bialystok, Bialystok, Poland;

31. Bioinformatics Interdepartmental Program, UCLA, Los Angeles, CA, USA;

32. Department of Pathology and Laboratory Medicine, David Geffen School of Medicine at UCLA, Los Angeles, CA, USA;

33. Department of Human Genetics, David Geffen School of Medicine at UCLA, Los Angeles, CA, USA;

34. Institute for Precision Health, UCLA, Los Angeles, CA, USA;

35. University Center for Primary Care and Public Health, Lausanne, Switzerland;

36. Department of Computational Biology, University of Lausanne, Lausanne, Switzerland;

37. Swiss Institute of Bioinformatics, Lausanne, Switzerland;

38. Estonian Genome Centre, Institute of Genomics, Faculty of Science and Technology, University of Tartu, Tartu, Estonia ;

39. Department of Neurology and Neurosurgery, Faculty of Medicine and Health Sciences, McGill University, Canada;

40. Analytic & Translational Genetics Unit, Massachusetts General Hospital, Harvard Medical School, Boston, MA, USA;

Correspondence to Andrea Ganna (andrea.ganna@helsinki.fi)

Abstract

Obesity is a significant public health concern. GLP-1 receptor agonists (GLP1-RA), predominantly in use as a type 2 diabetes treatment, are a promising pharmacological approach for weight loss, while bariatric surgery (BS) remains a durable, but invasive, intervention. Despite observed heterogeneity in weight loss effects, the genetic effects on weight loss from GLP1-RA and BS have not been extensively explored in large sample sizes, and most studies have focused on differences in race and ethnicity, rather than genetic ancestry. We studied whether genetic factors, previously shown to affect body weight, impact weight loss due to GLP1-RA therapy or BS in 10,960 individuals from 9 multi-ancestry biobank studies in 6 countries. The average weight change between 6 and 12 months from therapy initiation was -3.93% for GLP1-RA users, with marginal differences across genetic ancestries. For BS patients the weight change between 6 and 48 months from the operation was -21.17%. There were no significant associations between weight loss due to GLP1-RA and polygenic scores for BMI or type 2 diabetes or specific missense variants in the *GLP1R, PCSK1 and APOE* genes, after multiple-testing correction. A higher polygenic score for BMI was significantly linked to lower weight loss after BS (+0.7% for 1 standard deviation change in the polygenic score, $P = 1.24\times10^{-4}$), but the effect was modest and further reduced in sensitivity analyses. Our findings suggest that existing polygenic scores related to weight and type 2 diabetes and missense variants in the drug target gene do not have a large impact on GLP1-RA effectiveness. Our results also confirm the effectiveness of these treatments across all major continental ancestry groups considered.

Introduction

The obesity epidemic presents a significant public health burden, driving the need for effective treatment options.[1,2] Despite 75% of adults with obesity having attempted to lose weight, most have not achieved lasting success.[3] Amid these challenges, glucagon-like peptide-1 receptor agonists (GLP1-RA) have emerged as a promising solution. As first discovered in the late 1980s, glucagon-like peptide-1 (GLP-1) is linked to the incretin effect, which promotes insulin release from pancreatic beta cells, thus playing a crucial role in regulating blood sugar levels.[4,5] This evidence led to the approval of the first GLP1-RA for treatment of type 2 diabetes in 2005.[6–8] Nine years later, an association between GLP-1 and reduced food intake paved the way for the development and approval of the first GLP1-RA for obesity treatment.[5,9]

Clinical trials conducted in the following years demonstrated weight loss effects ranging from 6% to 16% for liraglutide and semaglutide, leading to a substantial increase in GLP1-RA usage worldwide[10–12]. Additional trials have further revealed cardiovascular[13,14] and renal benefits[15], with ongoing trials currently underway.[16] Overall, anticipated trends suggest that by 2030, the US will witness approximately 30 million patients using GLP1-RA drugs, representing roughly 9% of its population, underscoring their potential in combating the obesity crisis and exploring new therapeutic applications.[17–19]

Meanwhile, bariatric surgery (BS) stands as a durable and effective treatment for individuals with severe obesity (insurance payer threshold in the US: BMI ≥ 40 kg/m²)[20,21] and metabolic comorbidities (BMI ≥ 35 kg/m²), with some clinical guidelines now recommending its use for patients with a BMI ≥ 30 kg/m².[22,23] Meta-analyses combining results from randomized controlled trials and observational studies found that, on average, individuals undergoing BS experienced a weight loss of 26 kg. Separately, a randomized controlled trial reported a total weight loss of 25% after 5 years in patients undergoing sleeve gastrectomy and Roux-en-Y gastric bypass.[24,25] Although metabolic surgery carries typical risks for such interventions, including nutrient malabsorption, large-scale observational studies have demonstrated remarkable long-term benefits, such as reduced cardiovascular outcomes, lower all-cause mortality, and higher remission rates of type 2 diabetes, hypertension, and hyperlipidemia.[26–32]

Despite the robust weight loss effects observed with both GLP1-RA treatment and BS, there
exists heterogeneity in their effectiveness. One out of every five patients undergoing BS may not
achieve the desired weight loss within the first year or may experience weight regain within two
years.[33–36] The variability of GLP1-RA treatment effects on glycemic outcomes has been widely
investigated and can be partially explained by clinical indicators like low β-cell function or C-
peptide levels as well as genetic variants.[37–42] However, with the exception of type 2 diabetes,
sex, and baseline weight, the factors influencing the diversity in weight loss effects of these drugs
are still poorly understood.[43–45] Additionally, differences in patient adherence to treatment
regimens may further contribute to this variability.[46]
138
Genetics has been considered a potential factor for heterogeneity in weight loss treatments, but
its role remains largely unexplored.[43] A previous study has examined the role of a common
missense variant in the *GLP1R* genes in 57 women and found a weak link with GLP1-RA
treatment response.[39,47] A greater number and better-powered studies have been conducted in
individuals with BS [48,49], however, a consistent genome-wide association for weight loss has yet
to be identified. A previous study has shown that a polygenic score for BMI had a significant
impact on weight loss after biliopancreatic diversion with duodenal switch in 865 patients.[50]
146
147   Finally, there has been limited exploration into the effectiveness of these obesity treatments
148   across different genetic ancestries and across various healthcare settings.[10,51]
149
In our study we seek to i) characterize the observed GLP1-RA and BS body weight-lowering
effects using real-world data from 10 960 individuals across 9 biobanks in 6 countries ,with 6
major continental ancestry groups; ii) identify whether plausible genetic factors (i.e. polygenic
scores for BMI and T2D and coding variants in *GLP1R*, *PCSK1* and *APOE* gene) associate with
heterogeneity in GLP1-RA and BS weight loss effects, and thus identify patient groups that will
benefit most from such treatments; iii) compare the effects of GLP1-RA treatment and BS on
weight loss across major continental ancestry groups.
157
158

Results

GLP1-RA and bariatric surgery associated weight loss in 10 960 individuals from 9 cohorts and 6 major continental ancestry groups

We investigated changes in body weight associated with GLP1-RA treatment and BS across 9 biobank studies: Helsinki University Hospital (HUS, Finland; N = 633), Estonian Biobank (ESTBB; N = 464), UK Biobank (UKBB, United Kingdom; N = 810), All of Us (AoU, USA; N = 559), BioMe Biobank (BioMe, USA; N = 2,170), Mass General Brigham Biobank (MGBB, USA; N = 2,141), Atlas Biobank (UCLA-ATLAS, USA; N = 1,445), Qatar Biobank (QBB, Qatar; N = 2,383), Bialystok Bariatric Surgery Study (BBSS, Poland, N = 355). We included individuals that were at least 18 years of age when starting a GLP1-RA treatment or undergoing BS. Individuals undergo GLP1-RA treatment for different medical reasons, not only for weight loss. We defined GLP1-RA usage based on prescription or purchase data (ATC codes starting with A10B*; including exenatide, liraglutide, lixisenatide, albiglutide, dulaglutide, semaglutide, and beinaglutide) covering at least 12 months, and excluding individuals with indication of treatment discontinuation within this time window. For BS, we considered individuals undergoing the following procedures: Roux-en-Y gastric bypass, sleeve gastrectomy, adjustable gastric band, vertical banded gastroplasty, biliopancreatic diversion with duodenal switch. More information can be found in the **Methods** section.

We included only individuals who had an initial body weight measurement recorded at most 12 months prior to the baseline (defined as treatment initiation or surgery) and a follow-up measurement taken at least 6 but no more than 12 months after the baseline for GLP1-RA treatments, or within 48 months after baseline for BS. In case of multiple measurements within this time frame, we considered the median of such measurements (**Supplementary Figure 1** for a schematic of the approach used). We defined our main outcome of interest as the percentage change in body weight, calculated as the difference between the second body weight measurement (or median of multiple measurements) and the initial one, divided by the initial measurement .

Our study population included a total of 6,750 GLP1-RA users and 4,210 individuals undergoing BS (baseline characteristics are reported in **Table 1**). Overall, GLP1-RA users 1) had a lower

proportion of women (61.0% weighted average across studies) when compared to BS patients (73.1% weighted average across studies), 2) were on average older (57.30 *vs* 46.50), and 3) had a lower initial body weight (99.80 kg *vs* 116.96 kg). The proportion of individuals diagnosed with type 2 diabetes at time of GLP1-RA initiation was at least 48% across all studies and major continental ancestry groups.

Study	Ancestry	Number of individuals	Proportion of women (%)	Mean body weight (in kg) at baseline (SD)	Mean BMI (in kg/m2) at baseline (SD)	Mean age at baseline (SD)	T2D prevalence (%)	Most common GLP1-RA type	Proportion of semaglutide (%)
GLP1-RA									
HUS	EUR	255	68.00	113.70 (22.90)	39.56 (7.93)	53.79 (13.52)	65.49	semaglutide	60.40
ESTBB	EUR	191	62.30	109.18 (20.30)	37.68 (6.40)	56.95 (10.96)	89.01	semaglutide	72.77
UKBB	EUR	614	44.30	105.00 (19.90)	36.70 (6.14)	60.50 (7.15)	99.80	liraglutide	0.00
AoU	EUR	117	64.96	106.66 (24.18)	37.23 (7.84)	57.01 (11.03)	82.91	liraglutide	11.97
	AFR	91	83.52	104.8 (25.01)	37.94 (8.77)	54.57 (11.93)	92.31	liraglutide	18.68
MGBB	EUR	857	57.01	99.99 (21.48)	35.26 (6.79)	59.88 (12.17)	56.33	semaglutide	40.72
	AFR	156	54.32	104.65 (23.80)	37.25 (7.85)	54.31 (11.91)	60.24	dulaglutide	36.14
BioMe	AMR	614	66.45	91.19 (22.45)	34.34 (7.19)	59.07 (11.29)	67.75	dulaglutide	29.32
	AFR	728	69.78	101.49 (25.42)	36.15 (8.18)	57.00 (12.24)	69.50	dulaglutide	29.12
	EUR	369	44.44	98.26 (21.93)	33.93 (6.74)	60.32 (11.57)	47.97	semaglutide	37.13
	SAS	77	50.65	83.79 (20.10)	30.61 (6.70)	56.48 (12.59)	70.13	semaglutide	41.56
UCLA-ATLAS	AFR	133	64.00	101.6 (23.20)	35.50 (7.10)	61.00 (13.00)	91.00	semaglutide	57.10
	AMR	336	66.00	94.40 (22.60)	35.10 (7.10)	56.00 (13.00)	84.00	liraglutide	26.50
	EAS	120	47.00	81.50 (16.30)	29.70 (5.20)	60.00 (13.00)	92.00	semaglutide	61.70
	EUR	856	53.00	101.60 (22.00)	34.80 (6.90)	63.00 (13.00)	80.00	semaglutide	67.80
QBB	MID	1236	72.17	97.68 (24.54)	37.67 (8.90)	49.52 (10.75)	75,97	liraglutide	3.64
Bariatric surgery									
HUS	EUR	378	81.00	113.63 (19.50)	40.00 (5.71)	49.90 (9.39)	37.04		
ESTBB	EUR	273	76.56	121.24 (20.57)	41.80 (5.58)	44.72 (10.07)	28.21		
UKBB	EUR	196	69.40	121.00 (29.70)	44.20 (9.74)	55.50 (8.21)	50.00		
AoU	EUR	168	81.55	121.32 (29.88)	43.48 (9.85)	50.74 (12.62)	18.45		
	AFR	183	89.62	127.23 (32.55)	49.33 (27.56)	46.09 (11.67)	48.63		
MGBB	EUR	978	71.06	121.21 (26.52)	42.93 (7.90)	49.37 (12.16)	39.98		
	AFR	150	86.67	122.89 (23.92)	45.16 (8.27)	43.68 (11.67)	42.67		
BioMe	AMR	143	79.72	112.36 (20.81)	42.55 (6.13)	44.62 (11.48)	30.07		
	AFR	188	82.98	124.88 (25.63)	45.45 (7.90)	45.35 (12.16)	29.26		
	EUR	51	70.59	120.23 (25.29)	43.03 (7.95)	49.51 (13.34)	17.65		
QBB	MID	1147	69.49	102.14 (23.18)	38.55 (7.87)	41.66 (10.66)	40.02		
BBSS	EUR	355	56.06	138.48 (27.07)	47.10 (7.11)	47.10 (7.11)	35.21		

Table 1 | Baseline characteristics for 6 750 GLP1-RA users and 4 210 patients undergoing bariatric surgery included in our study population. For each cohort and major continental ancestry group (AFR: African, AMR: admixed American, EAS: East Asian, EUR: European, MID: Middle Eastern, SAS: South Asian), we report the total number of individuals eligible for analysis, the proportion of females, the mean (SD) body weight (in Kg) at baseline, the mean (SD) age (in years) at baseline, the prevalence of T2D, and (for GLP1-RA users only) the most commonly used type of medication and the proportion of semaglutide users.
HUS, Helsinki University Hospital; ESTBB, Estonian Biobank; UKBB, UK Biobank; AoU, All of US; MGBB, Mass General Brigham Biobank; BioMe, BioMe; UCLA-ATLAS, Atlas Biobank; QBB, Qatar Biobank; BBSS, Bialystok Bariatric Surgery Study

197 Among GLP1-RA users, we observed an average body weight change (**Figure 1**) across studies

198 of -3.93% (ranging from -1.08% to -7.07%), in line with results from observational studies.[52,53]

199 Among BS patients, we observed an average change of -21.17% (-15.00% to -27.72%).

Figure 1 | Average percentage change in body weight for 6 750 GLP1-RA users and 4 210 bariatric surgery patients. Bars represent the mean percentage change in body weight within each study and ancestry, and the overall combined weighted mean change. Error bars represent standard errors.

Demographic factors associated with GLP1-RA and bariatric surgery weight change

We explored the effect of baseline characteristics on body weight changes associated with GLP1-

RA treatment and BS. We fitted a multivariable linear model with percentage change in body

weight as outcome and baseline body weight, sex, age at baseline, the first 20 principal genetic

components and medication type (for GLP1-RA users only) as predictors. Each cohort was

analyzed separately, stratifying by major continental ancestry group, and the effects were then

meta-analyzed (**Table 2**). Baseline body weight was significantly associated with greater weight

loss for both GLP1-RA and BS treatments ($\beta_{GLP1-RA}$ = -0.05 % weight change compared to

baseline, $P = 1.63 \times 10^{-35}$; β_{BS} = -0.14 % weight change compared to baseline, $P = 5.49 \times 10^{-69}$)

and women had a significantly greater weight loss for both treatments ($\beta_{GLP1-RA}$ = -1.54 % weight

change compared to baseline, $P = 2.47 \times 10^{-16}$; β_{BS} = -2.66 % weight change compared to

baseline, $P = 6.83 \times 10^{-10}$). Being older at treatment initiation was only associated with less weight

loss for bariatric surgery (β_{BS} = 0.15 % weight change compared to baseline, $P = 2.54 \times 10^{-15}$).

The effects of baseline weight and sex on weight loss were highly heterogeneous across

ancestries and studies (I^2 for GLP1-RA = 0.99 and 0.83 for baseline weight and sex, respectively)

(**Supplementary Figures 2-4 and Supplementary Tables 1-2**).

	GLP1-RA			Bariatric surgery		
Variable	**Effect on percentage weight change**	**Standard error**	**P value**	**Effect on percentage weight change**	**Standard error**	**P value**
Baseline weight (kg)	-0.05	0.004	1.63×10^{-35}	-0.14	0.008	5.49×10^{-69}
Sex (female)	-1.54	0.188	2.74×10^{-16}	-2.66	0.431	6.83×10^{-10}
Age at baseline (years)	-0.02	0.008	0.052	0.15	0.016	9.85×10^{-19}

Table 2 | Effect of baseline weight, sex and age at baseline of percentage weight changes associated with GLP1-RA and bariatric surgery. Multivariable model adjusting for the three variables described in the table and the first 20 principal genetic components and medication type (for GLP1-RA users only). Negative effects indicate larger weight loss.

Marginal effect of ancestry on GLP1-RA and bariatric surgery associated weight loss

We evaluate whether the effect of GLP1-RA and BS associated weight loss was significantly

different across genetically-defined continental ancestry groups after accounting for differences

in weight at baseline, age, sex and, only for GLP1-RA, medication type (**Table 3**). Only multi-

ancestry biobanks (AoU, BioMe, MGBB, UCLA-ATLAS) were included in this analysis (N = 6 315).

Compared to individuals of European ancestry, all other ancestry groups had lower change in

body weight. However, this effect was statistically significant only among GLP1-RA users of

African (β = 0.76 % weight change from baseline weight, $P = 0.02$) and admixed American (β =

0.87 % weight change from baseline weight, $P = 3 \times 10^{-3}$) ancestries.

GLP1-RA			
Major continental ancestry group	**Effect on percentage weight change**	**Standard error**	**P value**
AFR	0.758	0.327	0.021
AMR	0.874	0.295	0.003
EAS	1.093	0.664	0.100
SAS	0.118	0.759	0.876
Bariatric surgery			
Major continental ancestry group	**Effect on percentage weight change**	**Standard error**	**P value**
AFR	1.688	1.308	0.197
AMR	-0.542	1.428	0.704

Table 3 | Associations between weight change and major continental ancestry groups. Meta-analysis effect sizes, standard errors and P values for the association between weight change and major continental ancestry groups (AFR: African, AMR: admixed American, EAS: East Asian, EUR: European, SAS: South Asian). Coefficients are derived from a multivariable model including percentage weight change as outcome and ancestry as categorical variable (with EUR as reference category), and adjusting for sex, age and weight at baseline and, for GLP1-RA only, medication type. Negative effects indicate larger weight loss

Plausible genetic factors do not associate with weight change due to GLP1-RA

We investigated the effect of genetic exposures with plausible effects on body weight changes. Genetic exposures were selected upon discussion between authors and before initiating analyses. First, we considered polygenic scores for BMI based on previous observation that a polygenic score was associated with BMI trajectories.[54,55] We also consider the only replicated genome-wide variant associated with BMI change in the largest study of BMI trajectories (*rs429358* in *APOE*).[54] Second, we considered a polygenic score for T2D based on the observation that biomarkers of type 2 diabetes were associated with effectiveness of GLP1-RA.[38,40] We confirmed that both polygenic scores for BMI and T2Ds were significantly associated with their respective traits across all major continental ancestry groups (**Supplementary Table 3**) although the effects were larger in individuals of European ancestry.

Third, we considered 13 SNPs in the *GLP1R* and *PCSK1* genes. We considered all missense

variants in the *GLP1R* gene with at least 1% minor allele frequency in at least one of major

continental ancestry group, as per gnomAD (version 4.1)[56], three of which have been functional

characterized as gain or loss of function (**Supplementary Table 4**).[57] We also considered two

non-synonymous variants in the *PCSK1* gene that have been associated with BMI variation in the

population.[58] Finally, we sought to replicate the effect of two variants associated with weight loss

in one of the largest studies of BS[49](**Methods**). After analyses, two missense variants in the

GLP1R genes, *rs2295006* and *rs201672448* were too rare to be tested across studies and were

not considered further.

We estimated the effect of each genetic exposure on weight change in each cohort and major

continental ancestry groups separately, fitting a linear model adjusted for baseline body weight,

sex, age at baseline, first 20 principal genetic components, and medication type (for GLP1-RA

users only), and we subsequently combined the effects through ancestry-specific and multi-

ancestry meta-analyses (**Figure 2, Supplementary Tables 5-6**). We did not adjust for T2D

diagnosis because we considered it a potential mediator between the genetic exposure and

weight change outcome.

Among GLP1-RA users, we did not observe any significant genetic exposure associated with

weight loss after multiple testing correction ($P < 0.004$, Bonferroni correction for 13 exposures

tested).

We further tested whether the lack of associations observed was possibly due to high effect-size

heterogeneity across ancestries, or due to lack of statistical power. We did not observe any

significant heterogeneity across ancestries for any of the genetic exposure (average I^2 across

genetic exposures for GLP1-RA was 0.06 **- Supplementary Tables 7-8**). Moreover, given the

current sample size for GLP1-RA users (N = 6 750) our power calculation showed we would have

at least 80% power to detect a statistically significant effect (at $P < 0.05$) of at least a 0.3% change

in body weight for 1 standard deviation (SD) change in the polygenic score, or for SNPs with a

minor allele frequency of at least 1% (**Methods**).

Figure 2 | Effect of 15 genetic exposures on body weight changes associated with GLP1-RA treatment and bariatric surgery.
a. Ancestry-specific and multi-ancestry meta-analysis effect sizes for association between percentage change in body weight and PGS for BMI and type 2 diabetes. Dots represent the percentage change in body weight per one standard deviation change in PGS, error bars represent the 95% confidence interval. Full dots represent statistical significance at P < 0.004.
b. Multi-ancestry meta-analysis effect sizes for the association between percentage change in body weight and genotype at each locus. Dots represent the effect size, error bars represent the 95% confidence interval. PGS, polygenic score.

277
278

A higher polygenic score for BMI associates with lower weight loss due to BS

Among BS patients, a higher polygenic score for BMI was significantly associated with lower weight loss ($\beta_{BMI\ PGS}$ = 0.70 % weight change compared to baseline for 1 SD change in the polygenic score, P = 1.24 × 10^{-4}) after multiple testing correction (P < 0.01, Bonferroni correction for 5 exposures tested), in the opposite direction than observed for weight at baseline (**Figure 2, Table 2**). A higher polygenic score for T2D was nominally associated with lower weight loss ($\beta_{T2D\ PGS}$ = 0.45 % weight change compared to baseline for 1 SD change in the polygenic score, P = 0.03).
We observed significant heterogeneity in the association between BMI PGS and weight loss across studies (I2 = 0.64, **Supplementary Table 8**).

291
292

Sensitivity analyses

To assess the robustness of our findings, we conducted a series of sensitivity analyses.

First, we consider different outcome definitions. We shortened the follow-up period for BS to 12 months to make it comparable to that of GLP1-RA, we considered only participants with weight measurements available at least 30 days prior to BS to account for the possibility of significant weight loss due to preoperative low-calorie diets and we used minimum rather than median weight during the follow-up period. This latter definition resulted in significant higher weight loss for GLP1-RA (-6.00% compared to -3.93% when using the median, **Supplementary Tables 9-10** and **Supplementary Figures 5-6**). All analyses confirmed the associations observed in primary analysis (**Supplementary Tables 11-19** and **Supplementary Figures 7-14**), although the effect of BMI PGS on weight loss due to BS was slightly reduced when considering a shortened follow-up period (**Supplementary Table 11** and **Supplementary Figure 8**).

Second, we considered different covariate adjustment and performed sub-group analyses. We adjusted the analyses for T2D status and observed similar results than primary analyses (**Supplementary Tables 12-13** and **Supplementary Figure 9)**. We considered only semaglutide and liraglutide as GLP1-RA treatments to capture their specific use for weight loss. While we observed a higher weight loss than considering all GLP1-RA treatments (-4.69% vs -3.93% **Supplementary Figure 10**), we confirmed the lack of effect of BMI and T2D PGS on weight loss observed in primary analyses (**Supplementary Table 14 and Supplementary Figure 11**).

Third, we considered different statistical models. Previous works have argued that using percentage changes in weight loss pre vs post therapy is statistically inappropriate and inefficient.[59–61] Thus, we considered a model using post-treatment weight as outcome, including the same predictors as in the main model and an additional term modeling the interaction between baseline weight and each genetic exposure. We observed similar results as in the primary model (**Supplementary Tables 16-17**). A second potential issue with the primary model we used relates with adjusting longitudinal change phenotypes for the baseline trait.[62,63] In particular, when testing for the effect of genetic exposure on drug response outcomes, adjusting phenotype changes (e.g. weight loss) for baseline levels (e.g. weight at treatment initiation) can result in false positive association when genetic variants are also associated with baseline levels[64.] Removing baseline weight from the model led to a reduction in the effect of the BMI PGS on weight loss due to BS (0.15% per 1 SD change in the polygenic score, P = 0.420 compared to 0.70% in the primary model; **Supplementary Tables 18-19** and **Supplementary Figure 14**).

Discussion

In this study we provide a comprehensive assessment of major demographic factors and plausible

genetic factors on body weight changes associated with GLP1-RA use and BS.

In line with previous work,[43,65–68] our main outcome of interest was the percentage change in body

weight between 6 and 12 months after GLP1-RA treatment initiation as well as between 6 and 48

months after BS.

By considering studies from 6 different countries, we were able to highlight the heterogeneity in

use of GLP1-RA across different healthcare systems. GLP1-RA users consisted mostly of

individuals with type 2 diabetes, reflecting their original use for treatment of this condition. Biobank

studies covering Boston (MGBB), New York (BioMe), Los Angeles (UCLA-ATLAS) and Helsinki

(HUS) had overall lower rates of type 2 diabetes individuals compared to the other studies and

lower weight at baseline, which, together with the availability of more updated electronic health

records data, suggest a larger number of individuals being prescribed GLP1-RA for weight loss.

However, care should be taken in interpreting variation in rate of type 2 diabetes across biobanks

as this might reflect different approaches used for capturing this diagnosis from the electronic

health records as well as different sampling strategies of biobanks.

The comparison between GLP1-RA and BS is informative as it highlights the younger age, higher

weight and lower percentage of men undergoing BS compared to GLP1-RA users. The weight

reduction was also approximately 5 times higher among BS patients compared to GLP1-RA

users. The weight reduction among GLP1-RA users was also more heterogeneous across

biobanks compared to BS, but in line with what reported previously in the literature using real

world data[41] and lower than results from clinical trials.[11,69] The lower weight reduction observed

among GLP1-RA users is partially influenced by the use of the median postoperative or post-drug

therapy weight, while we observed larger weight reduction when using minimum weight. However,

our approach provides a more conservative and robust estimate, better accounting for the

variability and potential uncertainty in clinical measurement data. Our findings also highlight a

consistent effect of baseline body weight and sex on weight loss associated with both GLP1-RA

treatment and BS, suggesting that individuals with higher weight at baseline, as well as women
benefit more from both interventions. These findings underscore the importance of considering
individual characteristics when designing weight loss interventions.

The main aim of this study was to evaluate the impact of genetic exposures on body weight
changes induced by GLP1-RA or BS. At the current sample size, a genome-wide scan would
have limited power to identify potential new signals. With 6,750 individuals, we would have at
least 80% statistical power to identify, at a genome-wide significant threshold, a genetic variant
with an allele frequency of at least 1% and an effect larger than 0.3% weight change per change
in one SD, which is unlikely given previous effects of common variants on treatment response.[64]
Instead, we formulated a set of hypotheses based on the results from the largest genome-wide
association studies of BMI and type 2 diabetes, and, explored biologically plausible mutations in
several candidate genes. Polygenic scores represent the most powerful genetic predictors for
BMI and type 2 diabetes and we confirm their significant association with the corresponding trait
across all major continental ancestry groups. Thus, lack of associations with these polygenic
scores would suggest that GLP1-RA and BS-related weight loss is independent of one's genetic
predisposition to BMI or type 2 diabetes, as explained by common genetic variants. Moreover,
because the biological underpinning of weight change is partially distinct from weight at baseline,
we also consider the few replicated genetic signals for BMI variability and BMI trajectories.

For bariatric surgery we observed that genetically higher BMI was associated with weight gain, in
the opposite direction for the phenotypic association with initial body weight. This might suggest
that individuals with a higher genetic predisposition for elevated BMI might regain weight, after
the operation, faster than those with lower genetic predisposition. To maintain weight loss, they
may need additional lifestyle modifications, behavioral health therapy or adjuvant pharmacologic
therapies as a GLP1-RA treatment.[70,71]

On the contrary, none of the genetic exposures were associated with weight loss among GLP1-
RA users, after multiple-testing correction. These results were not attributable to high
heterogeneity across studies or across major continental ancestry groups. Whether we directly
tested the impact of genetic exposure on percentage body weight change or used post-treatment
weight as outcome and tested the interaction between baseline weight and the genetic exposure,
results did not significantly change.

In addition, we conducted extensive sensitivity analyses to further examine the effect of BMI and
T2D PGS on weight loss after BS and GLP1-RA treatment. These analyses included alternative
definitions of weight outcomes, a shorter follow-up period and stricter inclusion criteria for BS,
adjustments for T2D status, subgroup analysis for GLP1-RA medication, and testing different
statistical models. Across all these analyses, we did not observe a significant effect of BMI or T2D
PGS on weight reduction following GLP1-RA treatment, in concordance with our primary analysis.
Regarding the impact of BMI PGS on weight loss due BS, we observed a reduction of effect in
some of the sensitivity analyses, in particular when excluding baseline weight adjustment from
the model. In this case, the directionality of the effect remained consistent, however the
association was no longer statistically significant. Overall, our results showed that the considered
genetic influence on weight loss in response to GLP1-RA treatment and BS may be too small to
impact clinical decision-making or patient outcomes. This statement should be revisited either by
using more powerful polygenic scores or by considering individuals undergoing GLP1-RA
treatment only for weight loss.

Another aim of the study was to compare the effects of GLP1-RA treatment and BS on weight
loss across major continental ancestry groups. Ancestry can provide more useful information
about population health than racial and ethnic categories because it is not directly influenced by
geographic, cultural, and sociopolitical forces.[72] Nonetheless, ancestry is also partially correlated
with self-reported race and ethnicity and socio-economic characteristics do differ across racial
and ethnic groups. Overall, we did not find large differences in GLP1-RA and BS associated
weight loss between major continental ancestry groups. We do observe that GLP1-RA users of
African and admixed American ancestry had statistically significantly lower change in body weight
compared to Europeans, after accounting for differences in sex, age, weight at baseline and
medication type. The effects were consistent in BioMe and UCLA-ATLAS biobanks for admixed
American ancestry, and additionally in AoU and MGBB for African ancestry. This might reflect
different socio-economic characteristics between individuals of admixed American and European
ancestry resulting, for example, in differences in access to treatment.

The main strength of our study is its size and diversity across countries, healthcare systems and
ancestries. Our study presents various limitations. First, we cannot be sure that all GLP1-RA
users fulfilled their prescription. This is a potential issue only in studies with prescription
information but not in studies with purchase information. Nonetheless weight loss was comparable
across both study types. Moreover, we ensured that observed changes in body weight are due to

the treatment by considering only measurements within one month from the last purchase or
prescription. Second, we could not accurately estimate treatment adherence, but overcome this
limitation by considering treatments of maximum 12 months, a period for which adherence has
been observed to be around 65%.[65] Third, we used ancestry labels based on major continental
ancestry groups, while ancestry can be better characterized as a continuous quantity.[72] This
discretization reflects pragmatic considerations to enable comparable analyses across the
different studies. Fourth, the absence of a control group in the GLP1-RA analysis may limit
distinguishing intentional from unintentional weight loss. However, large-scale real-world data and
consistent sensitivity analysis support the robustness of our results. Fifth, although T2D
medications like metformin can influence weight loss, adjustments for T2D in our sensitivity
analyses likely accounted for first-line medications. Finally, we did not assess the impact of GLP1-
RA dosing, as dose adjustments vary over time, higher doses do not consistently correlate with
greater weight loss and inconsistent reporting across studies could introduce biases.
440
In conclusion, our study suggests that GLP1-RA treatments work equally well in individuals
carrying common non-synonymous mutations in the GLP1R gene and in individuals at high
genetic risk for BMI and type 2 diabetes. It also suggests that BS is less effective among
individuals with higher polygenic score for BMI, but the effect is not large enough to be clinically
relevant. In fact, for each additional SD in the polygenic score for BMI, individuals undergoing BS
have on average 0.7% lower weight loss from their initial body weight. Finally, these two major
weight loss interventions achieved sustained weight reduction regardless of an individual's
genetic ancestry. However, socio-economic characteristics correlating with ancestry should be
further studied to better understand some of the remaining differences observed in this study.

Methods

Study population

In the current study, we included samples from 10 960 individuals from the following 9 biobanks: Helsinki University Hospital (HUS, Finland; N = 633), Estonian Biobank (ESTBB, Estonia; N = 464), UK Biobank (UKBB, United Kingdom; N = 810), All of Us (AoU, USA; N = 559), BioMe (BioMe, USA; N = 2,170), Mass General Brigham Biobank (MGBB, USA; N = 2,141), Atlas Biobank (UCLA-ATLAS, USA; N = 1,445), Qatar Biobank (QBB, Qatar; N = 2,383), Bialystok Bariatric Surgery Study (BBSS, Poland, N = 355) . The biobank studies include samples from (hospital) biobanks, prospective epidemiological and disease-based cohorts. Follow-up covers a total of 20 years with the earliest study starting follow-up in 2004 (AoU) and the latest study ending follow-up in 2024 (BioMe). In the **Supplementary Material** we provide a detailed description of the population selected and ancestry definitions in each study.

Inclusion and exclusion criteria for GLP1-RA analysis

For our GLP1-RA analysis we implemented the following inclusion and exclusion criteria. We required an initiation of a GLP1-RA treatment (ATC codes: A10BJ*, A10BX04, A10BX10, A10BX13, A10BX14) defined as the date of first medication purchase or medication prescription of GLP1-RA. Only first-time (GLP1-RA naïve) users within the datasets were included. Individuals were required to be on a GLP1-RA treatment for at least twelve months. This was defined through prescription length, regular drug prescription refills or drug purchases, depending on the dataset and healthcare setting.

Individuals aged under 18 were excluded from the analysis. Furthermore, individuals were required to have at least one body weight measurement maximum one year prior to treatment initiation or within 14 days after initiation, if no measure before initiation was available. If we observed multiple weight measurements within this time frame, only the closest one to initiation was considered. This measurement defined our baseline weight variable. At least one body weight measurement between 26 weeks and 52 weeks was necessary for each individual to be included in the analysis. Individuals were excluded if they underwent BS prior to or during the first year of GLP1-RA treatment.

Inclusion and exclusion criteria for BS analysis

For our BS analysis we implemented the following inclusion and exclusion criteria. Individuals
were included if they underwent any type of BS. The respected code definitions for BS varied
across countries and healthcare systems and can be found in **Supplementary Table 20**. The
proportions of all procedures across cohorts can be found in **Supplementary Table 21**.
Individuals aged under 18 at baseline were excluded from the analysis. Analogously to the GLP1-
RA analysis, individuals were required to have at least one body weight measurement maximum
one year prior to surgery or within 14 days after it, if no measure before baseline was available.
If we observed multiple weight measurements within this time frame, only the closest one to
initiation was considered. This measurement defined our baseline weight variable. Additionally,
at least one body weight measurement between 26 weeks and 208 weeks was necessary for
each individual to be included in the analysis.

Outcome definitions

In our primary model, for both GLP1-RA and BS we defined the outcome to be the percentage
change in body weight from baseline. The baseline body weight measured closest to T_0 and
between -52 weeks and +2 weeks from T_0 was defined as W_0.
The second body weight (mW_1) was defined as the median of body weight measurements
between 26 weeks and 52 weeks from T_0 for the GLP1-RA analysis and between 180 days and
1460 days for the BS analysis. This was to enhance robustness against outlier measurements
occurring within these intervals.
Therefore, the percentage change in body weight from baseline was calculated using
$\% \; weight \; change \; = \frac{mW_1 - W_0}{W_0} * 100$.
In our secondary model, we used mW_1 as an outcome.

Genetic exposures

We identified 15 genetic variants (single nucleotide polymorphism, SNP) and two PGS of interest
for our analysis. Among the included SNPs 3 are functionally characterized *GLP1R* variants
(*rs10305492, rs146868158, rs6923761*) associated with random glucose levels that have shown

decreased/increased response to different endogenous and exogenous GLP1-RA.[57] Seven SNPs

are *GLP1R* missense variants (*rs1042044, rs10305420, rs3765467, rs10305421, rs2295006,*

rs10305510, rs201672448) with a frequency over 1% in at least one ancestry group in gnomAD

(version 4.1).[56] One SNP is the only robustly replicated lead genome-wide significant variant

(*rs429358*, *ApoE*) in a GWAS of BMI change over time.[54,74] Two SNPs are *PCSK1* missense

variants (*rs6232, rs6235*) that have been found associated with BMI variations.[58] And two SNPs

(*rs728996, rs17702901*) were associated with an effect on excess body mass index loss amongst

BS patients.[49] A list of all genetic exposures and their frequencies across all major continental

ancestry groups can be found in **Supplementary Table 4**. The included PGS are for body mass

index (BMI) and type 2 diabetes (T2D).[75] The PGS weights were taken from Weissbrod et al.

2022[75] and obtained from UK Biobank. The scores were computed in each study using PLINK

2.0.[76] Because the PGS scores were derived from UK Biobank, we couldn't use the same score

in UK Biobank, instead we used Thompson et al. 2022.[77]

Statistical model

We employed a linear regression model to investigate the effect of the chosen genetic exposures

on weight change after initiation of GLP1-RA treatment and BS, respectively. In the primary model

we adjusted for the baseline weight, sex, age at initiation of treatment or surgery in years, the first

20 genetic principal components and study-specific covariates, such as the genotyping batch. In

the GLP1-RA analysis we additionally adjusted for the medication type within the drug class.

We defined $\% \, weight \, change$ as the outcome variable in the primary model.

Primary model:

$\% \, weight \, change \; = \beta_0 + \beta_1 * genetic \, exposure \; + \beta_2 * W_0 + \beta_3 * sex + \beta_4 * age \, at \, initiation \; +$

$\beta_5 * medication \, (only \, for \, GLP1) + \sum_{k=6}^{25} \beta_k * PC1{:}20 \; + \sum_{i=26}^{n} \beta_i * study \, specific \, covariates$

Meta-analyses

We conducted ancestry-specific and multi-ancestry fixed-effect meta-analyses for each of the
genetic exposures. Effect sizes were combined using fixed-effect inverse-variance-weighting
(IVW), as implemented in the R package "meta".[78] For each exposure we reported the unadjusted
P values, but considered as threshold for statistical significance *P* < 0.004 for GLP1-RA (i.e. P <
0.05/13, Bonferroni correction for 13 exposures tested) and *P* < 0.01 for BS (i.e. P < 0.05/5,
Bonferroni correction for 5 exposures tested). We tested for heterogeneity in effect sizes between
ancestries using Cochran's Q-test[79] and by inspecting the I^2 statistics[80], which is defined as:
540

$$I^2 = \left(\frac{Q - df}{Q}\right) * 100\%$$

542
543 where df is equal to $number\ of\ studies - 1$.

Power calculation

545 To calculate the statistical power for the PGS effect sizes, we used a t-test for linear regression
546 coefficient as implemented in the R package pwrs[81] assuming a standard deviation for the
547 outcome variable SDy = 7.8 (weighted combine SD across studies), a sample size of n = 6 750,
548 a total number of predictors k = 25 and an adjusted r2 = 0.05, to calculate power at P < 0.05
549 significance level.
550

For the coefficients of association with single variants, we estimated statistical power via the non-
centrality parameter (NCP) of the chi-square distribution. We defined NCP = 2 f (1 − f) n β^2
where f is MAF, n is the effective sample size, and β is the expected effect size.[82]
We set a beta of 0.3 (% weight change from baseline) as the expected genetic effect on GLP1
response, based on findings from a recent large-scale study on genetic influences on drug
response.[64]
557

Code Availability

The study utilized previously published analysis tools as described in the **Methods** section. Additional code used for these analyses is available at https://github.com/dsgelab/glp1-bs-genetics/

Data Availability

The Helsinki biobank can provide access for research projects within the scope regulated by the Finnish Biobank Act, which is research utilizing the biobank samples or data for the purposes of promoting health, understanding the mechanisms of disease or developing products and treatment practices used in health and medical care.

De-identified data of the MGBB that supports this study is available from the MGB Biobankportal at https://www.massgeneralbrigham.org/en/research-and-innovation/participate-in-research/biobank/for-researchers. Restrictions apply to the availability of these data, which are available to MGB-affiliated researchers via a formal application.

UK Biobank data are available through a procedure described at http://www.ukbiobank.ac.uk/using-the-resource/.

Registered researchers whose institutions have Data Use and Registration Agreements in place with All of Us that include the Controlled Tier can access genomic data.

Clinical and genotype data from UCLA ATLAS Community Health Initiative patients, de-identified for research purposes, are accessible to UCLA-approved researchers through the Discovery Data Repository (DDR).

BioMe data is available through a process described at https://icahn.mssm.edu/research/ipm/programs/biome-biobank/researcher-faqs.

Pseudonymised data and/or biological samples can be accessed for research and development purposes in accordance with the Estonian Human Genome Research Act. To access data, the research proposal must be approved by the Scientific Advisory Committee of the Estonian Biobank as well as by the Estonian Committee on Bioethics and Human Research. For more details on data access and relevant documents, please see https://genomics.ut.ee/en/content/estonian-biobank#dataaccess.

For access to the Bialystok Bariatric Surgery Study please refer to the Medical University of Bialystok.

The data and biosamples collected or generated by QBB will be made available to researchers
employed within or otherwise contractually bound to public and private institutions that conduct
scientific research and that meet the requirements detailed in the Qatar Biobank Research Access
policy.
The PGS weights used in this study are available in Weissbrod et al. 2022 and Thompson et al.
2022.[75]

Acknowledgements

We gratefully acknowledge the resources provided by the Institute for Precision Health and
participating patients from the UCLA ATLAS Community Health Initiative
(https://www.uclahealth.org/international-services/consulting-services/clinical-research-
consultation/ucla-atlas-community-health-initiative). The UCLA ATLAS Community Health
Initiative in collaboration with UCLA ATLAS Precision Health Biobank is a program of the Institute
for Precision Health, which directs and supports the biobanking and genotyping of biospecimen
samples from participating patients from UCLA in collaboration with the David Geffen School of
Medicine, UCLA Clinical and Translational Science Institute and UCLA Health. The ATLAS
Community Health Initiative is supported by UCLA Health, the David Geffen School of Medicine
and a grant from the UCLA Clinical and Translational Science Institute (UL1TR001881). This work
was financially supported in part by National Institutes of Health awards U01HG011715,
R01HG009120 and R01MH115676. The content is solely the responsibility of the authors and
does not necessarily represent the official views of the National Institutes of Health.
We would furthermore like to thank all participants of HUS
(https://www.helsinginbiopankki.fi/en/Helsinki-Biobank; https://www.hus.fi/en), ESTBB
(https://genomics.ut.ee/en/content/estonian-biobank), BBSS
(https://classic.clinicaltrials.gov/ct2/show/NCT04634591), QBB (https://www.qphi.org.qa/), BioMe
(https://icahn.mssm.edu/research/ipm/programs/biome-biobank), UKBB
(https://www.ukbiobank.ac.uk/), MGBB (https://www.massgeneralbrigham.org/en/research-and-
innovation/participate-in-research/biobank) and AOU (https://allofus.nih.gov/) for their generous
contribution. This work was further supported in part by funding from the Eric and Wendy Schmidt
Center at the Broad Institute of MIT and Harvard.
U.V. and K.A. have been funded by Estonian Research Council's personal research funding start-
up grant PSG759 and European Research Council' starting grant under the grant agreement no
101117251 (OBECAUSE). J.H.L. is supported by NIDDK K23 DK131345.

The activities of the EstBB are regulated by the Human Genes Research Act, which was adopted

in 2000 specifically for the operations of the EstBB. Individual level data analysis in the EstBB

was carried out under ethical approvals of 1.1-12/1409 and 1.1-12/2161 from the Estonian

Committee on Bioethics and Human Research (Estonian Ministry of Social Affairs), using data

according to release application 6-7/GI/18857 from the Estonian Biobank.

References

1. Hruby, A. & Hu, F. B. The Epidemiology of Obesity: A Big Picture. *Pharmacoeconomics* **33**, 673–689 (2015).

2. Boutari, C. & Mantzoros, C. S. A 2022 update on the epidemiology of obesity and a call to action: as its twin COVID-19 pandemic appears to be receding, the obesity and dysmetabolism pandemic continues to rage on. *Metabolism* **133**, 155217 (2022).

3. EASO Secretariat. European survey finds over three-quarters of adults with obesity have attempted to lose weight in the past year, but most have been unsuccessful. *EASO* https://easo.org/european-survey-finds-over-three-quarters-of-adults-with-obesity-have-attempted-to-lose-weight-in-the-past-year-but-most-have-been-unsuccessful/ (2022).

4. Creutzfeldt, W. The incretin concept today. *Diabetologia* **16**, 75–85 (1979).

5. Kreymann, B., Williams, G., Ghatei, M. A. & Bloom, S. R. Glucagon-like peptide-1 7-36: a physiological incretin in man. *Lancet* **2**, 1300–1304 (1987).

6. Nauck, M. A. *et al.* Normalization of fasting glycaemia by intravenous GLP-1 ([7-36 amide] or [7-37]) in type 2 diabetic patients. *Diabet. Med.* **15**, 937–945 (1998).

7. Nauck, M. A. *et al.* Effects of glucagon-like peptide 1 on counterregulatory hormone responses, cognitive functions, and insulin secretion during hyperinsulinemic, stepped hypoglycemic clamp experiments in healthy volunteers. *J. Clin. Endocrinol. Metab.* **87**, 1239–1246 (2002).

8. Amylin and Lilly Announce FDA Approval of BYETTA(TM) (Exenatide) Injection. *Eli Lilly and Company* https://investor.lilly.com/news-releases/news-release-details/amylin-and-lilly-announce-fda-approval-byettatm-exenatide.

9. Secher, A. *et al.* The arcuate nucleus mediates GLP-1 receptor agonist liraglutide-dependent weight loss. *J. Clin. Invest.* **124**, 4473–4488 (2014).

10. Ryan, D. H. *et al.* Long-term weight loss effects of semaglutide in obesity without diabetes

in the SELECT trial. *Nat. Med.* (2024) doi:10.1038/s41591-024-02996-7.

11. Wilding, J. P. H. *et al.* Once-Weekly Semaglutide in Adults with Overweight or Obesity. *N.*

Engl. J. Med. **384**, 989–1002 (2021).

12. Rubino, D. M. *et al.* Effect of Weekly Subcutaneous Semaglutide vs Daily Liraglutide on

Body Weight in Adults With Overweight or Obesity Without Diabetes: The STEP 8

Randomized Clinical Trial. *JAMA* **327**, 138–150 (2022).

13. Marso, S. P. *et al.* Liraglutide and Cardiovascular Outcomes in Type 2 Diabetes. *N. Engl. J.*

Med. **375**, 311–322 (2016).

14. Marso, S. P. *et al.* Semaglutide and Cardiovascular Outcomes in Patients with Type 2

Diabetes. *N. Engl. J. Med.* **375**, 1834–1844 (2016).

15. Perkovic, V. *et al.* Effects of Semaglutide on Chronic Kidney Disease in Patients with Type

2 Diabetes. *N. Engl. J. Med.* **391**, 109–121 (2024).

16. Sattar, N. *et al.* Cardiovascular, mortality, and kidney outcomes with GLP-1 receptor

agonists in patients with type 2 diabetes: a systematic review and meta-analysis of

randomised trials. *Lancet Diabetes Endocrinol* **9**, 653–662 (2021).

17. Watanabe, J. H., Kwon, J., Nan, B. & Reikes, A. Trends in glucagon-like peptide 1 receptor

agonist use, 2014 to 2022. *J. Am. Pharm. Assoc.* **64**, 133–138 (2024).

18. Melson, E., Ashraf, U., Papamargaritis, D. & Davies, M. J. What is the pipeline for future

medications for obesity? *Int. J. Obes.* (2024) doi:10.1038/s41366-024-01473-y.

19. Morgan, J. P. The increase in appetite for obesity drugs.

https://www.jpmorgan.com/insights/global-research/current-events/obesity-drugs (2023).

20. Weight Loss Surgery Requirements. *Duke Health*

https://www.dukehealth.org/treatments/weight-loss-surgery-candidate.

21. Bariatric Surgery Insurance Coverage. *Temple Health*

https://www.templehealth.org/services/bariatric-surgery/for-physicians/insurance-coverage.

22. Clapp, B. *et al.* American Society for Metabolic and Bariatric Surgery 2020 estimate of

metabolic and bariatric procedures performed in the United States. *Surg. Obes. Relat. Dis.*

18, 1134–1140 (2022).

23. After 30 Years — New Guidelines For Weight-Loss Surgery. *American Society for*

Metabolic and Bariatric Surgery https://asmbs.org/news_releases/after-30-years-new-

guidelines-for-weight-loss-surgery/ (2022).

24. Gloy, V. L. *et al.* Bariatric surgery versus non-surgical treatment for obesity: a systematic

review and meta-analysis of randomised controlled trials. *BMJ* **347**, f5934 (2013).

25. Salminen, P. *et al.* Effect of Laparoscopic Sleeve Gastrectomy vs Roux-en-Y Gastric

Bypass on Weight Loss, Comorbidities, and Reflux at 10 Years in Adult Patients With

Obesity: The SLEEVEPASS Randomized Clinical Trial. *JAMA Surg.* **157**, 656–666 (2022).

26. Lim, R., Beekley, A., Johnson, D. C. & Davis, K. A. Early and late complications of bariatric

operation. *Trauma Surg Acute Care Open* **3**, e000219 (2018).

27. Ma, I. T. & Madura, J. A., 2nd. Gastrointestinal Complications After Bariatric Surgery.

Gastroenterol. Hepatol. **11**, 526–535 (2015).

28. Holmberg, D. & Lagergren, J. What do we know about long-term effects of bariatric

surgery? *Br. J. Surg.* **110**, 1–2 (2022).

29. Doumouras, A. G. *et al.* Bariatric surgery and cardiovascular outcomes in patients with

obesity and cardiovascular disease:: A population-based retrospective cohort study: A

population-based retrospective cohort study. *Circulation* **143**, 1468–1480 (2021).

30. Kuno, T., Tanimoto, E., Morita, S. & Shimada, Y. J. Effects of bariatric surgery on

cardiovascular disease: A concise update of recent advances. *Front. Cardiovasc. Med.* **6**,

94 (2019).

31. Elsaid, M. I. *et al.* Association of bariatric surgery with cardiovascular outcomes in adults

with severe obesity and nonalcoholic fatty liver disease. *JAMA Netw. Open* **5**, e2235003

(2022).

32. Adams, T. D. *et al.* Long-term all-cause and cause-specific mortality for four bariatric

surgery procedures. *Obesity (Silver Spring)* **31**, 574–585 (2023).

33. Azagury, D. *et al.* Heterogeneity of weight loss after gastric bypass, sleeve gastrectomy,

and adjustable gastric banding. *Surgery* **165**, 565–570 (2019).

34. Cadena-Obando, D. *et al.* Are there really any predictive factors for a successful weight

loss after bariatric surgery? *BMC Endocr. Disord.* **20**, 20 (2020).

35. Çalık Başaran, N., Dotan, I. & Dicker, D. Post metabolic bariatric surgery weight regain: the

importance of GLP-1 levels. *Int. J. Obes.* (2024) doi:10.1038/s41366-024-01461-2.

36. Lauti, M., Kularatna, M., Hill, A. G. & MacCormick, A. D. Weight Regain Following Sleeve

Gastrectomy-a Systematic Review. *Obes. Surg.* **26**, 1326–1334 (2016).

37. Young, K. G. *et al.* Treatment effect heterogeneity following type 2 diabetes treatment with

GLP1-receptor agonists and SGLT2-inhibitors: a systematic review. *Commun. Med.* **3**, 131

(2023).

38. Jones, A. G. *et al.* Markers of β-Cell Failure Predict Poor Glycemic Response to GLP-1

Receptor Agonist Therapy in Type 2 Diabetes. *Diabetes Care* **39**, 250–257 (2016).

39. Dawed, A. Y. *et al.* Pharmacogenomics of GLP-1 receptor agonists: a genome-wide

analysis of observational data and large randomised controlled trials. *Lancet Diabetes*

Endocrinol **11**, 33–41 (2023).

40. Cardoso, P. *et al.* Phenotype-based targeted treatment of SGLT2 inhibitors and GLP-1

receptor agonists in type 2 diabetes. *Diabetologia* **67**, 822–836 (2024).

41. Brown, E., Wilding, J. P. H., Barber, T. M., Alam, U. & Cuthbertson, D. J. Weight loss

variability with SGLT2 inhibitors and GLP-1 receptor agonists in type 2 diabetes mellitus

and obesity: Mechanistic possibilities. *Obes. Rev.* **20**, 816–828 (2019).

42. Umapathysivam, M. M. *et al.* Type 2 Diabetes risk alleles in Peptidyl-glycine Alpha-

amidating Monooxygenase influence GLP-1 levels and response to GLP-1 Receptor

Agonists. *medRxiv* (2023) doi:10.1101/2023.04.07.23288197.

43. Jensterle, M., Rizzo, M., Haluzík, M. & Janež, A. Efficacy of GLP-1 RA Approved for Weight

730  Management in Patients With or Without Diabetes: A Narrative Review. *Adv. Ther.* **39**,

731  2452–2467 (2022).

732 44. Nathan, B. M. *et al.* Predictors of weight-loss response with glucagon-like peptide-1

733  receptor agonist treatment among adolescents with severe obesity. *Clin. Obes.* **6**, 73–78

734  (2016).

735 45. Vosoughi, K. *et al.* Association of Glucagon-like Peptide 1 Analogs and Agonists

736  Administered for Obesity with Weight Loss and Adverse Events: A Systematic Review and

737  Network Meta-analysis. *EClinicalMedicine* **42**, 101213 (2021).

738 46. Segal, J. B. *et al.* Assessing Heterogeneity of Treatment Effect in Real-World Data. *Ann.*

739  *Intern. Med.* **176**, 536–544 (2023).

740 47. Jensterle, M., Pirš, B., Goričar, K., Dolžan, V. & Janež, A. Genetic variability in GLP-1

741  receptor is associated with inter-individual differences in weight lowering potential of

742  liraglutide in obese women with PCOS: a pilot study. *Eur. J. Clin. Pharmacol.* **71**, 817–824

743  (2015).

744 48. van der Meer, R. *et al.* Genetic variants associated with weight loss and metabolic

745  outcomes after bariatric surgery: A systematic review. *Obes. Rev.* **24**, e13626 (2023).

746 49. Aasbrenn, M. *et al.* Genetic Determinants of Weight Loss After Bariatric Surgery. *Obes.*

747  *Surg.* **29**, 2554–2561 (2019).

748 50. de Toro-Martín, J. *et al.* Polygenic risk score for predicting weight loss after bariatric

749  surgery. *JCI Insight* **3**, (2018).

750 51. NCD Risk Factor Collaboration (NCD-RisC). Worldwide trends in underweight and obesity

751  from 1990 to 2022: a pooled analysis of 3663 population-representative studies with 222

752  million children, adolescents, and adults. *Lancet* **403**, 1027–1050 (2024).

753 52. White, G. E. *et al.* Real-world weight-loss effectiveness of glucagon-like peptide-1 agonists

754  among patients with type 2 diabetes: A retrospective cohort study. *Obesity* **31**, 537–544

755  (2023).

53. Inoue, K. *et al.* Long-term impact of liraglutide, a glucagon-like peptide-1 (GLP-1) analogue,

on body weight and glycemic control in Japanese type 2 diabetes: an observational study.

Diabetol. Metab. Syndr. **6**, 95 (2014).

54. Venkatesh, S. S. *et al.* The genetic architecture of changes in adiposity during adulthood.

medRxiv (2023) doi:10.1101/2023.01.09.23284364.

55. Xu, J. *et al.* Exploring the clinical and genetic associations of adult weight trajectories using

electronic health records in a racially diverse biobank: a phenome-wide and polygenic risk

study. *Lancet Digit Health* **4**, e604–e614 (2022).

56. Karczewski, K. J. *et al.* The mutational constraint spectrum quantified from variation in

141,456 humans. *Nature* **581**, 434–443 (2020).

57. Lagou, V. *et al.* GWAS of random glucose in 476,326 individuals provide insights into

diabetes pathophysiology, complications and treatment stratification. *Nat. Genet.* **55**, 1448–

1461 (2023).

58. Nead, K. T. *et al.* Contribution of common non-synonymous variants in PCSK1 to body

mass index variation and risk of obesity: a systematic review and meta-analysis with

evidence from up to 331 175 individuals. *Hum. Mol. Genet.* **24**, 3582–3594 (2015).

59. Vickers, A. J. The use of percentage change from baseline as an outcome in a controlled

trial is statistically inefficient: a simulation study. *BMC Med. Res. Methodol.* **1**, 6 (2001).

60. Curran-Everett, D. Explorations in statistics: the analysis of ratios and normalized data.

Adv. Physiol. Educ. **37**, 213–219 (2013).

61. Berry, D. A. & Ayers, G. D. Symmetrized Percent Change for Treatment Comparisons. *The*

American Statistician (2006) doi:10.1198/000313006X90684.

62. Lepage, B., Lamy, S., Dedieu, D., Savy, N. & Lang, T. Estimating the causal effect of an

exposure on change from baseline using directed acyclic graphs and path analysis.

Epidemiology **26**, 122–129 (2015).

63. Glymour, M. M., Weuve, J., Berkman, L. F., Kawachi, I. & Robins, J. M. When is baseline

adjustment useful in analyses of change? An example with education and cognitive

change. *Am. J. Epidemiol.* **162**, 267–278 (2005).

64. Sadler, M. C. *et al.* Leveraging large-scale biobank EHRs to enhance pharmacogenetics of

cardiometabolic disease medications. *medRxiv* (2024) doi:10.1101/2024.04.06.24305415.

65. Weiss, T. *et al.* Real-world weight change, adherence, and discontinuation among patients

with type 2 diabetes initiating glucagon-like peptide-1 receptor agonists in the UK. *BMJ*

Open Diabetes Res Care **10**, (2022).

66. Ghusn, W. *et al.* Weight Loss Outcomes Associated With Semaglutide Treatment for

Patients With Overweight or Obesity. *JAMA Netw Open* **5**, e2231982 (2022).

67. Xu, T. *et al.* Timing of Maximal Weight Reduction Following Bariatric Surgery: A Study in

Chinese Patients. *Front. Endocrinol.* **11**, 615 (2020).

68. Monpellier, V. M., Janssen, I. M. C., Antoniou, E. E. & Jansen, A. T. M. Weight Change

After Roux-en Y Gastric Bypass, Physical Activity and Eating Style: Is There a

Relationship? *Obes. Surg.* **29**, 526–533 (2019).

69. Knop, F. K. *et al.* Oral semaglutide 50 mg taken once per day in adults with overweight or

obesity (OASIS 1): a randomised, double-blind, placebo-controlled, phase 3 trial. *Lancet*

402, 705–719 (2023).

70. Hall, K. D. & Kahan, S. Maintenance of Lost Weight and Long-Term Management of

Obesity. *Med. Clin. North Am.* **102**, 183–197 (2018).

71. Mok, J. *et al.* Safety and Efficacy of Liraglutide, 3.0 mg, Once Daily vs Placebo in Patients

With Poor Weight Loss Following Metabolic Surgery: The BARI-OPTIMISE Randomized

Clinical Trial. *JAMA Surg.* **158**, 1003–1011 (2023).

72. Borrell, L. N. *et al.* Race and Genetic Ancestry in Medicine - A Time for Reckoning with

Racism. *N. Engl. J. Med.* **384**, 474–480 (2021).

73. Akshatha, S. & Nayak, U. B. The psychological distress associated with type 2 diabetes

mellitus represents an unmet need for drug discovery. *Med. Drug Discov.* **23**, 100196

(2024).

74. Kemper, K. E. *et al.* Genetic influence on within-person longitudinal change in

anthropometric traits in the UK Biobank. *Nat. Commun.* **15**, 3776 (2024).

75. Weissbrod, O. *et al.* Leveraging fine-mapping and multipopulation training data to improve

cross-population polygenic risk scores. *Nat. Genet.* **54**, 450–458 (2022).

76. Chang, C. C. *et al.* Second-generation PLINK: rising to the challenge of larger and richer

datasets. *Gigascience* **4**, 7 (2015).

77. Thompson, D. J. *et al.* UK Biobank release and systematic evaluation of optimised

polygenic risk scores for 53 diseases and quantitative traits. *bioRxiv* (2022)

doi:10.1101/2022.06.16.22276246.

78. Schwarzer, G., Carpenter, J. R. & Rücker, G. *Meta-Analysis with R*. (Springer International

Publishing, 2015).

79. Cochran, W. G. The Combination of Estimates from Different Experiments. *Biometrics* **10**,

101–129 (1954).

80. Higgins, J. P. T. & Thompson, S. G. Quantifying heterogeneity in a meta-analysis. *Stat.*

Med. **21**, 1539–1558 (2002).

81. Bulus, M. A practical guide to statistical power and sample size calculations in R.

https://cran.r-project.org/web/packages/pwrss/vignettes/examples.html (2023).

82. Kanai, M. *et al.* Insights from complex trait fine-mapping across diverse populations.

bioRxiv (2021) doi:10.1101/2021.09.03.21262975.

828