Association between plausible genetic factors and weight loss from GLP1-RA and bariatric surgery: a multi-ancestry study in 10 960 individuals from 9 biobanks

Jakob German*[1,2], Mattia Cordioli*[1,3], Veronica Tozzo[4], Sarah Urbut[5,6,7], Kadri Arumäe[8], Roelof A.J. Smit[9,10,11,12], Jiwoo Lee[3,6], Josephine H. Li[6,13,14,15], Adrian Janucik[14,16], Yi Ding[4], Akintunde Akinkuolie[4], Henrike Heyne[17], Andrea Eoli[18,19], Chadi Saad[20], Yasser Al-Sarraj[20], Rania Abdel-latif[20], Alexandra Barry[6,14], Zhe Wang[9], Estonian Biobank research team[31], Pradeep Natarajan[6,14,21,22], Samuli Ripatti[1,14,23, 33], Anthony Philippakis[2,3], Lukasz Szczerbinski[6,13,24,25,26], Bogdan Pasaniuc[4,27,28,29,30,], Hamdi Mbarek[20], Ruth J.F. Loos[9,10,11,12], Uku Vainik[8,31,32,+], Andrea Ganna[1,33,+]

* These authors contributed equally
+ These authors jointly supervised the project

1. Institute for Molecular Medicine Finland (FIMM), University of Helsinki, Helsinki, Finland;
2. Eric and Wendy Schmidt Center, Broad Institute of MIT and Harvard, Cambridge, MA, USA, 02142;
3. Broad Institute of MIT and Harvard, Cambridge, MA, USA;
4. Department of Computational Medicine, David Geffen School of Medicine at UCLA, Los Angeles, CA, USA;
5. Division of Cardiovascular Medicine, Massachusetts General Hospital, Boston, MA;
6. Center for Genomic Medicine, Massachusetts General Hospital, Boston, MA;
7. Cardiovascular Research Center, Massachusetts General Hospital, Boston, MA;
8. Institute of Psychology, Faculty of Social Sciences, University of Tartu, Tartu, Estonia;
9. Charles Bronfman Institute for Personalized Medicine, Icahn School of Medicine at Mount Sinai, New York, NY, USA;
10. Department of Environmental Medicine and Climate Science, Icahn School of Medicine at Mount Sinai, New York, NY, USA;
11. Novo Nordisk Foundation Center for Basic Metabolic Research, Faculty of Health and Medical Sciences, University of Copenhagen, Copenhagen, Denmark;
12. Novo Nordisk Foundation Center for Genomic Mechanisms of Disease, Broad Institute of MIT and Harvard, Cambridge, MA, USA;
13. Diabetes Unit, Massachusetts General Hospital, Boston, MA, USA;
14. Program in Medical & Population Genetics, Broad Institute of MIT and Harvard, Cambridge, MA, USA;
15. Harvard Medical School, Boston, MA, USA;
16. Center for Digital Medicine, Medical University of Bialystok, Bialystok, Poland;
17. Hasso Plattner Institut, Potsdam, Germany;
18. Digital Engineering Faculty, University of Potsdam, Potsdam, Germany, Prof.-Dr.-Helmert-Str. 2-3, 14482;
19. Hasso Plattner Institute for Digital Health at Mount Sinai, Icahn School of Medicine at Mount Sinai, New York City, NY, USA;
20. Qatar Genome Program, Qatar Precision Health Institute, Qatar Foundation, Doha, Qatar;

21. Personalized Medicine, Mass General Brigham, Boston, MA, USA;

22. Department of Medicine, Harvard Medical School, Boston, MA, USA;

23. Department of Public Health, Clinicum, Faculty of Medicine, University of Helsinki,

Helsinki, Finland;

24. Programs in Metabolism and Medical & Population Genetics, Broad Institute of MIT and

Harvard, Cambridge, MA, USA;

25. Department of Endocrinology, Diabetology and Internal Medicine, Medical University of

Bialystok, Bialystok, Poland;

26. Clinical Research Centre, Medical University of Bialystok, Bialystok, Poland;

27. Bioinformatics Interdepartmental Program, UCLA, Los Angeles, CA, USA;

28. Department of Pathology and Laboratory Medicine, David Geffen School of Medicine at

UCLA, Los Angeles, CA, USA;

29. Department of Human Genetics, David Geffen School of Medicine at UCLA, Los

Angeles, CA, USA;

30. Institute for Precision Health, UCLA, Los Angeles, CA, USA;

31. Estonian Genome Centre, Institute of Genomics, Faculty of Science and Technology,

University of Tartu, Tartu, Estonia ;

32. Department of Neurology and Neurosurgery, Faculty of Medicine and Health Sciences,

McGill University, Canada;

33. Analytic & Translational Genetics Unit, Massachusetts General Hospital, Harvard

Medical School, Boston, MA, USA;

Correspondence to Andrea Ganna (andrea.ganna@helsinki.fi)

Abstract

Obesity is a significant public health concern. GLP-1 receptor agonists (GLP1-RA), predominantly in use as a type 2 diabetes treatment, are a promising pharmacological approach for weight loss, while bariatric surgery (BS) remains a durable, but invasive, intervention. Despite observed heterogeneity in weight loss effects, the genetic effects on weight loss from GLP1-RA and BS have not been extensively explored in large sample sizes, and most studies have focused on differences in race and ethnicity, rather than genetic ancestry. We studied whether genetic factors, previously shown to affect body weight, impact weight loss due to GLP1-RA therapy or BS in 10,960 individuals from 9 multi-ancestry biobank studies in 6 countries. The average weight change between 6 and 12 months from therapy initiation was -3.93% for GLP1-RA users, with marginal differences across genetic ancestries. For BS patients the weight change between 6 and 48 months from the operation was -21.17%. There were no significant associations between weight loss due to GLP1-RA and polygenic scores for BMI or type 2 diabetes or specific missense variants in the *GLP1R, PCSK1 and APOE* genes, after multiple-testing correction. However, a higher polygenic score for BMI was significantly linked to lower weight loss after BS (+0.7% for 1 standard deviation change in the polygenic score, $P = 1.24 \times 10^{-4}$). In contrast, higher weight at baseline was associated with greater weight loss. Our findings suggest that existing polygenic scores related to weight and type 2 diabetes and missense variants in the drug target gene do not have a large impact on GLP1-RA effectiveness. Our results also confirm the effectiveness of these treatments across all major continental ancestry groups considered.

Introduction

The obesity epidemic presents a significant public health burden, driving the need for effective treatment options.[1,2] Despite 75% of adults with obesity having attempted to lose weight, most have not achieved lasting success.[3] Amid these challenges, glucagon-like peptide-1 receptor agonists (GLP1-RA) have emerged as a promising solution. As first discovered in the late 1980s, glucagon-like peptide-1 (GLP-1) is linked to the incretin effect, which promotes insulin release from pancreatic beta cells, thus playing a crucial role in regulating blood sugar levels.[4,5] This evidence led to the approval of the first GLP1-RA for treatment of type 2 diabetes in 2005.[6–8] Nine years later, an association between GLP-1 and reduced food intake paved the way for the development and approval of the first GLP1-RA for obesity treatment.[5,9]

Clinical trials conducted in the following years demonstrated weight loss effects ranging from 6% to 16% for liraglutide and semaglutide, leading to a substantial increase in GLP1-RA usage worldwide[10–12]. Additional trials have further revealed cardiovascular[13,14] and renal benefits[15], with ongoing trials currently underway.[16] Overall, anticipated trends suggest that by 2030, the US will witness approximately 30 million patients using GLP1-RA drugs, representing roughly 9% of its population, underscoring their potential in combating the obesity crisis and exploring new therapeutic applications.[17–19]

Meanwhile, bariatric surgery (BS) stands as a durable and effective treatment for individuals with severe obesity (BMI >= 40 kg/m^2) and metabolic comorbidities, and consistent with its effectiveness, the threshold has been reduced to a BMI >=30 kg/m^2.[20,21] Meta-analyses combining results from randomized controlled trials and observational studies found that, on average, individuals undergoing BS experienced a weight loss of 26 kg.[22,23] However, such an invasive procedure inherently carries increased risks of severe adverse events.[24,25]

Despite the robust weight loss effects observed with both GLP1-RA treatment and BS, there exists heterogeneity in their effectiveness. One out of every five patients undergoing BS may not achieve the desired weight loss within the first year or may experience weight regain within two years.[26–29] The variability of GLP1-RA treatment effects on glycemic outcomes has been widely investigated and can be partially explained by clinical indicators like low β-cell function or C-peptide levels as well as genetic variants.[30–35] However, with the exception of type 2 diabetes,

sex, and baseline weight, the factors influencing the diversity in weight loss effects of these drugs

are still poorly understood.[36–38] Additionally, differences in patient adherence to treatment

regimens may further contribute to this variability.[39]

Genetics has been considered a potential factor for heterogeneity in weight loss treatments, but

its role remains largely unexplored.[36] A previous study has examined the role of a common

missense variant in the *GLP1R* genes in 57 women and found a weak link with GLP1-RA

treatment response.[32,40] A greater number and better-powered studies have been conducted in

individuals with BS [41,42], however, a consistent genome-wide association for weight loss has yet

to be identified.

Finally, there has been limited exploration into the effectiveness of these obesity treatments

across different genetic ancestries and across various healthcare settings.[10,43]

In our study we seek to i) characterize the observed GLP1-RA and BS body weight-lowering

effects using real-world data from 10 960 individuals across 9 biobanks in 6 countries ,with 6

major continental ancestry groups; ii) identify whether plausible genetic factors (i.e. polygenic

scores for BMI and T2D and coding variants in *GLP1R*, *PCSK1* and *APOE* gene) associate with

heterogeneity in GLP1-RA and BS weight loss effects, and thus identify patient groups that will

benefit most from such treatments; iii) compare the effects of GLP1-RA treatment and BS on

weight loss across major continental ancestry groups.

Results

GLP1-RA and bariatric surgery associated weight loss in 10 960 individuals from 9 cohorts and 6 major continental ancestry groups

We investigated changes in body weight associated with GLP1-RA treatment and BS across 9 biobank studies: Helsinki University Hospital (HUS, Finland; N = 633), Estonian Biobank (ESTBB; N = 464), UK Biobank (UKBB, United Kingdom; N = 810), All of Us (AoU, USA; N = 559), BioMe Biobank (BioMe, USA; N = 2,170), Mass General Brigham Biobank (MGBB, USA; N = 2,141), Atlas Biobank (UCLA-ATLAS, USA; N = 1,445), Qatar Biobank (QBB, Qatar; N = 2,383), Bialystok Bariatric Surgery Study (BBSS, Poland, N = 355). We included individuals that were at least 18 years of age when starting a GLP1-RA treatment or undergoing BS. Individuals undergo GLP1-RA treatment for different medical reasons, not only for weight loss. We defined GLP1-RA usage based on prescription or purchase data (ATC codes starting with A10B*; including exenatide, liraglutide, lixisenatide, albiglutide, dulaglutide, semaglutide, and beinaglutide) covering at least 12 months, and excluding individuals with indication of treatment discontinuation within this time window. For BS, we considered individuals undergoing the following procedures: Roux-en-Y gastric bypass, sleeve gastrectomy, adjustable gastric band, vertical banded gastroplasty, biliopancreatic diversion with duodenal switch. More information can be found in the **Methods** section.

We included only individuals who had an initial body weight measurement recorded at most 12 months prior to the baseline (defined as treatment initiation or surgery) and a follow-up measurement taken at least 6 but no more than 12 months after the baseline for GLP1-RA treatments, or within 48 months after baseline for BS. In case of multiple measurements within this time frame, we considered the median of such measurements (**Supplementary Figure 1** for a schematic of the approach used). We defined our main outcome of interest as the percentage change in body weight, calculated as the difference between the second body weight measurement (or median of multiple measurements) and the initial one, divided by the initial measurement .

Our study population included a total of 6,750 GLP1-RA users and 4,210 individuals undergoing BS (baseline characteristics are reported in **Table 1**). Overall, GLP1-RA users 1) had a lower

174 proportion of women (61.0% weighted average across studies) when compared to BS patients
175 (73.1% weighted average across studies), 2) were on average older (57.30 *vs* 46.50), and 3) had
176 a lower initial body weight (99.80 kg *vs* 116.96 kg). The proportion of individuals diagnosed with
177 type 2 diabetes at time of GLP1-RA initiation was at least 48% across all studies and major
178 continental ancestry groups.
179

Study	Ancestry	Number of individuals	Proportion of women (%)	Mean body weight (in Kg) at baseline (SD)	Mean age at baseline (SD)	T2D prevalence (%)	Most common GLP1-RA type	Proportion of semaglutide (%)
GLP1-RA								
HUS	EUR	255	68.00	113.70 (22.90)	53.79 (13.52)	65.49	semaglutide	60.40
ESTBB	EUR	191	62.30	109.18 (20.3)	56.95 (10.96)	89.01	semaglutide	72.77
UKBB	EUR	614	44.30	105.00 (19.90)	60.50 (7.15)	99.80	liraglutide	0.00
AoU	EUR	117	64.96	106.66 (24.18)	57.01 (11.03)	82.91	liraglutide	11.97
	AFR	91	83.52	104.8 (25.01)	54.57 (11.93)	92.31	liraglutide	18.68
MGBB	EUR	857	57.01	99.99 (21.48)	59.88 (12.17)	56.33	semaglutide	40.72
	AFR	156	54.32	104.65 (23.80)	54.31 (11.91)	60.24	dulaglutide	36.14
BioMe	AMR	614	66.45	91.19 (22.45)	59.07 (11.29)	67.75	dulaglutide	29.32
	AFR	728	69.78	101.49 (25.42)	57.00 (12.24)	69.50	dulaglutide	29.12
	EUR	369	44.44	98.26 (21.93)	60.32 (11.57)	47.97	semaglutide	37.13
	SAS	77	50.65	83.79 (20.10)	56.48 (12.59)	70.13	semaglutide	41.56
UCLA-ATLAS	AFR	133	64.00	101.6 (23.20)	61.00 (13.00)	91.00	semaglutide	57.10
	AMR	336	66.00	94.40 (22.60)	56.00 (13.00)	84.00	liraglutide	26.50
	EAS	120	47.00	81.50 (16.30)	60.00 (13.00)	92.00	semaglutide	61.70
	EUR	856	53.00	101.60 (22.00)	63.00 (13.00)	80.00	semaglutide	67.80
QBB	MID	1236	72.17	97.68 (24.54)	49.52 (10.75)	75,97	liraglutide	3.64
Bariatric surgery								
HUS	EUR	378	81.00	113.63 (19.5)	49.90 (9.39)	37.04		
ESTBB	EUR	273	76.56	121.24 (20.57)	44.72 (10.07)	28.21		
UKBB	EUR	196	69.40	121.00 (29.70)	55.50 (8.21)	50.00		
AoU	EUR	168	81.55	121.32 (29.88)	50.74 (12.62)	18.45		
	AFR	183	89.62	127.23 (32.55)	46.09 (11.67)	48.63		
MGBB	EUR	978	71.06	121.21 (26.52)	49.37 (12.16)	39.98		
	AFR	150	86.67	122.89 (23.92)	43.68 (11.67)	42.67		
BioMe	AMR	143	79.72	112.36 (20.81)	44.62 (11.48)	30.07		
	AFR	188	82.98	124.88 (25.63)	45.35 (12.16)	29.26		
	EUR	51	70.59	120.23 (25.29)	49.51 (13.34)	17.65		
QBB	MID	1147	69.49	102.14 (23.18)	41.66 (10.66)	40.02		
BBSS	EUR	355	56.06	138.48 (27.07)	47.10 (7.11)	35.21		

Table 1 | Baseline characteristics for 6 750 GLP1-RA users and 4 210 patients undergoing bariatric surgery included in our study population. For each cohort and major continental ancestry group (AFR: African, AMR: admixed American, EAS: East Asian, EUR: European, MID: Middle Eastern, SAS: South Asian), we report the total number of individuals eligible for analysis, the proportion of females, the mean (SD) body weight (in Kg) at baseline, the mean (SD) age (in years) at baseline, the prevalence of T2D, and (for GLP1-RA users only) the most commonly used type of medication and the proportion of semaglutide users.
HUS, Helsinki University Hospital; ESTBB, Estonian Biobank; UKBB, UK Biobank; AoU, All of US; MGBB, Mass General Brigham Biobank; BioMe, BioMe; UCLA-ATLAS, Atlas Biobank; QBB, Qatar Biobank; BBSS, Bialystok Bariatric Surgery Study

Among GLP1-RA users, we observed an average body weight change (**Figure 1**) across studies of -3.93% (ranging from -1.08% to -7.07%), in line with results from observational studies.[44,45] Among BS patients, we observed an average change of -21.17% (-15.00% to -27.72%).

Figure 1 | Average percentage change in body weight for 6 750 GLP1-RA users and 4 210 bariatric surgery patients. Bars represent the mean percentage change in body weight within each study and ancestry, and the overall combined weighted mean change. Error bars represent standard errors.

Demographic factors associated with GLP1-RA and bariatric surgery weight change

We explored the effect of baseline characteristics on body weight changes associated with GLP1-RA treatment and BS. We fitted a multivariable linear model with percentage change in body weight as outcome and baseline body weight, sex, age at baseline, the first 20 principal genetic components and medication type (for GLP1-RA users only) as predictors. Each cohort was analyzed separately, stratifying by major continental ancestry group, and the effects were then meta-analyzed (**Table 2**). Baseline body weight was significantly associated with greater weight loss for both GLP1-RA and BS treatments ($\beta_{GLP1-RA}$ = -0.05 % weight change compared to baseline, P = 1.63 × 10^{-35}; β_{BS} = -0.14 % weight change compared to baseline, P = 5.49 × 10^{-69}) and women had a significantly greater weight loss for both treatments ($\beta_{GLP1-RA}$ = -1.54 % weight

change compared to baseline, $P = 2.47 \times 10^{-16}$; β_{BS} = -2.66 % weight change compared to
baseline, $P = 6.83 \times 10^{-10}$). Being older at treatment initiation was only associated with less weight
loss for bariatric surgery (β_{BS} = 0.15 % weight change compared to baseline, $P = 2.54 \times 10^{-15}$).
The effects of baseline weight and sex on weight loss were highly heterogeneous across
ancestries and studies (I^2 for GLP1-RA= 0.99 and 0.83 for baseline weight and sex, respectively)
(**Supplementary Figures 2-4 and Supplementary Tables 1-2**).

203
204

Variable	GLP1-RA			Bariatric surgery		
	Effect on percentage weight change	Standard error	P value	Effect on percentage weight change	Standard error	P value
Baseline weight (kg)	-0.05	0.004	1.63×10^{-35}	-0.14	0.008	5.49×10^{-69}
Sex (female)	-1.54	0.188	2.74×10^{-16}	-2.66	0.431	6.83×10^{-10}
Age at baseline (years)	-0.02	0.008	0.052	0.15	0.016	9.85×10^{-19}

Table 2 | Effect of baseline weight, sex and age at baseline of percentage weight changes associated with GLP1-RA and bariatric surgery. Multivariable model adjusting for the three variables described in the table and the first 20 principal genetic components and medication type (for GLP1-RA users only). Negative effects indicate larger weight loss.

Marginal effect of ancestry on GLP1-RA and bariatric surgery associated weight loss

We evaluate whether the effect of GLP1-RA and BS associated weight loss was significantly
different across genetically-defined continental ancestry groups after accounting for differences
in weight at baseline, age, sex and, only for GLP1-RA, medication type (**Table 3**). Only multi-
ancestry biobanks (AoU, BioMe, MGBB, UCLA-ATLAS) were included in this analysis (N = 6 315).
Compared to individuals of European ancestry, all other ancestry groups had lower change in
body weight. However, this effect was statistically significant only among GLP1-RA users of
African (β = 0.76 % weight change from baseline weight, $P = 0.02$) and admixed American (β =
0.87 % weight change from baseline weight, $P = 3 \times 10^{-3}$) ancestries.

GLP1-RA			
Major continental ancestry group	**Effect on percentage weight change**	**Standard error**	**P value**
AFR	0.758	0.327	0.021
AMR	0.874	0.295	0.003
EAS	1.093	0.664	0.100
SAS	0.118	0.759	0.876
Bariatric surgery			
Major continental ancestry group	**Effect on percentage weight change**	**Standard error**	**P value**
AFR	1.688	1.308	0.197
AMR	-0.542	1.428	0.704

Table 3 | Associations between weight change and major continental ancestry groups. Meta-analysis effect sizes, standard errors and P values for the association between weight change and major continental ancestry groups (AFR: African, AMR: admixed American, EAS: East Asian, EUR: European, SAS: South Asian). Coefficients are derived from a multivariable model including percentage weight change as outcome and ancestry as categorical variable (with EUR as reference category), and adjusting for sex, age and weight at baseline and, for GLP1-RA only, medication type. Negative effects indicate larger weight loss

Plausible genetic factors do not associate with weight change due to GLP1-RA

We investigated the effect of genetic exposures with plausible effects on body weight changes. Genetic exposures were selected upon discussion between authors and before initiating analyses. First, we considered polygenic scores for BMI based on previous observation that a polygenic score was associated with BMI trajectories.[46,47] We also consider the only replicated genome-wide variant associated with BMI change in the largest study of BMI trajectories (*rs429358* in *APOE*).[46] Second, we considered a polygenic score for T2D based on the observation that biomarkers of type 2 diabetes were associated with effectiveness of GLP1-RA.[31,33] We confirmed that both polygenic scores for BMI and T2Ds were significantly associated with their respective traits across all major continental ancestry groups (**Supplementary Table 3**) although the effects were larger in individuals of European ancestry.

Third, we considered 13 SNPs in the *GLP1R* and *PCSK1* genes. We considered all missense

variants in the *GLP1R* gene with at least 1% minor allele frequency in at least one of major

continental ancestry group, as per gnomAD (version 4.1)[48], three of which have been functional

characterized as gain or loss of function (**Supplementary Table 4**).[49] We also considered two

non-synonymous variants in the *PCSK1* gene that have been associated with BMI variation in the

population.[50] Finally, we sought to replicate the effect of two variants associated with weight loss

in one of the largest studies of BS[42](**Methods**). After analyses, two missense variants in the

GLP1R genes, *rs2295006* and *rs201672448* were too rare to be tested across studies and were

not considered further.

We estimated the effect of each genetic exposure on weight change in each cohort and major

continental ancestry groups separately, fitting a linear model adjusted for baseline body weight,

sex, age at baseline, first 20 principal genetic components, and medication type (for GLP1-RA

users only), and we subsequently combined the effects through ancestry-specific and multi-

ancestry meta-analyses (**Figure 2, Supplementary Tables 5-6**). As the approach of adjusting

longitudinal change phenotypes for the baseline trait has been criticized for potentially inducing

biases for genetic variants associated with the baseline trait[51], we also considered a second model

using post-treatment weight as outcome, including the same predictors as in the model above

and an additional term modeling the interaction between baseline weight and each genetic

exposure (**Supplementary Tables 7-8**).

Among GLP1-RA users, we did not observe any significant genetic exposure associated with

weight loss after multiple testing correction ($P < 0.004$, Bonferroni correction for 13 exposures

tested). Using the alternative model, we identified one missense variant in the *GLP1R* genes,

rs3765467, for which we observed a statistically significant interaction with baseline weight

($\beta_{w0*rs3765467} = 0.09$ kg weight change from baseline weight, $P = 2.7 \times 10^{-4}$) on the weight measured

post GLP1-RA treatment after multiple testing correction. When considering nominal significance,

another missense variant in the *GLP1R* genes*, rs6923761*, had a statistically significant

interaction with baseline weight ($\beta_{w0*rs6923761} = 0.02$ kg weight change from baseline weight, $P =$

0.01) on the weight measured post GLP1-RA treatment.

We further tested whether the lack of associations observed was possibly due to high effect-size

heterogeneity across ancestries, or due to lack of statistical power. We did not observe any

significant heterogeneity across ancestries for any of the genetic exposure (average I^2 across

genetic exposures for GLP1-RA was 0.06 **- Supplementary Tables 9-10**). Moreover, given the

current sample size for GLP1-RA users (N = 6 750) our power calculation showed we would have

at least 80% power to detect a statistically significant effect (at *P* < 0.05) of at least a 0.3% change

in body weight for 1 standard deviation (SD) change in the polygenic score, or for SNPs with a

minor allele frequency of at least 1% (**Methods**).

Figure 2 | Effect of 15 genetic exposures on body weight changes associated with GLP1-RA treatment and bariatric surgery.
a. Ancestry-specific and multi-ancestry meta-analysis effect sizes for association between percentage change in body weight and PGS for BMI and type 2 diabetes. Dots represent the percentage change in body weight per one standard deviation change in PGS, error bars represent the 95% confidence interval. Full dots represent statistical significance at P < 0.004.
b. Multi-ancestry meta-analysis effect sizes for the association between percentage change in body weight and genotype at each locus. Dots represent the effect size, error bars represent the 95% confidence interval. PGS, polygenic score.

270
271

A polygenic score for BMI associates with weight change due to

BS

Among BS patients, a higher polygenic score for BMI was significantly associated with lower

weight loss ($\beta_{BMI\ PGS}$ = 0.70 % weight change compared to baseline for 1 SD change in the

polygenic score, *P* = 1.24 × 10⁻⁴) after multiple testing correction (*P* < 0.01, Bonferroni correction

for 5 exposures tested), in the opposite direction than observed for weight at baseline (**Figure 2**).

A higher polygenic score for T2D was nominally associated with lower weight loss ($\beta_{T2D\ PGS}$ = 0.45

% weight change compared to baseline for 1 SD change in the polygenic score, P = 0.03).

We observed significant heterogeneity in the association between BMI PGS and weight loss

across studies (I2 = 0.64, **Supplementary Table 10**).

Discussion

In this study we provide a comprehensive assessment of major demographic factors and plausible

genetic factors on body weight changes associated with GLP1-RA use and BS.

In line with previous work,[36,52–55] our main outcome of interest was the percentage change in body

weight between 6 and 12 months after GLP1-RA treatment initiation as well as between 6 and 48

months after BS.

By considering studies from 6 different countries, we were able to highlight the heterogeneity in

use of GLP1-RA across different healthcare systems. GLP1-RA users consisted mostly of

individuals with type 2 diabetes, reflecting their original use for treatment of this condition. Biobank

studies covering Boston (MGBB), New York (BioMe), Los Angeles (UCLA-ATLAS) and Helsinki

(HUS) had overall lower rates of type 2 diabetes individuals compared to the other studies and

lower weight at baseline, which, together with the availability of more updated electronic health

records data, suggest a larger number of individuals being prescribed GLP1-RA for weight loss.

However, care should be taken in interpreting variation in rate of type 2 diabetes across biobanks

as this might reflect different approaches used for capturing this diagnosis from the electronic

health records as well as different sampling strategies of biobanks.

The comparison between GLP1-RA and BS is informative as it highlights the younger age, higher

weight and lower percentage of men undergoing BS compared to GLP1-RA users. The weight

reduction was also approximately 5 times higher among BS patients compared to GLP1-RA

users. The weight reduction among GLP1-RA users was also more heterogeneous across

biobanks compared to BS, but in line with what reported previously in the literature using real

world data[34] and lower than results from clinical trials.[11,56] Our findings also highlight a consistent

effect of baseline body weight and sex on weight loss associated with both GLP1-RA treatment

and BS, suggesting that individuals with higher weight at baseline, as well as women benefit more

from both interventions. These findings underscore the importance of considering individual

characteristics when designing weight loss interventions.

The main aim of this study was to evaluate the impact of genetic exposures on body weight changes induced by GLP1-RA or BS. At the current sample size, a genome-wide scan would have limited power to identify potential new signals. With 6,750 individuals, we would have at least 80% statistical power to identify, at a genome-wide significant threshold, a genetic variant with an allele frequency of at least 1% and an effect larger than 0.3% weight change per change in one SD, which is unlikely given previous effects of common variants on treatment response.[51] Instead, we formulated a set of hypotheses based on the results from the largest genome-wide association studies of BMI and type 2 diabetes, and, explored biologically plausible mutations in several candidate genes. Polygenic scores represent the most powerful genetic predictors for BMI and type 2 diabetes and we confirm their significant association with the corresponding trait across all major continental ancestry groups. Thus, lack of associations with these polygenic scores would suggest that GLP1-RA and BS-related weight loss is independent of one's genetic predisposition to BMI or type 2 diabetes, as explained by common genetic variants. Moreover, because the biological underpinning of weight change is partially distinct from weight at baseline, we also consider the few replicated genetic signals for BMI variability and BMI trajectories.

For bariatric surgery we observed that genetically higher BMI was associated with weight gain, in the opposite direction for the phenotypic association with initial body weight. This might suggest that individuals with a higher genetic predisposition for elevated BMI might regain weight, after the operation, faster than those with lower genetic predisposition. To maintain weight loss, they may need additional lifestyle modifications, behavioral health therapy or adjuvant pharmacologic therapies as a GLP1-RA treatment.[57,58]

On the contrary, none of the genetic exposures were associated with weight loss among GLP1-RA users, after multiple-testing correction. These results were not attributable to high heterogeneity across studies or across major continental ancestry groups. Whether we directly tested the impact of genetic exposure on percentage body weight change or used post-treatment weight as outcome and tested the interaction between baseline weight and the genetic exposure, results did not significantly change. In light of the associations observed for BS and following the hypothesis that individuals with genetically higher BMI tend to re-gain weight faster, it would not be surprising to find, at larger sample size, a significant association between BMI polygenic score and weight loss due to GLP1-RA. However, our analyses exclude that this effect, should it exist, is unlikely to guide clinical treatment assignment. Our results showed that the considered genetic

influence on weight loss in response to GLP1-RA treatment may be too small to impact clinical

decision-making or patient outcomes. This statement should be revisited either by using more

powerful polygenic scores or by considering individuals undergoing GLP1-RA treatment only for

weight loss.

Another aim of the study was to compare the effects of GLP1-RA treatment and BS on weight

loss across major continental ancestry groups. Ancestry can provide more useful information

about population health than racial and ethnic categories because it is not directly influenced by

geographic, cultural, and sociopolitical forces.[59] Nonetheless, ancestry is also partially correlated

with self-reported race and ethnicity and socio-economic characteristics do differ across racial

and ethnic groups. Overall, we did not find large differences in GLP1-RA and BS associated

weight loss between major continental ancestry groups. We do observe that GLP1-RA users of

African and admixed American ancestry had statistically significantly lower change in body weight

compared to Europeans, after accounting for differences in sex, age, weight at baseline and

medication type. The effects were consistent in BioMe and UCLA-ATLAS biobanks for admixed

American ancestry, and additionally in AoU and MGBB for African ancestry. This might reflect

different socio-economic characteristics between individuals of admixed American and European

ancestry resulting, for example, in differences in access to treatment.

The main strength of our study is its size and diversity across countries, healthcare systems and

ancestries. Our study presents various limitations. First, we cannot be sure that all GLP1-RA

users fulfilled their prescription. This is a potential issue only in studies with prescription

information but not in studies with purchase information. Nonetheless weight loss was comparable

across both study types. Moreover, we ensured that observed changes in body weight are due to

the treatment by considering only measurements within one month from the last purchase or

prescription. Second, we could not accurately estimate treatment adherence, but overcome this

limitation by considering treatments of maximum 12 months, a period for which adherence has

been observed to be around 65%.[52] Third, we used ancestry labels based on major continental

ancestry groups, while ancestry can be better characterized as a continuous quantity.[59] This

discretization reflects pragmatic considerations to enable comparable analyses across the

different studies.

In conclusion, our study suggests that GLP1-RA treatments work equally well in individuals

carrying common non-synonymous mutations in the GLP1R gene and in individuals at high

genetic risk for BMI and type 2 diabetes. It also suggests that BS is less effective among

individuals with higher polygenic score for BMI, but the effect is not large enough to be clinically

relevant. In fact, for each additional SD in the polygenic score for BMI, individuals undergoing BS

have on average 0.7% lower weight loss from their initial body weight. Finally, these two major

weight-loss interventions achieved sustained weight reduction regardless of an individual's

genetic ancestry. However, socio-economic characteristics correlating with ancestry should be

further studied to better understand some of the remaining differences observed in this study.

Methods

Study population

In the current study, we included samples from 10 960 individuals from the following 9 biobanks: Helsinki University Hospital (HUS, Finland; N = 633), Estonian Biobank (ESTBB, Estonia; N = 464), UK Biobank (UKBB, United Kingdom; N = 810), All of Us (AoU, USA; N = 559), BioMe (BioMe, USA; N = 2,170), Mass General Brigham Biobank (MGBB, USA; N = 2,141), Atlas Biobank (UCLA-ATLAS, USA; N = 1,445), Qatar Biobank (QBB, Qatar; N = 2,383), Bialystok Bariatric Surgery Study (BBSS, Poland, N = 355) . The biobank studies include samples from (hospital) biobanks, prospective epidemiological and disease-based cohorts. Follow-up covers a total of 20 years with the earliest study starting follow-up in 2004 (AoU) and the latest study ending follow-up in 2024 (BioMe). In the **Supplementary Material** we provide a detailed description of the population selected in each study.

Inclusion and exclusion criteria for GLP1-RA analysis

For our GLP1-RA analysis we implemented the following inclusion and exclusion criteria. We required an initiation of a GLP1-RA treatment (ATC codes: A10BJ*, A10BX04, A10BX10, A10BX13, A10BX14) defined as the date of first medication purchase or medication prescription of GLP1-RA. Individuals were required to be on a GLP1-RA treatment for at least twelve months. This was defined through prescription length, regular drug prescription refills or drug purchases, depending on the dataset and healthcare setting.

Individuals aged under 18 were excluded from the analysis. Furthermore, individuals were required to have at least one body weight measurement maximum one year prior to treatment initiation or within 14 days after initiation, if no measure before initiation was available. If we observed multiple weight measurements within this time frame, only the closest one to initiation was considered. This measurement defined our baseline weight variable. At least one body weight measurement between 26 weeks and 52 weeks was necessary for each individual to be included in the analysis. Individuals were excluded if they underwent BS prior to or during the first year of GLP1-RA treatment.

Inclusion and exclusion criteria for BS analysis

For our BS analysis we implemented the following inclusion and exclusion criteria. Individuals were included if they underwent any type of BS. The respected code definitions for BS varied across countries and healthcare systems and can be found in **Supplementary Table 11**. Individuals aged under 18 at baseline were excluded from the analysis. Analogously to the GLP1-RA analysis, individuals were required to have at least one body weight measurement maximum one year prior to surgery or within 14 days after it, if no measure before baseline was available. If we observed multiple weight measurements within this time frame, only the closest one to initiation was considered. This measurement defined our baseline weight variable. Additionally, at least one body weight measurement between 26 weeks and 208 weeks was necessary for each individual to be included in the analysis.

Outcome definitions

In our primary model, for both GLP1-RA and BS we defined the outcome to be the percentage change in body weight from baseline The baseline body weight measured closest to T_0 and between -52 weeks and +2 weeks from T_0 was defined as W_0.

The second body weight (mW_1) was defined as the median of body weight measurements between 26 weeks and 52 weeks from T_0 for the GLP1-RA analysis and between 180 days and 1460 days for the BS analysis. This was to enhance robustness against outlier measurements occurring within these intervals.

Therefore, the percentage change in body weight from baseline was calculated using

$$\% \, weight \, change \; = \; \frac{mW_1 - W_0}{W_0} * 100 \; .$$

In our secondary model, we used mW_1 as an outcome.

Genetic exposures

We identified 15 genetic variants (single nucleotide polymorphism, SNP) and two PGS of interest for our analysis. Among the included SNPs 3 are functionally characterized *GLP1R* variants (*rs10305492*, *rs146868158*, *rs6923761*) associated with random glucose levels that have shown decreased/increased response to different endogenous and exogenous GLP1-RA.[49] Seven SNPs

are *GLP1R* missense variants (*rs1042044, rs10305420, rs3765467, rs10305421, rs2295006,*

rs10305510, rs201672448) with a frequency over 1% in at least one ancestry group in gnomAD

(version 4.1).[48] One SNP is the only robustly replicated lead genome-wide significant variant

(*rs429358, ApoE*) in a GWAS of BMI change over time.[46,60] Two SNPs are *PCSK1* missense

variants (*rs6232, rs6235*) that have been found associated with BMI variations.[50] And two SNPs

(*rs728996, rs17702901*) were associated with an effect on excess body mass index loss amongst

BS patients.[42] A list of all genetic exposures and their frequencies across all major continental

ancestry groups can be found in **Supplementary Table 4**. The included PGS are for body mass

index (BMI) and type 2 diabetes (T2D).[61] The PGS weights were taken from Weissbrod et al.

2022[61] and obtained from UK Biobank. The scores were computed in each study using PLINK

2.0.[62] Because the PGS scores were derived from UK Biobank, we couldn't use the same score

in UK Biobank, instead we used Thompson et al. 2022.[63]

Statistical models

We employed two linear regression models (A and B) to investigate the effect of the chosen

genetic exposures on weight change after initiation of GLP1-RA treatment and BS, respectively.

In both models we adjusted for the baseline weight, sex, age at initiation of treatment or surgery

in years, the first 20 genetic principal components and study-specific covariates, such as the

genotyping batch. In the GLP1-RA analysis we additionally adjusted for the medication type within

the drug class.

In model A $\% \, weight \, change$ was the outcome variable, while in model B the outcome was mW_1.

In model B we additionally included the interaction term between the genetic exposure variable

and the baseline weight.

Model A (primary model):

$\% \, weight \, change \; = \beta_0 + \beta_1 * genetic \, exposure \; + \beta_2 * W_0 + \beta_3 * sex + \beta_4 * age \, at \, initiation \; +$

$\beta_5 * medication \, (only \, for \, GLP1) + \sum_{k=6}^{25} \beta_k * PC1{:}20 \; + \sum_{i=26}^{n} \beta_i * study \, specific \, covariates$

Model B (secondary model):

$mW_1 \; = \beta_0 + \beta_1 * genetic \, exposure \; + \beta_2 * W_0 + \beta_3 * genetic \, exposure * W_0 \; +$

$\beta_4 * sex + \beta_5 * age \, at \, initiation \; + \beta_6 * medication \, (only \, for \, GLP1) + \sum_{k=7}^{26} \beta_k * PC1{:}20 \; +$

472
$$\sum_{i=27}^{n} \beta_i * study\ specific\ covariates$$

Meta-analyses

We conducted ancestry-specific and multi-ancestry fixed-effect meta-analyses for each of the
genetic exposures. Effect sizes were combined using fixed-effect inverse-variance-weighting
(IVW), as implemented in the R package "meta".[64] For each exposure we reported the unadjusted
P values, but considered as threshold for statistical significance *P* < 0.004 for GLP1-RA (i.e. P <
0.05/13, Bonferroni correction for 13 exposures tested) and *P* < 0.01 for BS (i.e. P < 0.05/5,
Bonferroni correction for 5 exposures tested). We tested for heterogeneity in effect sizes between
ancestries using Cochran's Q-test[65] and by inspecting the I^2 statistics[66], which is defined as:

483
$$I^2 = \left(\frac{Q - df}{Q}\right) * 100\%$$

484
485 where df is equal to $number\ of\ studies - 1$.

Power calculation

To calculate the statistical power for the PGS effect sizes, we used a t-test for linear regression
coefficient as implemented in the R package pwrs[67] assuming a standard deviation for the
outcome variable SDy = 7.8 (weighted combine SD across studies), a sample size of n = 6 750,
a total number of predictors k = 25 and an adjusted r2 = 0.05, to calculate power at P < 0.05
significance level.

For the coefficients of association with single variants, we estimated statistical power via the non-
centrality parameter (NCP) of the chi-square distribution. We defined NCP = 2 f (1 − f) n β^2
where f is MAF, n is the effective sample size, and β is the expected effect size.[68]
We set a beta of 0.3 (% weight change from baseline) as the expected genetic effect on GLP1
response, based on findings from a recent large-scale study on genetic influences on drug
response.[51]

Code Availability

501 The study utilized previously published analysis tools as described in the **Methods** section.
502 Additional code used for these analyses is available at https://github.com/dsgelab/glp1-bs-
503 genetics/
504

Data Availability

The Helsinki biobank can provide access for research projects within the scope regulated by the
Finnish Biobank Act, which is research utilizing the biobank samples or data for the purposes of
promoting health, understanding the mechanisms of disease or developing products and
treatment practices used in health and medical care.
De-identified data of the MGBB that supports this study is available from the MGB Biobankportal
at https://www.massgeneralbrigham.org/en/research-and-innovation/participate-in-
research/biobank/for-researchers. Restrictions apply to the availability of these data, which are
available to MGB-affiliated researchers via a formal application.
UK Biobank data are available through a procedure described at
http://www.ukbiobank.ac.uk/using-the-resource/.
Registered researchers whose institutions have Data Use and Registration Agreements in place
with All of Us that include the Controlled Tier can access genomic data.
Clinical and genotype data from UCLA ATLAS Community Health Initiative patients, de-identified
for research purposes, are accessible to UCLA-approved researchers through the Discovery Data
Repository (DDR).
BioMe data is available through a process described at
https://icahn.mssm.edu/research/ipm/programs/biome-biobank/researcher-faqs.
Pseudonymised data and/or biological samples can be accessed for research and development
purposes in accordance with the Estonian Human Genome Research Act. To access data, the
research proposal must be approved by the Scientific Advisory Committee of the Estonian
Biobank as well as by the Estonian Committee on Bioethics and Human Research. For more
details on data access and relevant documents, please see
https://genomics.ut.ee/en/content/estonian-biobank#dataaccess.
For access to the Bialystok Bariatric Surgery Study please refer to the Medical University of
Bialystok.

The data and biosamples collected or generated by QBB will be made available to researchers employed within or otherwise contractually bound to public and private institutions that conduct scientific research and that meet the requirements detailed in the Qatar Biobank Research Access policy.

The PGS weights used in this study are available in Weissbrod et al. 2022 and Thompson et al. 2022.[61]

Acknowledgements

We gratefully acknowledge the resources provided by the Institute for Precision Health and participating patients from the UCLA ATLAS Community Health Initiative (https://www.uclahealth.org/international-services/consulting-services/clinical-research-consultation/ucla-atlas-community-health-initiative). The UCLA ATLAS Community Health Initiative in collaboration with UCLA ATLAS Precision Health Biobank is a program of the Institute for Precision Health, which directs and supports the biobanking and genotyping of biospecimen samples from participating patients from UCLA in collaboration with the David Geffen School of Medicine, UCLA Clinical and Translational Science Institute and UCLA Health. The ATLAS Community Health Initiative is supported by UCLA Health, the David Geffen School of Medicine and a grant from the UCLA Clinical and Translational Science Institute (UL1TR001881). This work was financially supported in part by National Institutes of Health awards U01HG011715, R01HG009120 and R01MH115676. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

We would furthermore like to thank all participants of HUS (https://www.helsinginbiopankki.fi/en/Helsinki-Biobank; https://www.hus.fi/en), ESTBB (https://genomics.ut.ee/en/content/estonian-biobank), BBSS (https://classic.clinicaltrials.gov/ct2/show/NCT04634591), QBB (https://www.qphi.org.qa/), BioMe (https://icahn.mssm.edu/research/ipm/programs/biome-biobank), UKBB (https://www.ukbiobank.ac.uk/), MGBB (https://www.massgeneralbrigham.org/en/research-and-innovation/participate-in-research/biobank) and AOU (https://allofus.nih.gov/) for their generous contribution. This work was further supported in part by funding from the Eric and Wendy Schmidt Center at the Broad Institute of MIT and Harvard.

U.V. and K.A. have been funded by Estonian Research Council's personal research funding start-up grant PSG759 and European Research Council' starting grant under the grant agreement no 101117251 (OBECAUSE). J.H.L. is supported by NIDDK K23 DK131345.

563 The activities of the EstBB are regulated by the Human Genes Research Act, which was adopted
564 in 2000 specifically for the operations of the EstBB. Individual level data analysis in the EstBB
565 was carried out under ethical approvals of 1.1-12/1409 and 1.1-12/2161 from the Estonian
566 Committee on Bioethics and Human Research (Estonian Ministry of Social Affairs), using data
567 according to release application 6-7/GI/18857 from the Estonian Biobank.
568

References

1. Hruby, A. & Hu, F. B. The Epidemiology of Obesity: A Big Picture. *Pharmacoeconomics* **33**, 673–689 (2015).

2. Boutari, C. & Mantzoros, C. S. A 2022 update on the epidemiology of obesity and a call to action: as its twin COVID-19 pandemic appears to be receding, the obesity and dysmetabolism pandemic continues to rage on. *Metabolism* **133**, 155217 (2022).

3. EASO Secretariat. European survey finds over three-quarters of adults with obesity have attempted to lose weight in the past year, but most have been unsuccessful. *EASO* https://easo.org/european-survey-finds-over-three-quarters-of-adults-with-obesity-have-attempted-to-lose-weight-in-the-past-year-but-most-have-been-unsuccessful/ (2022).

4. Creutzfeldt, W. The incretin concept today. *Diabetologia* **16**, 75–85 (1979).

5. Kreymann, B., Williams, G., Ghatei, M. A. & Bloom, S. R. Glucagon-like peptide-1 7-36: a physiological incretin in man. *Lancet* **2**, 1300–1304 (1987).

6. Nauck, M. A. *et al.* Normalization of fasting glycaemia by intravenous GLP-1 ([7-36 amide] or [7-37]) in type 2 diabetic patients. *Diabet. Med.* **15**, 937–945 (1998).

7. Nauck, M. A. *et al.* Effects of glucagon-like peptide 1 on counterregulatory hormone responses, cognitive functions, and insulin secretion during hyperinsulinemic, stepped hypoglycemic clamp experiments in healthy volunteers. *J. Clin. Endocrinol. Metab.* **87**, 1239–1246 (2002).

8. Amylin and Lilly Announce FDA Approval of BYETTA(TM) (Exenatide) Injection. *Eli Lilly and Company* https://investor.lilly.com/news-releases/news-release-details/amylin-and-lilly-announce-fda-approval-byettatm-exenatide.

9. Secher, A. *et al.* The arcuate nucleus mediates GLP-1 receptor agonist liraglutide-dependent weight loss. *J. Clin. Invest.* **124**, 4473–4488 (2014).

10. Ryan, D. H. *et al.* Long-term weight loss effects of semaglutide in obesity without diabetes

in the SELECT trial. *Nat. Med.* (2024) doi:10.1038/s41591-024-02996-7.

11. Wilding, J. P. H. *et al.* Once-Weekly Semaglutide in Adults with Overweight or Obesity. *N.*

Engl. J. Med. **384**, 989–1002 (2021).

12. Rubino, D. M. *et al.* Effect of Weekly Subcutaneous Semaglutide vs Daily Liraglutide on

Body Weight in Adults With Overweight or Obesity Without Diabetes: The STEP 8

Randomized Clinical Trial. *JAMA* **327**, 138–150 (2022).

13. Marso, S. P. *et al.* Liraglutide and Cardiovascular Outcomes in Type 2 Diabetes. *N. Engl. J.*

Med. **375**, 311–322 (2016).

14. Marso, S. P. *et al.* Semaglutide and Cardiovascular Outcomes in Patients with Type 2

Diabetes. *N. Engl. J. Med.* **375**, 1834–1844 (2016).

15. Perkovic, V. *et al.* Effects of Semaglutide on Chronic Kidney Disease in Patients with Type

2 Diabetes. *N. Engl. J. Med.* **391**, 109–121 (2024).

16. Sattar, N. *et al.* Cardiovascular, mortality, and kidney outcomes with GLP-1 receptor

agonists in patients with type 2 diabetes: a systematic review and meta-analysis of

randomised trials. *Lancet Diabetes Endocrinol* **9**, 653–662 (2021).

17. Watanabe, J. H., Kwon, J., Nan, B. & Reikes, A. Trends in glucagon-like peptide 1 receptor

agonist use, 2014 to 2022. *J. Am. Pharm. Assoc.* **64**, 133–138 (2024).

18. Melson, E., Ashraf, U., Papamargaritis, D. & Davies, M. J. What is the pipeline for future

medications for obesity? *Int. J. Obes.* (2024) doi:10.1038/s41366-024-01473-y.

19. Morgan, J. P. The increase in appetite for obesity drugs.

https://www.jpmorgan.com/insights/global-research/current-events/obesity-drugs (2023).

20. Clapp, B. *et al.* American Society for Metabolic and Bariatric Surgery 2020 estimate of

metabolic and bariatric procedures performed in the United States. *Surg. Obes. Relat. Dis.*

18, 1134–1140 (2022).

21. After 30 Years — New Guidelines For Weight-Loss Surgery. *American Society for*

Metabolic and Bariatric Surgery https://asmbs.org/news_releases/after-30-years-new-

guidelines-for-weight-loss-surgery/ (2022).

22. Gloy, V. L. *et al.* Bariatric surgery versus non-surgical treatment for obesity: a systematic

review and meta-analysis of randomised controlled trials. *BMJ* **347**, f5934 (2013).

23. Salminen, P. *et al.* Effect of Laparoscopic Sleeve Gastrectomy vs Roux-en-Y Gastric

Bypass on Weight Loss, Comorbidities, and Reflux at 10 Years in Adult Patients With

Obesity: The SLEEVEPASS Randomized Clinical Trial. *JAMA Surg.* **157**, 656–666 (2022).

24. Lim, R., Beekley, A., Johnson, D. C. & Davis, K. A. Early and late complications of bariatric

operation. *Trauma Surg Acute Care Open* **3**, e000219 (2018).

25. Ma, I. T. & Madura, J. A., 2nd. Gastrointestinal Complications After Bariatric Surgery.

Gastroenterol. Hepatol. **11**, 526–535 (2015).

26. Azagury, D. *et al.* Heterogeneity of weight loss after gastric bypass, sleeve gastrectomy,

and adjustable gastric banding. *Surgery* **165**, 565–570 (2019).

27. Cadena-Obando, D. *et al.* Are there really any predictive factors for a successful weight

loss after bariatric surgery? *BMC Endocr. Disord.* **20**, 20 (2020).

28. Çalık Başaran, N., Dotan, I. & Dicker, D. Post metabolic bariatric surgery weight regain: the

importance of GLP-1 levels. *Int. J. Obes.* (2024) doi:10.1038/s41366-024-01461-2.

29. Lauti, M., Kularatna, M., Hill, A. G. & MacCormick, A. D. Weight Regain Following Sleeve

Gastrectomy-a Systematic Review. *Obes. Surg.* **26**, 1326–1334 (2016).

30. Young, K. G. *et al.* Treatment effect heterogeneity following type 2 diabetes treatment with

GLP1-receptor agonists and SGLT2-inhibitors: a systematic review. *Commun. Med.* **3**, 131

(2023).

31. Jones, A. G. *et al.* Markers of β-Cell Failure Predict Poor Glycemic Response to GLP-1

Receptor Agonist Therapy in Type 2 Diabetes. *Diabetes Care* **39**, 250–257 (2016).

32. Dawed, A. Y. *et al.* Pharmacogenomics of GLP-1 receptor agonists: a genome-wide

analysis of observational data and large randomised controlled trials. *Lancet Diabetes*

Endocrinol **11**, 33–41 (2023).

33. Cardoso, P. *et al.* Phenotype-based targeted treatment of SGLT2 inhibitors and GLP-1

receptor agonists in type 2 diabetes. *Diabetologia* **67**, 822–836 (2024).

34. Brown, E., Wilding, J. P. H., Barber, T. M., Alam, U. & Cuthbertson, D. J. Weight loss

variability with SGLT2 inhibitors and GLP-1 receptor agonists in type 2 diabetes mellitus

and obesity: Mechanistic possibilities. *Obes. Rev.* **20**, 816–828 (2019).

35. Umapathysivam, M. M. *et al.* Type 2 Diabetes risk alleles in Peptidyl-glycine Alpha-

amidating Monooxygenase influence GLP-1 levels and response to GLP-1 Receptor

Agonists. *medRxiv* (2023) doi:10.1101/2023.04.07.23288197.

36. Jensterle, M., Rizzo, M., Haluzík, M. & Janež, A. Efficacy of GLP-1 RA Approved for Weight

Management in Patients With or Without Diabetes: A Narrative Review. *Adv. Ther.* **39**,

2452–2467 (2022).

37. Nathan, B. M. *et al.* Predictors of weight-loss response with glucagon-like peptide-1

receptor agonist treatment among adolescents with severe obesity. *Clin. Obes.* **6**, 73–78

(2016).

38. Vosoughi, K. *et al.* Association of Glucagon-like Peptide 1 Analogs and Agonists

Administered for Obesity with Weight Loss and Adverse Events: A Systematic Review and

Network Meta-analysis. *EClinicalMedicine* **42**, 101213 (2021).

39. Segal, J. B. *et al.* Assessing Heterogeneity of Treatment Effect in Real-World Data. *Ann.*

Intern. Med. **176**, 536–544 (2023).

40. Jensterle, M., Pirš, B., Goričar, K., Dolžan, V. & Janež, A. Genetic variability in GLP-1

receptor is associated with inter-individual differences in weight lowering potential of

liraglutide in obese women with PCOS: a pilot study. *Eur. J. Clin. Pharmacol.* **71**, 817–824

(2015).

41. van der Meer, R. *et al.* Genetic variants associated with weight loss and metabolic

outcomes after bariatric surgery: A systematic review. *Obes. Rev.* **24**, e13626 (2023).

42. Aasbrenn, M. *et al.* Genetic Determinants of Weight Loss After Bariatric Surgery. *Obes.*

Surg. **29**, 2554–2561 (2019).

43. NCD Risk Factor Collaboration (NCD-RisC). Worldwide trends in underweight and obesity

from 1990 to 2022: a pooled analysis of 3663 population-representative studies with 222

million children, adolescents, and adults. *Lancet* **403**, 1027–1050 (2024).

44. White, G. E. *et al.* Real-world weight-loss effectiveness of glucagon-like peptide-1 agonists

among patients with type 2 diabetes: A retrospective cohort study. *Obesity* **31**, 537–544

(2023).

45. Inoue, K. *et al.* Long-term impact of liraglutide, a glucagon-like peptide-1 (GLP-1) analogue,

on body weight and glycemic control in Japanese type 2 diabetes: an observational study.

Diabetol. Metab. Syndr. **6**, 95 (2014).

46. Venkatesh, S. S. *et al.* The genetic architecture of changes in adiposity during adulthood.

medRxiv (2023) doi:10.1101/2023.01.09.23284364.

47. Xu, J. *et al.* Exploring the clinical and genetic associations of adult weight trajectories using

electronic health records in a racially diverse biobank: a phenome-wide and polygenic risk

study. *Lancet Digit Health* **4**, e604–e614 (2022).

48. Karczewski, K. J. *et al.* The mutational constraint spectrum quantified from variation in

141,456 humans. *Nature* **581**, 434–443 (2020).

49. Lagou, V. *et al.* GWAS of random glucose in 476,326 individuals provide insights into

diabetes pathophysiology, complications and treatment stratification. *Nat. Genet.* **55**, 1448–

1461 (2023).

50. Nead, K. T. *et al.* Contribution of common non-synonymous variants in PCSK1 to body

mass index variation and risk of obesity: a systematic review and meta-analysis with

evidence from up to 331 175 individuals. *Hum. Mol. Genet.* **24**, 3582–3594 (2015).

51. Sadler, M. C. *et al.* Leveraging large-scale biobank EHRs to enhance pharmacogenetics of

cardiometabolic disease medications. *medRxiv* (2024) doi:10.1101/2024.04.06.24305415.

52. Weiss, T. *et al.* Real-world weight change, adherence, and discontinuation among patients

with type 2 diabetes initiating glucagon-like peptide-1 receptor agonists in the UK. *BMJ*

Open Diabetes Res Care **10**, (2022).

53. Ghusn, W. *et al.* Weight Loss Outcomes Associated With Semaglutide Treatment for

Patients With Overweight or Obesity. *JAMA Netw Open* **5**, e2231982 (2022).

54. Xu, T. *et al.* Timing of Maximal Weight Reduction Following Bariatric Surgery: A Study in

Chinese Patients. *Front. Endocrinol.* **11**, 615 (2020).

55. Monpellier, V. M., Janssen, I. M. C., Antoniou, E. E. & Jansen, A. T. M. Weight Change

After Roux-en Y Gastric Bypass, Physical Activity and Eating Style: Is There a

Relationship? *Obes. Surg.* **29**, 526–533 (2019).

56. Knop, F. K. *et al.* Oral semaglutide 50 mg taken once per day in adults with overweight or

obesity (OASIS 1): a randomised, double-blind, placebo-controlled, phase 3 trial. *Lancet*

402, 705–719 (2023).

57. Hall, K. D. & Kahan, S. Maintenance of Lost Weight and Long-Term Management of

Obesity. *Med. Clin. North Am.* **102**, 183–197 (2018).

58. Mok, J. *et al.* Safety and Efficacy of Liraglutide, 3.0 mg, Once Daily vs Placebo in Patients

With Poor Weight Loss Following Metabolic Surgery: The BARI-OPTIMISE Randomized

Clinical Trial. *JAMA Surg.* **158**, 1003–1011 (2023).

59. Borrell, L. N. *et al.* Race and Genetic Ancestry in Medicine - A Time for Reckoning with

Racism. *N. Engl. J. Med.* **384**, 474–480 (2021).

60. Kemper, K. E. *et al.* Genetic influence on within-person longitudinal change in

anthropometric traits in the UK Biobank. *Nat. Commun.* **15**, 3776 (2024).

61. Weissbrod, O. *et al.* Leveraging fine-mapping and multipopulation training data to improve

cross-population polygenic risk scores. *Nat. Genet.* **54**, 450–458 (2022).

62. Chang, C. C. *et al.* Second-generation PLINK: rising to the challenge of larger and richer

datasets. *Gigascience* **4**, 7 (2015).

63. Thompson, D. J. *et al.* UK Biobank release and systematic evaluation of optimised

polygenic risk scores for 53 diseases and quantitative traits. *bioRxiv* (2022)

doi:10.1101/2022.06.16.22276246.

64. Schwarzer, G., Carpenter, J. R. & Rücker, G. *Meta-Analysis with R*. (Springer International

Publishing, 2015).

65. Cochran, W. G. The Combination of Estimates from Different Experiments. *Biometrics* **10**,

101–129 (1954).

66. Higgins, J. P. T. & Thompson, S. G. Quantifying heterogeneity in a meta-analysis. *Stat.*

Med. **21**, 1539–1558 (2002).

67. Bulus, M. A practical guide to statistical power and sample size calculations in R.

https://cran.r-project.org/web/packages/pwrss/vignettes/examples.html (2023).

68. Kanai, M. *et al.* Insights from complex trait fine-mapping across diverse populations.

bioRxiv (2021) doi:10.1101/2021.09.03.21262975.

736