Machine learning for the prediction of spontaneous preterm birth using early

second and third trimester maternal blood gene expression: A Cautionary Tale

Kylie K Hornaday*[1][¶], Ty Werbicki[1][¶], Suzanne C Tough[2,3], Stephen L Wood[4], David W

Anderson[5], Constance H Li[6][^], Donna M Slater[1,4][^]

[1]Department of Physiology and Pharmacology, Cumming School of Medicine, University of
Calgary, Calgary, Alberta, Canada
11
12   [2]Department of Community of Health Sciences, Cumming School of Medicine, University of
13   Calgary, Calgary, Alberta, Canada
14
15   [3]Department of Pediatrics, Cumming School of Medicine, University of Calgary, Calgary,
16   Alberta, Canada
17
18   [4]Department of Obstetrics and Gynaecology, Cumming School of Medicine, University of
19   Calgary, Calgary, Alberta, Canada
20
21   [5]Department of Science, Langara College, Vancouver, British Columbia, Canada
22
23   [6]Department of Biochemistry and Molecular Biology, Cumming School of Medicine, University
24   of Calgary, Alberta, Canada
25
26
27
28   *Corresponding author

29   E-mail: kylie.hornaday@ucalgary.ca (KKH)

30   [¶]These authors contributed equally to this work

31   [^]These authors contributed equally to this work

Abstract

Spontaneous preterm birth (sPTB) remains a significant global health challenge and a leading cause of neonatal mortality and morbidity. Despite advancements in neonatal care, the prediction of sPTB remains elusive, in part due to complex etiologies and heterogeneous patient populations. This study aimed to validate and extend information on gene expression biomarkers previously described for predicting sPTB using maternal whole blood from the All Our Families pregnancy cohort study based in Calgary, Canada. The results of this study are two-fold: first, using additional replicates of maternal blood samples from the All Our Families cohort, we were unable to repeat the findings of a 2016 study which identified top maternal gene expression predictors for sPTB. Second, we conducted a secondary analysis of the original gene expression dataset from the 2016 study using five modelling approaches (random forest, elastic net regression, unregularized logistic regression, L2-regularized logistic regression, and multilayer perceptron neural network) followed by external validation using a pregnancy cohort based in Detroit, USA. The top performing model (random forest classification) suggested promising performance (area under the receiver operating curve, AUROC 0.99 in the training set), but performance was significantly degraded on the test set (AUROC 0.54) and further degraded in external validation (AUROC 0.50), suggesting poor generalizability, likely due to overfitting exacerbated by a low feature-to-noise ratio. Similar performance was observed in the other four learning models. Prediction was not improved when using higher complexity machine learning (e.g. neural network) approaches over traditional statistical learning (e.g. logistic regression). These findings underscore the challenges in translating biomarker discovery into clinically useful predictive models for sPTB. This study highlights the critical need for rigorous methodological safeguards and external validation in biomarker research. It also emphasizes the impact of data

55 noise and overfitting on model performance, particularly in high-dimensional omics datasets.

56 Future research should prioritize robust validation strategies and explore mechanistic insights to

57 improve our understanding and prediction of sPTB.

Introduction

Preterm birth, defined as delivery of a live infant prior to 37 weeks of gestation, occurred

in approximately 13.4 million births worldwide in 2020, and is a significant contributor to

mortality and morbidity in neonates and children under five [1, 2]. Child mortality related to

PTB complications has declined since 2000, in part due to advancements in treatments for

neonatal complications of prematurity such as respiratory distress syndrome. However, an

estimated 900,000 PTB-associated deaths of children under five still occurred in 2019 worldwide

[3].

Approximately one third of preterm births occur because of known maternal or fetal

indications, while the remaining two thirds occur following spontaneous onset of labour and/or

premature rupture of the fetal membranes (collectively termed spontaneous or sPTB). Most

sPTB are without known indication, making prediction and subsequent clinical management of

risk, challenging [2]. As one of the great obstetrical syndromes, considerable efforts have aimed

to identify predictive biomarkers of sPTB, however none so far have emerged to have clinical

utility, possibly due to heterogeneity within both patient populations and preterm birth

phenotypes, as well as risk of bias within study design [4-7]. Methodological safeguarding and

appropriate validation of models is important to determine feasibility, repeatability, robustness,

and generalizability of prediction [8-11]. Best practices for prediction modelling are well defined

in the literature [12, 13], and primary research articles reporting external validation of prediction

models have been increasingly published over the last five years [14-17]. However, in the

reproductive field, studies externally validating prediction models are limited [10, 18, 19]. Thus,

we sought herein to repeat and validate previous findings on prediction of sPTB, which

identified a predictive relationship between gene expression biomarkers and sPTB.

Gene expression biomarkers have been identified in maternal whole blood for the

prediction of sPTB, which presents a promising avenue for minimally invasive prediction as

peripheral blood can reflect global and uterine physiological and immunological changes during

pregnancy [20]. One example includes eight genes, *LOC100128908, MIR3691, LOC101927441,*

CST13P, ACAP2, ZNF324, SH3PXD2B, TBX21 that were identified as significantly predictive of

sPTB (65% sensitivity and 88% specificity after adjusting for history of abortion and anaemia) in

a stepwise logistic regression model published in 2016 by Heng *et al.* [21]. These gene

expression biomarkers were originally identified using an Affymetrix chip microarray analysis of

maternal whole blood from the All Our Families pregnancy cohort based in Calgary, Canada

[21]. The All Our Families pregnancy cohort presents a rare opportunity for testing experimental

repeatability, as maternal blood samples were collected and stored in four separate PAXgene

RNA tubes, two were used for the original study (22), and one for validating RNA quality and

integrity [22], leaving a remaining fourth sample for experimental validation. The study herein

sought to use the additional PAXgene tube to repeat and validate this predictive model to test

feasibility for clinical use.

It is also important to note that prediction algorithms that match too closely to the

training data, in other words, suffer from overfitting, are not generalizable to other populations,

which is one of the major limitations of prediction modelling. This problem is exacerbated by

small or non-representative training sets, where patterns identified may not be meaningfully

associated with the outcome, or constitute "noise", and thus the prediction does not translate

effectively beyond the original training observations. This stresses the importance of external

validation in order to identify robust, generalizable models to meaningfully push forward the

prediction of preterm birth. Therefore, an external pregnancy cohort based in Detroit, USA,

which collected maternal blood gene expression data from comparable timepoints in pregnancy,

was identified, presenting a unique opportunity for external validation to determine the

generalizability of prediction [23].

The original 2016 study used a logistic regression-based model; however, we

hypothesized that machine learning approaches could improve predictive performance. Machine

learning and other complex data analysis methods are particularly well suited for mining high

dimensional datasets, such as transcriptomic datasets, as they do not require the data to adhere to

any *a priori* assumptions [24]. Machine learning allows for the identification of non-obvious,

interactive, complex, and/or non-linear patterns which can go undetected when using traditional

statistical linear models [24, 25]. These patterns can be leveraged, both toward outcome

prediction (something highly valuable for complex medical conditions such as preterm birth),

and characterizing underlying disease mechanisms [26, 27]. This is particularly enticing, as the

underlying causes of sPTB remain poorly understood.

The overarching aim is to explore the repeatability, generalization, and robustness of a

prediction model for sPTB using maternal blood gene expression biomarkers, with emphasis on

external validation of prediction models for sPTB. Additionally, given the high complexity of

preterm birth, we wanted to explore whether additional learning algorithms could improve

prediction over traditional statistical learning. Specific aims are as follows:

1. To test the predictive utility of the gene expression biomarkers, *LOC100128908,*

MIR3691, LOC101927441, CST13P, ACAP2, ZNF324, SH3PXD2B, TBX21 previously

identified in maternal blood as biomarkers of spontaneous preterm birth [21].

2. To identify a predictive biomarker(s) for spontaneous preterm birth using maternal blood

gene expression data using machine learning best practices and multiple learning

approaches.

Methods

Biological samples and validation of top biomarkers

To test the reproducibility of top biomarkers identified in the literature, historical

biological samples were collected from the All Our Families cohort [28-30]. In brief, participants

were recruited between May 1[st], 2008, and December 31[st], 2011, at <25 weeks gestation and

provided consent for blood sample collection, and completed questionnaires including

information related to demographics, emotional and physical health. Participants provided

informed written consent at the time of recruitment from healthcare offices, community and

through Calgary Laboratory services, and were provided copies of their consent forms for their

records. This study was approved by the Conjoint Health Research Ethics Board at the

University of Calgary #REB15-0248 Predicting Preterm Birth Study. Biological samples were

collected at two points in pregnancy, timepoint 1 (T1) at 17-23 weeks gestation and timepoint 2

(T2) 28-32 weeks gestation. Maternal whole blood was collected directly into four separate

PAXgene blood RNA tubes which were then stored at -80°C prior to RNA isolation (PAXgene

Blood RNA Kit, Qiagen). Samples were de-identified prior to data collection. Two tubes were

previously used for the original prediction modelling and biomarker identification by Heng et al.,

[21], a third tube was used to assess RNA integrity in long term storage [22], and a fourth was

collected from storage from August 16[th] to November 3[rd], 2018 for use in the current study. For

the current study, n=47 participants who subsequently had a sPTB (<37 weeks) were included

(n=44 T1, n=42 T2 samples), in addition to n=45 participants who had a healthy term (38-42

weeks) delivery (n=40 T1, n=44 T2). A total of n=13 samples were missing from storage or

insufficient sample remaining (n=3 T1 sPTB, n=3 T2 sPTB, n=5 T1 term, n=2 T2 term), and n=2

T2 samples in the sPTB group were not included as delivery occurred prior to the second sample

collection. Maternal blood samples were collected, and RNA was isolated according to

manufacturer's instructions (RNAeasy minikit, Qiagen). The following genes were measured

using a probe-based assay (Quantigene, Invitrogen, ThermoFisher Scientific), which uses

identical probes to an Affymetrix microarray chip: *LOC100128908 (LMLN2), LOC101927441,*

CST13P, ACAP2, ZNF324, SH3PXD2B, TBX21. Due to limitations with measuring microRNA

(miRNA) using the assay, *MIR3691* was not measured. The same genes were also validated

using bulk RNA sequencing (Illumina mRNAseq following poly-A capture, NovaSeq S4 200

cycle). Sequences were aligned to the human reference genome GrCH38 using STAR to produce

reads per gene tables [31].

Data availability

166   The accession number for the gene expression data from the Calgary cohort and Detroit

167   cohort described herein are Gene Expression Omnibus GSE59491 and GSE 1494490,

168   respectively. Scripts for data preprocessing, differential expression analysis, feature selection,

169   and model training are available at https://github.com/tywerbicki/Slater_Lab_SPTB.

Population and expression dataset

To test whether additional learning models could improve predictive performance, a secondary analysis of the previously published maternal blood microarray data [21], was conducted. Gene expression data was downloaded as log2 robust multi-array average (RMA) values from the National Center for Biotechnology Information Gene Expression Omnibus (accession number: GSE59491) (All Our Families, AOF- Calgary cohort) [21]. The RMA preprocessing, including background correction and normalization was previously conducted [21, 23]. The dataset used herein contains high-throughput expression data from n=165 subjects (n=51 sPTB, n=114 matched term delivery controls) nested within the Calgary AOF cohort. Matched gene expression data from two timepoints, 17-23 weeks (T1) and 28-33 weeks (T2) was available for each participant. Two observations from the sPTB group were removed from the dataset as these deliveries occurred prior to the T2 collection and therefore have only one expression dataset.

An external dataset was additionally identified from a pregnancy cohort based in Detroit, USA [23], which collected maternal blood samples at comparable timepoints for gene expression analysis, (accession number: GSE149440), and was used for external validation of the model. Participants that had at least two matched blood samples collected within the same two timeframes (T1 and T2) were selected from within the Detroit cohort dataset for analysis, for a total of n=98 subjects (n=34 sPTB and n=64 matched term delivery controls) included for external validation.

The Calgary dataset was randomly split into ten cross validation folds: each tenth acted as a as a test set for the preprocessing, differential expression analysis, feature selection, and model

193   training done on the remaining 90% of the data. While there is no true hold out set for the

194   Calgary cohort, models are externally validated on the Detroit dataset. A schematic outlining the

195   analytical pipeline is provided in Figure 1.

197   **Figure 1. Schematic of analytical pipeline.**

Differential expression analysis

Differential expression was performed on the Calgary dataset with 10-fold cross validation.

Each training fold contains 85 term and 35 sPTB labeled patients, and each testing fold contains

29 term and 12 sPTB labeled patients. For each fold, features with low and excessively high

variance were filtered from analysis (feature kept if standard deviation between [0.001, 3]).

Features were also removed if microarray expression was lower than a value of 5 in more than

35 samples. Because T1 and T2 are modelled together, only genes with measurements at both

time points were kept after filtering.

For each remaining gene, microarray expression was modelled against a variable combining

timepoint with birth outcome using a linear model (lmFit). The correlations of repeated

measures (duplicateCorrelation) were estimated and specified in the linear model. Further, an

empirical Bayes model was used to adjust the linear model estimates for more robust estimates.

Differential expression results were explored using the following comparisons:

• sPTB group compared to term group at T1☐

• sPTB group compared to term group at T2☐

• T1 compared to T2 in sPTB group☐

• T1 compared to T2 in term group▯

• dT (T2-T1) in sPTB compared to term▯

The Benjamini-Hochberg procedure was used to perform FDR correction. Genes with a

family-wise error rate less than 0.05 for any of the above comparisons were considered

differentially expressed and kept for downstream modelling.▯▯

Feature selection analysis

Feature selection was performed using a stepwise additive selective (SAS) approach with

a logistic regression model. A cross-validation area under the receiver operator curve (AUROC)

was used as the metric for evaluating model improvement. The full set of input features includes

three features describing each differentially expressed gene: the log2 intensity at T1 (T1), the

log2 intensity at T2 (T2), and the difference between these measurements (T2-T1, or dT). The

possibility of interactions between each of these features was included by using a polynomial

model to generate pairwise feature combinations. The number of interaction terms was restricted

to a maximum of five.

The algorithm used here starts feature selection with an empty base logistic regression

model. At each iteration, the addition of each available feature to the current model was assessed

using a nested five-fold cross-validation scheme. The validation AUROC was calculated for

each cross-validation fold and averaged across the folds to evaluate the performance for adding

that feature to the current model. The feature corresponding with the highest average AUROC

was added if the marginal improvement to the current model met a predetermined threshold (ε)

of 0.015. If this minimum improvement in AUROC was not met, the next iteration re-run with

the unchanged current model. If there was no change in the current model for three iterations,

feature selection terminated and returned the features in the base model. Otherwise, the current

model was updated with the selected feature for the next iteration, and feature selection

continued until a maximum number of 30 predictive features were selected.

Model training and testing

The identified predictive features in each fold were used to train three models: an

unregularized logistic regression (LR), an L2-regularized logistic regression (ridge regression,

LR-reg), and a multilayer perceptron (MLP) neural network. The resultant three models were

assessed for predictive performance by fitting them on the cross-validation test set and the

external (Detroit) dataset. The Python Scikit-Learn implementation of logistic regression was

used with a Newton-Conjugate gradient and a maximum of 10,000 iterations. For the L2-

regularized logistic regression, a nested 5-fold cross-validation was used for hyperparameter

tuning on the penalty value (regularization strength).

The PyTorch, Scikit-Learn and Ray Tune Python libraries were used for MLP

implementation. The MLP has three layers: two hidden layers and one output layer. We used a

Leaky Rectified Linear Unit (Leaky ReLU) activation function, dropout regularization, and

performed batch normalization for each hidden layer. Nested five-fold cross-validation and Ray

Tune was used to optimize the following hyperparameters: the number of units in the first and

second hidden layers (l1, l2), dropout probabilities for the first and second hidden layers (p1, p2),

the learning rate (lr), weight decay (L2 regularization), and the training batch size. We set Ray

Tune to explore 500 hyperparameter configurations (num_samples=500) and used a reduction

factor of four ($\eta =4$) for more aggressive pruning. To select the best trained model, we compared

validation loss and AUROC. Training for the optimized MLP model was limited to 250 epochs.

Additionally, we fit two machine learning models after differential expression analysis

but prior to formal feature selection: an elastic net logistic regression (LR-ELN) and a random

forest classifier (RF). These models have internal feature selection and provide comparison

feature sets for our stepwise additive feature selection approach. The elastic net model was fit

using the Scikit-Learn implementation with a saga solver and a maximum of 10,000 iterations.

The random forest model was fit using the Scikit-Learn implementation

(RandomForestClassifier). Hyperparameters for both models (elastic net: mixing parameter,

regularization strength; random forest: n trees, max depth, min samples to split, min samples per

leaf, feature selection at split) were optimized using nested 5-fold cross-validation.

Results

Population demographics

275   Demographic characteristics of the population used for biomarker validation are

276   described in Table 1. Participants with sPTB did not significantly differ from the term group in

277   age, smoking status, alcohol use during pregnancy, history of abortion, history of PTB, gravidity

278   or parity.

Table 1. Patient demographic and clinical characteristics

	sPTB (n=47)	Term (n=45)
Maternal age (years)	31.5[30.2-32.8]	31.8[30.6-33.0]
Maternal ethnicity		
White	37, 79[64-89]%	35, 78[63-89]%
Non-white	10, 21[11-36]%	10, 22[11-37]%
Smoking during pregnancy		
Yes	5, 11[4-23]%	10, 22[11-37]%

No	36, 77[62-88]%	34, 76[60-87]%
Unknown	6, 13[5-26]%	1, 2[0.06-12]%
Alcohol during pregnancy		
Yes	17, 36[23-52]%	24, 53[38-68]%
No	22, 47[32-62]%	18, 40[26-56]%
Unknown	8, 17[8-31]%	3, 7[1-18]%
History of abortion		
Yes	4, 9[2-20]%	8, 18[8-32]%
No	43, 91[80-98]%	37, 82[68-92]%
History of PTB		
Yes	8, 17[8-31]%	2, 4[0.5-15]%
No	39, 83[69-92]%	43, 96[85-99]%
Gravidity	1.9[1.6-2.2]	1.9[1.7-2.2]
Parity	0.6[0.3-0.8]	0.5[0.4-0.7]
Gestational age at delivery (weeks)	33.9[33.2-34.6]	39.2[38.9-39.4]

Values are represented as mean[95% confidence interval] for continuous variables or n, %[95% confidence interval] for categorical variables.

Validation of biomarkers of preterm birth

Of the biomarkers measured, only five of seven were detectable in the study population using both a probe-based assay (Table 2, S1 Table) and RNA sequencing (S1 Table). As both the probe-based assay and RNA sequencing exhibited similar patterns, the following analysis was conducted on the probe-based assay data. Samples were tested at four concentrations (1.875, 3.75, 6.25, 25ng/uL RNA standards). Biomarkers *CST13P* and *LMLN2* were below the limit of detection (<LOD) in over 50% of the population (68% and 56% respectively) at all concentrations and thus were excluded from further analysis. One sample (term T2) was <LOD across all biomarkers, which was likely a technical issue with sample processing and thus excluded. Levels of *SH3PXD2B* were <LOD in 22% of the population and those <LOD were

294   assigned as one half of the basement level (3 MFI, mean fluorescence index units). The

295   remaining four biomarkers were present above the limit of detection in all samples.

296   **Table 2. Biomarker levels in maternal blood**

		ACAP2	*LOC101927441*	*ZNF324*	*SH3PXD2B*	*TBX2*
TERM	**T1 (n=40)**	11649[10656-12641]	546[479-613]	572[513-631]	27[18-36]	1128[1
	T2 (n=43)	10711[9727-11695]	465[397-533]	484[432-536]	24[17-31]	989[85
SPTB	**T1 (n=44)**	9868[8744-10629]	422[364-480]	444[394-494]	21[17-25]	895[75
	T2 (n=42)	11827[10049-13605]	471[388-554]	515[443-587]	39[26-52]	1042[8

Values are represented as mean [95% confidence interval] of mean fluorescence index MFI.

Four of the five measured biomarkers, *ACAP2* (p=0.0068)*, LOC101927441* (p=0.0082),

ZNF324 (p=0.0019)*,* and *TBX21* (p=0.0182) exhibited significantly lower levels in the sPTB

group compared to the term group at T1, and not at T2. When assessing biomarkers as a

measurement of T2/T1 ratios, *ACAP2* (p=0.0074)*, LOC101927441* (p=0.0273), *ZNF324*

(p=0.0170), *TBX21* (p=0.0119) ratios were significantly higher in the sPTB group than the term

group, suggesting a greater trajectory of increased expression through gestation in those with

sPTB (Figure 2). Though we do observe some differences in biomarker levels between sPTB and

term samples, only five of the eight originally identified biomarkers [21] could be measured

using the same population and methodology, and only four exhibited significant differences

between term and preterm groups.

Figure 2. Biomarkers of preterm birth. Values are reported as mean fluorescence index (MFI)

at either timepoint or a ratio of MFI values at T2 over T1. Analysed by one-way ANOVA

followed by Dunnett correction for multiple comparisons. Box and whisker plots represent

minimum, maximum and median values. *p-value<0.05, **p-value<0.01. *ACAP2*: ArfGAP with

coiled-coil, ankyrin repeat and PH domains 2, *ZNF324*: zinc finger protein 324. *SH3PCD2B*:

SH3 and PX domains 2B, *TBX21*: T-box transcription factor 21.

Differential expression analysis

320 Differential expression analysis identified 1108 genes that were differentially expressed

321 in at least one comparison, for a total of 3324 gene features (T1, T2 and T2-T1 for each gene)

322 kept for downstream modelling (S2 Tables).

Feature selection

The stepwise additive selection algorithm (SAS) identified 73 gene features which were

selected for downstream modelling using unregularized logistic regression, an L2-regularized

logistic regression, and a multilayer perceptron (MLP) neural network (S3 Table). The top (rank

#1) gene features for each iteration of cross validation (two times five-fold cross validation for a

total of ten iterations), are represented in Table 3. Notably, the top predictive genes did not show

consistency in ranking across iterations, and no features were selected more than two times in ten

iterations (range 1-2). For example, the top-most predictive feature identified by the SAS

algorithm, *MLPL51,* which encodes a mitochondrial ribosomal protein, at T1 was ranked #1

most predictive feature in two iterations but was assigned a score of zero (uninformative) the

remaining eight iterations, indicating that top features are not robust to noise in the dataset.

Table 3. Top features identified by the stepwise additive feature selection algorithm

Feature	Number of times selected	Average Rank	Gene name
MRPL51 at T2	2	1	mitochondrial ribosomal protein L51
ZNF136 at T2	2	1.5	zinc finger protein 136
TRBJ2-6 at T1	1	1	T cell receptor beta joining 2-6
DHRS7B at T1	1	1	dehydrogenase/reductase member 7B
NSRP1 at T2	1	1	nuclear speckle splicing regulatory preotin 1
C2CD2 at T1	1	1	C2 calcium dependent domain containing 2
NCALD at T1	1	1	neurocalcin delta
ZFP14 at T2	1	1	zinc finger protein 14
NOLC1 at T1	1	1	nucleolar and coiled-body phosphoprotein 1

Rows represent individual gene features. Number of times selected indicates the number of times
each feature was selected by the algorithm across ten iterations (2 times five-fold cross
validation). Average rank indicates the average rank assigned to each feature each time it was
selected, with 1 being the highest possible rank.

The elastic net logistic regression model and random forest classifier were fit after

differential expression analysis but prior to formal feature selection with SAS, as these models

perform internal feature selection. We identified 45 gene features selected by the elastic net in >5

iterations out of 10 cross validation folds, only 10 of which were the same features identified by

SAS. The random forest classifier selected 77 gene features in >5 out of 10 iterations, only 6 of

which were the same features identified by SAS (Figure 3). Notably, only four features were

identified by all three feature selection approaches (Table 4). A full list of features identified is

available in S2 Tables.

Figure 3. Features selected by the stepwise additive selector, random forest classifier, and

elastic net. Venn diagram of features selected by each method. SAS – stepwise additive selector,

RF – random forest, LR-ELN – elastic net logistic regression.

Table 4. Common features identified by three feature selection approaches

	SAS		RF		LR-ELN		
	number of times selected	average rank	number of times selected	average rank	number of times selected	average rank	
TRBJ2-6 **at T1**	1	1	7	16.6	8	25.5	
SLAMF8 **T2-T1**	1	2	6	61.9	10	175.9	
DHRS7B **T2-T1**	1	2	6	68.6	6	242	d
CCND3 **at T2**	1	3	6	236.9	7	7.1	

357 Rows represent individual gene features. Number of times selected indicates the number of times
358 each feature was selected by the algorithm across ten iterations (2 times five-fold cross
359 validation). Average rank indicates the average rank assigned to each feature each time it was
360 selected, with 1 being the highest possible rank.
361

Model performance for prediction of sPTB

All models showed promising performance in the training set, with the top performing

model was the random forest classifier (AUROC=0.99) (Figure 4, Table 5). However, all five

learning models showed significant degradation in performance when applied to the Calgary

cohort test set (AUROC range 0.54-0.59) and further degraded when validated externally

(AUROC range 0.50-0.52), which indicates a high degree of overfitting.

Figure 4. Model performance. Model performance metrics, AUROC (area under the receiver

operator curve), F1, PPV (positive predictive value), and sensitivity of each model in the

training, internal test and external validation sets.

Table 5. Model performance

Model	Predicted on	AUROC	Sensitivity	Specificity	PPV	
LR	**Calgary training**	0.86(0.83-	0.75(0.70-	0.96(0.95-0.97)	0.88(0.85-	0.

	set	0.89)	0.80)		0.91)	
LR	**Calgary test set**	0.54(0.51-0.57)	0.28(0.20-0.36)	0.80(0.75-0.85)	0.35(0.26-0.44)	0.
LR	**Detroit set**	0.52(0.5-0.54)	0.25(0.21-0.29)	0.80(0.77-0.83)	0.60(0.56-0.64)	0.
LR-ELN	**Calgary training set**	0.90(0.72-1)	0.9(0.73-1.07)	0.91(0.73-1.09)	0.91(0.74-1.08)	0.
LR-ELN	**Calgary test set**	0.59(0.56-0.62)	0.33(0.27-0.39)	0.84(0.79-0.89)	0.50(0.42-0.58)	0.
LR-ELN	**Detroit set**	0.52(0.51-0.53)	0.18(0.12-0.24)	0.87(0.83-0.91)	0.63(0.59-0.67)	0.
LR-reg	**Calgary training set**	0.84(0.8-0.88)	0.73(0.66-0.80)	0.96(0.95-0.97)	0.88(0.85-0.91)	0.
LR-reg	**Calgary test set**	0.54(0.5-0.58)	0.27(0.19-0.35)	0.82(0.78-0.86)	0.35(0.25-0.45)	0.
LR-reg	**Detroit set**	0.52(0.51-0.53)	0.24(0.21-0.27)	0.81(0.78-0.84)	0.60(0.57-0.63)	0.
RF	**Calgary training set**	0.99(0.98-1)	0.99(0.98-1)	1(1-1)	1(1-1)	
RF	**Calgary test set**	0.54(0.51-0.57)	0.14(0.08-0.2)	0.93(0.89-0.97)	0.58(0.35-0.81)	0.
RF	**Detroit set**	0.50(0.49-0.51)	0.84(0.65-1.03)	0.17(-0.03-0.37)	0.57(0.55-0.59)	0.
MLP	**Calgary training set**	0.87(0.83-0.91)	0.79(0.73-0.85)	0.96(0.95-0.97)	0.88(0.85-0.91)	0.92
MLP	**Calgary test set**	0.55(0.52-0.58)	0.30(0.24-0.36)	0.80(0.77-0.83)	0.39(0.33-0.45)	0.
MLP	**Detroit set**	0.52(0.5-0.54)	0.24(0.20-0.28)	0.81(0.79-0.83)	0.60(0.56-0.64)	0.

376 LR – logistic regression, LR-ELN – logistic regression with elastic net, LR-reg – regularized
377 logistic regression, RF – random forest, MLP – multilayer perceptron artificial feedforward
378 neural network. AUROC – area under the receiver operating curve, PPV – positive predictive
379 value, NPV – negative predictive value, FPR – false positive rate, FNR – false negative rate.
380

Assessing overfitting

All models showed significant degradation of performance in both internal and external

test sets as compared to training performance, indicating a high degree of overfitting during

training. To test the degree of overfitting, the unregularized LR and MLP algorithms were

retrained using permuted data, as these represent the lowest and highest complexity models used.

In brief, target labels (sPTB or term) were randomly scrambled during the preprocessing stage to

remove any relationship between the ground truth and predictive gene features before proceeding

with differential expression analysis, feature selection, and model training as previously

described. Using scrambled data, high performance was still observed in the training set, with the

highest performance by the MLP algorithm (AUC 0.80). This high performance despite the

forced disassociation between true sPTB labels and gene expression suggests overfitting in both

these models. Unsurprisingly, model performance was degraded when applied to both the

internal and external test sets (Table 6). These results underscore the overfitting we observed for

all five machine learning approaches.

Table 6. Model performance using permuted data

Model	Training Set	Test Set	Training AUROC	Test AUROC
MLP	Calgary	Calgary	0.80	0.49
LR	Calgary	Calgary	0.75	0.50
MLP	Calgary	Detroit	0.72	0.52
LR	Calgary	Detroit	0.63	0.52

397 Model performance as reported by area under the receiver operating curve (AUC) for two

398 prediction algorithms, multilayer perceptron (MLP) and logistic regression (LR) tested internally

399 and validated externally.

Discussion

We were unable to repeat the findings of Heng et al., [21] to predict spontaneous preterm birth using maternal blood gene expression. Most alarmingly, two of the eight topmost predictive genes were not detectable in blood samples from the same patients, suggesting issues with repeatability in probe-based RNA array methods used here, despite validation of RNA integrity over long-term storage [22]. Indeed, array reliability may be particularly problematic in lowly expressed genes and certain genes may be more subject to poor probe specificity [32, 33], though we were unable to conduct assessment of gene-specific expression levels over long-term storage. Microarray platforms for detection of gene expression signatures may be limited by low specificity of probes, and are not robust to gene variants, leading to inconsistencies in detection which likely impacted reproducibility. Additionally, we were unable to produce a more generalizable model through secondary analysis of the microarray data using alternate learning algorithms, and our results suggest a high degree of overfitting following external validation. This highlights the importance of repeat and validation studies to meaningfully progress the field of preterm birth prediction.

Heterogeneity in validation cohorts

Technical and/or biological differences between the training cohort and the cohort used for external validation may have further contributed to the lack of generalizability. One of the strengths of this study was that we were able to identify an external cohort with maternal blood gene expression data collected at the same timepoints during gestation, and used comparable microarray methods for expression analysis, however, batch effects between cohorts may still contribute to the lack of generalizability. Additionally, while the Calgary based training cohort was primarily Caucasian, with a mean age of 31, the population in the external validation cohort

from Detroit were mainly African American (92%) and younger (mean age 24). Additionally, the

Detroit based cohort had a higher percentage of a prior preterm birth, at 15.5% in the term

control population (as compared to 4% in the Calgary cohort) and as high as 50% in the sPTB

population (as compared to 17% in the Calgary cohort), suggesting that the Detroit cohort

represents a higher risk population with respect to preterm birth. The contribution of the social

determinants of health, including systemic racism and unequal access to resources and quality

healthcare among different populations including African Americans living in the United States

of America, may present distinct mechanisms leading to preterm birth as compared to the

population represented in the Calgary cohort.

Impact of overfitting and data leakage

Reassessment of the prediction model as published by Heng *et al*., [21] suggests that the

original model may have been unintentionally impacted by data leakage. Many prediction

studies, at least in the field of reproduction, are preceded by observational experiments to explore

biomarker patterns as possible predictors and to reduce the number of features for subsequent

biomarker discovery. Differential gene expression analysis is frequently used as a filtering step to

identify genes associated with an outcome. Often, these observational experiments are conducted

on the whole dataset, not the training set. As such, prediction models trained on this data are

biased by patterns that exist in the test set. This phenomenon, where information from outside

the training dataset is used to create the prediction model, is known as data leakage.

Consequently, the training dataset contains information about the outcome that would not be

otherwise available when using the model for prediction, artificially overinflating the predictive

performance when the model is applied to the test set. We, the authors, would like to express that

we believe data leakage was conducted entirely unintentionally. Indeed, we only identified the

data leakage when it was identified by a reviewer in a previous, failed submission for

publication, in which we had unintentionally committed the same type of data leakage during our

validation study.

Validation with data leakage, in our dataset, demonstrated high predictive performance in

the training set (AUC 0.72 with LR, 0.79 with MLP), and was not significantly degraded in the

test set (AUC 0.65 with LR, 0.85 with MLP). Note that with the presence of data leakage, the

machine learning MLP model had substantially higher performance (AUC 0.85) as compared to

analysis without data leakage (AUC 0.54 for comparable dataset). These misleading performance

metrics are a result of conducting differential expression analysis prior to splitting the training

and test sets. Models were built by preselecting candidate genes identified in the differential

expression analysis, which was informed by both the training and test sets, thus, model

performance was ultimately biased by information it should not have 'seen'. This stresses the

importance of methodological safeguarding and careful study design, to avoid possible sources

of bias and data leakage, particularly in omics or similar datasets that are prone to a high degree

of noise. External validation is also a highly powerful tool for testing the generalizability of

models that may have been subject to data leakage, and unintentional data leakage likely

contributes to the lack of reproducibility in some prediction studies.

One of the primary limitations to prediction using maternal blood gene expression

encountered was overfitting and noise within the dataset, which significantly skewed

performance estimates. Microarray and similar expression datasets typically contain thousands of

features, significantly inflating the feature to observation ratios in this analysis. Feature selection

approaches were unable to effectively reduce the noise within this dataset to obtain clinically

useful patterns as markers for spontaneous preterm birth. Limited sample size, particularly in the

test set, may have impacted our ability to detect true associations. Nonetheless, the sample size

of n=165 represents one of the largest maternal blood gene expression datasets available in the

pregnant population, a population that is currently understudied. A noisy dataset with high

feature to observation ratio likely also contributed to the inability to repeat and/or validate

previous findings. Possible consequences of data noise are further exacerbated when using

advanced methods such as machine learning, and, as evidenced in the study herein, high

complexity/machine learning approaches often do not demonstrate improved predictive

performance over traditional methods [34].

Biological significance of biomarkers identified

We observed limited consistency in genes identified between three feature selection

methods, suggesting that their association with sPTB is not robust. Only four common genes

were identified by the SAS, elastic net, and random forest classifiers: *TRPJ2-6, SLAMF8,*

DHRS7B, and *CCND3.* These identified genes associated with immune responses and cell

cycling. The first, *TRPJ2-6,* encodes a protein predicted to be part of the T cell receptor complex

in the J region of the variable domain of the T cell receptor beta chain that is involved in antigen

recognition [35]. *SLAMF8* encodes a member of the CD2 family of cell surface proteins involved

in lymphocyte activation. *DHRS7B* encodes a protein predicted to be involved in lipid

biosynthesis, and *CCND3* belongs to the highly conserved cyclin family and regulates the cell

cycle during G1/S transition [36]. Taken together, this may suggest that sPTB is associated with

changes in immune responses and cell cycling that can be observed in whole blood from the

maternal circulation.

Suggestions for future biomarker discovery

While maternal blood presents an enticing opportunity for minimally invasive prediction, peripheral blood is subject to a high noise signal from various physiological processes occurring within the body unrelated to uterine function during pregnancy. This stresses the need for improved feature identification. Biological compartments including cervicovaginal fluid, amniotic fluid, and the vaginal microbiome may better reflect the physiology of pregnancy and the transition to parturition [37, 38] though sample availability of reproductive and gestational tissues for research purposes is limited. An emerging strategy involves the use of cell-free nucleic acid biomarkers, which can be utilized to identify biomarkers with uterine origin in maternal blood for improved prediction of adverse pregnancy outcomes [39, 40]. Considerable research has been conducted to review the most robust predictors for sPTB, including but not limited to inflammatory biomarkers, maternal characteristics and genetic contributions [41-43], yet the most frequently used risk factors in current literature show variable predictive performance and poor robustness [44]. A recent meta-analysis identified the most robust predictors of PTB, including low gestational weight gain, interpregnancy interval following miscarriage <6months, and sleep-disordered breathing [43], and it is likely that combined biomarker approaches are necessary for prediction [45, 46]. Additionally, current literature often does not distinguish those predictors for a medically indicated PTB from those for sPTB, despite likely distinct aetiologies. It is also worth noting that the pervasive use of convenience sampling in reproductive studies (e.g. secondary analysis of biosamples used for routine antenatal screening) are not necessarily performed proximal to the outcome of interest (PTB). For many subjects, the delay from testing to outcome may make identifying true associations difficult.

Strategies to improve prediction

Future research in identifying biomarkers for the mechanisms of preterm labour are

important and likely requires biologically informed selection over unbiased feature selection, as

these methods may suffer more from high noise to feature ratios, especially considering the

heterogeneity of PTB and biological variation between individuals. Recent systematic reviews

have highlighted that currently described biomarkers have yet to show promising associations for

utility in prediction [47]. Yet, recent advancements point to proteomic signatures as a promising

biomarker for preterm birth [48], and biomarker discovery targeted to subtypes of PTB such as

PPROM [49]. The best models combine an understanding of the features (such as genes,

proteins, or patient characteristics) that are most important for determining the outcome and

robust methodologies. Identification of robust predictors of sPTB likely requires a combination

of advanced selection techniques and an improved understanding of the physiological

mechanisms of labour (including known pathways of biomarkers involved in sPTB).

Looking ahead, our recommendations for future research include safeguarding against

sources of data leakage, implementing cross-validation techniques as a measure of robustness,

external validation as a measurement of overfitting, and prioritizing repeatability and

reproducibility of findings. Specific recommendations for addressing data leakage are to ensure

testing data has no influence on model training, including in any discovery analysis (such as

differential expression analysis) which may precede model training. Dimensionality reduction

techniques, such as principal component analysis (PCA), can mediate overfitting by reducing

overall dimensionality. However, one disadvantage of dimensionality reduction using techniques

like PCA is that they collapse predictors into groups of predictors, which is less enticing for

540   clinical application, as cost and feasibility of clinical testing scales with the number of predictors.

541   Other recommendations for addressing overfitting are to conduct permutation tests to distinguish

542   true patterns from noise in addition to external validation. This likely includes incentivizing

543   repeated studies in published literature and improving data management, storage, and sharing

544   infrastructure, particularly for larger, multicentre studies [10, 11].

Limitations

The strength of this study was to undertake a replication and external validation, which is,

unfortunately, lacking in many prediction studies for preterm birth. However, this study is not

without limitations. The current assay was unable to measure miRNA, preventing the

measurement of one biomarker *MIR3691*, further, two of the biomarkers were not found to be

expressed on repetition of the study. Together this impacted our ability to fully replicate the

prediction model . As commercially available microarray assays were used, it leads to important

considerations of assay reliability. Additionally, the biomarker *SH3PXD2B* was expressed below

the limit of detection in 22% of the samples, for which values were imputed at half the limit of

detection. Replacing values with half the limit of the detection has not been shown to

significantly impact means when the percentage of imputed values is low, less than 25% [50].

Nonetheless, imputation of missing values can contribute to bias within the measurement, which

may explain why were unable to repeat the finding that *SH3PXD2B* expression is different in the

preterm population compared to term.

A nested 2:1 case control study design was selected to address class imbalance, as the

typical rate of sPTB is ~10%. However, given the moderate class imbalance in this study design,

and the possible bias inherent to resampling or weighting methods, no additional methods to

address class imbalance was conducted. There is no consensus on what degree of class imbalance

is acceptable, though moderate class imbalances (e.g. 2:1) would have more limited impacts on

model training as compared to 10:1. However, it has been suggested error rates for moderate

class imbalances (between 1:1 and 1:3) are underestimated, and more problematic than

commonly acknowledged [51]. As such, future work in exploring prediction models for preterm

birth could consider class weighting or resampling techniques, such as SMOTE (synthetic

minority over-sampling) of preterm birth cases or under-sampling of term cases to address this

class imbalance, in addition to a case control study design.

Conclusion

Our study serves as a cautionary tale for researchers, emphasizing the need for

transparency, rigorous methodological standards, as well as not only repeating results but

validation in external cohorts to advance the field of spontaneous preterm birth prediction

responsibly. Our findings also underscore the broader implications for omics studies for

discovery analysis, where high feature-to-observation ratios are common, which exacerbates the

challenge for mitigating bias and ensuring the reliability of predictive models. For example,

differential expression analysis without appropriate training and testing sets for validation

introduces inherent bias and limits the generalizability of patterns identified. Testing on internal

test sets alone is insufficient for measuring generalizability, especially in the instance of data

leakage. Yet, assessments of overfitting and external validation are not standard practice in

preterm birth prediction, and the authors stress their importance for meaningful future work in

this field.

Current studies on the prediction of sPTB suffer from a lack of external validation, and

perhaps unintentional data leakage, leading to a lack of generalized models that would be

clinically useful. Recent publication studies have identified various potential predictors of sPTB

in both maternal and fetal compartments, though many have not demonstrated robust prediction

in external validation nor are repeat studies of promising predictors are not common [52-55].

Further, our results suggest that maternal blood gene expression may either not be predictive of

sPTB, or that a high degree of noise limits our ability to detect true associations, stressing the

need for a more robust biomarker discovery. Identification of a few potential cell cycling and

immune biomarkers presents novel areas for future biomarker discovery. A better understanding

of biomarkers associated with sPTB would not only contribute to insight on the mechanism of

sPTB, but also to improved prediction of sPTB, allowing for targeted intervention for those at

risk for poor maternal and neonatal health outcomes related to prematurity.

References

1.   Ohuma EO, Moller AB, Bradley E, Chakwera S, Hussain-Alkhateeb L, Lewin A, et al. National,
regional, and global estimates of preterm birth in 2020, with trends from 2010: a systematic analysis.
Lancet. 2023;402(10409):1261-71. doi: 10.1016/s0140-6736(23)00878-4. PubMed PMID: 37805217.

2.   Purisch SE, Gyamfi-Bannerman C. Epidemiology of preterm birth. Seminars in Perinatology.
2017;41(7):387-91. doi: https://doi.org/10.1053/j.semperi.2017.07.009.

3.   Perin J, Mulick A, Yeung D, Villavicencio F, Lopez G, Strong KL, et al. Global, regional, and
national causes of under-5 mortality in 2000-19: an updated systematic analysis with implications for the
Sustainable Development Goals. Lancet Child Adolesc Health. 2022;6(2):106-15. Epub 20211117. doi:
10.1016/s2352-4642(21)00311-4. PubMed PMID: 34800370; PubMed Central PMCID:
PMCPMC8786667.

4.   Hornaday KK, Wood EM, Slater DM. Is there a maternal blood biomarker that can predict
spontaneous preterm birth prior to labour onset? A systematic review. PLOS ONE. 2022;17(4):e0265853.
doi: 10.1371/journal.pone.0265853.

5.   Marić I, Stevenson DK, Aghaeepour N, Gaudillière B, Wong RJ, Angst MS. Predicting Preterm
Birth Using Proteomics. Clin Perinatol. 2024;51(2):391-409. Epub 20240323. doi:
10.1016/j.clp.2024.02.011. PubMed PMID: 38705648; PubMed Central PMCID: PMCPMC11186213.

6.   Ramachandran A, Clottey KD, Gordon A, Hyett JA. Prediction and prevention of preterm birth:
Quality assessment and systematic review of clinical practice guidelines using the AGREE II framework.
Int J Gynaecol Obstet. 2024. Epub 20240415. doi: 10.1002/ijgo.15514. PubMed PMID: 38619379.

7.   Yang Q, Fan X, Cao X, Hao W, Lu J, Wei J, et al. Reporting and risk of bias of prediction models
based on machine learning methods in preterm birth: A systematic review. Acta Obstet Gynecol Scand.
2023;102(1):7-14. Epub 20221117. doi: 10.1111/aogs.14475. PubMed PMID: 36397723; PubMed Central
PMCID: PMCPMC9780725.

8.   Staffa SJ, Zurakowski D. Statistical Development and Validation of Clinical Prediction Models.
Anesthesiology. 2021;135(3):396-405. doi: 10.1097/aln.0000000000003871. PubMed PMID: 34330146.

9.   Steyerberg EW, Harrell FE, Jr. Prediction models need appropriate internal, internal-external, and
external validation. J Clin Epidemiol. 2016;69:245-7. PubMed PMID: 25981519.

10.   Sharifi-Heris Z, Laitala J, Airola A, Rahmani AM, Bender M. Machine Learning Approach for
Preterm Birth Prediction Using Health Records: Systematic Review. JMIR Med Inform.
2022;10(4):e33875. Epub 20220420. doi: 10.2196/33875. PubMed PMID: 35442214; PubMed Central
PMCID: PMCPMC9069277.

11.   Mennickent D, Rodríguez A, Opazo MC, Riedel CA, Castro E, Eriz-Salinas A, et al. Machine
learning applied in maternal and fetal health: a narrative review focused on pregnancy diseases and
complications. Front Endocrinol (Lausanne). 2023;14:1130139. Epub 20230519. doi:
10.3389/fendo.2023.1130139. PubMed PMID: 37274341; PubMed Central PMCID: PMCPMC10235786.

12.   Leisman DE, Harhay MO, Lederer DJ, Abramson M, Adjei AA, Bakker J, et al. Development and
Reporting of Prediction Models: Guidance for Authors From Editors of Respiratory, Sleep, and Critical
Care Journals. Crit Care Med. 2020;48(5):623-33. PubMed PMID: 32141923.

13.   Collins GS, Dhiman P, Ma J, Schlussel MM, Archer L, Van Calster B, et al. Evaluation of clinical
prediction models (part 1): from development to external validation. Bmj. 2024;384:e074819. Epub
20240108. doi: 10.1136/bmj-2023-074819. PubMed PMID: 38191193; PubMed Central PMCID:
PMCPMC10772854.

14.   Lenain R, Dantan E, Giral M, Foucher Y, Asar Ö, Naesens M, et al. External Validation of the
DynPG for Kidney Transplant Recipients. Transplantation. 2021;105(2):396-403. doi:
10.1097/tp.0000000000003209. PubMed PMID: 32108750.

15.   Russell FM, Herbert A, Kennedy S, Nti B, Powell M, Davis J, et al. External validation of the
ultrasound competency assessment tool. AEM Educ Train. 2023;7(3):e10887. Epub 20230622. doi:
10.1002/aet2.10887. PubMed PMID: 37361190; PubMed Central PMCID: PMCPMC10288010.

16.   Yun JS, Han K, Choi SY, Cha SA, Ahn YB, Ko SH. External validation and clinical application of
the predictive model for severe hypoglycemia. Front Endocrinol (Lausanne). 2022;13:1006470. Epub
20220929. doi: 10.3389/fendo.2022.1006470. PubMed PMID: 36246915; PubMed Central PMCID:
PMCPMC9556834.

17.   Slieker RC, van der Heijden A, Siddiqui MK, Langendoen-Gort M, Nijpels G, Herings R, et al. Performance of prediction models for nephropathy in people with type 2 diabetes: systematic review and external validation study. Bmj. 2021;374:n2134. Epub 20210928. doi: 10.1136/bmj.n2134. PubMed PMID: 34583929; PubMed Central PMCID: PMCPMC8477272.

18.   Chaemsaithong P, Sahota DS, Poon LC. First trimester preeclampsia screening and prediction. Am J Obstet Gynecol. 2022;226(2s):S1071-S97.e2. Epub 20200716. doi: 10.1016/j.ajog.2020.07.020. PubMed PMID: 32682859.

19.   Neary C, Naheed S, McLernon DJ, Black M. Predicting risk of postpartum haemorrhage: a systematic review. Bjog. 2021;128(1):46-53. Epub 20200805. doi: 10.1111/1471-0528.16379. PubMed PMID: 32575159.

20.   Feyaerts D, Marić I, Arck PC, Prins JR, Gomez-Lopez N, Gaudillière B, et al. Predicting Spontaneous Preterm Birth Using the Immunome. Clin Perinatol. 2024;51(2):441-59. Epub 20240404. doi: 10.1016/j.clp.2024.02.013. PubMed PMID: 38705651.

21.   Heng YJ, Pennell CE, McDonald SW, Vinturache AE, Xu J, Lee MWF, et al. Maternal whole blood gene expression at 18 and 28 weeks of gestation associated with spontaneous preterm birth in asymptomatic women. PLoS ONE. 2016;11(6). PubMed PMID: 610977402.

22.   Stephenson NL, Hornaday KK, Doktorchik CTA, Lyon AW, Tough SC, Slater DM. Quality assessment of RNA in long-term storage: The All Our Families biorepository. PLoS One. 2020;15(12):e0242404. Epub 20201201. doi: 10.1371/journal.pone.0242404. PubMed PMID: 33259520; PubMed Central PMCID: PMCPMC7707483.

23.   Tarca AL, Pataki BÁ, Romero R, Sirota M, Guan Y, Kutum R, et al. Crowdsourcing assessment of maternal blood multi-omics for predicting gestational age and preterm birth. Cell Reports Medicine. 2021;2(6):100323-. doi: 10.1016/j.xcrm.2021.100323.

24.   Alpaydin E. Introduction to machine learning. Ieee Xplore d, Mit Press p, ebrary I, editors: Cambridge, Massachusetts : MIT Press; Third edition.; 2014.

25.   Theodoridis S. Machine learning : a Bayesian and optimization perspective: Amsterdam, Netherlands : Academic Press; First edition.; 2015.

26.   Dhar V. Data science and prediction. Commun ACM. 2013;56(12):64–73. doi: 10.1145/2500499.

27.   Arain Z, Iliodromiti S, Slabaugh G, David AL, Chowdhury TT. Machine learning and disease prediction in obstetrics. Curr Res Physiol. 2023;6:100099. Epub 20230519. doi: 10.1016/j.crphys.2023.100099. PubMed PMID: 37324652; PubMed Central PMCID: PMCPMC10265477.

28.   McDonald CR, Darling AM, Conroy AL, Tran V, Cabrera A, Liles WC, et al. Inflammatory and angiogenic factors at mid-pregnancy are associated with spontaneous preterm birth in a cohort of Tanzanian women. PLoS ONE. 2015;10(8). PubMed PMID: 606057658.

29.   Gracie SK, Lyon AW, Kehler HL, Pennell CE, Dolan SM, McNeil DA, et al. All Our Babies Cohort Study: recruitment of a cohort to predict women at risk of preterm birth through the examination of gene expression profiles and the environment. BioMed Central;http://www.biomedcentral.com/1471-2393/10/87;University of Calgary;Medicine; 2011.

30.   Tough SC, McDonald SW, Collisson BA, Graham SA, Kehler H, Kingston D, et al. Cohort
Profile: The All Our Babies pregnancy cohort (AOB). International Journal of Epidemiology.
2017;46(5):1389-90k. doi: 10.1093/ije/dyw363.

31.   Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, et al. STAR: ultrafast universal
RNA-seq aligner. Bioinformatics. 2012;29(1):15-21. doi: 10.1093/bioinformatics/bts635.

32.   Draghici S, Khatri P, Eklund AC, Szallasi Z. Reliability and reproducibility issues in DNA
microarray measurements. Trends Genet. 2006;22(2):101-9. Epub 20051227. doi:
10.1016/j.tig.2005.12.005. PubMed PMID: 16380191; PubMed Central PMCID: PMCPMC2386979.

33.   Kothapalli R, Yoder SJ, Mane S, Loughran TP, Jr. Microarray results: how accurate are they?
BMC Bioinformatics. 2002;3:22. Epub 20020823. doi: 10.1186/1471-2105-3-22. PubMed PMID:
12194703; PubMed Central PMCID: PMCPMC126254.

34.   Christodoulou E, Ma J, Collins GS, Steyerberg EW, Verbakel JY, Van Calster B. A systematic
review shows no performance benefit of machine learning over logistic regression for clinical prediction
models. J Clin Epidemiol. 2019;110:12-22. Epub 20190211. doi: 10.1016/j.jclinepi.2019.02.004. PubMed
PMID: 30763612.

35.   Lefranc M-P. Immunoglobulin and T Cell Receptor Genes: IMGT(®) and the Birth and Rise of
Immunoinformatics. Frontiers in immunology. 2014;5:22-. doi: 10.3389/fimmu.2014.00022.

36.   Meyerson M, Harlow E. Identification of G1 kinase activity for cdk6, a novel cyclin D partner.
Molecular and Cellular Biology. 1994;14(3):2077-86. doi: 10.1128/MCB.14.3.2077.

37.   Chakoory O, Barra V, Rochette E, Blanchon L, Sapin V, Merlin E, et al. DeepMPTB: a vaginal
microbiome-based deep neural network as artificial intelligence strategy for efficient preterm birth
prediction. Biomarker Research. 2024;12. doi: 10.1186/s40364-024-00557-1.

38.   Chang Y, Li W, Shen Y, Li S, Chen X. Association between interleukin-6 and preterm birth: a
meta-analysis. Ann Med. 2023;55(2):2284384. Epub 20231127. doi: 10.1080/07853890.2023.2284384.
PubMed PMID: 38010798; PubMed Central PMCID: PMCPMC10836263.

39.   Cowan AD, Rasmussen M, Jain M, Tribe RM. Predicting Preterm Birth Using Cell-Free
Ribonucleic Acid. Clin Perinatol. 2024;51(2):379-89. Epub 20240313. doi: 10.1016/j.clp.2024.02.008.
PubMed PMID: 38705647.

40.   Moufarrej MN, Vorperian SK, Wong RJ, Campos AA, Quaintance CC, Sit RV, et al. Early
prediction of preeclampsia in pregnancy with cell-free RNA. Nature. 2022;602(7898):689-94. Epub
20220209. doi: 10.1038/s41586-022-04410-z. PubMed PMID: 35140405; PubMed Central PMCID:
PMCPMC8971130.

41.   Tang ID, Mallia D, Yan Q, Pe'er I, Raja A, Salleb-Aouissi A, et al. A Scoping Review of Preterm
Birth Risk Factors. Am J Perinatol. 2024;41(S 01):e2804-e17. Epub 20230925. doi: 10.1055/s-0043-
1775564. PubMed PMID: 37748506.

42.   Li J, Ge J, Ran N, Zheng C, Fang Y, Fang D, et al. Finding the priority and cluster of
inflammatory biomarkers for infectious preterm birth: a systematic review. J Inflamm (Lond).
2023;20(1):25. Epub 20230724. doi: 10.1186/s12950-023-00351-0. PubMed PMID: 37488605; PubMed
Central PMCID: PMCPMC10367376.

43. Mitrogiannis I, Evangelou E, Efthymiou A, Kanavos T, Birbas E, Makrydimas G, et al. Risk
factors for preterm birth: an umbrella review of meta-analyses of observational studies. BMC Med.
2023;21(1):494. Epub 20231213. doi: 10.1186/s12916-023-03171-4. PubMed PMID: 38093369; PubMed
Central PMCID: PMCPMC10720103.

44. Ferreira A, Bernardes J, Gonçalves H. Risk Scoring Systems for Preterm Birth and Their
Performance: A Systematic Review. J Clin Med. 2023;12(13). Epub 20230628. doi:
10.3390/jcm12134360. PubMed PMID: 37445395; PubMed Central PMCID: PMCPMC10342801.

45. Mirzaei A, Hiller BC, Stelzer IA, Thiele K, Tan Y, Becker M. Computational Approaches for
Connecting Maternal Stress to Preterm Birth. Clin Perinatol. 2024;51(2):345-60. Epub 20240315. doi:
10.1016/j.clp.2024.02.003. PubMed PMID: 38705645.

46. Creswell L, Rolnik DL, Lindow SW, O'Gorman N. Preterm Birth: Screening and Prediction. Int J
Womens Health. 2023;15:1981-97. Epub 20231221. doi: 10.2147/ijwh.s436624. PubMed PMID:
38146587; PubMed Central PMCID: PMCPMC10749552.

47. Hornaday KK, Stephenson NL, Canning MT, Tough SC, Slater DM, Fujioka K. Maternal
cytokine profiles in second and early third trimester are not predictive of preterm birth. PloS one.
2024;19(12):e0311721. doi: 10.1371/journal.pone.0311721.

48. Elkahlout R, Mohammed SGAA, Najjar A, Farrell T, Rifai HA, Al-Dewik N, et al. Application of
Proteomics in Maternal and Neonatal Health: Advancements and Future Directions. Proteomics Clinical
applications. 2025:e70004. doi: 10.1002/prca.70004.

49. Kirk M, RothEkmann J, Overgaard M, Ekelund CK, Hegaard HK, Rode L. A systematic review
of first-trimester blood biomarkers associated with preterm prelabor rupture of the fetal membranes.
Biomarkers. 2025:1-21. doi: 10.1080/1354750X.2025.2475474.

50. Croghan W, Egeghy PP, editors. Methods of Dealing with Values Below the Limit of Detection
using SAS Carry2003.

51. Weiss GM, He H, Ma Y. Foundations of Imbalanced Learning. 1 ed. United States: Wiley; 2013.
p. 13-41.

52. Becking EC, Bekker MN, Henrichs J, Bax CJ, Sistermans EA, Henneman L, et al. Fetal Fraction
of Cell Free DNA in the Prediction of Adverse Pregnancy Outcomes: A Nationwide Retrospective
Cohort Study. BJOG : an international journal of obstetrics and gynaecology. 2025;132(3):318-25. doi:
10.1111/1471-0528.17978.

53. Chen Y, Shi X, Wang Z, Zhang L. Development and validation of a spontaneous preterm birth
risk prediction algorithm based on maternal bioinformatics: A single-center retrospective study. BMC
pregnancy and childbirth. 2024;24(1):763-12. doi: 10.1186/s12884-024-06933-x.

54. Zhang Y, Sylvester KG, Wong RJ, Blumenfeld YJ, Hwa KY, Chou CJ, et al. Prediction of risk for
early or very early preterm births using high-resolution urinary metabolomic profiling. BMC pregnancy
and childbirth. 2024;24(1):783-11. doi: 10.1186/s12884-024-06974-2.

55. Ramzan F, Rong J, Roberts CT, O'Sullivan JM, Perry JK, Taylor R, et al. Maternal Plasma
miRNAs as Early Biomarkers of Moderate-to-Late-Preterm Birth. International journal of molecular
sciences. 2024;25(17):9536. doi: 10.3390/ijms25179536.

Supporting Information

S1 Table. Raw fluorescence index and sequencing reads for predictive genes tested in maternal blood. Isolated RNA from whole maternal blood was analyzed for gene expression using a QuantiGene Plex custom assay (Qiagen) (S1.1) and using bulk RNA sequencing (S1.2).

S2 Tables. Differential expression analysis results. Fold differences in each comparison, average expression, F value, p-value and adjusted p-value for each gene identified as differentially expressed in each fold of 10-fold cross validation.

S3 Table. Feature selection. Features selected by the stepwise additive selector (SAS), elastic net regression model (LR-ELN) and random forest classifier.