1

A cross-sectional survey of General Practitioners' knowledge of the wait times for

adolescent mental health specialists and services in Australia

Bridianne O'Dea[1,2]

Mirjana Subotic-Kerry[2]

Thomas Borchard[2]

Belinda Parker[2]

Bojana Vilus[2]

Frank Iorfino[3]

Alexis E. Whitton[2]

Ben Harris-Roxas[4]

Tracey D. Wade[1]

Madelaine K. de Valle[1]

Nicholas Glozier[5,6]

Jennifer Nicholas[7,8]

Michelle Torok[2,3]

Taylor A. Braund[2,3]

Philip J. Batterham[9]

Author affiliations:

1. Flinders University Institute for Mental Health and Wellbeing, College of Education, Psychology and Social Work, Bedford Park, South Australia, Australia.

2. Black Dog Institute, Faculty of Medicine and Health, University of New South Wales, Sydney, New South Wales, Australia.

3. Brain and Mind Centre, University of Sydney, Sydney, New South Wales, Australia

4. School of Population Health, Faculty of Medicine and Health, University of New South Wales, Sydney, New South Wales, Australia

5. Sydney Medical School, University of Sydney, Sydney, New South Wales, Australia

6. Australian Research Council Centre of Excellence for Children and Families over the Life Course

7. Centre for Youth Mental Health, University of Melbourne, Melbourne, Victoria, Australia

8. Orygen, Melbourne, Victoria, Australia

9. Centre for Mental Health Research, Australian National University, Canberra, Australian Capital Territory, Australia.

Corresponding author:

Bridianne O'Dea

Flinders University Institute for Mental Health and Wellbeing, Flinders University, Adelaide, South Australia, Australia, 5042

Email: bridianne.odea@flinders.edu.au

ORCID: 0000-0003-1731-210X

Statements and Declarations

Competing Interests:

Ben Harris-Roxas is an Associate Editor of Australian Journal of Primary Health. To mitigate this potential conflict of interest they were blinded from the review process.

Declaration of Funding:

This research was supported by a donation from the Buxton Family Foundation, Australian Unity, the Frontiers Technology Clinical Academic Group Industry Connection Seed Funding Scheme, and the UNSW Medicine, Neuroscience, Mental Health and Addiction Theme and SPHERE Clinical Academic Group Collaborative Research Funding. BOD is supported by an NHMRC MRFF Investigator Fellowship (MRF1197249). AEW is supported by an NHMRC Investigator Fellowship (2017521). This work is supported by a NHMRC MRFF Million Minds Grant (MRF2035416).

Author contributions:

BOD conceived the project and led the funding acquisition with support from MSK, BP, AEW, BHR. BOD, BP, BHR, and TB led the development of the survey. BP, TB, MSK and EL provided research and operational support. MSK, TB, and BV analysed the data with statistical support from PJB. NG, JN, AEW, MT, TAB, and FI were chief investigators of some of the funding and provided scientific oversight. BOD and BV wrote the first draft of the manuscript with all authors providing feedback. All authors reviewed and approved the final manuscript.

ABSTRACT

Background: General Practitioners (GPs) play a key role in referring adolescents with

depression and/or anxiety to mental health specialists and services, but their capacity to do so

may be compromised by service wait times. It is unclear how GPs manage the mental

healthcare of adolescents when the choice of treatment is not available. This study aimed to

explore GPs' self-reported referral practices to mental health specialists and services for

adolescent depression and/or anxiety, as well as their perceived knowledge, acceptability, and

impacts of the wait times for these.

Methods: A cross-sectional online survey of 192 GPs in Australia who self-identified as

treating adolescents (12 to 17 years old) with depression and/or anxiety.

Results: GPs most frequently referred adolescents with depression and/or anxiety to

psychologists. However, the mean estimated wait time for psychologists was 57.26 days (SD:

47.91, Mdn: 45.0, range: 5-365), which was four times the proposed acceptable wait time (M:

14.66 days, SD: 8.70). Nearly all GPs (81.8%) had increased their level of care for

adolescents due to long waits but had limited training in and knowledge of strategies for

effective self-management.

Conclusions: GPs in Australia lack information on the wait times for adolescent mental

health specialists and services, despite frequent referrals. Greater knowledge of wait times,

training in wait time approaches, and self-directed digital interventions may help to enhance

the quality of primary care provided to adolescents.

Keywords: Primary care, General Practitioner, Mental Health Services, Adolescent,

Depression

INTRODUCTION

General Practitioners (GPs) are the first source of professional help for many

adolescents experiencing depression and anxiety and play a critical role in facilitating access

to specialist treatment (Crouch et al., 2019; Lawrence et al., 2024; MacDonald et al., 2018).

When treating mental illness, GPs are expected to follow clinical guidelines and escalate care

as required. Australia's clinical guidelines recommends specialists referrals for psychological

therapy as first-line treatment for youth depression (Malhi et al., 2021) and anxiety (Andrews

et al., 2018), with pharmacotherapies reserved for more severe cases with specialist

supervision due to potential side effects. Other public mental health services (e.g.

Headspace), community Child and Adolescent Mental Health Services (CAMHS) and

hospital inpatient centres support primary care. However, psychological presentations to GPs

have sharply increased over time, placing further strain on an already overburdened primary

care system (Daneshmand et al., 2023; John et al., 2023; Russell et al., 2024; Villas-Boas et

al., 2023). Mounting evidence suggests that GPs' capacity to provide appropriate care and

utilise referrals is increasingly limited by long wait times to mental health specialists and

services (Dumesnil et al., 2018; Fleury et al., 2012; Paton et al., 2021; Petrie et al., 2021).

Despite no nationally agree-upon standards, up to 40% of Australians mental help-

seekers waited 'longer than acceptable' for care (Mental Health Commission of New South

Wales, 2024). On average, adolescents wait more than 100 days (Edbrooke-Childs &

Deighton, 2020; Kowalewski et al., 2011; McNicholas et al., 2021; Punton et al., 2022;

Subotic-Kerry et al., 2025) with similar or longer delays for psychiatrists and paediatricians

globally (Edbrooke-Childs & Deighton, 2020; Kowalewski et al., 2011; Smith et al., 2018).

Comparatively, systems in the UK, Norway and Canada follow wait time benchmarks of 4-6

weeks for psychological therapy. Delays to care represent a period of elevated clinical risk

for adolescents and their GPs, as symptoms may worsen but treatment has not commenced.

GPs often receive no information about expected wait times (Fleury et al., 2012) despite

impacts on patient confidence and satisfaction (Bleustein et al., 2014). Faced with long waits,

many adolescents return to their GP (Subotic-Kerry et al., 2025), placing further strain on

practitioners who often lack the training, resources, and time to provide adequate interim

support (De Silva et al., 2017). Despite these challenges, clinical guidelines offer little advice

to GPs on how to proceed when first-line treatment is unavailable.

It is unclear how Australian GPs adapt to service shortages in youth mental health. A

recent systematic review identified only 18 waiting list interventions, and none were initiated

in primary care (Valentine et al., 2024). Some evidence suggests GPs have recommended

lifestyle changes (e.g. exercise) and digital mental health tools (Keyes et al., 2023), scheduled

follow-up appointments (Budd et al., 2021), delivered brief interventions through practice

nurses (Koet et al., 2024) or medication (Ball et al., 2023). However short consultation times

limit the extent of support that GPs can offer. Despite the known risks of delayed care, little

is understood about how GPs manage adolescent mental health in the context of prolonged

wait times.

Aims of current study

This study explored GPs' experiences of wait times for mental health specialists and

services for adolescents with depression and/or anxiety in Australia. We examined their

referral practices, awareness of wait times, perceptions of acceptable waits, and the impacts

on GPs, patients, and treatment. Findings will inform how service availability affects GPs'

capacity to deliver timely, high-quality care to adolescents.

142   **METHOD**

Design

This study utilised a national cross-sectional online survey (co-designed with GPs)

conducted in Australia between May 2022 to November 2022, at the end of the second wave

of the COVID-19 pandemic. The sample size target was estimated on the proportion of GPs

in Australia at the time of the survey (N=26, 599) who were likely to have mental health

presentations from adolescents (overestimated to be 90%, given lack of available data), a

95% Confidence Level and margin of error of 5%. Based on this, the target sample was 379.

The study was approved by the University of New South Wales Sydney Human Research

Ethics Committee (HC220107).

Participants, recruitment, and consent

Eligible participants were Australian GPs who self-reported treating adolescents (aged

12–17) with depression and/or anxiety and had verified registration with Allied Health

Practitioner Regulation Agency (AHPRA) and the Royal Australian College of General

Practitioners (RACGP) or the Australian College of Rural and Remote Medicine (ACRRM).

Screening included registration verification, correct reporting of valid Medicare Benefits

Schedule (MBS) item numbers and naming of their Primary Health Network (PHN). GPs

were recruited via articles, GP magazines, paid and unpaid social media posts, GP

conferences, PHN newsletters, and email lists from affiliated organisations. Advertisements

linked to an online survey that contained study information and consent. Eligible participants

received a $50 AUD voucher and were emailed study results.

Measures

Survey items were adapted from prior studies (Fleury et al., 2012; Steele et al., 2010).

See Supplementary Table S1 for full details.

GP Characteristics: Age, gender identity, work location (state/territory, metro vs

rural/regional), years of experience, employment status (full-/part-time), and number of

practices. GPs reported frequency of adolescent (12–17 years) consultations per week,

proportion of adolescent patients, and proportion with depression/anxiety. They indicated if

adolescent depression/anxiety presentations had increased, remained stable, or decreased over

12 months. Additional mental health training, specifically in adolescent mental health, and

frequency of parent/guardian attendance during consultations were recorded.

Referral and Wait Times: Using 5-point Likert scales (never=1 to always=5), GPs

rated how often they referred to mental health professionals/services or recommended other

interventions. They estimated wait times (days) for subsidised/public services (psychologists,

psychiatrists, headspace, CAMHS, inpatient centres), with an 'unsure' option. Acceptability

of these wait times was rated on a 0–5 scale, and GPs indicated what they considered

acceptable wait times in days. GPs reported whether they or their practice monitored wait

times, and how helpful this knowledge was for treatment decisions. Using 5-point Likert

scales (strongly disagree=1 to strongly agree=5), GPs rated agreement with statements on the

impact of wait times on their duties and patient outcomes. They also indicated how often they

referred adolescents elsewhere or increased their care due to long waits.

Care During Wait Times: GPs rated the importance of providing mental health

strategies/resources to waiting adolescents (1=not at all to 5=extremely important).

Frequency of recommending 13 listed interventions (e.g., sleep hygiene) was rated (never=1

to always=5), with free-text options for other interventions and suggestions on helpful

supports during waits. GPs rated the quality of clinical support provided to adolescents and

their parents/guardians (1=very poor to 5=excellent) and self-rated their knowledge of helpful

strategies/resources during wait times (1=no knowledge to 5=superior knowledge).

Data analyses

Qualtrics administered the survey. Invalid responses were determined by email/IP checks and response patterns (e.g., item responses, response speed). Suspected cases were reviewed by two researchers and discrepancies resolved by a third (see Supplementary Table S2). Analyses were conducted in SPSS. Participants' postcodes were used to classify Geographic area was classified using postcodes and the Australian Government's classification system. Wait times exceeding 12 months were excluded given referral expiry dates. For "select all that apply" items, percentages were based on total responses. To compare routine and wait-time care, mean scores were calculated and, where appropriate, combined into broader intervention types (e.g., diet and physical activity were grouped as lifestyle modifications). Free-text responses were analysed thematically by two independent raters, with disagreements resolved by a third. Associations between referral/recommendation frequencies and other variables were assessed using ordinal correlations (Spearman's Rho), with a Bonferroni correction applied ($p < .002$). This was used to examine associations between GPs' knowledge of wait times and background factors (e.g., experience, gender, weekly youth contact, adolescent patient load, mental health training) and referral frequency.

RESULTS

Sample characteristics

A total of 837 surveys were submitted, with 645 (77.1%) removed due to being invalid responses ($n=523$), ineligible ($n=62$), incomplete ($n=55$), or completed too quickly ($n=7$). The final sample consisted of 190 GPs. Most participants were female (71.6%, n=136), worked full-time (60.0%, n=114), and were based in a single practice (71.6%, n=136) in metropolitan areas (69.5%, n=132; see Supplementary Table S3 for state

breakdown). GPs had a mean age of 42.38 years (SD = 10.25, range = 27–73) and 11.69

years of experience (SD = 9.93, range = 0.5–50). Two-thirds (66.8%, n=127) frequently

treated adolescents (12–17 years), with most reporting adolescents made up a moderate

(60.0%, n=114) to very high (12.1%, n=20) share of their caseload. The majority reported a

moderate (46.3%, n=88) to high/very high (45.2%, n=86) prevalence of adolescent

depression and/or anxiety, which had increased over the past year (80.0%, n=152). Parents

were often or always present during consultations (57.9%, n=110). While 63.7% (n=121) had

completed additional mental health training, fewer (28.9%, n=55) had specific training in

youth mental health.

GPs' referrals and recommendations for adolescent depression and/or anxiety.

As shown in Table 1, referral and recommendation frequency varied among GPs

treating adolescent depression and/or anxiety. Most GPs (92.1%, n=175) "always" or "often"

referred to psychologists (M = 4.24, SD = 0.67), while 39.5% (n=75) did so for headspace

centres (M = 3.23, SD = 0.88). All GPs recommended ongoing GP care, lifestyle changes,

and sleep hygiene to some extent, with moderate to strong positive correlations between these

(r = 0.35–0.68, $p < .001$). Fewer than 1 in 10 GPs (7.9%, n=14) frequently prescribed

medication. In free-text responses, some GPs recommended social/family/peer support

(12.3%), mindfulness or meditation (8.2%), or further medical investigations (5.5%).

Table 1. Frequency of GP referrals and/or recommendations to mental health specialists

and services for adolescent depression and/or anxiety in Australia (N=190).

Type of specialist and/or service provider	M	SD	Mdn	Range
Referrals to health professionals				

Psychologist	4.24	0.67	4.0	1-5
School counsellor/psychologist	3.32	0.79	3.0	1-5
Psychiatrist	2.75	0.67	3.0	1-5
Paediatrician	2.75	0.77	3.0	1-5
Any other allied health care provider	2.59	0.94	3.0	1-5
Referrals to services or programs				
Headspace centres	3.23	0.88	3.0	1-5
Local Child and Adolescent Mental Health Service (CAMHS)	2.60	0.76	3.0	1-5
Structured in-person therapeutic programs (e.g., Cool Kids)	2.48	1.12	2.0	1-5
Support groups (e.g., a group of people meeting to share experiences)	2.04	1.01	2.0	1-5
Inpatient centres	1.99	0.66	2.0	1-5
Other GP-led interventions				
Ongoing appointments with GP	4.40	0.83	5.0	1-5
Medication (e.g. anti-depressants)	2.72	0.66	3.0	1-5
Other recommendations				
Lifestyle modification (e.g., diet and/or exercise, physical activity)	4.29	0.76	4.0	2-5
Sleep hygiene	4.26	0.71	4.0	2-5
Digital psychoeducation (e.g., websites)	3.81	0.93	4.0	1-5
Telephone helplines (e.g., Kids Helpline)	3.56	1.12	4.0	1-5
Digital mental health interventions (e.g.	3.44	1.05	4.0	1-5

smartphone apps, web-based programs)				
Psycho-education handouts (e.g. brochures)	3.02	1.16	3.0	1-5

Table notes: Frequency reported on 5-point Likert scale ranging from never (0) to always (5).

Several associations were observed between referral and recommendation types. Referrals to psychologists were linked only to ongoing care ($r = .37$, $p < .001$). Referrals to school counsellors were associated with headspace centres, ongoing care, and lifestyle changes ($r = .25–.28$, p < .001). Psychiatrist referrals correlated with paediatricians, CAMHS, and inpatient centres ($r = .23–.37$, $p < .001$). Referring to headspace was linked with CAMHS and allied health ($r = .21–.23$, $p < .002$). Medication recommendations were associated with CAMHS and inpatient referrals ($r = .22–.24$, $p < .002$), which may reflect variations in patient symptom severity.

GPs' knowledge of wait times for mental health specialists and services.

Only 10 GPs actively tracked wait times (5.3%) and did so by contacting providers directly or by asking patients in follow up consultations. All but one participant ($n=189$, 99.5%) reported that up-to-date information on wait times would be helpful to some extent for informing their treatment decisions (M: 4.30, SD: 0.87, Mdn: 5.0, Range: 1 to 5).

As shown in Table 2, 27–37% of GPs were unsure about wait times for public mental health services (e.g., headspace, CAMHS, inpatient centres), despite referring to them, often for severe cases. No GP characteristics were associated with knowledge of wait times for psychologists, psychiatrists, headspace, or inpatient centres ($p = .022–.707$). Knowledge of CAMHS wait times was linked to a higher adolescent mental health caseload ($r = –.24$, $p <$ .001). Wait time knowledge was generally unrelated to referral frequency, except for

inpatient centres, where uncertainty was associated with higher referral rates ($r = .24$, $p <$

.001).

GPs estimated the longest wait times for psychologists, headspace centres, and

psychiatrists, ranging from 56 to 113 days (see Table 2). Metropolitan GPs reported

significantly longer waits for psychiatrists (+39.48 days, $p < .001$) and CAMHS (+35.4 days,

$p < .002$) than regional/rural GPs; no other geographic differences were found ($p = .118$–

.378; see Table S4). Wait time estimates were unrelated to caseload factors ($p = .080$–.843).

Most GPs deemed wait times for psychologists (86.7%) and psychiatrists (89.4%)

unacceptable; over half said the same for headspace (57.9%) and CAMHS (56.4%). Across

services, GPs considered acceptable wait times to range from 6.61 to 26.10 days—

significantly shorter than their estimates ($p < .001$).

Table 2. The mean estimated wait times for the frequently referred mental health treatment and services for adolescents and the acceptability of wait times among GPs in Australia (N=190).

	Estimated wait time (days)				Acceptability ratings for wait times						Proposed acceptable wait time (days)
	M (SD)	Mdn	Range	Unsure n (%)	M (SD)	Mdn	Range	n (%) Unacceptable	n (%) Neutral	n (%) Acceptable	M (SD, range, Mdn)
Psychologist	57.26 (47.91)	45.0	5-365	12 (6.3)	1.64 (0.87)	1.0	1-5	163 (86.7)	16 (8.5)	9 (4.8)	14.66 (8.70, 1-60, 14.0)
Psychiatrist	113.22 (79.05)	90.0	4-365	15 (7.9)	1.44 (0.88)	1.0	1-5	169 (89.4)	11 (5.8)	9 (4.7)	26.1 (16.65, 1-90, 28.0)
Headspace centres	56.35 (51.10)	43.5	1-365	51 (26.8)	2.20 (1.15)	2.0	1-5	106 (57.9)	59 (32.2)	18 (9.8)	13.95 (9.15, 1-60, 14.0)
CAMHS	52.41	30.0	0-365	58	2.31	2.0	1-5	101 (56.4)	52	26 (14.5)	11.45 (11.30, 0-

	(76.71)			(30.5)	(1.23)				(27.6)	90, 7.0)
Inpatient centres	24.33 (33.51)	14.0	0-180	70 (36.8)	2.64 (1.21)	3.0	1-5	60 (38.9)	71 (46.1)	23 (14.9)	6.61 (8.88, 0-60, 3.0)

Note: *n* varies (range: 83 to 190) for each provider/service because estimations are only asked if they endorsed referring to that provider/service.

Total estimated wait times were capped at 12 months. Higher acceptability ratings indicate greater acceptability.

The impact of wait times on GPs and their patients.

Over two-thirds of GPs (70.0%, *n*=133) often or always referred elsewhere due to

long wait times, and most (82.1%, *n*=156) reported increasing their level of care. As shown in

Table 3, nearly all GPs agreed that long waits negatively affected their duty of care (96.3%),

increased adolescent feelings of abandonment (94.2%), risk of dropout (95.7%), self-harm or

suicide (90.5%), and reduced help-seeking (90.5%). In free-text responses, GPs also noted

impacts on family/carer stress (10.5%), negative perceptions of the healthcare system (7.3%),

and increased patient costs (5.2%).

Table 3. Level of agreement on the impacts of wait times on GPs and their adolescent patients (N=190).

	M (SD)	Mdn
Impact on GPs		
Increased professional responsibility and duty of care	4.74 (0.57)	5.0
Feelings of systemic neglect and abandonment	4.20 (0.91)	4.0
Increased prescribing of mental health medication	3.85 (0.97)	4.0
Impact on adolescent patients		
Increased feelings of abandonment and neglect	4.54 (0.65)	5.0
Increased treatment drop-out/discontinuation	4.47 (0.70)	5.0
Exacerbated self-harm and suicide risk	4.44 (0.73)	5.0
Reduced likelihood of future help-seeking	4.44 (0.73)	5.0
Increased social withdrawal and dysfunction	4.42 (0.64)	4.0
Delayed remission/recovery	4.42 (0.63)	4.0
Negatively impact adherence to treatment	4.41 (0.78)	4.0

Greater deterioration of symptoms or illness	4.39 (0.70)	4.0
Increased use of unhelpful coping strategies and behaviours	4.39 (0.71)	4.5
Greater chances of relapse	4.36 (0.66)	4.0
Increased school disengagement/absenteeism	4.33 (0.76)	4.0
Left feeling that their issues are not important	4.21 (0.80)	4.0
Increased rates of hospitalisation	3.95 (0.87)	4.0

Table note: Scores ranged from 1 to 5, with higher scores indicating greater levels of agreement.

GPs rated the quality of support provided to adolescents during wait times as fair (35.3%), good (50.5%), or excellent (6.3%) with a mean score of 3.53 (SD:0.79). Similar ratings were reported for care provided to parents/guardians (fair: 43.7%, good: 41.6%, excellent: 4.7%; M:3.39, SD:0.78). Nearly all GPs (95.8%) viewed it as very or extremely important to offer mental health strategies and resources during wait periods. However, most described their knowledge as minimal (8.4%), basic (37.9%), or adequate (48.4%), with only 5.3% reporting superior knowledge.

Commonly recommended interventions included ongoing GP appointments (88.9%), sleep hygiene (85.2%), exercise (84.7%), mindfulness (74.7%), and diet changes (66.8%). Over half also recommended mental health websites (67.8%), apps (55.2%), and web-based programs (50.5%). In free-text responses, GPs cited social support (21.1%) and relaxation activities (19.7%). As shown in Table 4, the frequency of interventions did not differ between routine care and wait-time care.

26 **Table 4. A comparison of the frequency of interventions recommended by GPs in**

27 **routine care for adolescent depression and/or anxiety with those recommended during**

28 **the wait time (N=190).**

Type of intervention	Routine care M (SD)	Wait time M (SD)	Wait time Mdn (Range)
Ongoing appointments with GP	4.40 (0.83)	4.33 (0.70)	4.0 (2-5)
Sleep hygiene	4.26 (0.72)	4.15 (0.72)	4.0 (2-5)
Lifestyle modification (e.g., diet and/or exercise, physical activity)	4.29 (0.76)	4.08 (0.81)	4.0 (1-5)
Digital psychoeducation (e.g. mental health websites)	3.81 (0.93)	3.92 (0.92)	4.0 (1-5)
Digital mental health interventions (e.g. smartphone apps, web-based programs)	3.44 (0.93)	3.47 (1.11)	4.0 (1-5)
Telephone helplines (e.g., Kids Helpline)	3.56 (1.12)	3.56 (1.12)	4.0 (1-5)
School counsellor/psychologist	3.32 (0.79)	3.49 (0.80)	3.0 (1-5)
Psycho-education handouts (e.g. brochures)	3.02 (1.16)	3.30 (1.15)	3.0 (1-5)
Medication prescribing (e.g. anti-depressants)	2.72 (0.66)	2.2 (0.75)	3.0 (1-5)

Table note: Scores ranged from 1 to 5, with higher scores indicating greater levels of

agreement.

When describing ideal interventions for adolescents awaiting mental health treatment,

GPs most frequently mentioned ongoing GP appointments (30.7%), digital interventions

(28.1%), lifestyle changes (14.6%), multidisciplinary support (13.5%), and regular check-ins

(13.0%).

DISCUSSION

This study explored GPs' self-reported referrals to mental health specialists and

services for adolescent depression and/or anxiety in Australia, their estimates and

acceptability of the wait times, and perceived impacts on treatment decisions. This study

confirmed that GPs routinely referred adolescents with anxiety and/depression to specialist

services with limited insight into anticipated wait times. GPs' knowledge of wait times was

not influenced by how frequently they referred patients, patient caseload, location, or

training. Given that timely access to care is a key determinant of outcomes for adolescents,

there is an urgent need to provide GPs with more precise estimates of expected wait times to

enhance their referral decision-making and patient care. Overall, GPs felt that the wait times

for almost all mental health specialists/services in Australia were unacceptable and should be

less than one month. Information-sharing from psychologists who can accept new referrals

should be prioritised, given they were the most common referral and often at capacity.

GPs agreed they were an important source of support in the wait time and had

increased their involvement in care, consistent with reports by Australian adolescents

(Subotic-Kerry et al., 2025). However, many interventions recommended by GPs during the

wait time were the same interventions recommended in routine care. This suggests that GPs

were not introducing supports specifically to address the unique needs of adolescents during

the wait period. This may contribute to the sense of abandonment and neglect reported by

youth (Subotic-Kerry et al., 2025). Digital resources warrant further investigation, as these

can reduce symptoms without increasing service demand. Improving the digital literacy of

GPs may increase the adoption of this approach to patient care during the wait time. Although

lifestyle interventions are included in clinical guidelines for depression (Malhi et al., 2021),

future research would benefit from clarifying whether GPs carefully assess the suitability of

this self-management approach for adolescents. Physical activity is often the least preferred

brief intervention among youth (Schley et al., 2019) with low-quality evidence to support its

use (Oberste et al., 2020). The increased rates of disordered eating among adolescents (John

et al., 2023) may also mean that some lifestyle modifications may be contraindicated.

Screening tools may help GPs to tailor self-management recommendations while also helping

to direct referrals more appropriately, ensuring patients are indeed waiting for the right

provider.

GPs agreed that long wait times had a negative impact on their professional

wellbeing. Specialised training in youth mental health and single session therapy (SST)

approaches may increase GPs' confidence and skills in managing the wait time. SSTs may be

especially useful for GPs looking to improve the support provided to patients in follow-up

appointments (Perreault et al., 2023). Family carers may be an underutilised support for GPs

and adolescents during the wait time, given that many parents are present at appointments

(Cronin et al., 2023). While parents have reported their own preferences for care during the

wait time (Cunningham et al., 2013), future work should endeavour to bring GPs,

adolescents, and their carers together to design a cooperative approach to wait time care that

not only meets the psychological needs of adolescents but also strengthens family support

and the capacity of GPs to provide high quality care.

Limitations

Despite extensive recruitment efforts, our sample target was not reached. While larger

than many prior studies, participants were younger and more likely to be female than national

averages, aligning with data showing female GPs manage more youth mental health (Royal

Australian College of General Practitioners, 2022). Referral patterns, wait time estimates, and

acceptability ratings likely vary with case complexity, which was not assessed. Future studies

should explore patient characteristics linked to increased risk during wait periods—such as

developmental stage, socio-economic disadvantage, and parental separation. Qualitative

methods (e.g. semi-structured interviews) could offer deeper insight into GP decision-

making.

Conclusion

Australian GPs have limited awareness of mental health service wait times but

recognise that access to this information could improve referral and treatment decisions.

While systemic efforts to reduce wait times are essential, GPs would benefit from training in

strategies to support adolescents during these delays. Further research is needed to strengthen

the supportive role GPs play in adolescent help-seeking.

Data availability statement: A deidentified dataset is available by contacting the

corresponding author bridianne.odea@flinders.edu.au.

References
99
Andrews, G., Bell, C., Boyce, P., Gale, C., Lampe, L., Marwat, O., Rapee, R., & Wilkins, G.

(2018). Royal Australian and New Zealand College of Psychiatrists clinical practice

guidelines for the treatment of panic disorder, social anxiety disorder and generalised

anxiety disorder. *Aust N Z J Psychiatry*, *52*(12), 1109-1172.

https://doi.org/10.1177/0004867418799453

Ball, W. P., Black, C., Gordon, S., Ostrovska, B., Paranjothy, S., Rasalam, A., Ritchie, D.,

Rowlands, H., Rzewuska, M., Thompson, E., Wilde, K., & Butler, J. E. (2023).

Inequalities in children's mental health care: analysis of routinely collected data on

prescribing and referrals to secondary care. *BMC Psychiatry*, *23*(1), 22.

https://doi.org/10.1186/s12888-022-04438-5

Bleustein, C., Rothschild, D. B., Valen, A., Valatis, E., Schweitzer, L., & Jones, R. (2014).
Wait times, patient satisfaction scores, and the perception of care. *American Journal of Managed Care*, *20*(5), 393-400.

Budd, M., Iqbal, A., Harding, C., Rees, E., & Bhutani, G. (2021). Mental health promotion and prevention in primary care: What should we be doing vs. what are we actually doing? *Mental Health and Prevention*, *21*. https://doi.org/10.1016/j.mhp.2020.200195

Cronin, M., McLoughlin, K., Foley, T., & McGilloway, S. (2023). Supporting family carers in general practice: a scoping review of clinical guidelines and recommendations. *BMC Primary Care*, *24*(1), 234. https://doi.org/10.1186/s12875-023-02188-1

Crouch, L., Reardon, T., Farrington, A., Glover, F., & Creswell, C. (2019). "Just keep pushing": Parents' experiences of accessing child and adolescent mental health services for child anxiety problems. *Child Care Health Dev*, *45*(4), 491-499. https://doi.org/10.1111/cch.12672

Cunningham, C. E., Chen, Y., Deal, K., Rimas, H., McGrath, P., Reid, G., Lipman, E., & Corkum, P. (2013). The Interim Service Preferences of Parents Waiting for Children's Mental Health Treatment: A Discrete Choice Conjoint Experiment. *Journal of Abnormal Child Psychology*, *41*(6), 865-877. https://doi.org/10.1007/s10802-013-9728-x

Daneshmand, R., Acharya, S., Zelek, B., Cotterill, M., & Wood, B. (2023). Changes in Children and Youth's Mental Health Presentations during COVID-19: A Study of Primary Care Practices in Northern Ontario, Canada. *Int. J. Environ. Res. Public Health*, *20*(16), 6588. https://www.mdpi.com/1660-4601/20/16/6588

De Silva, T., Prakash, A., Yarlagadda, S., Johns, M. D., Sandy, K., Hansen, V., Phelan, S., & Pit, S. (2017). General practitioners' experiences and perceptions of mild moderate depression management and factors influencing effective service delivery in rural

Australian communities: a qualitative study. *Int J Ment Health Syst*, *11*, 54.

https://doi.org/10.1186/s13033-017-0159-x

Dumesnil, H., Apostolidis, T., & Verger, P. (2018). Opinions of general practitioners about

psychotherapy and their relationships with mental health professionals in the

management of major depression: A qualitative survey. *PloS One*, *13*(1), e0190565.

https://doi.org/10.1371/journal.pone.0190565

Edbrooke-Childs, J., & Deighton, J. (2020). Problem severity and waiting times for young

people accessing mental health services. *BJPsych Open*, *6*(6), e118.

https://doi.org/10.1192/bjo.2020.103

Fleury, M.-J., Imboua, A., Aubé, D., Farand, L., & Lambert, Y. (2012). General practitioners'

management of mental disorders: A rewarding practice with considerable obstacles.

BMC Family Practice, *13*(1), 19. https://doi.org/10.1186/1471-2296-13-19

John, J. R., Khan, J. R., Lin, P. I., Jonnagaddala, J., Hu, N., Belcher, J., Liaw, S. T., Lingam,

R., & Eapen, V. (2023). A nationwide study of COVID-19 impact on mental health-

related presentations among children and adolescents to primary care practices in

Australia. *Psychiatry Research*, *326*, 115332.

https://doi.org/10.1016/j.psychres.2023.115332

Keyes, B., McCombe, G., Broughan, J., Frawley, T., Guerandel, A., Gulati, G., Kelly, B. D.,

Osborne, B., O'Connor, K., & Cullen, W. (2023). Enhancing GP care of mental health

disorders post-COVID-19: a scoping review of interventions and outcomes. *Irish*

Journal of Psychological Medicine, *40*(3), 470-486.

https://doi.org/10.1017/ipm.2022.17

Koet, L. B. M., Gerger, H., Jansen, W., Bindels, P. J. E., & de Schepper, E. I. T. (2024).

Evaluation of practice nurses' management of paediatric psychosocial problems in

general practice. *J Public Health*, *46*(2), e261-e268.

https://doi.org/10.1093/pubmed/fdae008

Kowalewski, K., McLennan, J. D., & McGrath, P. J. (2011). A preliminary investigation of

wait times for child and adolescent mental health services in Canada. *Journal of the*

Canadian Academy of Child and Adolescent Psychiatry. Journal de l'Académie

Canadienne de Psychiatrie de L'enfant et de l'Adolescent, *20*(2), 112-119.

Lawrence, S. M., Saab, M. M., Savage, E., Hegarty, J., & FitzGerald, S. (2024). Adolescents'

perspectives and experiences of accessing general practitioner services: A systematic

review. *Journal of Child Health Care*, *0*(0), 13674935241239837.

https://doi.org/10.1177/13674935241239837

MacDonald, K., Fainman-Adelman, N., Anderson, K. K., & Iyer, S. N. (2018). Pathways to

mental health services for young people: a systematic review. *Social Psychiatry and*

Psychiatric Epidemiology, *53*(10), 1005-1038. https://doi.org/10.1007/s00127-018-

1578-y

Malhi, G. S., Bell, E., Bassett, D., Boyce, P., Bryant, R., Hazell, P., Hopwood, M., Lyndon,

B., Mulder, R., Porter, R., Singh, A. B., & Murray, G. (2021). The 2020 Royal

Australian and New Zealand College of Psychiatrists clinical practice guidelines for

mood disorders. *Aust N Z J Psychiatry*, *55*(1), 7-117.

https://doi.org/10.1177/0004867420979353

McNicholas, F., Kelleher, I., Hedderman, E., Lynch, F., Healy, E., Thornton, T., Barry, E.,

Kelly, L., McDonald, J., Holmes, K., Kavanagh, G., & Migone, M. (2021). Referral

patterns for specialist child and adolescent mental health services in the Republic of

Ireland during the COVID-19 pandemic compared with 2019 and 2018. *BJPsych*

Open, *7*(3), e91. https://doi.org/10.1192/bjo.2021.48

Mental Health Commission of New South Wales. (2024). *Waited longer than acceptable for*

an appointment with a health professional. Retrieved 7th June from

https://www.nswmentalhealthcommission.com.au/measuring-change-

indicator/waited-longer-acceptable-appointment-health-professional

Oberste, M., Medele, M., Javelle, F., Lioba Wunram, H., Walter, D., Bloch, W., Bender, S.,

Fricke, O., Joisten, N., Walzik, D., Großheinrich, N., & Zimmer, P. (2020). Physical

Activity for the Treatment of Adolescent Depression: A Systematic Review and

Meta-Analysis [Systematic Review]. *Frontiers in Physiology*, *11*.

https://doi.org/10.3389/fphys.2020.00185

Paton, K., Gillam, L., Warren, H., Mulraney, M., Coghill, D., Efron, D., Sawyer, M., &

Hiscock, H. (2021). Clinicians' perceptions of the Australian Paediatric Mental

Health Service System: Problems and solutions. *Australian and New Zealand Journal*

of Psychiatry, *55*(5), 494-505. https://doi.org/10.1177/0004867420984242

Perreault, K., Breton, M., & Berbiche, D. (2023). An innovative model of psychological

service delivery in primary healthcare: the Single-Session Intervention. *BMC Primary*

Care, *24*(1), 1. https://doi.org/10.1186/s12875-022-01949-8

Petrie, K., Baldwin, P., Crawford, J., & Harvey, S. B. (2021). The voice of mental health

practice in Australia: a mixed-method cross-sectional study of gaps and areas of need.

Australian and New Zealand Journal of Public Health, *45*(4), 318-324.

https://doi.org/10.1111/1753-6405.13078

Punton, G., Dodd, A. L., & McNeill, A. (2022). 'You're on the waiting list': An interpretive

phenomenological analysis of young adults' experiences of waiting lists within mental

health services in the UK. *PloS One*, *17*(3), e0265542.

https://doi.org/10.1371/journal.pone.0265542

Royal Australian College of General Practitioners. (2022). *General Practice Health of the*

Nation 2022. https://www.racgp.org.au/getmedia/80c8bdc9-8886-4055-8a8d-

ea793b088e5a/Health-of-the-Nation.pdf

Russell, M. J., Urichuk, L., Parker, N., Agyapong, V. I. O., Rittenbach, K., Dyson, M. P., &

Hilario, C. (2024). Youth mental health care use during the COVID-19 pandemic in

Alberta, Canada: an interrupted time series, population-based study. *Child Adolesc*

Psychiatry Ment Health, *18*(1), 101. https://doi.org/10.1186/s13034-024-00785-1

Schley, C., Pace, N., Mann, R., McKenzie, C., McRoberts, A., & Parker, A. (2019). The

headspace Brief Interventions Clinic: Increasing timely access to effective treatments

for young people with early signs of mental health problems. *Early Interv Psychiatry*,

13(5), 1073-1082. https://doi.org/10.1111/eip.12729

Smith, J., Kyle, R. G., Daniel, B., & Hubbard, G. (2018). Patterns of referral and waiting

times for specialist Child and Adolescent Mental Health Services. *Child Adolesc Ment*

Health, *23*(1), 41-49. https://doi.org/10.1111/camh.12207

Steele, M., Shapiro, J., Davidson, B., Floyd, G., Johnston, J., Stretch, N., & Mohammed, A.

(2010). Survey Comparing Criteria Used by Rural and Urban Primary Care

Physicians for Referrals to Child and Adolescent Psychiatrists and Children's Mental

Health Agencies in Ontario. *Journal of the Canadian Academy of Child and*

Adolescent Psychiatry. Journal de l'Académie Canadienne de Psychiatrie de L'enfant

et de l'Adolescent, *19*(4), 284-289.

Subotic-Kerry, M., Borchard, T., Parker, B., Li, S. H., Choi, J., Long, E. V., Batterham, P. J.,

Whitton, A. E., Gokiert, A., Spencer, L., & O'Dea, B. (2025). While they wait: A

cross-sectional survey on wait times for mental health treatment for anxiety and

depression for Australian adolescents. *BMJ Open*, *15*, e087342.

https://doi.org/10.1136/bmjopen-2024-087342

Valentine, A. Z., Hall, S. S., Sayal, K., & Hall, C. L. (2024). Waiting-list interventions for

children and young people using child and adolescent mental health services: a

systematic review. *BMJ Ment Health*, *27*(1). https://doi.org/10.1136/bmjment-2023-

300844

Villas-Boas, S. B., Kaplan, S., White, J. S., & Hsia, R. Y. (2023). Adolescent Total and

Mental Health-Related Emergency Department Visits During the COVID-19

Pandemic. *JAMA Netw Open*, *6*(10), e2336463.

https://doi.org/10.1001/jamanetworkopen.2023.36463