A Framework to Assess Clinical Safety and Hallucination Rates of LLMs for Medical Text Summarisation

Authors: Elham Asgari[1,2], Saleh Khalil[1], Nina Montaña-Brown[1], Magda Dubois[1], Jasmine Balloch[1], Joshua Au Yeung[1], Dominic Pimenta[1]

Affiliation: 1-Tortus AI, London, UK, 2- Guy's and St Thomas NHS Trust, London, UK
Corresponding author: asgelham@gmail.com

Abstract

Integrating large language models (LLMs) into healthcare settings can improve workflow efficiency and patient care by automating tasks such as summarising consultations. However, ensuring the fidelity between LLM outputs and ground truth information is crucial, as errors can lead to miscommunication between patients and clinicians, resulting in incorrect diagnoses, treatment decisions and compromised patient safety. We introduce a clinician-in-the-loop framework with: 1) a clinically and technically-informed error taxonomy to classify LLM outputs, 2) an experimental structure to comprehensively and iteratively compare outputs within our LLM document generation pipeline, 3) a clinical safety framework to assess potential harms of errors in LLM outputs, and 4) an encompassing graphical user interface (GUI), CREOLA, to perform and assess all previous steps. Our clinical error metrics were derived from 18 experimental configurations involving LLMs for clinical note generation consisting of 49,590 transcript and 12,999 clinical note sentences. Overall, we observed a 1.47% hallucination rate (44% rated 'major') and a 3.45% omission rate (17% 'major'). Through iterative prompts and workflow refinements, we reduced major errors below previously reported human note-taking error rates, underscoring the potential of our framework to enable safer clinical documentation.

Introduction

One of the most appealing applications of LLMs in healthcare is for administrative tasks [1]. Clinicians devote a substantial amount of time to documentation [2], and prolonged interaction with electronic health records, where clinical documentation is logged, has been demonstrated to raise cognitive load and lead to burnout [3]. In fact, the use of LLMs for clinical documentation, especially clinical note generation [4] or consultation summarisation [5,6], is an active area of research.

However, LLMs are known to produce errors in many settings, from document summarisation [7], to general reasoning tasks as well as more clinically relevant tasks [8]. These errors can be categorised as "hallucinations" [9]: known as an event where LLMs generate information that is not present in the input data, or omissions: the event where LLMs miss relevant information from the original document. Errors in clinical documentation generation can lead to inaccurate recording and communication of facts [10,11]. Inaccuracies in the document summarisation task can introduce misleading details [8] into transcribed conversations or summaries, potentially delaying diagnoses [12] and causing unnecessary patient anxiety.

The problem of hallucinations poses a significant challenge to date [1,13]. The occurrence of hallucinations has previously been attributed to the data quality during model training [14,15], the type of model training methodology [16] and prompting strategies [17].

Recent work has established that hallucination may be an intrinsic, theoretical property of all LLMs [9]. Consequently, there is a growing body of work focused on the technical evaluation of LLM accuracy and the detection and mitigation of hallucinations in LLMs [18]. However, the prevalence, causation, and evaluation of hallucinations in a clinical context, as well as their subsequent impact on clinical safety, remains an open question.

Clinical documentation can be variable in quality [19,20], and studies estimate that human-generated clinical notes have, on average, at least 1 error and 4 omissions [21]. Given the increased usage of LLMs for clinical documentation [22,23], several methods have been proposed for evaluating clinical documentation generated using LLMs.

Relevant clinical evaluation frameworks typically include categorising clinical errors for downstream analysis. Typically, these differ from traditional natural language processing (NLP) taxonomies [16], which have separated hallucination types into distinct categories, for example, into "intrinsic" and "extrinsic" [24], "factuality" and "faithfulness" [16], "factual mirage" and "silver lining" [25] errors. The differences between general and clinical taxonomies arise from the necessity of increased granularity of clinical error types, which are not captured by the broader, general methods.

For example, Tierney et al. [26] propose using a modified version of the Physician Documentation Quality Instrument-9, accounting for hallucinations and bias, while Abacha et al. [23] propose evaluating clinical note quality using automated metrics. However, these relevant clinical categorisations have not assessed the implications of the mistakes for downstream harm.

Automated metrics, such as Recall-oriented Understudy for Gisting Evaluation (ROUGE)[27], Bilingual Evaluation Understudy (BLEU) [28] and Bidirectional Encoder Representations from Transformers (BERT)-score [29], while useful for comparing model-generated text with expert-written examples, exhibit significant limitations when applied to the evaluation of healthcare-related content. These metrics, primarily focused on surface-level textual similarity, fail to capture the semantic nuances, contextual dependencies, and domain-specific knowledge crucial for accurate medical discourse [30]. This deficiency is particularly problematic in healthcare settings, where understanding complex medical concepts (e.g., symptoms, diagnoses, treatments) and their interrelationships is paramount for patient well-being and effective decision-making.

Despite the exponential growth in benchmarks for model reasoning abilities [31], the evaluation of LLMs on clinical tasks has typically been carried out via "question-answering" (QA) benchmarks [5,8,32]. These tasks assess models' accuracy over various clinical questions, typically derived from licensing exams. While these methods offer insights into the factual knowledge and reasoning abilities of LLMs, they do not assess clinical or medical capabilities such as medical text summarisation.

Singhal et al. [33] have outlined the challenges of evaluating LLMs in various medical contexts, including medical exams, research and consumer queries. They have proposed a human evaluation model for the answers provided by different LLMs that checks on factuality, precision, possible harm and bias. Other evaluation factors such as fairness, transparency, trustworthiness and accountability have been suggested in using LLMs in healthcare [34]. In a more recent study, Tang et al assessed human evaluation based on metrics such as coherence, factual consistency, comprehensiveness and potential harm. Interestingly, they assessed the clinician's preference for different outputs [35]. More recently, Tam et al [36] have introduced QUEST as a framework for human evaluation of LLMs in healthcare following a comprehensive literature review on the topic. QUEST includes five principles for human evaluation of LLMs, including Quality of information, Understanding and reasoning, Expression style and persona, Safety and harm, and Trust and confidence.

Multiple benchmarks have been proposed to evaluate model summarisation capabilities in the biomedical domain, including over biomedical literature [37–40], medical forum conversations [41], and radiology reports [22,42]. However, these benchmarks do not capture the nuances of patient-facing clinical interactions, where LLM-documentation holds most promise.

Recently, Umapathi et al. [43] have assessed models' tendency towards hallucination. They reported that LLMs were significantly variable in their accuracy depending on the prompts used. However, the MedHALT benchmark is limited to assessing LLM's reasoning capabilities over the medical domain in a QA format. Most relevantly, Moramarco et al. [21] benchmark BART models on the PriMock dataset and find that they produce 3.9 errors and 6.6 omissions on average per note. However, they did not assess the model's impact or human errors on patient safety as part of their study.

This study aims to contribute to the ongoing effort to ensure clinical safety in using LLMs for note generation by introducing a framework which has four components: 1) a clinically and technically-informed error taxonomy to classify LLM outputs, 2) an experiment structure to comprehensively and iteratively compare outputs within our LLM document generation pipeline, 3) a clinical safety framework to assess potential harms of errors in LLM outputs, and 4) an encompassing graphical user interface (GUI), CREOLA, to perform and assess all previous steps. Figure 1 shows our workflow based on the framework. We present our findings and insights from applying our framework, which, to our knowledge, represents the largest manual evaluation of LLM clinical note generation to date.

Our objective is to promote the efficient, reliable, and confident use of LLMs for clinical documentation, thus supporting healthcare providers in delivering high-quality care and overall reducing the administrative workload for clinicians.

Figure 1: Our workflow for the assessment of LLM output using CREOLA platform

This diagram illustrates the process we followed using the available dataset for various experiments, including the input from clinicians for labelling and consolidating reviews, followed by a safety analysis.

Our objective is to promote the efficient, reliable, and confident use of LLMs for clinical documentation, thus supporting healthcare providers in delivering high-quality care and overall reducing the administrative workload for clinicians.

Results

Dataset

We conducted a series of 18 experiments, each consisting of 25 primary care consultation transcripts from the PriMock dataset [44] For each transcript, we generated paired clinical documentation using an LLM, resulting in 450 consultation transcript-note pairs. This was a total of 49,590 transcript sentences and 12,999 clinical note sentences that were manually evaluated and labelled for any hallucinations or omissions.

Experiments

Our experiments were guided by our framework, which provides a systematic approach to evaluate LLM outputs quantitatively. Using a baseline LLM prompt and workflow, we generated 25 transcript-note pairs. We recruited 50 medical doctors for manual evaluation. For each paired transcript-note, we had two clinician reviewers evaluate each sentence in the clinical note to ensure it was evidenced in the transcript; sentences that were not evidenced were labelled as hallucinations. We also highlighted each sentence in the transcript for review, checking if it was present in the output note if it was clinically relevant; and if not, they were labelled as an omission. If the hallucination or omission could change the diagnosis or management of the patient (if left uncorrected), it was marked as 'Major', otherwise, they would be labelled as 'minor'. In cases of discrepancy between the two reviewers, consolidation was performed by a senior clinician with over 20 years of clinical experience.

We additionally identified the specific sections of the notes where the hallucinations occur (main history, examination, discussion, symptoms assessment, and plan). The result of each experiment informed our subsequent experiment approach to analyse how prompt/ workflow/ engineering changes affect the hallucinatory potential for clinical note generation.

All experiments were conducted on CREOLA, our in-house platform, designed to allow clinicians to identify and label relevant hallucinations and omissions in clinical text. Using this platform, we were able to implement our framework to quantify and track changes in our prompts and model configurations and iteratively modify our approach to ensure the safe integration of LLM-generated summaries into clinical practice.

Hallucinations

Of 12,999 sentences in 450 clinical notes, 191 sentences had hallucinations (1.47%), of which 84 sentences (44%) were major (could impact patient diagnosis and management if left uncorrected). Of the hallucination types, 82 (43%) were fabricated, 56 (30%) were negations, 33 (17%) were contextual, and 20 (10%) were related to causality.

Major hallucinations occurred in all sections, but most commonly in Plan (21%), Assessment (10.5%), and Symptoms (5.2%) sections. The most common hallucination type were fabrications and primarily appeared in the planning section of the clinic note (Figure 2). Examples of the various hallucination types are available in supplementary materials.

Figure 2. Incidence of hallucinations, their types, the section of the note they appear in, and clinical risk.

The figure illustrates the occurrence of hallucinations in the generated sentences based on different prompts (a) and their corresponding percentages (b). It also highlights the type and clinical severity of hallucinations (c) and the specific sections of the note where they appeared (d).

Omissions

Of the 49,590 sentences from our consultation transcripts, 1,712 sentences were omitted (3.45%), of which 286 (16.7%) of which were classified as major and 1,426 (83.3%) as minor. Figure 3 shows the number of omissions and the percentage of omitted sentences based on different prompts.

Figure 3: Number of omitted sentences based on different prompts

The figure illustrates the number of omitted sentences using different prompts (a) and their respective percentage (b).

Grading Hallucinations and Omissions by Clinical Safety Impact

Inspired by protocols in medical device certifications, we applied the clinical safety assessment of our framework described in the methods section. We classified the clinical risk (Major or Minor) and evaluated the risk severity of all identified major hallucinations as depicted in Figure 4. We also determined the risk of hallucinations based on their type and where in the sentence they occurred (Figure 5).

Figure 4: Severity of risk in major hallucinations

We assessed the clinical risk resulting from major hallucinations based on our suggested framework inspired by protocols in medical device certifications.

Figure 5: Hallucination risk assessment

We have categorised the clinical severity of hallucination risks based on their type (a) and the section of the clinical notes where they occurred (b).

We conducted the risk assessment on the omissions (Figure 6 A) and detailed the note section that would likely have been affected (Figure 6 B). Major omissions were most common in current issues, followed by PMFS, and Info and Plan sections (54.5%. 35.0%, 10.5% respectively).

Figure 6: Severity of clinical risk for major omissions and the section of the clinical note where they most occurred

This figure illustrates the clinical risk severity due to major omissions (a) and indicates the sections of the notes where they are most likely to occur (b).

Examples of hallucinations and omissions are available in supplementary materials.

Iterative experiments can significantly reduce hallucination and omission rates in LLM-generated clinical notes

Through a series of 18 iterative experiments, we tested a combination of prompting and workflow strategies, including structured prompting, atomisation, function calls and JSON-based outputs, an additional LLM revision step, and templating (SOAP -Subjective, Objective, Assessment, plan- note), which are explained in more detail in the methods section.

Modifying the prompt from the baseline used in Experiment 1 to include a style update used in Experiment 8, resulted in a reduction of both major and minor omissions. Although there was a slight increase in hallucinations in Experiment 8, these were mostly minor. Figure 7 illustrates the number of hallucinations and omissions recorded in Experiments 1 and 8.

Figure 7. A comparison of omissions and hallucinations between experiment 1 and experiment 8. Comparing omissions (a) and hallucinations (b) using the base prompt from experiment 1 and the updated style prompt from experiment 8.

Each connected dot represents the change in hallucinations or omissions for a given document.

We then compared the outputs using various structured prompts in experiments 3 and 8 illustrated in Table 1.

Experiment 3	Experiment 8
You are a highly accurate medical office assistant drafting documentation for a physician. Every decision you take is life or death and must be 100% accurate. DO NOT ADD any content that isn't specifically mentioned IN THE TRANSCRIPT. From the attached transcript generate a clinical note based on the below template format for the physician to review, include all the relevant information and do not include any information that isn't explicitly mentioned in the transcript. If nothing is mentioned just return [NOT MENTIONED]. It is vital that all the information in the note is as accurate as possible. Avoid repeating the same information in different sections where possible. Write the note from the perspective of the physician. DO NOT add associate or relate causes for medical conditions unless explicitly specified by the Physician. See below for a template to outline the structure of the output and style preferences to follow.	You are a medical office assistant drafting documentation for a physician. DO NOT ADD any content that isn't specifically mentioned IN THE TRANSCRIPT. From the attached transcript generate a SOAP note based on the below template format for the physician to review, include all the relevant information and do not include any information that isn't explicitly mentioned in the transcript. If nothing is mentioned just return [NOT MENTIONED]. It is VITAL that all the information in the note is as accurate as possible. Avoid repeating the same information in different sections where possible. Write the note from the perspective of the physician. Only include any section of the template if there is information from the transcript, otherwise omit it.
Template: Referral Reason / reason for appointment: History - Allergies - Medications - History of presenting complaint - Past Medical History - Family/Social History - Sensitive information Observations: - Examination findings - Investigation results - Impression or clinical assessment Plan: - Planned investigations - Follow up - New prescribed medication or therapies - Communication, reassurance & patient understanding of care - Actions for referrer/GP Clarity: - Explained medical terms	Template for Clinical SOAP Note Format: Subjective: - HPI: [include here any mentioned symptoms, chronological narrative of patients complaints, information obtained from other sources(always identify source if not the patient).] - Past medical history. [include here all of the patients past conditions, treatments and encounters, also include relevant social history here including smoking, alcohol, drug use and occupation/travel history] - Review of systems [include here any additioinal symptoms in other organs that is relevant to the initial presentation] - Current medications [list medicines out each on a seperate line, in a standard format where the information is mentioned: [DRUG NAME][DRUG DOSE][DRUG FREQUENCY][INDICATION]] Objective: - Vital signs [including any mentioned blood pressure, pulse rate, oxygen saturation, temperature] - Physical exam [the examination findings from the physical exam, if mentioned] - Test Results [include in this section any lab test results or imaging reports] Assessment / Problem List: - Assessment: [A one sentence description of the patient and major problem as described by the physician, including the diagnosis the physician has identified] - Problem list: [A numerical list of clinical problems arising from this encounter and active ongoing medical problems the patient has. Present each problem as [Condition][Status:active/suspected/confirmed/past/unknown], list each problem on a separate line, leave status as unknown if not mentioned in the transcript] Plan: [include here any management plan mentioned in the transcript, including patient education, prescriptions, tests, referrals or other plans.]
Style preferences: - Write from the perspective of the physician (first person) - Be ultra concise - Use bullet points and broken sentences	Please adhere to the following style guidelines: - Write from the perspective of the physician (first person) - Be ultra-concise - Be ultra-precise, do not use generalising terms - Be highly detailed - Include ALL important negations in the relevant sections (e.g. the patient has no fever) the clinician has elicited as well as all positive findings. - Use bullet points and single words, not sentences. - Always list medications in a list in the following format for each one: medicine, dose, frequency, indication - Always document if drug allergies are present or not - Examination findings always refer to a physical exam, only include signs here, not symptoms - Preserve quantities if mentioned in the text

Table 1: Prompt changes that led to decreased hallucinations and omissions

The change in the prompt from experiments 3 to 8 reduced the incidence of major hallucinations by 75% (from 4 to 1), major omissions by 58% (24 to 10), and minor omissions by 35% (114 to 74) (Figure 8). By following structured prompting, including a style update and instructing the model to output the status "unknown" for instances where information was missing from the transcript, we significantly improved performance.

Figure 8: Comparison of hallucination and omission counts between two experiments, assessing differences in prompt engineering effect on the quality of outputs

The figure shows the changes in the counts of hallucinations (a) and omissions (b) between experiments 3 and 8 which have resulted from the changes in the prompts.

In a subsequent experiment (experiment 5), we found that incorporating a chain-of-thought prompt (Table 1, supplementary material), to extract facts from the transcript - a process referred to as atomisation - before generating the clinical note, led to an increase in major hallucinations and omissions. Figure 9 shows the comparison between omissions and hallucinations between this experiment and our base Experiment (Experiment 1).

Figure 9. A comparison of omissions and hallucinations between experiment 1 and experiment 5.

Comparing omissions (a) and hallucinations (b) in our base experiment (experiment 1) with experiment 5 where the transcript was broken down into concise facts before being passed to the LLM for note generation. Each connected dot represents the change in hallucinations or omissions for a given document.

We also compared the output of experiment 5 to experiment 3, which used structured prompts. We found that major hallucinations increased from 4 to 25, minor hallucinations from 5 to 29, major omissions from 24 to 47, and minor omissions from 114 to 188 (Fig 10). This result precluded the new

change from being evaluated for clinical safety, as the increase in hallucinations and omissions was considered too large to be considered useful.

Figure 10: Comparison of hallucination and omission counts between two experiments, assessing the efficacy of a data-extraction intermediate step versus a normal note-generation step.

We found that using prompts with function calling was useful in ensuring that the outputs adhered to a specific structure required for different electronic health records. Utilising our framework, we iteratively improved the performance of the structured notes across several experiments (6, 9, 10, and 11). From the first to the last iteration (experiments 6 to 11), we made meaningful improvements to the prompts, including instructions on adherence to subheadings and the addition of a writing style guidance (e.g., a list of writing rules to follow). Table 2 shows the prompts used in the four function call experiments.

Eperiment 6	Experiment 9
Please use the function below to generate a customised output. provide your output in json format. For each parameter value you provide, make sure to include all properties defined in the schema. If a parameter is an array please try to separate ideas into separate items. You are a highly accurate medical officer drafting documentation for a physician. You will receive a transcript of a medical consultation between a patient and a clinician. Your task is to identify the patient's key problems in that encounter, and then extract information provided to you in the format outlined in a JSON schema for each individual problem. A problem is a single discrete issue for the patient, encompassing presenting complaint, associated symptoms, and relevant history - e.g. shortness of breath, needs new housing, recent bereavement etc. It is VITAL that you include all properties mentioned in the schema, if there is a field that is not mentioned in the transcript just write [NOT MENTIONED]. Only include information strictly mentioned in the transcript. Failure to do so may cause harm to the patient. DO NOT duplicate information in more than one problem. Aim to capture all problems mentioned in the transcript into their own array items.	Please use the function below to generate a customised output. provide your output in json format. For each parameter value you provide, make sure to include all properties defined in the schema. If a parameter is an array please try to separate ideas into separate items. You are a highly accurate medical officer drafting documentation for a physician. You will receive a transcript of a medical consultation between a patient and a clinician. Your task is to identify the patient's key problems in that encounter, and then extract information provided to you in the format outlined in a JSON schema for each individual problem. A problem is a single discrete issue for the patient, encompassing presenting complaint, associated symptoms, and relevant history - e.g. shortness of breath, needs new housing, recent bereavement etc. It is VITAL that you include all properties mentioned in the schema, if there is a field that is not mentioned in the transcript just write [NOT MENTIONED]. Only include information strictly mentioned in the transcript. Failure to do so may cause harm to the patient. DO NOT duplicate information in more than one problem. For each parameter you provide in the tool call, please adhere to the following style guidelines: - Write from the perspective of the physician (first person) - Be ultra-concise - Be ultra-precise, do not use generalising terms - Be highly detailed - Include ALL important negations in the relevant sections (e.g. the patient has no fever) the clinician has elicited as well as all positive findings - Use bullet points and single words, not sentences - Always list medications in a list in the following format for each one: medicine, dose, frequency, indication - Always document if drug allergies are present or not - Examination findings always refer to a physical exam, only include signs here, not symptoms - Preserve quantities if mentioned in the text - Avoid repeating the same information in different sections where possible
Experiment 10	**Experiment 11**
Please use the function below to generate a customised output. provide your output in json format. For each parameter value you provide, make sure to include all properties defined in the schema. If a parameter is an array please try to separate ideas into separate items. You are a highly accurate medical officer drafting documentation for a physician. You will receive a transcript of a medical consultation between a patient and a clinician. Your task is to identify the patients key problems in that encounter, and then extract information provided to you in the format outlined in a JSON schema for each individual problem. A problem is a single discrete issue for the patient, encompassing presenting complaint, associated symptoms, and relevant history - e.g. shortness of breath, needs new housing, recent bereavement etc. It is VITAL that you include all properties mentioned in the schema, if there is a field that is not mentioned in the transcript just write [NOT MENTIONED]. Only include information strictly mentioned in the transcript. Failure to do so may cause harm to the patient. DO NOT duplicate information in more than one problem. For each parameter you provide in the tool call, please adhere to the following style guidelines: - Write from the perspective of the physician (first person) - Be ultra-concise - Be ultra-precise, do not use generalising terms - Be highly detailed - Include ALL important negations in the relevant sections (e.g. the patient has no fever) the clinician has elicited as well as all positive findings - Use bullet points and single words, not sentences - Always list medications in a list in the following format for each one: medicine, dose, frequency, indication - Always document if drug allergies are present or not - Examination findings always refer to a physical exam, only include signs here, not symptoms - Preserve quantities if mentioned in the text - Avoid repeating the same information in different sections where possible	Please use the function below to generate a customised output. provide your output in json format. For each parameter value you provide, make sure to include all properties defined in the schema. If a parameter is an array please try to separate ideas into separate items. You are a highly accurate medical officer drafting documentation for a physician. You will receive a transcript of a medical consultation between a patient and a clinician. Your task is to identify the patients key clinical problems (the presenting complaint or complaints) in that encounter, and then extract information provided to you in the format outlined in a JSON schema for each individual problem. A problem is a single discrete issue for the patient, encompassing presenting complaint, associated symptoms, and relevant history - e.g. shortness of breath, needs new housing, recent bereavement etc. It is VITAL that you include all properties mentioned in the schema, if there is a field that is not mentioned in the transcript just write [NOT MENTIONED]. Only include information strictly mentioned in the transcript. Failure to do so may cause harm to the patient. DO NOT duplicate information in more than one problem. For each parameter you provide in the tool call, please adhere to the following style guidelines: - Write from the perspective of the physician (first person) - Be ultra-concise - Be ultra-precise, do not use generalising terms - Be highly detailed - Include ALL important negations in the relevant sections (e.g. the patient has no fever) the clinician has elicited as well as all positive findings. - Use bullet points and single words, not sentences. - Always list medications in a list in the following format for each one: medicine, dose, frequency, indication - Always document if drug allergies are present or not - Examination findings always refer to a physical exam, only include signs here, not symptoms - Preserve quantities if mentioned in the text - Avoid repeating the same information in different sections where possible. - Ensure the output only has the headings Description of Problem, History, Examination and Comments - and no other headings

Table 2. Prompts used in experiments 6, 9, 10 and 11

As a result of these changes, we eliminated major omissions completely, decreasing them from 61 to 0, and reduced minor omissions by 58%, from 130 to 54. Additionally, we lowered the total number of hallucinations by 25%, reducing them from 4 to 3 (Figure 11).

Figure 11: number of hallucinations and omissions in the function call experiments.

This figure illustrates how the change in prompts within the function call experiments can change the number of major and minor hallucinations and omissions in the output

We then examined the two best-performing experiments with the fewest hallucinations and omissions (experiments 8 and 11) for the type of hallucinations they produced and where they were more likely to appear in the sentence (Figure 12).

Figure 12: Clinical risk assessment for hallucinations in experiments 8 and 11

The figures illustrate the type and severity of clinical risk resulting from major hallucinations in Experiments 8 (a) and 11 (c) and the section of the notes they occur (b and d).

Experiment 11 did not have any major omissions, and the risk assessment of major omissions for experiment 8 is shown in Figure 13.

Figure 13: Assessment of clinical risk resulting from major omissions

The figure illustrates the clinical risk from major omissions in experiment 8 and the section of the clinical notes it has occurred.

Discussion

Our study supported that hallucinations and omissions may be intrinsic theoretical properties of current LLMs [9]. LLMs can output unfactual or unfaithful text with high degrees of confidence [45] which can be particularly dangerous in a high-stakes environment such as healthcare. Our framework quantifies the clinical impact and implications that LLM omissions and hallucinations may lead to if unchecked or uncorrected; only then can clinical safety be meaningfully addressed. Once LLM errors are identified and quantified, we can make iterations on LLM prompts, workflows or engineering design to reduce or eliminate these errors. By concentrating on minimising errors that could significantly impact patient care, we can align LLMs with clinical safety standards and regulations.

a, we have conducted the largest manual evaluation on the task of LLM clinical note generation to date. To implement our framework at scale we built CREOLA, an in-house platform, designed to enable clinicians to identify and label relevant hallucinations and omissions in clinical text and inform future experiments. However, similar publicly available platforms (e.g. labelbox, explosion AI, autoblocks etc.) may be used for this task. Experiments within CREOLA can validate or discredit architectures and prompt approaches in a safe sandbox environment before clinical deployment.

Our experiment results show that omissions are more common than hallucinations (3.45% to 1.47%, respectively). However, hallucinations were much more likely to be classified as a "major" error compared to omissions (44% to 16.7%, respectively). This means that hallucinations are more likely to lead to downstream harm and impact clinical care if left uncorrected. Hallucinations occurred in all

sections, but most commonly (20%) in the Plan section of the clinical note. This is an important finding as this section often contains direct instructions or actions to colleagues or patients that can impact clinical safety. The most concerning hallucinations were the negation type (30% of total hallucinations). These mostly appeared in the planning section and contradicted what was said during the consultation. Negation hallucinations can lead to significant confusion and harm, and without the full context of the consultation, readers may struggle to discern which negation is true and which is false.

By designing prompts that addressed specific aspects of the notes (base, template and style), we were able to focus our iterations to achieve the best results. However, it is important to note that summarisation tasks require the ablation of certain data from the original text to make it a concise, relevant, and useful summary artefact. Optimal omission rates depend significantly on the context, the quality of input and the reviewer receiving output. Our framework allows us to focus on the clinical impact of omissions and hallucinations quantitatively. Overall, our hallucination rates are similar to those reported in the literature for generalist tasks [46]. For the clinical summarisation task, experiment 8 achieved 1 major hallucination and 10 major omissions, whilst experiment 11 achieved 2 major hallucinations and 0 major omissions over 25 notes. These results are highly encouraging, as our iterative experiment process has resulted in fewer errors per note than those reported in the clinical literature. Moramarco et al. [21] reported 3.9 errors and 6.6 omissions per note as produced by a BART model and 1 error and 4 omissions per human-written note. This improvement is likely due to the large parameter sizes of modern large language models in combination with our framework. Although this rate is subject to change depending on the text and experiment, our results suggest that we can achieve state-of-the-art, sub-human clinical error rates by carefully engineering and subsequently validating LLMs to produce safe outputs.

Our study is limited in several ways. Firstly, the sample size of medical transcripts used was relatively small; the sample size was chosen to balance the trade-off of annotation volume required for the comparison of different experiments against sample size and number of experiments performed. Additionally, we only evaluated one LLM (GPT-4), selected due to its established performance in text summarisation at the time of our experiments. The ongoing development and enhancement of open-source large language models (LLMs) are likely to boost their application in medicine [47]. Recently, Zhang et al. reviewed how fine-tuning open-source LLMs such as PRIMERA, LongT5, and Llama-2 can enhance their ability to summarise medical evidence effectively [48]. Furthermore, our experiments use a direct prompting scheme (Supplementary data 1). Newer methods such as (but not limited to) Retrieval-Augmented Generation (RAG) [49], Chain of Thought (CoT) [50], or the use of knowledge graphs [51] have recently been used to enhance the performance of LLMs. For example, by equipping LLMs with domain-specific knowledge, RAG enables the models to generate more precise and pertinent results [52,53], whilst CoT generally enhances model reasoning abilities. A straightforward extension of this work is using this framework over different experimental configurations, such as using different models or prompting techniques, and comparing the impact on reported performance to clinical safety metrics.

Finally, using human annotators to evaluate large amounts of data is expensive and unsustainable. In the long run, the automated evaluation of model output [54] is a consequential future direction which will enable the scalable assessment of a larger volume of information, with clinicians remaining in the loop by "supervising" evaluator models via the inspection of a sub-sample of the outputs. 'LLM-as-a-Judge', which refers to an LLM tasked with evaluating, scoring, or assessing the quality, correctness, or appropriateness of outputs, has recently been described [55] and its applicability has been discussed [56]. Utilising the capabilities of LLMs for initial screening can significantly reduce the time and resource demands on human evaluators. This can make the CREOLA framework more scalable and efficient over time. This hybrid approach aligns with ongoing advancements in AI and has the potential to maintain rigorous oversight while ensuring scalability.

In this work, we present a framework for the clinical safety assessment of LLMs in clinical documentation scenarios. Using the CREOLA platform, we analyse the impact of prompting techniques on the safety of LLM outputs. Our iterative modification process allows us to reach new low hallucination and omission rates - our best-performing experiments outperform previously reported model and human error rates - facilitating confident deployment of our solutions to end clinical users. Additionally, CREOLA provides a sandbox environment which buffers users and patients from harm in case the iteration leads to higher clinical error rates. The addition of clinical safety assessment to prompt evaluation creates a valuable framework for implementing note summarisation tools in clinical practice. We propose that our suggested framework, which combines the assessment of hallucinations and omissions with an evaluation of their impact on clinical safety, can serve as a governance and clinical safety assessment template for various organisations. This approach aims to empower clinicians to become key stakeholders in the deployment of large LLMs in clinical settings.

Methods

We propose a multi-component framework to evaluate hallucinations and omissions in clinical documentation generated by LLMs. Central to our approach is the concept of "clinician in the loop". Given their expertise, clinicians are uniquely positioned to identify clinical errors made by the models, making their involvement essential. A specialised annotation platform (CREOLA) was developed to facilitate clinician labelling for each experimental dataset.

Experimental design: Our experiments systematically assessed how various prompting techniques and workflow structures influenced the accuracy and reliability of clinical notes derived from primary care consultation transcripts. Typical parameters varied in our experiments included the complexity and specificity of prompts (such as the addition of structured sections, negations, or perspective changes), as well as the number of LLM calls, such as introducing an additional revision step through a secondary LLM call. For consistency and reproducibility, we used OpenAI's GPT-4 (GPT-4-32k-0613), setting the seed to 210, temperature to 0, and a top-p value of 0.95 to accommodate clinical language complexities.

All prompts used are detailed in Supplementary Materials, Table 4. To achieve a meaningful clinical comparison of efficacy and safety in a data-driven way, our framework relies on the definition of a 'baseline' experiment against which to compare results. The baseline experiment must use the same input data points as the new experiment. To clearly attribute an experiment's results to a specific change, we aim only to alter one parameter from the baseline experiment configuration at a time.

We used different methodologies to assess and improve model output as described below:

Model Improvements: This outlines modifications to individual LLM calls within our workflow while preserving the same overall structure. Common modifications include the prompt, the model used for the call, or the model hyperparameters, such as maximum output tokens (Table 1).

Workflow Improvements: We implemented changes to explore new methods for generating a specific type of output. For instance, in our clinical note generation based on a transcript, we decided to extract a list of facts from the transcript before making a single call to the LLM for the final note (Supplementary Table 1). We then evaluated how this approach affected the frequency of hallucinations and omissions (Figures 9 and 10). Additionally, we included an extra LLM call in some experiments to improve the quality of the output (Supplementary Figure 1).

Clinician vs LLM generated notes: Several members of our clinical team were tasked with creating notes based on consultation transcripts. We then utilised the framework to identify any hallucinations

and omissions in these notes, which allowed us to compare the clinician-created notes with those generated by the language model. Results shown in supplementary Figure 3, Experiment 17.

Summary of Experimental Approaches in LLM-Based Note Summarisation

Our study evaluated various approaches to prompt design and output structuring for LLM-based clinical notes. The experiments were designed to iteratively refine the model's ability to produce accurate, structured, and clinically relevant summaries. The key methodological approaches across our 18 experiments are summarised below:

Baseline Prompts (Experiments 1 & 2): These were the initial prompts we had in our product prior to adopting the framework. We used these as a benchmark against which the later prompts were compared.

Structured Prompts (Experiments 3, 7, 8, and 4): Prompts were organised into three components: base (context and goal setting), template (content and structure), and style (formatting). Experiment 3 was a customisation experiment to test a new prompt structure (base, template, style preferences) for generating custom notes based on a transcript. Experiment 7 introduced a first-person perspective in generated notes. Experiment 8 refined the style section by incorporating negations and an "unknown" category for problems not explicitly mentioned in transcripts. Experiment 4 tested an enhancement to the baseline SOAP (Subjective, Objective, Assessment, Plan) note by improving medication record representation.

Atomisation (Experiment 5): This method used a chain-of-thought prompt to extract atomic facts from transcripts to ensure the precise organisation of clinical details. The approach facilitated structured extraction, breaking down information into fundamental components.

Function Calls & JSON-Based Output (Experiments 6, 9, 10, 11): LLMs were instructed to generate responses in structured JSON format instead of free text. This structured format was optimised for integration with primary care electronic health record systems. Successive experiments refined style handling, negation accuracy, and clinical specificity, progressively reducing hallucinations.

Structured Prompt + LLM Revision Step (Experiments 14 & 15): We added a second LLM pass to review and refine outputs based on structured prompting. Experiment 14 built on Experiment 11 with a revision step to improve SOAP notes and introduce an "unknown" option for missing details. Experiment 15 applied this process to a 'Bad SOAP' note, containing hallucinations and omissions, to evaluate how well errors could be mitigated.

New Note Generation Approach (Experiment 16): A novel template-driven method was introduced for generating customised outputs. However, comparison with baseline results (Experiment 8) revealed an increase in major hallucinations and minor omissions, highlighting potential trade-offs.

Clinician vs. LLM Comparison (Experiment 17): In this experiment, clinicians manually created notes based on medical transcripts. These notes were then assessed for hallucinations and omissions, providing a benchmark against LLM-generated content. Interestingly, findings suggested slightly more hallucinations in clinician-written notes but fewer omissions, highlighting key differences in human vs. LLM-generated summaries.

Experiment 18: In this experiment, we assessed the performance of the notes within the publicly available ACI Bench dataset.

Hallucination and Omission Taxonomy: We follow the conventional AI literature and taxonomise LLM errors into two types 1) hallucinations, which are instances of text unsupported by the associated clinical documentation, and 2) omissions [21], which are instances where relevant details are missed in the supporting evidence. Furthermore, inspired by protocols in medical device certifications [57,58], we categorise errors as either 'major' or 'minor', where major errors can impact on the diagnosis or the management of the patient if not corrected.

To make our categories more granular, we propose to divide hallucinations into four categories: 1) fabrication, occurring when the model produced information that was not evidenced in the text, 2) negation, occurring when the model output negates a clinically relevant fact, 3) causality, occurring when a model speculates the cause of a given condition without explicit support from the text, and 4) contextual, occurring when the model mixes topics otherwise not related to the given context.

In the case of omissions, we further divide them into sections: 1) current issues, occurring when details about the current presentation were omitted, 2) PMFS (past medical history, medication history, family and social history), occurring when details about the past medical history, medications including allergies, family and social history, including drinking and smoking, were omitted, and 3) information and plan: when discussions and explanations of the condition and management plans were omitted. Examples of each of the sub-categories are provided in the Supplementary Materials.

Experimental Structure and Annotation Protocol: Here, we define a process to assess how model parameters affect the model outputs and clinical safety. To do this, we define "experiments", which are parametrised by: 1) the number of data points processed by the LLM, 2) the type of data the LLM will ingest, 3) the model configuration (type of model, random seed, temperature,...), 4) the prompt used to obtain an LLM output, and 5) the number of clinicians which must review the data point for clinical errors.

Given an experiment configuration, we extract model outputs from the input data and store the results in a database with the associated experiment metadata. We task annotators to classify whether given sub-sections of the output contain hallucinations or omissions according to our taxonomy, and explain in free text the reason for classification. The annotators were volunteer doctors who were paid £5 per note for annotations. Recognising the subjectivity inherent in annotation, we require annotation by at least two clinicians for each input-output pair. This step is followed by a consolidation step, i.e. a detailed review by our internal team of senior clinicians, ensuring a consistent evaluation of all annotations.

Clinical Safety Assessment: Recognising that safety assessment is a crucial part of using any medical technology, we designed a safety evaluation framework of the LLM outputs based on the framework used for evaluating a medical device [57,58]. Overall, this assessment involves estimating the likelihood of an error happening (Table 3) in conjunction with the potential impact of the error on the clinical outcome if it does occur. Table 4 shows the classification of the level of harm, and Figure 14 presents the estimation of risk based on the likelihood and consequences of an event.

Likelihood Category	Interpretation
Very high	Certain or almost certain; highly likely to occur
High	Not certain but very possible: reasonably expected to occur in the majority of cases
Medium	Possible

Low	Could occur but in the great majority of occasions will not
Very low	Negligible or nearly negligible possibility of occurring

Table 3. The likelihood of a hazard occurring

Consequence classification	Interpretation	Number of patients affected
Catastrophic	Death	Multiple
	Permanent life-changing incapacity and any condition for which the prognosis is death or permanent life-changing incapacity, severe injury or severe incapacity from which recovery is not expected in the short term	Multiple
Major	Death	Single
	Permanent life-changing incapacity and any condition for which the prognosis is death or permanent life-changing incapacity, severe injury or severe incapacity from which recovery is not expected in the short term	Single
	Severe injury or severe incapacity from which recovery is expected in the short term	Multiple
	Severe psychological trauma	Multiple
Considerable	Severe injury or severe incapacity from which recovery is expected in the short term	Single
	Severe psychological trauma	Single
	Minor injury or injuries from which recovery is not expected in the short term	Multiple
	Significant psychological trauma	Multiple
Significant	Minor injury or injuries from which recovery is not expected in the short term	Single

	Significant psychological trauma	Single
	Minor injury from which recovery is expected in the short term	Multiple
	Minor psychological upset; inconvenience	Multiple
Minor	Minor injury from which recovery is expected in the short term; Minor psychological upset; inconvenience; any negligible severity	Single

Table 4: Guidance for assessing the level of harm

Likelihood		Minor	Significant	Considerable	Major	Catastrophic
	Very high	3	4	4	5	5
	High	2	3	3	4	5
	Medium	2	2	3	3	4
	Low	1	2	2	3	4
	Very low	1	1	2	2	3
		Consequence				

Figure 14. Risk estimation based on the likelihood and consequence of harm occurrence

To maintain consistency in assessing the likelihood of hallucinations and omissions in each experiment, we created a percentage-based metric for their occurrence across experiments, as detailed in Supplementary data 1. 'Very High' likelihood represents scenarios where errors were very common (>90%), whereas 'Very Low' likelihood was associated with situations where errors were rare (<1%). For the 'Medium' likelihood category, which covers 10–60%, we used a broader range to accommodate the output variability and the understanding that some errors may be less predictable or depend on context (Table 5).

	Per 25 examples	Possibility
Very High	22.5	90%
High	15	60%
Medium	7	10-60%
Low	2.5	10%
Very Low	0.5	1%

Table 5. Calculating the likelihood of an error occurring in the text output

CREOLA, Clinical Review of LLMs and AI: We combine the experiment design, hallucination and omission taxonomy, and clinical safety evaluation in a platform we denote CREOLA, short for Clinical Review of LLMs and A1 (pays tribute to Creola Katherine Johnson [59], a pioneering human computer at NASA. Just as human computers were integral to the safe landing of Apollo moon missions, clinicians play a vital role in safely integrating AI technologies into clinical practice).

The platform is used to identify resultant changes in generated clinical documentation arising from changes to processes in LLM architecture. As illustrated in the "experimental structure", these changes could involve - but are not limited to - the type of model used or prompts used to obtain outputs. The platform was hosted as a Streamlit web application (https://creola.tortus.ai/); the annotation user interface is displayed in Figure 15.

A

B

Figure 15: The annotation user interface used to identify hallucinations (a) and omissions (b), as well as categorise them into major and minor categories.

To facilitate clinician review, the closest sentence matches (highlighted in yellow) for each portion of the text under review were extracted from the counterpart document. In the case of hallucinations, portions of text in the note were compared to the consultation (a), whereas for omissions, portions of the transcript were compared against the note (b).

Annotator recruitment: As outlined earlier, our framework requires annotators to review model outputs. Clinicians are uniquely skilled in critically assessing the veracity of clinical facts in the text. Therefore, we ask clinicians to annotate errors for our experiments. Annotators could register to contribute to the annotation through the CREOLA platform. To ensure annotators had a good understanding of the process, one-to-one tuition was initially provided by the study team. As the number of annotators grew, a short online course was developed to explain the annotation process, followed by a questionnaire to ensure a comprehensive understanding of the material. The annotators were only able to participate if they completed the questionnaire correctly. The annotators could contact the study teams with any questions through the CREOLA platform in order to ensure any problems in the platform were dealt with promptly.

Acknowledgement
We would like to thank the many physicians who helped with our CREOLA experiments and the annotation of the clinical transcripts and notes. No funding has been received for this study.

Author contributions

DP, EA, MD, NM, SK and JB contributed to the concept, design and execution of the study. MD and SK built the CREOLA platform and with NM designed the various prompts and experiments. MD analysed all the results and prepared the figures. EA and NM wrote, reviewed and revised the paper. EA and DP

designed the clinical safety framework, reviewed all the annotations and scored the impact of errors on clinical safety. JAY provided expert advice on paper writing and direction, and contributed to significant paper revision and rewriting. EA, JAY, MD, NM, SK, JB and DP contributed to the review and revision of the paper.

Competing interests

DM is the CEO of the company Tortus AI and all authors were employees of Tortus AI at the time of the writing the paper (MD, NM, SK, and JB as full time and EA part time, JAY full time employee at the time of revision). The authors declare no competing interests.

Data availability

We have used data from Primock and ACI bench which are publicly available clinical transcripts and notes (references in the main text).

Code availability

We have added all our prompts used in the supplementary materials of the article. As explained in the methods section, we used OpenAI's GPT-4 (GPT-4-32k-0613) as the LLM for all our experiments.

References

1. Clusmann, J. *et al.* The future landscape of large language models in medicine. *Communications Medicine* **3**, 141 (2023).
2. Becker, G. *et al.* Four minutes for a patient, twenty seconds for a relative - an observational study at a university hospital. *BMC Health Serv Res* **10**, 94 (2010).
3. Asgari, E. *et al.* Impact of Electronic Health Record Use on Cognitive Load and Burnout Among Clinicians: Narrative Review. *JMIR Med Inform* **12**, e55499 (2024).
4. Ali, S. R., Dobbs, T. D., Hutchings, H. A. & Whitaker, I. S. Using ChatGPT to write patient clinic letters. *Lancet Digit Health* **5**, e179–e181 (2023).
5. Patel, S. B. & Lam, K. ChatGPT: the future of discharge summaries? *Lancet Digit Health* **5**, e107–e108 (2023).
6. Van Veen, D. *et al.* Adapted large language models can outperform medical experts in clinical text summarization. *Nat Med* **30**, 1134–1142 (2024).
7. Zhang, T. *et al.* Benchmarking Large Language Models for News Summarization. *Trans Assoc Comput Linguist* **12**, 39–57 (2024).
8. Ahmad, M. A., Yaramis, I. & Roy, T. D. Creating Trustworthy LLMs: Dealing with Hallucinations in Healthcare AI. (2023).
9. Xu, Z., Jain, S. & Kankanhalli, M. Hallucination is Inevitable: An Innate Limitation of Large Language Models. (2024).
10. Kripalani, S. *et al.* Deficits in Communication and Information Transfer Between Hospital-Based and Primary Care Physicians. *JAMA* **297**, 831 (2007).
11. Adane, K., Gizachew, M. & Kendie, S. The role of medical data in efficient patient care delivery: a review. *Risk Manag Healthc Policy* **Volume 12**, 67–73 (2019).
12. Schiff, G. D. Diagnostic Error in Medicine. *Arch Intern Med* **169**, 1881 (2009).
13. Kumar, S., Balachandran, V., Njoo, L., Anastasopoulos, A. & Tsvetkov, Y. Language Generation Models Can Cause Harm: So What Can We Do About It? An Actionable Survey. in *Proceedings of the 17th Conference of the European Chapter of the*

Association for Computational Linguistics 3299–3321 (Association for Computational Linguistics, Stroudsburg, PA, USA, 2023). doi:10.18653/v1/2023.eacl-main.241.

14. Bender, E. M., Gebru, T., McMillan-Major, A. & Shmitchell, S. On the Dangers of Stochastic Parrots. in *Proceedings of the 2021 ACM Conference on Fairness, Accountability, and Transparency* 610–623 (ACM, New York, NY, USA, 2021). doi:10.1145/3442188.3445922.

15. Rando, J. & Tramèr, F. Universal Jailbreak Backdoors from Poisoned Human Feedback. (2023).

16. Huang, L. *et al.* A Survey on Hallucination in Large Language Models: Principles, Taxonomy, Challenges, and Open Questions. (2023).

17. Rawte, V. *et al.* Exploring the Relationship between LLM Hallucinations and Prompt Linguistic Nuances: Readability, Formality, and Concreteness. (2023).

18. S.M Towhidul Islam Tonmoy, S. M. M. Z. V. J. A. R. V. R. A. C. A. Das. A Comprehensive Survey of Hallucination Mitigation Techniques in Large Language Models. *Computer Science, Computation and Language* (2024).

19. Overhage, J. M., Qeadan, F., Choi, E. H. E., Vos, D. & Kroth, P. J. Explaining Variability in Electronic Health Record Effort in Primary Care Ambulatory Encounters. *Appl Clin Inform* **15**, 212–219 (2024).

20. Shahbodaghi, A., Moghaddasi, H., Asadi, F. & Hosseini, A. Documentation Errors and Deficiencies in Medical Records: A Systematic Review. *J Health Manag* **26**, 351–368 (2024).

21. Moramarco, F. *et al.* Human Evaluation and Correlation with Automatic Metrics in Consultation Note Generation. in *Proceedings of the 60th Annual Meeting of the Association for Computational Linguistics (Volume 1: Long Papers)* 5739–5754 (Association for Computational Linguistics, Stroudsburg, PA, USA, 2022). doi:10.18653/v1/2022.acl-long.394.

22. Demner-Fushman, D. *et al.* Preparing a collection of radiology examinations for distribution and retrieval. *Journal of the American Medical Informatics Association* **23**, 304–310 (2016).

23. Abacha, A. Ben, Yim, W., Michalopoulos, G. & Lin, T. An Investigation of Evaluation Metrics for Automated Medical Note Generation. (2023).

24. Ji, Z. *et al.* Survey of Hallucination in Natural Language Generation. *ACM Comput Surv* **55**, 1–38 (2023).

25. Huang, Y., Tang, K., Chen, M. & Wang, B. A Comprehensive Survey on Evaluating Large Language Model Applications in the Medical Industry. (2024).

26. Tierney, A. A. *et al.* Ambient Artificial Intelligence Scribes to Alleviate the Burden of Clinical Documentation. *NEJM Catal* **5**, (2024).

27. Lin C-Y. *ROUGE: A Package for Automatic Evaluation of Summaries. Text Summarization Branches Out.* . (Association for Computational Linguistics, 2004).

28. Papineni, K. , R. S. , W. T. & Z. W. A method for automatic evaluation of machine translation. *The Proceedings of ACL-2002, ACL, Philadelphia, PA* (2002).

29. Zhang, T., Kishore, V., Wu, F., Weinberger, K. Q. & Artzi, Y. BERTScore: Evaluating Text Generation with BERT. (2019).

30. Moreno, A. C. & Bitterman, D. S. Toward Clinical-Grade Evaluation of Large Language Models. *International Journal of Radiation Oncology*Biology*Physics* **118**, 916–920 (2024).

31. Minaee, S. *et al.* Large Language Models: A Survey. (2024).

32. Evans, O. *et al.* Truthful AI: Developing and governing AI that does not lie. (2021).

33. Singhal, K. *et al.* Large Language Models Encode Clinical Knowledge. (2022).

34. Reddy, S. Evaluating large language models for use in healthcare: A framework for translational value assessment. *Inform Med Unlocked* **41**, 101304 (2023).

35. Tang, L. *et al.* Evaluating large language models on medical evidence summarization. *NPJ Digit Med* **6**, 158 (2023).

36. Tam, T. Y. C. *et al.* A framework for human evaluation of large language models in healthcare derived from literature review. *NPJ Digit Med* **7**, 258 (2024).

37. Cohan, A. *et al.* A Discourse-Aware Attention Model for Abstractive Summarization of Long Documents. (2018).

38. Gupta, V., Bharti, P., Nokhiz, P. & Karnick, H. SumPubMed: Summarization Dataset of PubMed Scientific Articles. in *Proceedings of the 59th Annual Meeting of the Association for Computational Linguistics and the 11th International Joint Conference on Natural Language Processing: Student Research Workshop* 292–303 (Association for Computational Linguistics, Stroudsburg, PA, USA, 2021). doi:10.18653/v1/2021.acl-srw.30.

39. Luo, Z., Xie, Q. & Ananiadou, S. CitationSum: Citation-aware Graph Contrastive Learning for Scientific Paper Summarization. (2023).

40. DeYoung, J., Beltagy, I., van Zuylen, M., Kuehl, B. & Wang, L. L. MS2: Multi-Document Summarization of Medical Studies. (2021).

41. Song, Y., Tian, Y., Wang, N. & Xia, F. Summarizing Medical Conversations via Identifying Important Utterances. in *Proceedings of the 28th International Conference on Computational Linguistics* 717–729 (International Committee on Computational Linguistics, Stroudsburg, PA, USA, 2020). doi:10.18653/v1/2020.coling-main.63.

42. Johnson, A. E. W. *et al.* MIMIC-CXR, a de-identified publicly available database of chest radiographs with free-text reports. *Sci Data* **6**, 317 (2019).

43. Pal, A., Umapathi, L. K. & Sankarasubbu, M. Med-HALT: Medical Domain Hallucination Test for Large Language Models. (2023).

44. Papadopoulos Korfiatis, A. and M. F. and S. R. and S. A. *PriMock57: A Dataset of Primary Care Mock Consultations. Https://Github.Com/Babylonhealth/Primock57*. (2022).

45. Farquhar, S., Kossen, J., Kuhn, L. & Gal, Y. Detecting hallucinations in large language models using semantic entropy. *Nature* **630**, 625–630 (2024).

46. Simon Hughes, M. B. Vectara Hallucination Leaderboard comparing LLM performance at maintaining factual consistency when summarizing a set of facts. *Vectara, Inc. https://huggingface.co/spaces/vectara/leaderboard* (2023).

47. Riedemann, L., Labonne, M. & Gilbert, S. The path forward for large language models in medicine is open. *NPJ Digit Med* **7**, 339 (2024).

48. Zhang, G. *et al.* Closing the gap between open source and commercial large language models for medical evidence summarization. *NPJ Digit Med* **7**, 239 (2024).

49. Lewis, P. *et al.* Retrieval-Augmented Generation for Knowledge-Intensive NLP Tasks. *arXiv:2005.11401* (2020).

50. Wei, J. *et al.* Chain-of-Thought Prompting Elicits Reasoning in Large Language Models. *arXiv:2201.11903* (2022).

51. Jia, M., Duan, J., Song, Y. & Wang, J. medIKAL: Integrating Knowledge Graphs as Assistants of LLMs for Enhanced Clinical Diagnosis on EMRs. *arXiv:2406.14326* (2024).

52. Wang, Y., Ma, X. & Chen, W. Augmenting Black-box LLMs with Medical Textbooks for Clinical Question Answering. *arXiv:2309.02233* (2023).

53. Gilbert, S., Kather, J. N. & Hogan, A. Augmented non-hallucinating large language models as medical information curators. *NPJ Digit Med* **7**, 100 (2024).

54. Desmond, M., Ashktorab, Z., Pan, Q., Dugan, C. & Johnson, J. M. EvaluLLM: LLM assisted evaluation of generative outputs. in *Companion Proceedings of the 29th International Conference on Intelligent User Interfaces* 30–32 (ACM, New York, NY, USA, 2024). doi:10.1145/3640544.3645216.

55. Zheng, L. *et al.* Judging LLM-as-a-Judge with MT-Bench and Chatbot Arena. *Computer Science: Computation and Language* https://arxiv.org/abs/2306.05685 (2023).

56. Gu, J. *et al.* A Survey on LLM-as-a-Judge. *Computer Science: Computation and Language* arXiv:2411.15594 (2024).

57. International Organization for Standardization. *Https://Www.Iso.Org/Standard/59752.Html*.

58. International Organization for Standardization. *Https://Www.Iso.Org/Standard/72704.Html*.

59. https://en.wikipedia.org/wiki/Katherine_Johnson. Katherine Johnson.

Supplementary materials

Examples of hallucinations in the medical output text:

Fabricated Facts/Negations

Facts (diagnostics, plan, communication, …) that are completely fabricated (were never mentioned)

Type	Reason	Sentence
Minor	Clinical assessment not explicitly mentioned but accurate	Impression or clinical assessment: Chronic cough with sputum production and exertional dyspnea, possibly exacerbated by smoking, with a history of asthma.
Major	This is not in the input text shown	Follow up: Urgent referral to emergency department for further evaluation and management.
Minor	Not in the transcript	Treatment plan: Rest, continue with Advil as needed for pain relief.
Minor	Only neck mentioned, no other joints	Pertinent review of systems mentioned: No stiffness or loss of motion in joints.

Negated Events

Clinical events (e.g., symptoms, negations, surgeries, diagnostics) that were mentioned but not recorded

Type	Reason	Sentence
Major	Patient reports tiredness and feeling hot	No other significant symptoms.
Minor	Has had attendance due to peanut allergy	She has not had any surgeries or hospitalisations.

Context Conflation

Topics that were mentioned but mixed up in the note.

Type	Reason	Sentence
Minor	Exposure to mold and dust is at home and asbestos exposure in living and workplace.	Family/Social History: Works in crowded, poorly conditioned workplace with potential mold, dust, and asbestos exposure.
Minor	The cough preceded the shortness of breath and has been over three weeks.	History of presenting complaint: Shortness of breath and coughing started two weeks ago, worsening over time.
Minor	The past medical history is epilepsy. The missed medication is related to the current presentation.	Past Medical History: Epilepsy, seizure disorder, infrequent seizures; missed medication dose.

Assumed causality

Made-up causality between two things that were mentioned independently

Type	Reason	Sentence
Major	I can't see in the text that the physician ascribes the pain as 'stress related'.	Impression or clinical assessment: Chest pain of unknown origin, possibly stress related.
Major	Shouldn't make associations without being explicitly mentioned in the transcript	Impression or clinical assessment: Possible upper respiratory tract infection or COPD exacerbation, considering significant smoking history and environmental exposures.

Minor	No mention of smoking exacerbating dyspnoea	Impression or clinical assessment: Chronic cough with sputum production and exertional dyspnea, possibly exacerbated by smoking, with a history of asthma.

Examples of omissions in the medical output text and the section it occurred:

Current Issue

Type	Reason	Sentence
Major	Palpitations not mentioned. This points the diagnosis towards Pulmonary embolism/infective cause rather than cardiac	Yes, since this morning, it's felt like my heart is racing. Okay. Have you noticed any skin changes at all? Any rashes? No rashes. Okay.
Minor	Blood in sputum important	And has there been any kind of green, slimy mucus in your cough at all? Uh, no, it's been a dry cough. Dry cough. Okay. And have you noticed at all coughing up blood?
Minor	Need to mention if any change in smell/taste as may indicate COVID infection	Uh, no, I haven't noticed that. What about your sense of taste? Are you being, are you able to taste things? Yeah, I think so. Okay.

Information and Plan

Type	Reason	Sentence
Major	Advice: if the pregnancy test is positive, call straight back to organise be seen. Important safety net	So if you can do that today and if it's positive we need to see you straight away so give us a call back. Okay. Have you got any other questions? Right.
Minor	Explanation of shortness of breath important to include to gauge severity of this	Oh yeah, it's like when I'm doing walking, and I do exercise now and then, so it hasn't been... Super bad, like it hasn't stopped me from doing any of those things, but it's been pretty unusual and I'm a little bit worried.
Minor	Discussing diagnosis	So yeah, based on what we discussed today, it seems that you may have an upper respiratory tract infection.

PMFS Issues

Type	Reason	Sentence
Major	No allergies is an important negation to record	Okay. Right. Any allergies? Nothing. Yeah. Nothing. No allergies at all. Right here. Fine.
Major	Need to mention smoking history as this is important in someone presenting with chest pain. Is a risk factor for myocardial infarction etc	And do you use any recreational drugs like marijuana? Never. Okay. And do you smoke? Yes, I smoke about two packs of cigarettes per day.
Minor	Smoking status in an asthmatic should be recorded	No, I don't smoke. Do you drink any alcohol regularly? Yeah, one glass. How often is it? One glass of what? One glass of wine. Every... how often did you say?

Prompts and note generation

We used various approaches and prompts in our experiments. Each approach contributed valuable insights into refining both the process and output of LLM-based summarisation, allowing us to use the model with the best output for our clinical product. All prompts we used are listed in supplementary data 1.

Initial prompts
Experiments 1 and 2 were the initial prompts we had prior to utilising the CREOLA framework provided in the table below. These prompts served as baselines to our experiments before they were replaced.

Structured prompts
These were organised into three components: the base (core query), the template (context or format instructions), and the style (tone or level of detail). The purpose of this structured design was to address feedback on our outputs by changing specific sections of our prompt.

Experiments 3, 7, and 8 utilised the structured prompt. In experiment 7, we improved the base note to generate content from a first-person perspective. In experiment 8, we revised the style section of our SOAP note by incorporating negations and other style modifications. Additionally, we included "unknown" as an option for the status of problems not described in the transcript. Experiment 4 used the baseline SOAP note with an additional sentence to improve medication records.

Atomisation
In this model, atomic facts were extracted from our input text, which enabled precise extraction and organisation of clinical details. In experiment 5, atomisation was carried out to extract facts from the transcript. This process involved breaking the transcript down into individual atomic facts, which were then used to create a clinical note. The description did not provide a specific example. The procedure combined an adverb system prompt with instructions written in the third person.

Supplementary Table 1 shows the extraction of atomic facts from the transcript, followed by note generation by the LLM.

Atomisation of clinical facts		LLM note generation
Facts: - the patient has a cough. - the patient's cough is productive. - the patient brings up gunk when coughing. - the gunk is green. - the gunk is sometimes yellowish. - the patient has no blood in the cough. - the patient uses a couple of Kleenexes a day for the sputum. - the patient has had a cough for the past few years. - the patient feels the cough is getting worse. - the patient coughs throughout the day. - the patient's cough is worse in the morning. - the patient cannot think of anything that triggers the cough. - the cough has been getting worse over the last few years. - the cough's worsening has been slow. - the patient has had episodes of cough with sputum for the last few years. - taking a deep breath makes the patient's cough worse. - nothing helps the cough go away. - the patient has no runny nose. - the patient has no sore throat. - the patient has no difficulty breathing. - the patient has not traveled outside of the country. - the patient has no fever. - the patient has no chills. - the patient has no night sweats. - the patient has increased fatigue. - the patient has been feeling more tired than usual in the last few months. - the patient's hands are always a little swollen. - the patient's feet are always a little swollen. - the patient's ankles are always a little swollen. - the patient has gained a few pounds over the years. - the patient has no chest pain. - the patient has trouble breathing when walking a few blocks or going up a few flights of stairs. - the patient has urinary issues at nighttime.	- the patient has to get up a lot at night to use the bathroom. - the patient has no bowel issues. - the patient's sleep has been poor. - the patient does not wake up during the night to take a breath. - the patient uses one pillow. - the patient has lost five pounds in the last few months. - the patient has not been trying to lose weight. - the patient's appetite has been good. - the patient has no recent infections. - the patient has no runny nose. - the patient has no sore throat. - the patient has not been exposed to anyone with tuberculosis. - the patient had a normal tuberculosis test for work. - the patient's blood pressure has been high. - the patient was diagnosed with high blood pressure in the last few years. - the patient takes Lisinopril for blood pressure. - the patient has no known medication allergies. - the patient has not been hospitalized. - the patient had a c-section for the birth of her daughter. - the patient was in the hospital for a few days for the c-section. - the patient has had no other surgeries besides the c-section. - the patient's father had a heart attack at age 78. - the patient's father passed away from the heart attack. - the patient has no family history of cancer. - the patient has no known family history of asthma. - the patient lives in an apartment. - the patient lives with her husband and daughter. - the patient works at the local grocery store. - the patient eats one or two meals a day. - the patient sometimes snacks throughout the day. - the patient's diet includes a variety of meats and vegetables. - the patient does not exercise currently. - the patient does not use recreational drugs. - the patient drinks a glass of wine three or four nights a week. - the patient drinks five or six ounces of wine per glass. - the patient smokes cigarettes. - the patient smokes one to two packs of cigarettes per day. - the patient has been smoking for nearly 40 years.	History. - Allergies: No known medication allergies. - Medications: Lisinopril for high blood pressure. - History of presenting complaint: Chronic productive cough for years, worsening slowly, worse in the morning, exacerbated by deep breaths, no identifiable triggers, associated with increased fatigue and weight loss without trying. - Past Medical History: High blood pressure, c-section. - Family/Social History: Father had a heart attack and passed away at 78, no family history of cancer or asthma, lives with husband and daughter, works at a grocery store, diet includes meats and vegetables, no exercise, smokes 1-2 packs of cigarettes daily for nearly 40 years, drinks wine 3-4 nights a week. - Sensitive information: [NOT MENTIONED]. Observations: - Examination findings: [NOT MENTIONED]. - Investigation results: [NOT MENTIONED]. - Impression or clinical assessment: Chronic cough likely related to smoking, with symptoms suggestive of chronic bronchitis or early chronic obstructive pulmonary disease (COPD), considering the history of smoking and productive cough. Differential diagnosis includes heart failure, given peripheral edema, weight gain, and dyspnea on exertion. Plan: - Planned investigations: Chest X-ray, pulmonary function tests, complete blood count, basic metabolic panel, echocardiogram to assess for heart failure. - Follow up: Schedule follow-up appointment after investigations are complete. - New prescribed medication or therapies: [NOT MENTIONED]. - Communication, reassurance & patient understanding of care: Explained the need for investigations to determine the cause of the cough and associated symptoms. Discussed the impact of smoking on health and the potential benefits of cessation. - Actions for referrer/GP: [NOT MENTIONED]

Supplementary Table 1: Atomisation of clinical facts (a) and LLM note generation (b)

Figures 9 and 10 illustrate how the use of atomic facts prior to note generation leads to a significant increase in hallucinations and omissions compared to our base and structured prompts (experiments 1 and 3, respectively)

Function call and updated style block
A function call is the process of executing a specific function in a program by providing its name and the required input arguments. When called, the function performs its defined task and returns a result or performs an action.

In these experiments, the LLM was instructed to generate responses in a structured JSON format rather than free text. This output format was chosen due to its compatibility with clinician note summarisation in primary care. Experiments 6, 9, 10, and 11 were conducted using this format.

Experiment 9 aimed to test an updated style block for our EMIS (Educational Management Information System, which is an electronic health record system used widely in primary care settings in the United Kingdom) note, incorporating negations and other stylistic enhancements. We also revised several descriptions in the function schema. The block featured an EMIS note with the new style, which included negations and improvements such as conciseness. In experiment 10, we tested the EMIS function call with enhanced descriptions for examination and comments. This refinement aimed to ensure that only information present in the transcript was included. Experiment 11 was a continuation of experiment 10, and utilised the EMIS function call with improved descriptions and additional style guidance, specifically to avoid including information outside of the transcript.

Structured prompt + LLM revision step
In these experiments, we added an additional LLM call in order to review and refine its initial output to meet the requirements of a structured prompt, aiming to enhance the overall quality of the output.

In experiments 14 and 15, we applied this method. In experiment 14, we built upon the baseline established in Experiment 11 by adding a revision step after updating the state-of-the-art (SOTA) SOAP block. This revision included modifications to incorporate negations and other style enhancements, such as being more concise. Additionally, we introduced "unknown" as an option for the status of problems not described in the transcript. Experiment 15 involved adding a checking step to the 'Bad SOAP' note block, which was a SOAP note containing numerous hallucinations and omissions.

As noted in the supplementary Figure 1, experiment 14 had a slightly lower number of major omissions and minor hallucinations compared to experiment 8. Experiment 15 had fewer major hallucinations and omissions compared with experiment 3.

Supplementary Figure 1. Impact of adding an LLM revision step on the incidence of omissions and hallucinations

The figure illustrates the comparison between experiments 14 and 8 on the occurrence of omissions (a) and hallucinations (b), similar to the comparison between omissions (c) and hallucinations (d) between experiments 15 and 3.

New approach to note generation

In experiment 16, we developed a new method for generating customised outputs based on a specific template. We used our SOAP (Subjective, Objective, Assessment, Plan) note as the template and compared it to the baseline prompt output (from experiment 8). Interestingly, this comparison resulted in an increase in the number of major hallucinations and minor omissions, as shown in Supplementary Figure 2.

Supplementary Figure 2: Comparing hallucinations and omissions between our baseline prompt and customised output
This figure illustrates the changes in omissions (a) and hallucinations (b) between using the baseline prompt and a prompt designed to generate a customised output

Information gathering experiments

In experiment 17, we asked several clinicians to create notes based on medical transcripts. We then evaluated these notes for the presence of hallucinations and omissions. The goal was to understand the extent of inaccuracies in human-generated notes and to compare them with those generated by LLM. The clinician-generated notes exhibited slightly more major hallucinations and fewer omissions, as shown in Supplementary Figure 3.

Supplementary Figure 3. Comparing hallucinations and omissions between LLM-generated notes and human-generated notes

Figure 3 illustrates the number of major and minor omissions (a) and hallucinations (b) between LLM -generated notes in experiment 8 and human-generated notes in experiment 17