**Clinical management, epidemiology, and recurrence of human cystic**

**echinococcosis in a secondary care level hospital in an endemic area of the**

**Andes in Sicuani, Cusco, Peru**

Roberto Pineda-Reyes[1], Miguel M. Cabada[2,3], Maria L. Morales[3], Jan Hattendorf[1], Paola

Vergaray[4], Ruben Bascope[5], Jakob Zinsstag[1*]

[1] One Health Group, Human and Animal Health Unit, Department of Epidemiology and

Public Health, Swiss Tropical and Public Health Institute, associated to University of Basel,

Allschwil, Switzerland

[2] Division of Infectious Diseases, Department of Internal Medicine, University of Texas

Medical Branch, Galveston, TX, USA

[3] Cusco Branch – Alexander von Humboldt Tropical Medicine Institute, Universidad Peruana

Cayetano Heredia, Cusco, Peru

[4] Departamento de Cirugia, Hospital Alfredo Callo Rodriguez, Sicuani, Cusco, Peru

[5] Programa de Control de Enfermedades Zoonoticas, Gerencia Regional de Salud de Cusco,

Cusco, Peru

* Corresponding author (JZ): E-mail: jakob.zinsstag@swisstph.ch   Phone: +41 61 284 81 39

*Keywords: recurrence, hydatid disease, echinococcosis, Andes, South America*

*Word count: Abstract, 263 words; manuscript text, 3763.*

*Running head: Recurrence of cystic echinococcosis in Sicuani, Cusco, Peru*

*Tables and figures: 3 tables, 1 figure.*

**Abstract**

Peru has the highest incidence of human cystic echinococcosis (CE) in South America and most cases are reported in the Central and Southern Peruvian Andes, including the Cusco region. We reviewed medical records of patients with CE admitted between 2010-2019 to a level-2 hospital in Sicuani, which serves the Canchis province of Cusco. We collected data on surgical management and disease recurrence. Records from 115 patients were included, their median age was 25 years (IQR, 16-46) and 60% were female. Most patients (68.7%) presented with a single liver cyst. The Gharbi classification was used in 75/107 (70.1%) patients with an ultrasound and in 91/142 cysts (64.8%). Thirty-one (34.1%) were classified as Gharbi I, 33 (36.3%) Gharbi II, 21 (23.1%) Gharbi III, and 6 (6.6%) Gharbi IV. Pre-surgical complications were reported in 41.7%. One hundred two patients underwent surgery. In 39.2% a secondary CE prevention measure was documented, 5.9% had two, and one had the three recommended measures. Post-surgical cyst recurrence was reported in 16.5% at a median 32.3 months (IQR, 3.9-46.6) and readmissions in 12.7%. In the multivariable analysis, having ≥3 cysts (OR 6.26, 95% CI 1.10-39.85) was associated with post-surgical cyst recurrence. Advanced and complicated CE and post-surgical recurrence were common in Sicuani. Standardization of ultrasound staging-guided clinical management along with secondary CE prevention measures could effectively decrease the disease burden linked to clinical care, improving outcomes and decreasing costs. Ultimately, only integrated control at the source of infection including hand hygiene, secure disposal of offal, avoidance of dog feeding offal, and deworming of dogs can reduce transmission and lead to elimination.

**Introduction**

Cystic echinococcosis (CE) is a zoonosis caused by the metacestode stage of the dog tapeworm *Echinococcus granulosus* (1). *E. granulosus* has a global distribution, being endemic in South America, North and East Africa, Central Asia and China (2). Peru has the highest incidence of CE in South America, accounting for 84.8% of all reported cases in year 2018 (3). The district of Sicuani in the Cusco region is a highly endemic area in the Southern Peruvian Andes. Up to 45% of CE cases referred to tertiary care hospitals in Cusco are from Sicuani (4). A population survey using focus ultrasound to detect abdominal CE in four communities of the Cusco region found that 78% of patients diagnosed with CE were from Sicuani (5). The economic and urban expansion of the Sicuani city have created areas where unregulated livestock slaughter activities, stray dogs, and humans interact closely. Prompt diagnosis and appropriate treatment of cases in Sicuani are crucial to address the burden of CE in the healthcare system.

The WHO-Informal Working Group on Echinococcosis (WHO-IWGE) ultrasound staging system is used to diagnose and guide the clinical management of liver CE (6). However, many low- and middle-income countries (LMIC) have not implemented the proposed WHO ultrasound staging which may expose patients to unnecessary procedures (7). In developing countries, CE has a high post-surgical recurrence risk, with rates that range between 4.7% and 16.4% (8-11). Clinical characteristics such as the cyst size, number of cysts, and presence of complications have been associated with disease severity and recurrence (10, 12, 13). Treatment precautions to avoid cyst rupture and intraabdominal seeding may decrease the risk of recurrence related to surgical care.

The recommendations to avoid cyst rupture and secondary CE include the use of surgical field protection with hypertonic saline-soaked drapes, injection of a scolicidal agent into the cyst, and prescription of pre-surgical albendazole (ABZ) (6). Increasing evidence shows that appropriate standardization of surgical procedures may be associated with lower risk of recurrence (14-16). A prospective study found no mortality and zero recurrence at 24

months after standardized partial cystectomy, which was in part attributed to careful

avoidance of intraabdominal spillage and prescription of perioperative ABZ.(16) In addition,

the "watch-and-wait" approach is recommended as the standard care for inactive cysts to

spare asymptomatic patients from unnecessary treatment (17, 18).

We conducted a review of medical records of patients admitted during a 10-year period to a

low-complexity hospital in the city of Sicuani in Cusco, Peru, to describe the CE

presentation, management, and post-surgical recurrence in a highly endemic area.

**Methods**

*Study design and setting*

This is a review of medical records from patients hospitalized at the Alfredo Callo Rodriguez

(ACR) Hospital in Sicuani. This hospital is a level-2 public healthcare center that provides

specialty medical care to patients from the Canchis and Canas provinces of Cusco (19).

During March 2024, we collected data from all patients admitted to the hospital for CE

between January 1, 2010, and December 31, 2019. The pre-COVID-19 period was selected

to avoid collecting information that could reflect gaps in care or practice of CE treatment in

Sicuani because of the pandemic public health emergency response. This timeframe also

coincides with the renovation and expansion of surgical services in the hospital in 2010.

The city of Sicuani is the capital of the Canchis province in the Cusco region at an elevation

of 3,550 meters above sea level. The Sicuani district has an urban population of 47,400 and

a rural population of 10,400 habitants (20). Sicuani is a commerce center for farmers of the

surrounding areas and is the second largest city in the Cusco region.

*Study procedures*

Medical record numbers of patients with CE were identified using ICD-10 codes (B67.0-

B67.9) assigned as the discharge diagnosis through the hospital's statistics department.

Surgical logbooks were handsearched to ensure all surgical CE cases were retrieved.

Demographic and clinical data were collected via ODK software (© 2022 Get ODK Inc.,

California, USA) using an electronic case report form, which was piloted at the start of data

collection to ensure data accuracy.

*Definitions*

*Pre-surgical complications* were defined as documented evidence of cyst-related infectious

processes, fistulas, mass effect, allergic manifestations, hemorrhage, or trauma before

surgical treatment. One patient who presented to the hospital with abdominal pain and a 13-

week pregnancy was also considered complicated before surgery. The three assessed

*secondary CE prevention measures* included surgical field protection with drapes soaked in

hypertonic saline, injection of hypertonic saline into the cyst prior to evacuation, and the

prescription of pre-surgical ABZ (6). Hemoglobin values were adjusted for age, sex,

pregnancy status, and district of residence elevation in order to define *mild, moderate* or

*severe anemia* (21). *Post-surgical recurrence* was defined as the development of a new cyst

at or near the site of the previously removed cyst and/or within the cavity (e.g., peritoneum,

pleura) where cyst(s) were operated on. The *time-to-recurrence* was estimated as the time

from the surgery date until the date of recurrence diagnosis. In one case the date of

recurrence was documented only as "2024" and it was assigned "January-1-2024" as the

corresponding diagnosis date. *New cyst* was defined as a newly diagnosed cyst in a location

different from other operated cysts. *Readmission* was defined as return to inpatient care

related to CE after discharge. *Readmission rate* was calculated by dividing the number

patients with at least one readmission by the total number of hospitalization discharges.

*Annual cumulative incidence calculation*

The annual cumulative incidence of new hospital admissions was calculated dividing the

number of new CE hospitalizations per year by the total population at risk each year. The

population at risk was defined as the total population of the Canas and Canchis provinces

per year. Population data for years 2015 to 2019 was extracted from a repository database

of the Peruvian Ministry of Health and data for years 2009 to 2014 was extracted from a third

party vendor database, CPI Peru (22, 23).

*Statistical analysis*

Data analysis and visualization were performed using the R software, version 4.3.2 (R

Foundation for Statistical Computing, Vienna, Austria). Means with standard deviation (SD),

medians with interquartile range (IQR), and frequencies were used to assess the

distributions of demographic, epidemiological, and clinical characteristics of patients. A

multivariable logistic regression analysis was performed to determine independent

associations with the outcome variable "post-surgical recurrence". We pre-selected the

variables "age", "sex", and added further variables to the multivariable model based on

p<0.15 on univariable analyses with the outcome variable. A *P* value of <0.05 was

considered statistically significant. Confidence intervals were calculated at the 95% level.

*Ethical Statement*

This study was reviewed and approved by the Ethics Committee of Northern and Central

Switzerland (Ethikkommission Nordwest- und Zentralschweiz, EKNZ, No. AO_2023-00096)

and the Institutional Ethics Committee (Comite Institucional de Etica en Investigacion, No.

213622) of the Universidad Peruana Cayetano Heredia in Lima, Peru.

**Results**

*Demographic and clinical characteristics*

One hundred and twenty-five medical record numbers were identified. Six were duplicated, 2 were misrecorded as CE, and 2 had missing charts. In total, 115 unique medical records from patients with CE were included in the study (**Table 1**). The median age was 25 years (IQR, 16-46) and 60% were female. Most patients (103, 89.6%) were residents of the Canchis province and 73.9% lived in the Sicuani district. Fifty-six patients (48.7%) were referred from primary healthcare centers and 57.1% of these were referred from the Pampaphalla health center.

One hundred eighty-five echinococcal cysts were diagnosed in the 115 patients, 169 were diagnosed upon presentation to the hospital and 16 during readmissions. These involved the liver, lungs, stomach, spleen, duodenum, gallbladder, diaphragm, adnexa, bladder, and colon. Most (142, 84%) were located in the liver, 16 within the abdominal cavity, and 11 in the lungs. Four patients had disseminated liver and intraabdominal cysts. The maximum number of lung cysts per patient was 2. The median diameter of the largest liver cyst per patient was 12.8 cm (IQR, 9.2-15.0) and for lung cysts was 8.5 cm (IQR, 4.5-12.6). Seventy-seven (67%) patients had a liver cyst of ≥10 cm in diameter.

Most patients (98, 85.2%) presented with cysts only in the liver, 7 (6.1%) with both liver and intraabdominal cysts, 5 (4.3%) with liver and lung cysts, and 5 (4.3%) with only lung cysts. Most patients presented with a single liver cyst (79, 68.7%). Forty-eight (41.7%) cases presented with complicated cysts before surgery, 31 (27%) had one pre-surgical complication, and 17 (14.8%) had two or more. The most common pre-surgical complications were infection of the cyst content (17, 14.9%), cysto-biliary fistulas (9, 7.8%), liver abscess (9, 7.8%), cyst rupture (8, 7%), and peritonitis (5, 4.3%) (**Table 2**). A summary of the clinical presentation characteristics and laboratory results is shown in **Supplementary Table 1S**.

*Ultrasound diagnosis*

One hundred seven patients (93%) had a diagnostic ultrasound. In 75 (70.1%), a liver ultrasound was performed and used for staging according to the Gharbi classification (6), and in 20 (18.7%) a liver ultrasound was performed but no staging was reported. Twelve patients had lung cysts, disseminated disease, complicated cysts, or unclear or missing ultrasound reports precluding the use of a cyst diagnostic and management classification. In total, 91 (64.8%) of the 142 liver cysts were staged using the Gharbi classification. Thirty-one (34.1%) were Gharbi I, 33 (36.3%) Gharbi II, 21 (23.1%) Gharbi III, and 6 (6.6%) Gharbi IV. Seven (6.1%) patients were diagnosed with inactive cysts, including one who had a Gharbi IV cyst diagnosed intraoperatively.

*Treatment, secondary CE prevention measures, and post-surgical complications*

Only patients with liver and intraabdominal cysts underwent surgery in Sicuani (102, 88.7%). Ninety-five of the 102 patients (93.1%) underwent open surgery, 6 (5.9%) laparoscopic surgery, and one laparoscopic surgery converted to open laparotomy. The most common surgical techniques were cystectomy plus drainage (74, 72.5%) and pericystectomy plus drainage (9, 8.8%) (**Supplementary Table 2S**). In one case a Gharbi IV cyst was found and not excised and two patients with inactive cysts (2%) underwent excision. In one case the liver cyst was not found during the surgery. None of the patients had a percutaneous procedure performed. In 18 (17.6%) patients, the surgical report did not describe the location of the cyst in the liver. Seventy (68.6%) patients had right liver lobe cysts, 26 (25.5%) left liver lobe cysts, and 12 (11.8%) had cysts in both liver lobes. In 46 (45.1%) patients the liver segment VIII was involved, in 32 (31.4%) the segment VII, in 24 (23.5%) the VI, in 17 (16.7%) the IV, in 12 (11.8%) the V, in 10 (9.8%) the II, in 8 (7.8%) the III, and in one patient the segment I.

Forty (39.2%) patients had one secondary CE prevention measure documented, 6 (5.9%) had two, and only one had the three recommended measures. Pre-surgical ABZ was used in

33 (32.4%) cases, injection of hypertonic saline before cyst opening was documented in 16

(15.7%) cases, and surgical field protection with hypertonic saline soaked drapes was used

in 6 (5.9%) cases.

ABZ was the only antiparasitic prescribed. In 57 (55.9%) patients, ABZ was prescribed after

surgery, in 27 (26.5%) before and after surgery, in 7 (6.9%) no ABZ was prescribed, in 6

(5.9%) ABZ was prescribed only before surgery, and in 5 cases the timing of the prescription

was unknown. The median post-surgical duration of ABZ treatment prescribed was 30 days

(IQR, 22-34) and the median pre-surgical duration of ABZ treatment prescribed was 2 days

(IQR, 1-4).

Post-surgical complications occurred in 22/96 (22.9%) patients with complete post-operative

documentation. The most common were the development of residual fluid collections (14,

14.6%), post-surgical abscess (3, 3.1%), and biliary leakage (3, 3.1%) (**Table 2**). The three

patients with post-surgical liver abscess were readmitted to the hospital.

*Post-surgical recurrence, readmission, and annual cumulative incidence*

One patient required a second surgical intervention within the same hospitalization for post-

surgical biliary leak, and one required intensive care due to post-surgical hemodynamic

instability. Eleven (9.6%) patients were referred to a higher level of care, 9 for treatment of

pulmonary CE, 1 for post-surgical shock, and 1 for potential need of post-surgical intensive

care. Of these, 10 patients were transferred to the Hospital Regional del Cusco and 1 patient

to the Hospital Antonio Lorena in Cusco. Post-surgical recurrence was documented in 16/97

(16.5%) patients. The median time to recurrence after surgery was estimated at 32.3 months

(IQR, 3.9-46.6). The multivariable logistic regression analysis showed that having ≥3 cysts

compared to a single cyst (50% vs 15.1%, OR=6.26, 95% CI 1.10-39.85) was associated

with post-surgical recurrence (**Figure 1 and Supplementary Table 3S**).

Fourteen (14/110, 12.7%) patients were readmitted to the hospital at least once, 6 did for

cyst recurrence, 2 for treatment complications, 2 for complications of a previous cyst plus

cyst recurrence, 3 for elective two-stage treatment, and one for a new cyst. One patient had

2 readmissions. Twelve out of the 14 patients with readmissions had active echinococcal

cysts. The median age was 40 years (IQR, 26.5-62), 7 (58.3%) were female, and 5 (41.7%)

were referred from the Pampaphalla health center. Single liver cyst was the presentation in

9/12 (75%) of patients with readmissions. Readmissions contributed with 16 additional cysts

to the overall count for the 10-year study period. The annual cumulative incidence of new

hospital admissions for CE at the ACR Hospital of Sicuani are reported in **Table 3**.

**Discussion**

Our study describes the clinical characteristics and management of CE at a low-complexity

hospital in an endemic area of the Southern Peruvian Andes. Most CE patients requiring

admission to this hospital were part of the economically active population and presented

often with advanced disease (cysts ≥10 cm in diameter) and/or pre-surgical complications.

Despite the common use of ultrasound for the evaluation of CE, the WHO-IWGE system was

not used for the diagnosis and management of liver CE in any of the cases. Instead, the

Gharbi classification was used on a limited number of cysts. Secondary CE prevention

measures were documented in less than half of the surgical procedures. One in six patients

had recurrence of CE after surgery and one in eight patients were readmitted to the hospital

at least once. Our data suggest that educational interventions could improve the medical and

surgical management of CE at the ACR Hospital of Sicuani, thus decreasing the morbidity

associated with disease recurrence and surgical procedures.

Between 2016-2018, the Peruvian Ministry of Health reported a CE cumulative incidence of

29.2-45 cases per 100,000 in the Cusco region, which is one of the highest in the country

(24). During the same period, our cumulative incidence calculation ranged between 8.4-11.2

new admissions to the ACR Hospital per 100,000. These numbers are also higher than most

other regions in Peru, even though they do not include outpatient consultations, cases

diagnosed in primary healthcare centers, and cases from the Social Security Health System

(EsSalud) or private healthcare centers. The ACR Hospital of Sicuani serves the populations

of the Canas and Canchis provinces. These provinces have large young age populations,

which are affected by CE. The period between infection and clinical presentation, or 'latent

period', typically lasts many years in CE. This suggests that active transmission occurs in

Sicuani and that it happens early in life. A large case series of hospitalized patients in two

regions of Spain showed a distribution skewed towards the older age groups with only 2%

cases in patients younger than 19 years (25, 26). The authors attributed this age distribution

in part to the successful interruption of transmission by control programs based on stray

dogs elimination and safe disposal of livestock offal (27). A similar scenario was described in

Chile, where most reported cases between 2018-2022 were diagnosed among patients

between 45-74 years (28). Efforts to control CE transmission in the Canas and Canchis

provinces have been limited to the treatment of dogs in rural areas and sporadic screening

of children. A concerted effort with a One Health approach is crucial to formulate effective

control strategies in the area.

In our study, most patients had liver cysts and 41.7% presented to the hospital with

complications. This is a high proportion of complications, similar to the 40.3% reported in a

study in Salamanca, Spain (29). Compared to the latter, our study had slightly lower

percentage of mechanical complications (20% vs 25%); however, we encountered a higher

proportion of infectious complications (25.2% vs 17.8%), cyst rupture (8.7% vs 2.9%), and

advanced disease (cysts ≥10 cm in 68.7% versus 45%) (29). These discrepancies could be

due to a selection bias related to the hospital level of complexity, our lower proportion of lung

cysts, or due to differences in healthcare access. Another retrospective series in the Junin

region in the Central Peruvian Andes also reported over 50% cysts ≥10 cm, but their

proportion of pre-surgical complications was significantly lower (4%) likely due to including

cases from outpatient services and private hospitals (30).

We could not establish a significant association between pre-surgical complications and CE

recurrence. Other studies with larger samples were also unable to demonstrate associations

between infectious, mechanical, or allergic complications and CE recurrence (9, 29). This

could indicate that the risk of recurrence depends on other clinical variables associated with

surgical or post-surgical care (9). Large cyst diameter has been proposed as a marker of

severity and risk factor for recurrence (12). Despite the large cyst diameters found in our

patients compared to other studies, an association between this variable and pre-surgical

complications, recurrence, or readmission was not detected (11, 13, 29). We did observe an

association between higher number of cysts and risk of post-surgical recurrence. Number of

cysts has been identified as risk factor for complications and recurrence (10, 13). Pacifico

and colleagues found that patients with ≥3 cysts had a four-fold higher risk for pre-surgical

complications in a secondary case-control analysis of a study in the Central Andes of Peru

(13). Similarly, having ≥3 cysts was an independent risk factor for recurrence in a

retrospective study of 672 surgically treated patients in Morocco (10). A large parasite

burden may lead to subclinical cyst rupture and seeding in some patients before hospital

admission or during surgical interventions.

Ultrasound staging was documented in two-thirds of those with liver cysts. Ultrasound CE

diagnosis and staging is not yet implemented in many LMIC (7). The WHO-IWGE staging

system was not used for the management of CE in our study sample. This classification

serves as a clinical management guide to consider medical and invasive (percutaneous,

surgical) interventions, or a 'watch and wait' approach (6). Patients with liver cysts who

underwent surgery presented with Gharbi I or II in our study, and some could have been

candidates for percutaneous treatment (31). However, this treatment option is not available

in endemic areas of Peru. Patients with Gharbi III cysts in our study had a greater proportion

of recurrence compared to the other stages. Surgery is the treatment of choice for Gharbi III

cysts due to high recurrence rates reported in percutaneous treatment studies (31, 32).

Multiple daughter cysts within a solid matrix, as seen in Gharbi III, may be more difficult to

access and drain likely missing smaller cysts which may not get sufficient scolicidal drug

exposure within the matrix, increasing the risk of treatment failure (32, 33). Early ultrasound

screening along with implementation of WHO-IWGE staging and percutaneous procedures

may decrease costs, unnecessary procedures, and hospitalization time in Sicuani.

We found a higher post-surgical recurrence rate in comparison to other studies from different

global regions (8, 10, 11, 34-37). A minority of patients had documentation of recommended

interventions to prevent spillage of cyst contents during surgery. In some cases, surgical

field drapes were soaked on normal saline or iodine rather than hypertonic saline. In others,

the instillation of a scolicidal agent into the cyst was performed after opening and

evacuation. An association between spillage-prevention measures and post-surgical

recurrence could not be determined. In addition, patients with documented adequate

spillage-prevention measures had a higher proportion of recurrence. This could be explained

by differences in documentation or care in patients with more complicated disease. In these

patients, the surgeons may have performed a more detailed documentation of interventions

or used spillage prevention more often. Although careful implementation of these measures

has been associated with no recurrence at 24 months in prospective studies (16), it is

unclear whether other measures used at the ACR Hospital are effective. We could not find

an association between recurrence and the timing of ABZ prescription (e.g., pre-surgical,

post-surgical, or both) or the duration of ABZ treatment but our power to find these

association was probably limited by our sample size. In addition, there was no

documentation of medication adherence, and some patients might have had problems to

afford or take antiparasitics consistently. Pre-surgical ABZ is a core measure for secondary

prevention in liver CE (31). Prospective studies assessing the timing and dosing of

antiparasitics as well as the appropriateness of spillage-prevention measures during surgery

may help close this information gap.

Our study had limitations that should be acknowledged. First, this is a retrospective review of

medical charts with a small sample size. Missing information in certain variables such as

ultrasound staging did not allow testing their association to the outcome. Second, as the

hospital had a digitalized database of medical record numbers only since 2017, cases

between 2010-2016 were manually searched in the surgical logbooks. Therefore, patients

hospitalized during this period who did not undergo surgery are missing in our cohort. Third,

the description of symptoms, complications, and management of patients in our study

represent the clinical judgement and documentation of different surgeons in Sicuani during a

long time span. These may not concur with the standard of care in other contexts and

generalizations should be carefully assessed. Additionally, surgical reports documentation

differed among surgeons, which precluded an analysis on the association of surgical

techniques and recurrence. Finally, the lack of follow-up in some patients may have caused

an underestimation of cyst recurrence.

In conclusion, advanced and complicated liver CE and post-surgical recurrence were

common among hospitalized patients at the ACR Hospital in the city of Sicuani in Cusco,

Peru. Practical measures including standardization of ultrasound-guided clinical

management, secondary CE prevention measures, and follow-up could decrease the CE

burden linked to clinical care, improving outcomes and decreasing costs. Ultimately, the

contextual ecological cycle of transmission must be established in Sicuani, as only an

integrated control at the source of infection involving interventions such as hand hygiene,

secure disposal of offal, avoidance of dog feeding offal, and deworming of dogs can reduce

transmission and lead to elimination. This requires a One Health approach and a socio-

ecological strategy analysis (38), similar to rabies control (39).

**Acknowledgements**

We thank the Alfredo Callo Rodriguez Hospital's Administration Office as well as the personnel of the Statistics and Archives Department for their assistance during the study period.

**Financial support**

This study was supported by the University of Basel "Bologna Master" thesis funding from the Department of Education and Training of the Swiss Tropical and Public Health Institute, Allschwil, Switzerland. Logistic and field activities were funded by the Cusco Branch – Alexander von Humboldt Tropical Medicine Institute at Universidad Peruana Cayetano Heredia, Cusco, Peru.

**Conflicts of interest**

The authors have no competing interests.

**Authors' details**

**RP-R:** Swiss Tropical and Public Health Institute, associated to University of Basel, Allschwil, Switzerland, Kreuzstrasse 2, 4123 Allschwil, Switzerland, repineda1@gmail.com; **MMC:** Division of Infectious Diseases, University of Texas Medical Branch, Galveston Texas, 301 University Boulevard, Rte. 0435, Galveston, TX 77555-0435, USA, micabada@utmb.edu and Cusco Branch – Alexander von Humboldt Tropical Medicine Institute, Universidad Peruana Cayetano Heredia, Jr. Jose Carlos Mariategui J-6, Wanchaq 08002, Cusco, Peru, miguel.cabada.s@upch.pe; **MLM:** Cusco Branch – Alexander von Humboldt Tropical Medicine Institute, Universidad Peruana Cayetano Heredia, Jr. Jose Carlos Mariategui J6, Wanchaq 08002, Cusco, Peru, maria.morales.f@upch.pe; **JH:** Swiss Tropical and Public Health Institute, associated to University of Basel, Allschwil, Switzerland, Kreuzstrasse 2, 4123 Allschwil, Switzerland, jan.hattendorf@swisstph.ch; **PV:** Departamento de Cirugia, Hospital Alfredo Callo Rodriguez – Sicuani, Av. Manuel Callo Zevallos Nro. 519, Sicuani, Cusco, Peru, miqxonc21@gmail.com; **RB:** Programa de Control de Enfermedades Zoonoticas, Gerencia Regional de Salud de Cusco, Prolongacion Avenida de la Cultura 1525, Cusco 08003, Peru, bascopecusco@gmail.com; **JZ:** Swiss Tropical and Public Health Institute, associated to University of Basel, Allschwil, Switzerland, Kreuzstrasse 2, 4123 Allschwil, Switzerland, jakob.zinsstag@swisstph.ch.

**References**

1.      Craig PS, McManus DP, Lightowlers MW, Chabalgoity JA, Garcia HH, Gavidia CM,

et al. Prevention and control of cystic echinococcosis. Lancet Infect Dis. 2007;7(6):385-94.

2.      Deplazes P, Rinaldi L, Alvarez Rojas CA, Torgerson PR, Harandi MF, Romig T, et al.

Global Distribution of Alveolar and Cystic Echinococcosis. Adv Parasitol. 2017;95:315-493.

3.      PAHO/WHO. Equinococosis: informe epidemiológico en la región de América del Sur

- 2018, n.4, 2020: Veterinary Public Health - PANAFTOSA - PAHO/WHO; 2020 [Available

from: https://iris.paho.org/handle/10665.2/51942.

4.      Flores Quispe O. Equinococosis quística en niños y adolescentes en los hospitalales

Regional del Cusco y Adolfo Guevara Velasco, Cusco a 3400 msnm, 2010-2014. Cusco,

Peru.: San Antonio Abad del Cusco.; 2015.

5.      Tamarozzi F, Hou A, Morales ML, Giordani MT, Vilca F, Mozo K, et al. Prevalence

and Risk Factors for Human Cystic Echinococcosis in the Cusco Region of the Peruvian

Highlands Diagnosed Using Focused Abdominal Ultrasound. Am J Trop Med Hyg.

2017;96(6):1472-7.

6.      Brunetti E, Kern P, Vuitton DA, Writing Panel for the W-I. Expert consensus for the

diagnosis and treatment of cystic and alveolar echinococcosis in humans. Acta Trop.

2010;114(1):1-16.

7.      Bold B, Hattendorf J, Shagj A, Tserendovdon B, Ayushkhuu T, Luvsandorj A, et al.

Patients with cystic echinococcosis in the three national referral centers of Mongolia: A

model for CE management assessment. PLoS neglected tropical diseases.

2018;12(8):e0006686.

8.      Kapan M, Kapan S, Goksoy E, Perek S, Kol E. Postoperative recurrence in hepatic

hydatid disease. J Gastrointest Surg. 2006;10(5):734-9.

9.      Velasco-Tirado V, Romero-Alegría Á, Belhassen-García M, Alonso-Sardón M,

Esteban-Velasco C, López-Bernús A, et al. Recurrence of cystic echinococcosis in an

endemic area: a retrospective study. BMC Infect Dis. 2017;17(1):455.

454 10. El Malki HO, El Mejdoubi Y, Souadka A, Zakri B, Mohsine R, Ifrine L, et al. Does

455 primary surgical management of liver hydatid cyst influence recurrence? J Gastrointest Surg.

456 2010;14(7):1121-7.

457 11. Yosra K, Ghada H, Bochra A, Fatma T, Said J, Riadh J. Predictive Factors of Liver

458 Hydatid Cyst Recurrence in Children. J Pediatr Surg. 2023;58(12):2362-7.

459 12. Bedioui H, Ayari H, Bouslama K, Maghrebi H, Hsairi H, Jouini M, et al. [Recurrence of

460 hydatid cyst of liver: predictive factors: Tunisian experience]. Bull Soc Pathol Exot.

461 2012;105(4):265-9.

462 13. Pacifico J, Cabrera R, Salgado-Diaz S, Mormontoy W, Suarez-Ognio L. Factors

463 associated with pre-surgical complications in patients with cystic echinococcosis from

464 endemic areas of Peru. Rev Peru Med Exp Salud Publica. 2021;38(1):33-40.

465 14. Al-Saeedi M, Ramouz A, Khajeh E, El Rafidi A, Ghamarnejad O, Shafiei S, et al.

466 Endocystectomy as a conservative surgical treatment for hepatic cystic echinococcosis: A

467 systematic review with single-arm meta-analysis. PLoS Negl Trop Dis.

468 2021;15(5):e0009365.

469 15. Abbasi Dezfouli S, El Rafidi A, Aminizadeh E, Ramouz A, Al-Saeedi M, Khajeh E, et

470 al. Risk factors and management of biliary leakage after Endocystectomy for hepatic cystic

471 echinococcosis. PLoS Negl Trop Dis. 2023;17(10):e0011724.

472 16. Al-Saeedi M, Khajeh E, Hoffmann K, Ghamarnejad O, Stojkovic M, Weber TF, et al.

473 Standardized endocystectomy technique for surgical treatment of uncomplicated hepatic

474 cystic echinococcosis. PLoS Negl Trop Dis. 2019;13(6):e0007516.

475 17. Lissandrin R, Tamarozzi F, Mariconti M, Manciulli T, Brunetti E, Vola A. Watch and

476 Wait Approach for Inactive Echinococcal Cyst of the Liver: An Update. Am J Trop Med Hyg.

477 2018;99(2):375-9.

478 18. Junghanss T, da Silva AM, Horton J, Chiodini PL, Brunetti E. Clinical management of

479 cystic echinococcosis: state of the art, problems, and perspectives. Am J Trop Med Hyg.

480 2008;79(3):301-11.

19.     Gerencia Regional de Salud Cusco. Analisis de Situacion de Salud - Canchis Cusco,

Peru: Red Nacional de Epidemiologia - RENACE Peru; 2020 [Available from:

https://www.redsaludcce.gob.pe/Modernidad/archivos/epidemiologia/ASIS/ASIS_Canchis_2

020.pdf.

20.     Instituto Nacional de Estadistica e Informatica (INEI). Censo 2018, Resultados

Definitivos, Region Cusco. Lima, Peru: Gobierno del Peru; 2018.

21.     Guideline on haemoglobin cutoffs to define anaemia in individuals and populations.

WHO Guidelines Approved by the Guidelines Review Committee. Geneva2024.

22.     Ministerio de Salud del Peru. Estadistica Poblacional Lima: Ministerio de Salud del

Peru.; 2023 [Available from: https://www.minsa.gob.pe/reunis/data/poblacion_estimada.asp.

23.     Compania Peruana de Estudios de Mercado y Opinion Publica (CPI Peru).

Estadisticas Poblacionales Peru: CPI Peru; [Available from:

https://cpi.pe/banco/estadisticas-poblacionales.html.

24.     Centro Nacional de Epidemiologia Prevencion y Control de Enfermedades. Boletin

Epidemiologico del Peru: Ministerio de Salud del Peru; 2021 [Available from:

https://www.dge.gob.pe/epipublic/uploads/boletin/boletin_202136_25_150233.pdf.

25.     Lopez-Bernus A, Belhassen-Garcia M, Prieto-Vicente AJ, Alonso-Sardon M, Carpio-

Perez A, Velasco-Tirado V, et al. [Epidemiological update of hydatid disease in hospitals in

the public health system of Extremadura (2003-2012)]. Enferm Infecc Microbiol Clin.

2016;34(4):232-6.

26.     Lopez-Bernus A, Belhassen-Garcia M, Alonso-Sardon M, Carpio-Perez A, Velasco-

Tirado V, Romero-Alegria A, et al. Surveillance of Human Echinococcosis in Castilla-Leon

(Spain) between 2000-2012. PLoS Negl Trop Dis. 2015;9(10):e0004154.

27.     Jimenez S, Perez A, Gil H, Schantz P, Ramalle E, Juste R. Progress in control of

cystic echinococcosis in La Rioja, Spain: decline in infection prevalences in human and

animal hosts and economic costs and benefits. Acta Trop. 2002;83(3):213-21.

28.     Ministerio de Salud - Gobierno de Chile. Informe: Situación de la Equinococosis

Quística/Hidatidosis en Chile, 2018-2022. Chile: Ministerio de Salud - Gobierno de Chile;

2023 [Available from: https://diprece.minsal.cl/wp-content/uploads/2024/03/INFORME-

HIDATIDOSIS-2018-2022.pdf.

29.    Collado-Aliaga J, Romero-Alegria A, Alonso-Sardon M, Muro A, Lopez-Bernus A,

Velasco-Tirado V, et al. Complications Associated with Initial Clinical Presentation of Cystic

Echinococcosis: A 20-year Cohort Analysis. Am J Trop Med Hyg. 2019;101(3):628-35.

30.    Salgado Díaz S, Suarez-Ognio L,  Cabrera R. Características clínicas y

epidemiológicas de la equinococosis quística registradas en un área endémica en los Andes

centrales del Perú. Neotropical Helminthology. 2007;1(2):69-83.

31.    Kuehn R, Uchiumi LJ, Tamarozzi F. Treatment of uncomplicated hepatic cystic

echinococcosis (hydatid disease). Cochrane Database Syst Rev. 2024;7(7):CD015573.

32.    Kabaalioglu A, Ceken K, Alimoglu E, Apaydin A. Percutaneous imaging-guided

treatment of hydatid liver cysts: do long-term results make it a first choice? Eur J Radiol.

2006;59(1):65-73.

33.    Men S, Hekimoglu B, Yucesoy C, Arda IS, Baran I. Percutaneous treatment of

hepatic hydatid cysts: an alternative to surgery. AJR Am J Roentgenol. 1999;172(1):83-9.

34.    Velasco-Tirado V, Romero-Alegria A, Belhassen-Garcia M, Alonso-Sardon M,

Esteban-Velasco C, Lopez-Bernus A, et al. Recurrence of cystic echinococcosis in an

endemic area: a retrospective study. BMC Infect Dis. 2017;17(1):455.

35.    Alzoubi M, Daradkeh S, Daradka K, Shattarat LN, Al-Zyoud A, Al-Qalqili LA, et al.

The recurrence rate after primary resection cystic echinococcosis: A meta-analysis and

systematic literature review. Asian J Surg. 2024.

36.    Azizoglu M, Aydogdu B, Kamci TO, Arslan S, Basuguy E, Bilici S, et al. Analysis of

Complications of a Neglected Disease: 13 Years of Experience with Liver Hydatid Cysts in a

High-Volume Hospital. Medicina (Kaunas). 2024;60(10).

37.    Manterola C, Otzen T, Urrutia S, Group M. Risk factors of postoperative morbidity in

patients with uncomplicated liver hydatid cyst. Int J Surg. 2014;12(7):695-9.

38.    Zinsstag J, Meyer, J.M., Bonfoh, B., Fink, G., Dimov, A. One Health in Human-

Environment Systems: Linking Health and the Sustainable use of Natural Resources. CABI

One Health. 2024;3: 1:1-11.

39.    Bucher A, Dimov, A., Fink, G., Chitnis, N., Bonfoh, B., Zinsstag, J. Benefit-cost

analysis of coordinated strategies for control of rabies in Africa. Nature communications.

2023;14:5370.

546 **Table 1.** General characteristics of 115 patients hospitalized for cystic echinococcosis between
547 2010 and 2019 at the Alfredo Callo Rodriguez Hospital of Sicuani, Cusco, Peru.
548

| Variable[*] – no. (%) | All patients (n=115) | Pre-surgical complications[a] | |
|---|---|---|---|
| | | No (n=63) | Yes (n=48) |
| Age group | | | |
|   <18 years | 34 (29.6) | 18 (28.6) | 15 (31.3) |
|   18-64 years | 67 (58.3) | 41 (65.1) | 25 (52.1) |
|   ≥65 years | 14 (12.2) | 4 (6.3) | 8 (16.7) |
| Female | 69 (60) | 35 (55.6) | 31 (64.6) |
| District of residence | | | |
|   Sicuani | 85 (73.9) | 47 (74.6) | 38 (79.2) |
|   Marangani | 7 (6.1) | 6 (9.5) | 1 (2.1) |
|   Other | 18 (15.7) | 10 (15.9) | 8 (16.7) |
| Referred from primary care center | 56 (48.7) | 38 (60.3) | 18 (37.5) |
| Location of cyst | | | |
|   Only liver | 98 (85.2) | 57 (90.5) | 38 (79.2) |
|   Liver and other intraabdominal | 7 (6.1) | 4 (6.3) | 2 (4.2) |
|   Only lung | 5 (4.4) | 1 (1.6) | 4 (8.3) |
|   Liver and lung | 5 (4.4) | 1 (1.6) | 4 (8.3) |
| Number of cysts | | | |
|   Single cyst | 84 (73) | 49 (77.8) | 33 (68.8) |
|   2 cysts | 20 (17.4) | 9 (14.3) | 9 (18.8) |
|   ≥3 cysts | 11 (9.6) | 5 (7.9) | 6 (12.5) |
| Largest cyst diameter | | | |
|   <5 cm | 3 (2.6) | 1 (1.6) | 2 (4.2) |
|   5-10 cm | 28 (24.3) | 16 (25.4) | 12 (25) |
|   ≥10 cm | 79 (68.7) | 45 (71.4) | 31 (64.6) |
| Anemia severity | | | |
|   No anemia | 57 (49.6) | 33 (52.4) | 24 (50) |
|   Mild | 27 (23.5) | 19 (30.2) | 8 (16.7) |
|   Moderate | 22 (19.1) | 10 (15.9) | 12 (25) |
|   Severe | 3 (2.6) | 0 (0) | 3 (6.3) |
| Ultrasound performed | 107 (93) | 62 (98.4) | 45 (93.8) |
|   Ultrasound staging | | | |
|     Gharbi staging | 75 (65.2) | 48 (76.2) | 27 (56.3) |
|     No staging method | 20 (17.4) | 8 (12.7) | 12 (25) |
|     Other | 12 (10.4) | 6 (9.5) | 6 (12.5) |
| Treatment approach | | | |
|   Surgery + post-ABZ | 57 (49.6) | 37 (58.7) | 20 (41.7) |
|   Surgery + pre- & post-ABZ | 27 (23.5) | 15 (23.8) | 12 (25) |
|   No treatment | 10 (8.7) | 2 (3.2) | 8 (16.7) |
|   Only surgery | 7 (6.1) | 4 (6.3) | 3 (6.3) |
|   Surgery + pre-ABZ | 6 (5.2) | 3 (4.8) | 3 (6.3) |
|   Only ABZ | 3 (2.6) | 2 (3.2) | 1 (2.1) |
| Post-surgical complications | 22 (22.9)[b] | 12 (19) | 10 (20.8) |
| Referred to higher care center | 11 (9.6) | 2 (3.2) | 9 (18.8) |
| Post-surgical recurrence | 16 (16.5)[c] | 8 (12.7) | 6 (12.5) |
| Readmission | 14 (12.7)[d] | 9 (14.3) | 5 (10.4) |

[*] Subjects with no data available in variables "sex" (n=1), "district of residence" (n=5), "largest
cyst diameter" (n=5), "anemia severity" (n=6), "ultrasound performed" (n=5), and "treatment
approach" (n=5). *ABZ,* albendazole, *pre-,* pre-surgical, *post-,* post-surgical.
[a] No data available in 4 patients.
[b] 22/96 surgically treated patients with complete post-surgical documentation; in 5/102 patients
the post-surgical follow-up was incomplete, and in 1/102 the immediate post-surgical follow-up
was incomplete.
[c] 16/97 surgically treated patients with complete charts.
[d] 14/110 discharged patients with complete charts.

**Table 2.** Pre-surgical, intraoperative, and post-surgical complications by anatomical

location of cysts.

| Complication type[*] | n (%) |
|---|---|
| **Pre-surgical (n=48, 41.7%[a])** | |
| Infectious (n=29, 25.2%) | |
|    Purulent cyst fluid | 17 (14.8) |
|    Abscess | 9 (7.8) |
|    Peritonitis | 5 (4.4) |
|    Pneumonia | 5 (4.4) |
|    Cholecystitis | 2 (1.7) |
|    Cholangitis | 1 (0.9) |
| Mechanical (n=23, 20%) | |
|    Fistula | 11 (9.6) |
|      Cysto-biliary | 9 (7.8) |
|      Cysto-pleural-bronchial | 1 (0.9) |
|      Cholecysto-gastric | 1 (0.9) |
|    Cyst rupture or vomica | 10 (8.7) |
|    Mass effect[b] | 5 (4.4) |
| Respiratory (n=5, 4.4%) | |
|    Respiratory insufficiency | 5 (4.4) |
| Hemorrhagic (n=3, 2.6%) | |
|    Hemoptysis | 2 (1.7) |
|    Hemorrhagic cyst fluid | 1 (0.9) |
| Allergic (n=3, 2.6%) | |
|    Rash/urticaria | 3 (2.6) |
| Other (n=3, 2.6%) | |
|    Pregnancy | 1 (0.9) |
|    Abdominal trauma | 1 (0.9) |
| **Intraoperative (n=2, 2%[c])** | |
| Anaphylactic shock | 1 (1.0) |
| Cardiac arrest | 1 (1.0) |
| Hemoperitoneum | 1 (1.0) |
| Spleen laceration | 1 (1.0) |
| **Post-surgical (n=22, 22.9%[d])** | |
| Residual collection | 14 (14.6) |
| Abscess | 3 (3.1) |
| Biliary leakage | 3 (3.1) |
| Cutaneous fistula | 2 (2.1) |
| Adhesions | 1 (1.0) |
| Hemorrhage | 1 (1.0) |
| Post-cardiac arrest | 1 (1.0) |
| Sepsis | 1 (1.0) |
| Wound infection | 1 (1.0) |

[*] Patients may have more than one complication.

[a] 48/115 total patients.

[b] Biliary obstruction (n=3), bowel obstruction (n=1), abdominal

compartment syndrome (n=1).

[c] 2/102 surgically treated patients.

[d] 22/96 surgically treated patients with complete post-operative

documentation.

**Figure 1.** Forest plot showing the univariable and multivariable logistic regression analyses
of variables associated with post-surgical recurrence. Odds ratios and confidence intervals
in black represent univariable logistic regression models for each independent variable.
Odds ratios and confidence intervals in blue represent the multivariable analysis. *ABZ,*
albendazole, *CI,* confidence interval, *npos/N,* number with recurrence/total per category,
*OR,* odds ratio, *Ref.,* reference category. *Including pre- and post-surgical treatment.

**Table 3.** Annual cumulative incidence of new hospital admissions for CE at the Alfredo Callo Rodriguez Hospital of Sicuani during the period 2010-2019.

| Year | Hospital admissions for new CE diagnosis | | | All hospital admissions due to CE[*] | | | Population at risk[§] | Cumulative incidence ×100,000 |
| | Patients | Echinococcal cysts | | Admissions | Echinococcal cysts | | | |
| | n=115 (%) | n=169 (%) | Mean number of cysts per patient (SD) | n=126 (%) | n=185 (%) | Mean number of cysts per patient (SD) | | |
|---|---|---|---|---|---|---|---|---|
| 2019 | 21 (18.3) | 33 (19.5) | 1.57 (1.12) | 22 (17.5) | 35 (18.9) | 1.59 (1.10) | 144,607 | 14.5 |
| 2018 | 16 (13.9) | 21 (12.4) | 1.31 (0.79) | 17 (13.5) | 22 (11.9) | 1.29 (0.77) | 143,826 | 11.1 |
| 2017 | 16 (13.9) | 27 (16.0) | 1.69 (1.78) | 19 (15.1) | 30 (16.2) | 1.58 (1.64) | 143,059 | 11.2 |
| 2016 | 12 (10.4) | 17 (10.1) | 1.42 (1.17) | 15 (11.9) | 20 (10.8) | 1.33 (1.05) | 142,266 | 8.4 |
| 2015 | 20 (17.4) | 31 (18.3) | 1.55 (0.83) | 23 (18.2) | 38 (20.5) | 1.65 (1.07) | 141,444 | 14.1 |
| 2014 | 9 (7.8) | 11 (6.5) | 1.22 (0.44) | 9 (7.1) | 11 (6.0) | 1.22 (0.44) | 141,900 | 6.3 |
| 2013 | 5 (4.4) | 9 (5.3) | 1.8 (0.84) | 5 (4.0) | 9 (4.9) | 1.8 (0.84) | 142,300 | 3.5 |
| 2012 | 9 (7.8) | 11 (6.5) | 1.22 (0.44) | 9 (7.1) | 11 (6.0) | 1.22 (0.44) | 142,800 | 6.3 |
| 2011 | 5 (4.4) | 7 (4.1) | 1.4 (0.89) | 5 (4.0) | 7 (3.8) | 1.4 (0.89) | 143,200 | 3.5 |
| 2010 | 2 (1.7) | 2 (1.2) | 1.0 (0.00) | 2 (1.6) | 2 (1.1) | 1.0 (0.00) | 143,600 | 1.4 |

[*]Including readmissions, but not counting one patient readmitted in 2020 for a liver cyst, one patient readmitted in 2020 for multiple liver and intraabdominal cysts, and two patients readmitted for drainage of post-surgical liver abscesses.
[§]Combining the Canas and Canchis provinces total population.
*CE,* cystic echinococcosis, *SD,* standard deviation.