Improving Access and Efficiency of Acute Ischemic Stroke Treatment Across Four Canadian Provinces: A Stepped-Wedge Trial

Noreen Kamal, PhD[1,2,3], Elena A. Cora, MD PhD[4,5], Simone Alim BSc (In Progress)[6], Judah Goldstien PhD[7,8], David Volders MD[4,5], Shadi Aljendi, PhD[10], Heather Williams MD[11], Patrick T. Fok MDCM PhD[4,7,9] Etienne van der Linde MD[12], Trish Helm-Neima BSc[11], Tania Chandler MD[13], Alix Carter MD[4,7,8], Renee Cashin BSc[14], Brian Metcalfe MD[15], Julie Savoie BNSc[16], Wendy Simpkin BN[4], Fraser Clift MD[14,17], Cassie Chisholm MSc[14], Michael D Hill MD MSc[18], Bijoy K. Menon MD MSc[18], Stephen Phillips MD[3,4] and the ACTEAST Collaborators

Department of Industrial Engineering, Faculty of Engineering, Dalhousie University, Halifax, Canada
Department of Community Health and Epidemiology, Faculty of Medicine, Dalhousie University, Halifax, Canada
Department of Medicine (Division of Neurology), Faculty of Medicine, Dalhousie University, Halifax, Canada
Nova Scotia Health, Halifax, Canada
Department of Diagnostic Radiology, Faculty of Medicine, Dalhousie University, Halifax, Canada
Department of Mathematics and Statistics, Faculty of Science, Dalhousie University, Halifax, Canada
Department of Emergency Medicine, Faculty of Medicine, Dalhousie University, Halifax, Canada
Emergency Health Services, Halifax, Canada
EHS LifeFlight, Nova Scotia, Canada
Faculty of Computer Science, University of New Brunswick, Fredericton, Canada
Health PEI, Charlottetown, Canada
Dr. GB Cross Memorial Hospital, NL Health Services, Clarenville, Canada
Horizon Health Network, Saint John, Canada
Discipline of Emergency Medicine, Faculty of Medicine, Memorial University, St. John's, Canada
Medflight NL, St. John's, Canada
Vitalité Health Network, Moncton, Canada
Department of Medicine, Faculty of Medicine, Memorial University, St. John's, Canada
Department of Clinical Neurosciences, Radiology and Community Health Sciences, Cumming School of Medicine, University of Calgary, Calgary, Canada

Short Title: Improving Stroke Treatment

Corresponding Author:
Noreen Kamal, PhD PEng
Associate Professor
Dalhousie University

Department of Industrial Engineering
5269 Morris Street, Room 100
PO Box 15000
Halifax NS B3H 4R2 Canada
Noreen.Kamal@dal.ca
Phone: +1-902-494-3293

Tables: 3
Figures: 2
Word Count: 6307 (including Title Page, Abstract, Text, References, Tables and Figures Legends)

ACTEAST Collaborators:
Shahram Abootalebi
Brian Archer
Leslie Austin-Smith
Dylan Blacquiere
Ashley Boyce
Greg Browne
Jamie Drapeau
Charles Duffy
Marsha Eustace
Rachel Goss
Kelly Goudey
Edgar Goulette
Gordon Gubitz
Jens Heidenreich
Milan Jurga
Nadia LeBreton
Michelle MacGrath
Carolyn MacPhail
Alier Marrero
Gregory McNally
Jennifer McVey
Matthew Murphy
Susannah Piercey
Amer Qureshi
Nicole Tupper
Scott Theriault
Lindsay Richards
Shauna Sommerville
Janel Swain

Abstract [300 words]

Background: An evidence-to-practice gap exists in acute ischemic stroke treatment, and improvements in access and efficiency of treatment with iv thrombolysis and endovascular thrombectomy (EVT) is needed. The objective of this study was to improve access and efficiency of ischemic stroke treatment across four Atlantic Canadian Provinces.

Methods: A stepped-wedge cluster trial was conducted over 30 months with 3 clusters covering 34 sites. The trial was conducted across all 4 Atlantic Canadian provinces: Nova Scotia (NS), New Brunswick (NB), Prince Edward Island (PE), and Newfoundland and Labrador (NL). The design was quasi-randomized, with each cluster associated with one or more provinces: cluster 1 – NS; cluster 2 – NB and PE; and cluster 3 – NL. The patient population was all ischemic stroke patients across all 4 provinces. The intervention was a 6-month modified Quality Improvement Collaborative (mQIC), which was modified from the Breakthrough Series Collaborative to be half of the 1-year period and conducted virtually. The intervention consisted of assembling an interdisciplinary improvement team, 2 full-day workshops, webinars, and virtual site visits. Suggested changes included 6 process improvement strategies.

Results: The proportion of patients that received treatment did not increase significantly with the intervention [0.4% increase for patients that received thrombolysis and/or EVT (p=0.68)]. Median door-to-needle time was reduced by 9.2 minutes with the intervention (p=0.01). Cluster 3 saw the greatest improvements in both access and efficiency.

Conclusions: A mQIC intervention resulted in improvement of process measures like door-to-needle time. Quality improvement initiatives may need to be longer to see improvements in proportion of patients treated. Tailored interventions for each health system can ensure that each system sees improvement. In-person activities might be critical to ensure fidelity of the intervention.

Introduction

Revolutionary advances in stroke treatment have occurred over the past two decades from thrombolysis using intravenous (IV) alteplase in the mid-1990's[1,2,3] to the most recent ground-breaking trials in 2015 showing the high efficacy of endovascular thrombectomy (EVT).[4,5] These treatments can reduce disability due to stroke, thus reducing the need for assistance with daily living tasks,[7] increase return to the workplace,[8] and reduce hospital costs. [9]

Although treatment of acute ischemic stroke patients with alteplase and EVT are part of guideline care in Canada and around the world,[10,11] less than optimal utilization rates for both of these treatments are observed. This evidence-to-practice gap is not a new phenomenon;[12,13] however, the substantial economic and societal benefits of these therapies continue to make it critical to pursue optimal uptake. This evidence-to-practice gap is exacerbated by both geo-political and socio-political divides. There is a split between urban and rural access to treatment; access to stroke physicians or neurologists is limited in rural settings. Furthermore, access to EVT in rural areas is even more challenging because the treatment is relatively new and requires specialised equipment and personnel. Although both of these therapies mitigate disability, their effectiveness has been shown to be highly time dependent. In stroke treatment, minutes matter,[5,14] bringing to light the mantra, "time is brain".[15] Therefore, there have been calls to reduce hospital thrombolysis treatment times to 30 minutes from hospital arrival,[10,16] and to create seamless transfer processes for more efficient access to EVT for patients living outside major urban centers.[17,18] Atlantic Canada is comprised of four provinces: Nova Scotia (NS), New Brunswick (NB), Prince Edward Island (PE), and Newfoundland and Labrador (NL).

Stroke treatment in Atlantic Canada is particularly challenging because of small populations with a significant proportion that live in rural areas.

Using a stepped-wedge cluster trial design, this study [ACTEAST (Atlantic Canada Together Enhancing Acute Stroke Treatment)] aimed to evaluate if access to and efficiency of acute ischemic stroke treatment with iv thrombolysis and EVT across Atlantic Canada would improve with a *modified Quality Improvement Collaborative (mQIC) intervention*.

Methods

The ACTEAST Project was a 30-month stepped-wedge cluster trial. The trial was quasi-randomised in-so-far as the sites were not randomized to a cluster for pragmatic reasons; rather, each cluster was associated with a province. A quasi-randomisation methodology was used because of the close network of all stroke centers across each province, which would make it difficult to not contaminate each cluster if sites were to be randomised to different clusters. Each cluster had a single center that was capable of EVT treatment with the remainder of sites only capable of thrombolysis treatment; patients at these thrombolysis-only sites who were eligible for EVT were transferred to the EVT-capable center in their province. There were a total of 34 sites with 10 sites in cluster 1, 12 sites in cluster 2, and 12 sites in cluster 3. The stepped-wedge cluster trial design is shown in Figure 1. This trial took place entirely during the COVID-19 pandemic where restriction and additional processes were required including testing all patients for COVID-19. These restrictions also limited the administration of intervention activities to being entirely virtual.

Setting:

ACTEAST was conducted across all four Atlantic Canadian provinces, which are smaller and less populated provinces as compared to the other Canadian provinces and have a combined population of approximately 2.5 million people. The Atlantic provinces have a larger percent of its population living in rural areas: in NS, 43% of its population live in rural areas, 48% in NB; 53% in PE; and 41% in NL. This is more than double other Canadian provinces; for example, only 14%, 17%, 14%, and 19% live in rural areas for Ontario, Alberta, British Columbia, and Quebec, respectively.[19] Atlantic Canada also has an older population.[20] The age and risk standardized stroke incidence rate is greater in Atlantic Canada than in other provinces: up to 140 strokes/100,000 people/year in the Atlantic provinces compared to 113 strokes/100,000 people/year in Ontario.[21] All provinces have a provincially administered universal healthcare system. Each province has developed a provincial stroke system, where specific hospitals have been designated as stroke centers. These designated stroke centers have CT (computed tomography) scanners and expertise to treat patients with thrombolysis. They have access to stroke neurology expertise through telehealth. Ambulances bypass closer hospitals to take patients directly to a designated stroke center, which are distributed across each province to ensure optimal access for the population in each province.

EVT services were available in NS and NB in the cities of Halifax and Saint John respectively. PE has set-up cross-provincial transfer agreement with the EVT-capable hospital in Halifax, NS. NL did not have EVT services until June 20, 2022, which was completed during the trial period. Initially, only local patients had access to EVT, so there were no patients transferred for EVT in NL during the trial period.

Site Enrolment and Patient Population:

Enrolment into the ACTEAST trial was done at a site level. The Principal Investigator (NK) developed relationships with key medical and administrative personnel in all provinces to ensure that all designated stroke centers enrolled. All enrolled sites were required to set-up an interdisciplinary improvement team that consisted of paramedics, emergency department nurses, emergency physicians, stroke physicians (if applicable), stroke nurses (if applicable), CT technologists, radiologists, and administrators. Each team member was enrolled into the study and consented their participation.

Measures:

To measure access to treatment, the proportion of all ischemic stroke patients who received treatment was used. There were 3 separate measures for this: proportion that received iv thrombolysis only; proportion that received EVT only; and proportion that received iv thrombolysis and/or EVT (any treatment). We also assessed the proportion of patients that were transferred for EVT from a thrombolysis-only center.

To assess speed at which treatment was administered (treatment efficiency), all community onset ischemic stroke patients that received treatment were included. Thrombolysis efficiency was assessed using door-to-needle time (time from hospital arrival to the start of thrombolysis treatment) and 911-to-needle time (time from 911 call to start of thrombolysis treatment). EVT efficiency was assessed using door-to-arterial-access time (time from hospital arrival to the start of EVT treatment) and 911-to-arterial-access time (time from 911 call to start of EVT treatment). Transfer efficiency was assessed using door-in-door-out

time, which is the time from arrival at the thrombolysis-only center to departure from this center to the EVT-capable center.

Patient outcomes were measured using a pragmatic approach by utilizing hospital discharge disposition. The outcome measures included proportion of patients discharged home and proportion of patients that died in-hospital. This was done for all ischemic stroke patients and for all treated stroke patients (excluding in-hospital stroke patients).

Intervention:

The intervention used a mQIC methodology adapted from the Institute for Healthcare Improvement's Breakthrough Series Collaborative.[22] The mQIC began with each site formally enrolling and assembling an interdisciplinary team, as described in the *Site Enrolment and Patient Population* section. The mQIC consisted of two full-day workshops, where the enrolled team from each site attended. The first workshop highlighted evidence on treatment of ischemic stroke and strategies for improving efficiency of treatment. The afternoon consisted of each team planning their improvements and reporting it back to the entire group. The second workshop was conducted 6-8 weeks after the first, where teams presented their changes, and the afternoon also had a planning period with report back. During the 6-month intervention period, the sites were supported with virtual site visits and webinars. Full details of the mQIC design and engagement of all sites and personnel have previously been reported.[23]

All enrolled sites and their improvement teams were provided with a change package. This outlined the following changes for each team to trial, adapt, and implement at their site:

1. *Prenotification and stroke team activation*: paramedics pre-notify the hospital of an incoming stroke patient, and paramedics ensure two IV lines are in place; the hospital creates a single call activation of the stroke team.

2. *Rapid registration process*: the hospital adopts a "registration as *unknown*" (similar for trauma patients), pre-registration process by using information from paramedics, or a quick registration process where only minimal information is entered initially.

3. *Patient moved to CT scanner on Emergency Medical Services (EMS) stretcher*: this includes having a thrombolysis kit ready and conducting a quick neurological exam on the way to the scanner.

4. *Not waiting for bloodwork when not indicated for the patient*

5. *Thrombolysis administered in imaging area*

6. *Rapid transfer process for EVT*: adopting a heads-up call to the transport system when presented with a severe stroke.

Data Collection:

Data were collected by different approaches for each province. NS and PE have provincial registries that contained all the necessary data. Data is collected for the registries by a stroke coordinator through chart reviews. EMS data were linked to the registry data to obtain the date and time of the 911 call. Data from NB was collected by conducting chart reviews, which were done either by the data analyst or by the health authority's research office. Data from NL was collected by chart review conducted either by the local stroke coordinator or through Memorial University's Quality of Care NL organization.

Statistical Analysis:

The data for this trial includes all ischemic stroke patients in the entire province regardless of the number of sites that formally enrolled. The trial is therefore a population based study; no sample size was calculated. All missing values for age, sex, and National Institutes of Health Stroke Scale (NIHSS) were imputed using the K-Nearest Neighbour algorithm (k-NN). For the comparison analysis related to all ischemic stroke patients, unadjusted proportion comparison is conducted using χ^2 analysis, while a mixed effects logistic regression with cluster as random effects variable is used for adjusted analysis. For analysis related to all treated patients, unadjusted analysis comparison is conducted using χ^2 for proportions and Wilcoxon rank-sum test for continuous variables. The comparison analysis uses a mixed effects logistic regression for binary/categorical variables and a mixed effects quantile regression for continuous variables. In these analyses, age, sex, and NIHSS are fixed effects variables while cluster was a random effects variable. We did not include the time of the intervention as cluster captures the temporality of the stepped-wedge trial, and the overall intervention period is short. When analyzing results by cluster, we used a simple quantile regression model.

Ethical Considerations and Trial Registration:

Ethical approval for this study was obtained from the NS Health Research Ethics Board (REB# 1025460), Health PEI Ethics Board, Horizon Health Network Research Ethics Board (ROMEO File #: 100906 and RS#: 2020-2893), Vitalité Health Network Ethics Office (ROMEO file number: 101305), and Newfoundland and Labrador Health Research Ethics Board (Researcher Portal File #: 20210255 and Reference #: 2020.074). The trial is registered with the ISRCTN registry (ISRCTN11109800, https://www.isrctn.com/ISRCTN11109800).

Results

Over 30 months from May 1 2020 to October 31 2022, 8594 ischemic stroke patients were included in the analysis for this trial, out of which 1576 (18.3%) patients received acute treatment. There were 3065 ischemic stroke patients in the pre-intervention period of whom 547 (17.8%) received treatment, and 3806 ischemic stroke patients were in the post-intervention period of whom 718 (18.9%) patients were treated. The full details of patient allocation by cluster and period are shown in Figures 1 and 2.

The results for all ischemic stroke patients are shown in Table 1. The adjusted proportion of patients that received treatment did not increase significantly with a 0.4% increase for patients that received thrombolysis and/or EVT (p=0.68). Cluster 3 showed the most significant improvements in proportion of patients treated with a 6.52% increase (p=0.003) in the proportion receiving treatment. Cluster 3 also saw a 39.35% increase (p<0.0001) in the proportion of patients discharged home.

For the results of all acutely treated ischemic stroke patients, Table 2 shows the comparison of baseline characteristics. Overall the pre-intervention and post-intervention periods are similar except the median time from onset to 911 call was 5 minutes faster (p=0.001) in the post-intervention period.

The process and outcomes results for all community onset ischemic patients treated with thrombolysis and/or EVT are shown in Table 3. The only time measure to see a significant reduction is door-to-needle time, which went down to a median of 62.0 minutes (IQR: 38.7-85.9 minutes) from 71.2 minutes (IQR: 43.5-99.5 minutes) (p=0.01). Cluster 1 showed no significant improvements. Cluster 2 showed a 7.5 minute reduction in the median door-to-CT time

(p=0.009), and a 12.6 minute reduction in the median door-to-needle time (p=0.01). Cluster 3 showed the most improvement, and their post-intervention median door-to-needle times was the lowest of all 3 clusters; they went from a median door-to-needle time of 54.5 minutes (IQR: 44.0-67.0 minutes) to a median of 49.0 minutes (IQR: 41.7-56.3 minutes) (p=0.05). This improvement also carried through to the median 911-to-needle time, where the median time dropped by 12.7 minutes (p=0.05). Cluster 3 also saw a 5.81% increase (p=0.0081) in the proportion of ischemic stroke patients that received thrombolysis, and a 6.52% increase (p=0.003) in the proportion of ischemic stroke patients that received thrombolysis and/or EVT.

Discussion

These results show that improvements were minimal and highly dependent on the cluster. Although there were improvements to the door-to-needle times, these improvements are only present in clusters 2 and 3. Furthermore, the overall reduction in median door-to-needle time was only 9.2 minutes, which is much less compared to previous studies that used a similar intervention.[24-26] The door-to-needle initiative in Alberta saw approximately 30-minute reduction in median times;[24] the California initiative resulted in approximately 20-minute reduction;[25] and the Chicago initiative saw a 15.5 minute reduction.[26] None of these studies used a stepped-wedge trial design, which may account for some of the lower impact, as their improvements may have been overstated due to secular trends and other factors. The settings in Alberta, California, and Chicago studies are much more urban than Atlantic Canada; even in the population study for Alberta Canada, where 17% of their population lives in rural areas compared to approximately 50% for Altantic Canada.[19] Additionally, the intervention period was much shorter in this study, which may suggest that more time is needed to allow sites to

implement their changes especially if more patients are being seen outside of urban centers. Furthermore, the entire trial took place during a time of COVID-19 restrictions. Hospitals had additional tasks such as conducting a COVID-19 test on all patients; furthermore, personnel were participating in projects related to mitigating spread of COVID-19, which created a significant distraction from implementing changes to improve acute stroke treatment.[27] In addition, the mQIC was held entirely virtually due to the pandemic, and these results may signal that virtual Improvement Collaboratives are less efficacious, and there is a need for face-to-face workshops and site visits. Previous studies have noted that virtual quality improvement require a longer lead time,[28] so a shorter implementation period combined with virtual delivery in our study may have had a compounding effect on the impact of the intervention.

These results show that the intervention also needs to be tailored for each health system. This study shows that cluster 3 was able to improve the proportion of patients that receive treatment and efficiency of treatment, cluster 2 was able to improve door-to-needle times, and cluster 1 was not able to make any changes. Since this stepped-wedge trial was designed such that each intervention was similar with limited customization, this may have had variable impact on each health system. This shows the need to assess each health system and tailor the implementation strategy for each health system based on their barriers.[29] Furthermore, additional customized supports to each site may have been necessary, which was not done in this intervention.

The process changes were only observed in the door-to-needle metric. This is likely because the improvement strategies were based on improving this single metric. Additional evidence-based strategies to improve door-to-arterial-access and door-in-door-out times are

needed to see improvements in these metrics. Furthermore, Improvement Collaboratives that focus on a single metric are shown to be more successful than those that focus on multiple metrics and improvements.[24, 30]

Although the results showed significant improvements to the proportion of patients discharged home, it is unlikely that the intervention had this level of impact on this pragmatic outcome measures. Although there was an increase in the proportion of patients that received treatment in cluster 3 and improvement in thrombolysis time for clusters 2 and 3, these improvements are unlikely to have seen a 13% improvement in proportion of ischemic stroke patients discharged home. This shows the need to have true outcomes measures such as 90-day mRS (modified Rankin Score) even for quality improvement initiatives.

Conclusions

The intervention to improve access and efficiency of acute ischemic stroke treatment in four Atlantic Canadian provinces made a 9.2-minute improvement in median door-to-needle times. A longer intervention period that is 1-year long and tailored to each health system with in-person workshops and sites visits may result in more improvements.

Author Conflicts of Interest and Disclosures:
NK: reports funding from CIHR, NSERC, Medtronic, Mitacs, NS Health, and NS Department of Education and Early Childhood Development. She has received speaker honorarium for Heart & Stroke Foundation of NB. She is part-owner of DESTINE Health Inc.

Sources of Funding and Role of Funder Statement

This study was funded by the Canadian Institutes for Health Research (CIHR) Project Grant (PJT-169124). A small grant was also received from Medtronic. The study funders had no influence in the design and conduct of the study; collection, management, analysis, and interpretation of

the data; preparation, review, or approval of the manuscript; and decision to submit the manuscript for publication.

Acknowledgements

Owen Parfrey at Memorial University's Quality of Care NL assisted with facilitating data procurement and collection across NL. Jody Rhodenizer at Horizon Health Network was key in obtaining data for all Horizon hospital in NB. We thank the research office at Vitalité Health Network for visiting hospitals in the Vitalité Health Authority in NB. Dr. Jessalyn Holodinsky guided us in understanding the best methodology for statistical analysis.

References

1. Wardlaw JM, Murray V, Berge E, Del Zoppo GJ. Thrombolysis for acute ischaemic stroke. *Cochrane database of systematic reviews*. 2009;(4): 1-142. doi: 10.1002/14651858.CD000213.pub3.

2. National Institute of Neurological Disorders and Stroke rt-PA Stroke Study Group. Tissue plasminogen activator for acute ischemic stroke. *New England Journal of Medicine*. 1995;333(24):1581-1588. doi: 10.1056/NEJM199512143332401.

3. Emberson J, Lees KR, Lyden P, Blackwell L, Albers G, et al. Effect of treatment delay, age, and stroke severity on the effects of intravenous thrombolysis with alteplase for acute ischaemic stroke: a meta-analysis of individual patient data from randomised trials. *The Lancet*. 2014;384(9958):1929-35. doi: 10.1016/S0140-6736(14)60584-5.

4. Goyal M, Menon BK, Van Zwam WH, Dippel DW, Mitchell PJ, Demchuk AM, Dávalos A, Majoie CB, van der Lugt A, De Miquel MA, Donnan GA, et al. Endovascular thrombectomy after large-vessel ischaemic stroke: a meta-analysis of individual patient data from five randomised trials. *The Lancet*. 2016;387(10029):1723-31. doi: 10.1016/S0140-6736(16)00163-X.

5. Goyal M, Demchuk AM, Menon BK, Eesa M, Rempel JL, Thornton J, Roy D, Jovin TG, Willinsky RA, Sapkota BL, Dowlatshahi D., et al. Randomized assessment of rapid endovascular treatment of ischemic stroke. *New England Journal of Medicine*. 2015;372(11):1019-30. doi: 0.1056/NEJMoa1414905.

6. Adamson J, Beswick A, Ebrahim S. Is stroke the most common cause of disability?. *Journal of Stroke and Cerebrovascular Diseases*. 2004;13(4):171-7. doi: 10.1016/j.jstrokecerebrovasdis.2004.06.003.

7. Claesson L, Lindén T, Skoog I, Blomstrand C. Cognitive impairment after stroke–impact on activities of daily living and costs of care for elderly people. Cerebrovascular Diseases. 2005;19(2):102-9. doi: 10.1159/000082787.

8. Daniel K, Wolfe CD, Busch MA, McKevitt C. What are the social consequences of stroke for working-aged adults?: A systematic review. *Stroke*. 2009;40(6):e431-40. doi: 10.1161/STROKEAHA.108.534487.

9. Mittmann N, Seung SJ, Coté R, Hill MD, Mackey A, et al. Impact of disability status on ischemic stroke costs in Canada in the first year. *CJNS*. 2012; 39(6): 739-800. doi: 10.1017/S0317167100015638.

10. Boulanger JM, Lindsay MP, Gubitz G, et al. Canadian stroke best practice recommendations for acute stroke management: prehospital, emergency department, and acute inpatient stroke care, 6th edition, update 2018. *International Journal of Stroke*. 2018; 13: 949-984. doi: 10.1177/1747493018786616.

11. Powers WJ, Rabinstein AA, Ackerson T, Adeoye OM, Bambakidis NC, et al. 2018 guidelines for the early management of patients with acute ischemic stroke: a guideline for healthcare professionals from the American Heart Association/American Stroke Association. *Stroke*. 2018;49(3):e46-99. doi: 10.1161/STR.0000000000000158.

12. Lang ES, Wyer PC, Haynes RB. Knowledge translation: closing the evidence-to-practice gap. Annals of emergency medicine. 2007;49(3):355-63. doi: 10.1016/j.annemergmed.2006.08.022.

13. Mallonee S, Fowler C, Istre GR. Bridging the gap between research and practice: a continuing challenge. Injury prevention. 2006;12(6):357-9. doi:10.1136/ip.2006.014159.

14. Saver JL, Goyal M, Van der Lugt AA, Menon BK, Majoie CB, Dippel DW, Campbell BC, et al. Time to treatment with endovascular thrombectomy and outcomes from ischemic stroke: a meta-analysis. *JAMA*. 2016;316(12):1279-89. doi:10.1001/jama.2016.13647.

15. Saver JL. Time is brain—quantified. *Stroke*. 2006 Jan 1;37(1):263-6. doi:10.1161/01.STR.0000196957.55928.ab.

16. Kamal N, Benavente O, Boyle K, Buck B, Butcher K, Casaubon LK, Côté R, Demchuk AM, Deschaintre Y, Dowlatshahi D, Gubitz GJ, et al.. Good is not good enough: the benchmark stroke door-to-needle time should be 30 minutes. *CJNS*. 2014;41(6):694-6. doi:10.1017/cjn.2014.41.

17. Kamal N, Jeerakathil T, Demchuk AM, Smith EE, Mann B, Buck B, Hill MD on behalf of the QuICR Alberta Stroke Program. Door-in-door-out times across a Canadian Province for Optimal Access to endovascular therapy. *European Stroke Journal.* 2018; 3(1_suppl): 3 – 586. doi:10.1177/2396987318770127.

18. Kamal N, Demchuk AM, Hill MD, Yu AY, Stotts G, et al. Canadian trends in door-in-door-out times for endovascular treatment. *International Journal of Stroke.* 2017;12 Issue 4_suppl:4-84. 2017. doi:10.1177/1747493017721569.

19. Statistics Canada. Canada Goes Urban. https://www150.statcan.gc.ca/n1/pub/11-630-x/11-630-x2015004-eng.htm. Accessed July 29, 2019.

20. Statistics Canada. Chart 3 Proportion of the population aged 15 to 64 within the total population, Canada, provinces and territories, 2016. https://www150.statcan.gc.ca/n1/daily-quotidien/170503/cg-a003-eng.htm

21. Kamal N, Lindsay MP, Côté R, Fang J, Kapral MK, Hill MD. Ten-year trends in stroke admissions and outcomes in Canada. *Canadian Journal of Neurological Sciences*. 2015;42(3):168-75. doi:10.1017/cjn.2015.20.

22. Institute for Healthcare Improvement. The Breakthrough Series: IHI's Collaborative Model for Achieving Breakthrough Improvement. *IHI Innovation Series white paper*. Boston: Institute for Healthcare Improvement; 2003.

23. Kamal N, Aljendi S, Carter A, Cora EA, Chandler T, Clift F, Fok PT, Goldstein J, Gubitz G, Hill MD, Menon BK. Improving access and efficiency of ischemic stroke treatment across four Canadian provinces using a stepped wedge trial: Methodology. *Frontiers in Stroke*. 2022;1:1014480. doi:10.3389/fstro.2022.1014480.

24. Kamal N, Jeerakathil T, Stang J, Liu M, Rogers E, Smith EE, Demchuk AM, Siddiqui M, Mann B, Bestard J, Lang E. Provincial door-to-needle improvement initiative results in improved patient outcomes across an entire population. *Stroke*. 2020;51(8):2339-46. doi:10.1161/STROKEAHA.120.029734.

25. Nguyen-Huynh MN, Klingman JG, Avins AL, Rao VA, Eaton A, Bhopale S, et al; KPNC Stroke FORCE Team. Novel Telestroke Program improves thrombolysis for acute stroke across 21

hospitals of an Integrated Healthcare System. *Stroke*. 2018;49:133–139. doi:10.1161/STROKEAHA.117.018413

26. Prabhakaran S, Lee J, O'Neill K. Regional learning collaboratives produce rapid and sustainable improvements in stroke thrombolysis times. *Circ Cardiovasc Qual Outcomes*. 2016;9:585–592. doi:10.1161/CIRCOUTCOMES.116.003222.

27. Aljendi S, Mrklas KJ, Kamal N. Qualitative Evaluation of a Quality Improvement Collaborative Implementation to Improve Acute Ischemic Stroke Treatment in Nova Scotia, Canada. *Preprints*. doi:10.20944/preprints202407.2211.v1

28. Khurshid Z, De Brún A, Moore G, McAuliffe E. Virtual adaptation of traditional healthcare quality improvement training in response to COVID-19: a rapid narrative review. *Human resources for health*. 2020;18:1-8. doi:10.1186/s12960-020-00527-2.

29. Powell BJ, Beidas RS, Lewis CC, Aarons GA, McMillen JC, Proctor EK, Mandell DS. Methods to improve the selection and tailoring of implementation strategies. *The journal of behavioral health services & research*. 2017;44:177-94. doi:10.1007/s11414-015-9475-6.

30. Schouten LM, Hulscher ME, van Everdingen JJ, Huijsman R, Grol RP. Evidence for the impact of quality improvement collaboratives: systematic review. *Bmj*. 2008;26;336(7659):1491-4. doi:10.1136/bmj.39570.749884.BE.

Figures:

Cluster	Lead in Phase (May/20-Oct/20)	Phase 1 (Nov/20-Apr/21)	Phase 2 (May/21-Oct21)	Phase 3 (Nov/21-Apr/22)	Phase 4 (May/22-Oct/22)	Total
Cluster 1 [NS; n=10]	N(IS)= 663 N(treated) = 145	N(IS) = 664 N(treated) = 124	N(IS) = 699 N(treated) = 129	N(IS) = 682 N(treated) = 137	N(IS) = 621 N(treated) = 135	N(IS)= 3329 N(treated)= 670
Cluster 2 [NB,PE; n=12]	N(IS) = 597 N(treated) = 114	N(IS) = 657 N(treated) = 99	N(IS) = 654 N(treated) = 106	N(IS) = 670 N(treated) = 89	N(IS) = 707 N(treated) = 90	N(IS) = 3285 N(treated) = 498
Cluster 3 [NL; n=12]	N(IS) = 354 N(treated) = 41	N(IS) = 403 N(treated) = 64	N(IS) = 391 N(treated) = 65	N(IS) = 408 N(treated) = 69	N(IS) = 424 N(treated) = 93	N(IS) = 1980 N(treated) = 332
Total	N(IS) = 1614 N(treated) = 300	N(IS) = 1724 N(treated) = 287	N(IS) = 1744 N(treated) = 300	N(IS) = 1760 N(treated) = 295	N(IS) = 1752 N(treated) = 318	N(IS) = 8594 N(treated)=1500

□ Pre-Intervention ▨ Post-Intervention ■ Intervention

Figure 1: Stepped-wedge trial design and timeline. NS: Nova Scotia, NB: New Brunswick, PE: Prince Edward Island, NL: Newfoundland and Labrador; n denotes the number of sites in each cluster; N(IS): number of all ischemic stroke patients; N(treated): number of community onset ischemic stroke patients treated with thrombolysis and/or Endovascular Thrombectomy.

Figure 2: Flow diagram of patient allocation across each period and cluster. DTN: Door-to-needle time, DTAA: Door-to-arterial-access time, DIDO: Door-in-door-out time.

Tables:

Table 1: Proportion of patients treated and discharge disposition for all ischemic stroke patients; adjustments are only made for cluster allocation; treated patients include both community and hospital onset stroke patients.

Entire Cohort	Pre period (Control) n = 3065	Post period (Intervention) n = 3803	p (significance)
	N (%)	**N (%)**	
Received Thrombolysis	474 (15.46%)	620 (16.30%)	0.36
Received Thrombolysis (adjusted)	495 (16.15%)	647 (17.02%)	0.3
Received EVT	115 (3.75%)	196 (5.15%)	0.006
Received EVT (adjusted)	123 (4.00%)	151 (3.98%)	0.9
Received either thrombolysis and/or EVT	547 (17.85%)	718 (18.87%)	0.28
Received either thrombolysis and/or EVT (Adjusted)	570 (18.60%)	723 (19.00%)	0.68
Transferred for EVT	118 (3.84%)	193 (5.07%)	0.017
Transferred for EVT (Adjusted)	127 (4.15%)	145 (3.80%)	0.14
Discharged home	1015 (33.10%)	1569 (41.25%)	< 0.0001
Discharged home (Adjusted)	*1016 (33.15%)*	*1768 (46.50%)*	*< 0.0001*
Died	324 (10.57%)	570 (14.98%)	< 0.0001
Died (Adjusted)	349 (11.39%)	464 (12.20%)	0.3
Cluster 1	**N=663**	**N=2002**	
Received Thrombolysis	132 (19.90%)	383 (19.10%)	0.70
Received EVT	40 (6.03%)	122 (6.09%)	1
Received either thrombolysis and/or EVT	155 (23.37%)	444 (22.17%)	0.55
Transferred for EVT[@]	43 (6.40%)	114 (5.60%)	0.51
Discharged home	236 (35.59%)	699 (34.91%)	0.78
Died	117 (17.64%)	360 (17.98%)	0.89
Cluster 2	**N=1254**	**N-1377**	
Received Thrombolysis	165 (13.15%)	147 (10.67%)	0.056
Received EVT	75 (5.98%)	71 (5.15%)	0.40
Received either thrombolysis and/or EVT	215 (17.14%)	181 (13.14%)	0.004
Transferred for EVT[@]	75 (5.98%)	79 (5.73%)	0.85
Discharged home	543 (43.30%)	616 (44.73%)	0.48
Died	165 (13.15%)	189 (13.72%)	0.71
Cluster 3	**N=1148**	**N=424**	
Received Thrombolysis	*177 (15.41%)*	*90 (21.22%)*	*0.0081*
Received EVT	*0*	*3 (0.70%)*	*0.02*
Received either thrombolysis and/or EVT	*177 (15.41%)*	*93 (21.93%)*	*0.003*
Transferred for EVT	0	0	-
Discharged home	*236 (20.55%)*	*254 (59.90%)*	*< 0.0001*
Died	42 (3.65%)	21 (4.95%)	0.30

Table 2: Baseline characteristics for all community onset ischemic stroke patients that received treatment with thrombolysis and/or endovascular thrombectomy

	Control		Intervention		Significance
All Clusters	n	value	n	Value	
Age mean (SD)	528	70.8 (13.2)	669	70.4 (13.6)	0.64
Sex, women (%)	528	236 (44.7%)	670	303 (45.2%)	0.90
NIHSS, median (IQR)	230	10 (6-14)	242	10 (5-15)	0.60
Onset to door, median (IQR)	493	85.0 (52.0-146.0)	629	85.0 (57.0-130.0)	0.70
Onset to 911, median (IQR)	*364*	*22.0 (8.7-68.0)*	*434*	*17.0 (6.7-45.0)*	*0.001*
Cluster 1					
Age, mean (SD)	145	73.0 (13.3)	402	71.01 (13.0)	0.13
Sex, women (%)	145	68 (46.8%)	402	188 (46.8%)	1
NIHSS, median (IQR)	73	9 (5-12)	147	8 (5-13)	0.84
Onset to door, median (IQR)	144	84.8 (54.5-134.2)	381	85.1 (60.0-127.0)	0.94
Onset to 911, median (IQR)	99	14.2 (7.6-35.7)	279	13.2 (6.0-35.1)	0.40
Cluster 2					
Age, mean (SD)	213	69.8 (13.7)	179	68.8 (14.2)	0.48
Sex, women (%)	213	95 (44.6%)	179	67 (37.4%)	0.18
NIHSS, median (IQR)	102	11 (7-15)	90	12 (7-17)	0.62
Onset to door, median (IQR)	192	82.5 (52.0-142.8)	159	88.0 (55.0-135.0)	0.71
Onset to 911, median (IQR)	167	25.0 (7.0-92.5)	116	20.0 (7.0-63.25)	0.36
Cluster 3					
Age, mean (SD)	170	70.5 (12.2)	89	70.8 (14.3)	0.64
Sex, women (%)	170	73 (42.94%)	90	48 (53.33%)	0.14
NIHSS, median (IQR)	55	10 (6-14)	6	5 (5-9.75)	0.20
Onset to door, median (IQR)	157	88.0 (51.0-150.0)	90	78.5 (45.0-134.0)	0.19
Onset to 911, median (IQR)	98	37.5 (13.0-85.0)	39	32.0 (19.0-74.0)	0.88

Table 3: Treatment times and outcome measures for all community onset ischemic stroke patients treated with thrombolysis and/or endovascular thrombectomy. Adjustments were made for cluster, age, sex and NIHSS. IQR: interquartile range.

	Control		Intervention		Significance
	n	Values	n	Value	
Door to CT, median (IQR) (unadjusted)	464	17.0 (9.0-36.2)	580	17.0 (10.0-28.0)	0.80
Door to CT, median (IQR) (adjusted)	464	21.9 (8.2-40.0)	580	20.6 (11.6-29.3)	0.70
Door to Needle, median (IQR) (unadjusted)	*416*	*73.0 (49.0-109.0)*	*566*	*62.5 (40.0-93.0)*	*< 0.0001*
Door to Needle, median (IQR) (adjusted)	*416*	*71.2 (43.5-99.5)*	*566*	*62.0 (38.7-85.9)*	*0.01*
Door-in-door-out, median (IQR) (unadjusted)	55	148.0 (116.0-189.5)	91	155.5 (127.0-188.5)	0.62
Door-in-door-out, median (IQR) (adjusted)	55	151.0 (100.0-202.0)	91	160.0 (147.0-173.0)	0.80
Door-to-arterial-access, median (IQR) (unadjusted)	106	58 (11.2-101.7)	178	70.0 (30.0-100.0)	0.11
Door-to-arterial-access, median (IQR) (adjusted)	106	54.5(8.2-101.0)	178	50.0 (11.0-88.0)	0.50
911-to-needle, median (IQR) (unadjusted)	330	129.6 (97.0-164.6)	413	125.0 (93.0-157.0)	0.14
911-to-needle, median (IQR) (adjusted)	330	152.0 (44.2-259.8)	413	144.0 (75.4-212.6)	0.62
911-to-arterial-access, median (IQR) (unadjusted)	69	270.0 (147.0-356.0)	88	273.0 (160.0-349.0)	0.96
911-to-arterial-access, median (IQR) (adjusted)	69	256.0 (232.0-280.0)	88	239.0 (191.0-319.0)	0.97
Length of Stay[&] median (IQR)	423	6.0 (3.0-13.0)	520	6.0 (3.0-14.0)	0.30
Length of Stay[&] median (IQR) (adjusted)	423	5.75 (4.8-6.6)	520	6.9 (4.8-9.0)	0.16
Discharged home, n (%)	528	272 (51.5%)	673	282 (41.9%)	0.001
Discharged home, n (%) (adjusted)	528	274 (52.0%)	673	284 (42.1%)	0.002
Mortality at discharge, n (%)	528	84 (15.9%)	673	121 (17.9%)	0.40
Mortality at discharge, n(%) (adjusted)	528	84 (16.0%)	673	119 (17.7%)	0.46
Cluster 1					
Door to CT, median (IQR) (unadjusted)	125	13.0 (8.0-21.0)	341	15 .0 (8.0-22.0)	0.23
Door to CT, median (IQR) (adjusted)	125	13.0 (10.8-15.2)	341	14.8 (13.5-16.0)	0.18
Door to Needle, median (IQR) (unadjusted)	122	57.0 (41.5-83.5)	346	58.0 (37.0-83.0)	0.79
Door to Needle, median (IQR) (adjusted)	122	57.2 (50.7-63.8)	346	58.5 (54.5-62.5)	0.70
Door-in-door-out, median (IQR) (unadjusted)	19	137.0 (114.18-212)	49	153.0 (134.0-187.0)	0.62
Door-in-door-out, median (IQR) (adjusted)	19	136.8 (103.6-169.9)	49	155.0 (134.6-175.6)	0.36
Door-to-arterial-access, median (IQR) (unadjusted)	37	86.0 (56.0-123.0)	106	77.5 (63.0-101.8)	0.42
Door-to-arterial-access, median (IQR) (adjusted)	37	85.7 (69.6-101.7)	106	77.0 (67.7-86.5)	0.37
911-to-needle, median (IQR) (unadjusted)	102	113.9 (89.5-145.3)	279	119.2 (93.8-153.3)	0.30
911-to-needle, median (IQR) (adjusted)	102	113.4 (101.0-125.6)	279	118.5 (111.0-125.9)	0.40
911-to-arterial-access, median (IQR) (unadjusted)	15	213.0 (146.1-332.1)	43	180.0 (135.9-349.1)	0.67

911-to-arterial-access, median (IQR) (adjusted)	15	236.0 (131.0-341.0)	43	203.0 (165.0-241.0)	0.42
Length of Day, median (IQR)[&] (unadjusted)	114	6 (3-14)	300	6 (3-13)	0.67
Length of Day, median (IQR)[&] (adjusted)	114	6.7 (4.7-8.6)	300	6.6 (5.5-7.8)	0.90
Discharged home, n (%) (unadjusted)	145	65 (44.8%)	401	177 (44.1%)	0.80
Discharged home, n (%) (adjusted)	145	63 (43.7%)	401	179 (44.7%)	0.35
Mortality at discharge, n (%) (unadjusted)	145	27 (18.6%)	401	81(20.1%)	0.68
Mortality at discharge, n (%) (adjusted)	145	27 (18.9%)	401	80 (20.4%)	0.70
Cluster 2					
Door to CT, median (IQR) (unadjusted)	188	33.0 (22.8-50.2)	156	26.0 (15.0-42.0)	0.001
Door to CT, median (IQR) (adjusted)	*188*	*32.5 (28.6-36.2)*	*156*	*25.0 (22.4-29.6)*	*0.009*
Door to Needle, median (IQR) (unadjusted)	149	102.0 (82.0-123.0)	131	90.0 (67.0-113.0)	0.005
Door to Needle, median (IQR) (adjusted)	*149*	*102.2 (95.0-109.0)*	*131*	*89.6 (83.0-97.0)*	*0.01*
Door-in-door-out, median (IQR) (unadjusted)	36	148.0 (123.0-194.0)	42	157.0 (119.0-189.0)	0.55
Door-in-door-out, median (IQR) (adjusted)	36	146.8 (127.0-167.0)	42	162.6 (144.0-181.0)	0.26
Door-to-arterial-access, median (IQR) (unadjusted)	69	23.0 (9.0-53.0)	69	22.0 (10.0-79.0)	0.61
Door-to-arterial-access, median (IQR) (adjusted)	69	27.2 (10.0-44.0)	69	21.6 (4.0-38.0)	0.65
911-to-needle, median (IQR) (unadjusted)	129	154.0 (128.0-180.0)	94	151.0 (120.8-189.5)	0.37
911-to-needle, median (IQR) (adjusted)	129	151.6 (143.4-159.7)	94	146.0 (136.5-155.6)	0.38
911-to-arterial-access, median (IQR) (unadjusted)	54	281.0 (149.0-356.0)	45	309.0 (224.0-349.0)	0.36
911-to-arterial-access, median (IQR) (adjusted)	54	280.4 (240.7-320.1)	45	307.1 (263.6-350.6)	0.37
Length of Stay[&] median (IQR) (unadjusted)	171	6 (3-11)	145	6 (3-18)	0.55
Length of Stay[&] median (IQR) (adjusted)	171	6.9 (5.5-8.3)	145	7.6 (5.7-9.4)	0.60
Discharged home, n (%) (unadjusted)	*213*	*108 (50.7%)*	*179*	*47 (26.2%)*	*< 0.0001*
Discharged home, n (%) (adjusted)	*213*	*110 (51.6%)*	*179*	*49 (27.3%)*	*0.01*
Mortality at discharge, n (%) (unadjusted)	213	38 (17.8%)	179	32 (17.8%)	0.95
Mortality at discharge, n (%) (adjusted)	213	36 (16.9%)	179	32 (17.8%)	0.90
Cluster 3					
Door to CT, median (IQR) (unadjusted)	151	9.0 (4.0-17.0)	83	12.0 (6.0-21.0)	0.13
Door to CT, median (IQR) (adjusted)	151	9.6 (8.0-12.0)	78	11.8 (9.0-15.0)	0.24
Door to Needle, median (IQR) (unadjusted)	*145*	*54.0 (40.0-75.0)*	*89*	*50.0 (31.0-73.0)*	*0.05*
Door to Needle, median (IQR) (adjusted)	*145*	*54.5 (44.0-67.0)*	*89*	*49.0 (41.7-56.3)*	*0.05*
Door-in-door-out, median (IQR) (unadjusted)	0	-	0	-	
Door-in-door-out, median (IQR) (adjusted)	0	-	0	-	
Door-to-arterial-access, median (IQR) (unadjusted)	0	-	3	72.0 (69.5-91.0)	
Door-to-arterial-access, median (IQR) (adjusted)	0	-	3	72.0 (69.5-91.0)	
911-to-needle, median (IQR) (unadjusted)	*99*	*103.0 (70.0-146.0)*	*40*	*87.0 (66.0-121.0)*	*0.004*
911-to-needle, median (IQR) (adjusted)	*99*	*102.8 (90.9-114.7)*	*40*	*90.1 (70.9-109.3)*	*0.05*

911-to-arterial-access, median (IQR)	0	-	0	-	
Length of Stay[&] median (IQR) (unadjusted)	138	5 (3-12.50)	84	7 (3-22)	0.16
Length of Stay[&] median (IQR) (adjusted)	138	4.7 (3-6)	75	7.5 (5-9)	0.08
Discharged home, n (%) (unadjusted)	170	99 (58.2%)	93	58 (62.3%)	0.60
Discharged home, n (%) (adjusted)	170	100 (59.0%)	93	57 (61.2%)	0.79
Mortality at discharge, n (%) (unadjusted)	170	19 (11.17%)	93	9 (9.6%)	0.86
Mortality at discharge, n (%) (adjusted)	170	17 (10.2%)	93	11 (11.8%)	0.80

[&]LOS does **not include** patients that died in hospital

Cluster	Lead in Phase (May/20-Oct/20)	Phase 1 (Nov/20-Apr/21)	Phase 2 (May/21-Oct21)	Phase 3 (Nov/21-Apr/22)	Phase 4 (May/22-Oct/22)	Total
Cluster 1 [NS; n=10]	N(IS)= 663 N(treated) = 145	N(IS) = 664 N(treated) = 124	N(IS) = 699 N(treated) = 129	N(IS) = 682 N(treated) = 137	N(IS) = 621 N(treated) = 135	N(IS)= 3329 N(treated)= 670
Cluster 2 [NB,PE; n=12]	N(IS) = 597 N(treated) = 114	N(IS) = 657 N(treated) = 99	N(IS) = 654 N(treated) = 106	N(IS) = 670 N(treated) = 89	N(IS) = 707 N(treated) = 90	N(IS) = 3285 N(treated) = 498
Cluster 3 [NL; n=12]	N(IS) = 354 N(treated) = 41	N(IS) = 403 N(treated) = 64	N(IS) = 391 N(treated) = 65	N(IS) = 408 N(treated) = 69	N(IS) = 424 N(treated) = 93	N(IS) = 1980 N(treated) = 332
Total	N(IS) = 1614 N(treated) = 300	N(IS) = 1724 N(treated) = 287	N(IS) = 1744 N(treated) = 300	N(IS) = 1760 N(treated) = 295	N(IS) = 1752 N(treated) = 318	N(IS) = 8594 N(treated)=1500

Pre-Intervention   Post-Intervention   Intervention