Health professionals' beliefs and attitudes towards preconception care: A systematic review

Authors

Cherie Caut[1], Danielle Schoenaker[2,3,4], Erica McIntyre[5,6], Amie Steel[1]

1. Australian Research Centre in Complementary and Integrative Medicine, School of Public Health, Faculty of Health, University of Technology Sydney, Building 10, Level 8, 235-253 Jones Street, Ultimo NSW 2007 Australia.
2. School of Human Development and Health, Faculty of Medicine, University of Southampton, Southampton, UK.
3. MRC Lifecourse Epidemiology Centre, University of Southampton, Southampton, UK.
4. NIHR Southampton Biomedical Research Centre, University of Southampton and University Hospital Southampton NHS Foundation Trust, Southampton, UK.
5. Institute for Sustainable Futures, University of Technology Sydney, Building 10, Level 10, 235-253 Jones Street, Ultimo NSW 2007 Australia. Visiting Fellow
6. Research Institute for Innovative Solutions for Well-being and Health, University of Technology Sydney, Ultimo NSW, Australia.

Abstract

Background

Health professionals have previously identified various barriers and factors that would help facilitate preconception care services in healthcare settings. Clinically relevant preconception information and clinical practice guidelines have since been developed to aid health professionals in preconception care delivery. This systematic review aimed to 1) synthesise recent literature (past five years) describing health professionals' beliefs and attitudes towards preconception care services or programmes and 2) determine if the experience of health professionals providing preconception care has changed compared to literature reviews conducted more than five years ago.

Methods

Five databases were searched between 27/01/2016 and 27/01/2022. Primary quantitative and qualitative research studies were included if they examined health professionals' beliefs and attitudes towards delivering preconception care services or programmes. Study quality was assessed using the CASP Checklist (qualitative studies) and AXIS tool (quantitative studies). Data synthesis used thematic categorisation adapted from the framework approach.

Results

Twenty-one studies were included (n=8 qualitative, n=11 quantitative, n=2 mixed-

methods studies). Methodological quality was generally good for qualitative studies

but varied for quantitative studies. The results covered three categories: 1)

addressing preconception care health literacy (i.e. lack of knowledge, awareness,

training and resources), 2) *clinical practicalities of preconception car*e (i.e. need for

coordination of care and clarity on role responsibility), and 3) *the role of the patient*

(i.e. need for public health education to support patient-led conversations).

Conclusions

Little has changed regarding the barriers and facilitators to providing preconception

care reported by health professionals. To improve the provision of preconception

care, there is a need to co-develop professional and public preconception health

education, clinical resources, and a coordinated preconception healthcare service

model.

Keywords

preconception care, health professionals, beliefs and attitudes, systematic review

59
60

Introduction

Women's and men's modifiable preconception health (PCH) and health behaviours—

such as body composition, lifestyle behaviours, nutrition, environmental exposures,

and birth spacing—can affect maternal and child health[1-3]. The need to address

such risk factors has led to clinical and public health measures that aim to screen for

PCH risks and implement intervention strategies to optimise PCH and the health

behaviours of prospective parents before conception, known as preconception care

(PCC)[4]. The importance of PCC is highlighted by health policies and strategies

produced by a number of countries around the world[5]. It has received dedicated

attention from the World Health Organization[4] as a critically important

component of healthcare that can impact multiple generations.

While PCC is valuable and important for prospective parents, healthcare providers

experience a range of barriers to implementing PCC services or programmes[6, 7].

Previous systematic reviews that included studies conducted more than five years

ago have reported health professionals' (HPs) experiences providing PCC and

describe the types of barriers they experience, including but not limited to poor

interprofessional communication, insufficient clinical time, funding, clinically

relevant information, and public and HP awareness of the benefits of PCC[6, 7].

Conversely, HPs reported that adequate knowledge of PCC, prospective parents

discussing their intention to become pregnant or requesting PCC, and clinical PCC

resources would enable them to facilitate PCC[7]. To address some of these barriers

and promote PCC services, clinician-focussed PCC information and guidelines have

been developed[8, 9], and relevant information for implementing PCC programmes

in organisations[10]. A recent systematic review, however, identified that existing

clinical practice guidelines (CPGs) on PCC supported by high-quality evidence are

lacking[11]. Emerging research efforts aim to establish population PCH priorities and

co-develop strategies to address them at a healthcare services level[12].

While these guidelines aim to assist HPs providing PCC, an up-to-date

understanding of HPs' views towards PCC delivery is needed to identify if the

increasing availability of guidelines has reduced or changed previously reported

barriers and if HPs also require other types of support to implement PCC

meaningfully. Therefore, this systematic review aimed to 1) synthesise recent

literature describing health professionals' beliefs and attitudes towards

preconception care services or programmes and 2) compare these experiences to

previous systematic review findings[6, 7].

Methods

100   This systematic review is reported following the Preferred Reporting Items for

101   Systematic Reviews and Meta-analyses 2020[13] and was prospectively registered on

102   the International Prospective Register of Systematic Reviews (PROSPERO;

103   CRD42021249386).

Eligibility criteria

106   Original primary research (quantitative and qualitative studies) published between

107   27/01/2016 and 27/01/2022 that sampled HPs (including, but not limited to, general

108   practitioners, midwives, and obstetricians) and examined their beliefs and/or

109   attitudes towards delivering PCC services or programmes were eligible for inclusion.

Information sources

Keyword and MeSH terms were employed in the databases MEDLINE (OVID), EMBASE (OVID), Maternity and Infant Care (OVID), CINAHL (EBSCO), and PsycINFO (EBSCO), with the following limits: title and abstract, studies in humans, published within a five-year limit up to 27/01/2022. No limits to language were applied. The search strategy is presented in Supplementary File 1.

Selection process

The search was conducted on 27 January 2022. CC completed electronic database searches and downloaded citations and abstracts into EndNoteX9 citation management software. Duplication screening occurred before citations were exported into Covidence systematic review software[14]. Articles were screened by title and abstract by CC, with 10% of studies screened for reliability by DS. Full-text articles were downloaded and screened by CC, with 10% of studies screened for reliability by AS before final inclusion for review. Disagreements were discussed until a consensus was reached. A third reviewer (DS, AS or EM) was invited to adjudicate if unresolved. References lists of the included studies were searched for additional eligible studies. Article exclusion reasons from the full-text screening stage were recorded.

Data collection process

CC extracted data from eligible studies into a customised form, with data from 10% of studies extracted by AS to establish reliability.

135 Data items

136 Data extracted included the study reference, title, aims, type of PCC service, location,

137 population, study design, data collection method(s), sample size, and findings on HP

138 beliefs and attitudes towards PCC.

139
Critical appraisal

The Critical Appraisal Skills Programme (CASP) Qualitative Checklist[15] evaluated

study reporting and methodological quality. Quantitative studies were assessed via

the Appraisal tool for Cross-Sectional Studies (AXIS)[16]. CC critically appraised

included studies, and AS assessed 10% of included studies for reliability.

Disagreements were discussed until a consensus was reached. A third reviewer (EM

or DS) was invited to adjudicate if unresolved.

147
Synthesis methods

A framework approach for applied and policy-relevant research was employed to

analyse the data, identify common themes across the studies, and then categorise

these findings to determine the key themes[17]. Findings on beliefs and attitudes

towards PCC from the included studies were extracted by CC reading and re-reading

each paper. Common themes were then identified across the studies and further

defined into categories. Each study was assigned as many categories as relevant to

their reported findings.

Results

157 Study selection

158 Searches retrieved 394 articles. After removing duplicates, 237 titles and abstracts

159 were screened (Figure 1). Thirty-one full-text articles were checked for eligibility; 21

160   met the inclusion criteria and were included in this review. Reasons for exclusion at

161   full-text screening are presented in Supplementary File 2.

162
163   **[Insert Figure 1. Prisma flow diagram]**

164
Study characteristics

All articles (n=21) were published in English between 2017 and 2022 (Table 1).

Eleven studies[18-28] employed a cross-sectional study design and survey method

for data collection. Eight qualitative studies were identified utilising focus groups

(n=2)[29, 30] or interviews (n=6)[31-36]. Two studies[37, 38] collected data using

mixed methods; Devido et al. [37] employed a focus group and survey, and Mass et

al. [38] a survey and interactive workshop. The included studies were conducted in

the United States (US; n=4)[18, 25, 36, 37]; Australia (n=4)[21-24], the Netherlands

(n=3)[30, 32, 38]; and one study each in New Zealand[19], India[20], Europe[31],

Indonesia[29], Canada[26], United Kingdom (UK)[33], Malawi[34], South

Africa[27], Nigeria[35], and China[28]. A range of HPs were represented, including

but not limited to nurses, midwives, general practitioners, obstetricians and

gynaecologists. Sample size varied among the qualitative (n=16 to 92), quantitative

(n=77 to 992) and mixed-methods (n=48 to 299) studies.

179
180   **[Insert Table 1. Characteristics of included studies]**

181
Critical appraisal of included studies

Quality assessment details are presented in Table 2 and Table 3 for qualitative and

quantitative studies, respectively.

Most of the qualitative studies fulfilled all the CASP checklist[15] criteria for

assessing methodological quality. One study[34] needed to be more transparent on

the appropriateness of the recruitment strategy, consideration of researcher bias and

rigour of data analysis. The methodological quality of the quantitative studies was

varied[16]. Sample size justification, non-response bias and funding sources or

conflicts of interest were the most poorly reported criteria, followed by the use of

instruments or measurements that had been trialled, piloted, or published

previously.

[Item Table 2. CASP Quality Assessment]

[Item Table 3. AXIS Quality Assessment]

197
198 Results of syntheses

199 Three common thematic categories across the findings from the reviewed studies

200 were identified and defined: *addressing preconception care health literacy, clinical*

201 *practicalities of preconception care*, and *the role of the patient*.

202
203 **Addressing preconception care health literacy**

204 Seventeen studies reported findings related to addressing PCC health literacy[18-26,

205 28-31, 33, 34, 36, 37]. These studies addressed PCH knowledge, clinical training

206 needs and access to resources.

207
Preconception health knowledge gaps

Insufficient knowledge about PCH as a barrier to HPs' PCC practices was commonly

reported. For example, an Australian cross-sectional study of GPs (n=110) found

their lack of knowledge of PCC guidelines to be one of the most common barriers to

their delivery of PCC[24]; this view was shared by 40% of midwives in a second

Australian study[21] and a broader group of HPs from a UK qualitative study.[33]

The lack of knowledge as a barrier to PCC provision also applied to GPs (n=304) in

Australia providing care to men[23], of whom almost all respondents (90%)

indicated that they did not feel confident about their knowledge of the factors

affecting male fertility[23]. In contrast, a study[28] of HPs in China found that

26.9% of those providing care to women with type 1 diabetes felt they lacked the

knowledge to deliver PCC to this population. At the same time, a US study reported a

significant positive association between teaching self-efficacy and preconception

counselling knowledge among parish nurses (n=48) providing diabetes education to

women with diabetes[37]. Only one cross-sectional study sampling dentists and

gynaecologists (n=300) from India investigated clinician characteristics associated

with PCH knowledge[20]. It was found in logistic regression analyses that older age

doctors (Adjusted odds ratio [AOR]=1.055 (95% Confidence interval [CI]: 1.055-

1.092, p=0.002), the field of specialisation - dentists (AOR=1.635, 95% CI: 1.064-

2.511, p=0.002), and years of practicing experience > 5 years (AOR=2.254, 95% CI:

1.46-3.45, p=0.001) were significantly associated with higher knowledge levels of

periodontal health and adverse pregnancy outcomes[20].

Awareness of preconception risks and guidelines

HPs reported needing to be made aware of PCH risks and of existing CPGs designed

to assist them in screening for PCH risks and delivering PCH interventions[19-21, 23,

24, 26, 37]. This issue was highlighted in a study involving dentists and

gynaecologists (n=300), where only 79% were aware of the association between

periodontal health and preterm low birth weight[20]. In Australia, a study of GPs

(n=110) found that only 53% were aware of PCC-CPGs[24]. Even where HPs reported

being aware of guidelines, familiarity with the recommendations may pose another

barrier, as indicated in an Australian cross-sectional study where fewer HPs

(including GPs, obstetricians and gynaecologists, midwives and dietitians) were

aware of the recommended dose (38%) or duration (44%) of preconception iodine

supplementation[21]. This was despite 71% of those HPs being aware that the

National Health and Medical Research Council recommends this intervention[21]. In

this Australian study, awareness of the recommendation was positively associated

with recommending iodine supplements[21]. In contrast, in a cross-sectional study

of GPs (n=200) in New Zealand (NZ), most GPs reported being aware of the risks of

obesity in pregnancy, over 50% of these GPs reported practice that was inconsistent

with guidelines, and only 12% of these same GPs were aware of the New Zealand

Ministry of Health Guidance for Healthy Weight in Pregnancy[19]. A similar finding

was found in a Canadian cross-sectional study assessing periconceptional folic acid

recommendations of GPs (n=77), where only half knew the correct dose and duration

of folic acid for low-risk women[26].

253
The desire for further training

HPs reported wanting more information and education on PCH risks and

interventions[21-23, 30, 34] to improve their confidence in delivering PCC[22, 23].

For example, in a cross-sectional study of Australian health professionals (n=396),

HPs (including GPs, obstetricians and gynaecologists, midwives and dietitians)

indicated they would be interested in receiving new information relating to

iodine[21], and in another Australian cross-sectional study almost all maternal child

and family health nurses (MCaFHN) (n=192) agreed that more information and

formal education opportunities on the topic would increase their confidence to

discuss PCH[22]. A majority (54.6%) of Australian GPs (n=304) suggested that more

information and education about factors that affect male fertility would make them

more confident to talk to male patients about fertility[23]. Some studies specifically

identified lack of training as the barrier to providing PCC and an increased likelihood

of HPs providing PCH information and PCC to their patients if they had received

further training[18, 25, 28, 29, 31, 36]. For example, in a cross-sectional study from

the US, 15% of GPs (n=443) reported they needed training before implementing

PCC[25]. In another cross-sectional study from the US, obstetricians and

gynaecologists (n=297) reported offering expanded carrier screening more

commonly if they were fellowship-trained (80%) compared to those who were not

(70%)[18].

274
Access to resources

Nine studies reported findings related to clinician access to resources supporting

PCC delivery[21-24, 33, 36, 37]. The types of resources participants suggested

include sample meal plans[21], screening guidelines, relevant research findings[21],

online tools[22], factsheets, trustworthy websites[22, 23], waiting room posters[23],

checklists, brochures[24], expert clinician champions, and medical record

prompts[36]. Two other qualitative studies reported structural resource

requirements such as PCC policy and procedures used to support them to implement

PCC programs into clinical practice settings, as reported in a study of HPs (n=92)

from the US[36] and a funding model through third-party reimbursement to

improve the delivery of PCC as described by HPs (n=30) in a study from the

Netherlands[30].

Clinical practicalities of preconception care

Fifteen studies reported findings related to the clinical practicalities of PCC[20-25, 27-30, 32, 34-36, 38].

Coordination of care

In some studies, HPs reported that a lack of interprofessional referral and care coordination was a barrier to providing PCC[20, 28-30, 32, 36, 38]. One cross-sectional study of dentists and gynaecologists (n=300) from India, for example, found that gynaecologists rarely refer to dentists despite being aware of the link between periodontal health and preterm low birth weight, and only 12% of gynaecologists referred patients to the dentists in the preconception period.*[20]* Several studies[20, 28-30] reported HPs describing a desire for an integrated, multi-disciplinary approach to providing PCC services, greater coordination and referral networks between HPs[30, 32, 36, 38] and emphasised the importance of patient follow-up[36]. In a study from the Netherlands, HPs (n=299) suggested that while the responsibility for providing PCC consultations is best suited to primary care, many other HPs involved may act as referrers towards PCC[38].

Role responsibility

Most HPs in the included studies believed it to be part of their role to provide preconception risk screening, PCH promotion and provision or referral for PCH interventions. These include GPs[23, 24, 38], MCaFHNs[22], midwives[38], nurses[27] and specialist physicians[35]. In a qualitative study from the Netherlands, HPs (including midwives, obstetricians and gynaecologists, fertility specialists, GPs, preventive child health care workers, maternity health care providers, physiotherapists, pharmacists and dieticians) (n=30) expressed that the provision of

PCC is challenging due to unclear allocation of responsibilities[30]. Although HPs

(including midwives, nurses, obstetricians and gynaecologists and GPs)

acknowledged that the responsibility to provide PCC consultations lies with all

professions[34] in an Australian cross-sectional study, 84% of GPs (n=110) reported

that they should be the primary providers of PCC[24]. In a mixed-methods study

from the Netherlands, HPs (including nurses, midwives, GPs, physiotherapists,

preventive child healthcare professionals, dieticians, policy officers, maternity care

assistants, and gynaecologists) (n=299) suggested that the responsibility for

providing PCC consultations lies within primary care, mainly GPs (95.6%) and

midwives (94.4%)[38]. In this same study, HPs found it significantly more

challenging to start a conversation about a wish to conceive than midwives (26.8%

versus 20.2%, p=0.006)[38]; they felt less competent to provide preconception

information (32.3% versus 15.1%, p=<0.001)[38]. Similar findings were reported in a

cross-sectional study from the US (n=443), where most GPs (88%) felt pregnancy

intention screening should be routinely included in primary care[25]. In contrast, in

a qualitative study from Nigeria, HPs (including nurses and specialist physicians)

(n=26) stated that PCC services should be offered at all levels of health care with

referral when needed[35]. Specialist physicians from this Nigerian study also

identified the relevance of PCC to their practice, stating that those with chronic

diseases would benefit more[35].

333
Clinician time

HPs report that one of the barriers to providing effective PCC is an insufficient

amount of consultation time[21, 22, 24, 29]. For example, in an Australian cross-

sectional study[21], the main reason for not discussing dietary sources of iodine with

women was insufficient time[21]. Time constraints were also reported by MCaFHNs

(n=192)[22], GPs (n=110)[24] and a broader range of HPs (n=32)[29], as the most

frequently endorsed barrier to promoting PCH. Several of the MCaFHNs (n=192)

suggested that adding a scheduled visit dedicated to interconception health advice to

those who want it would be helpful[22].

The role of the patient

Seven studies reported findings related to the role of the patient[18, 23, 32-34, 36,

38].

347
Community education and health promotion

Three studies indicated that HPs would like to see an increase in public health

education that improves their patients' knowledge of PCH risk factors and the

benefits of PCC[18, 32, 38] and perceive the absence of such community awareness

as a barrier to PCC[32]. Most (56%) obstetricians and gynaecologists (n=297)

believed that offering expanded carrier screening to their patients should be

restricted to those diseases important to the couple, and 52% believed that screening

should only occur when the clinical significance of each disease being screened for is

understood by the couple[18]. In a qualitative study from the Netherlands, HPs

(including midwives, GPs, obstetricians and gynaecologists, cardiologists and

gastroenterologists) (n=20) stated that barriers affecting the uptake and delivery of

PCC included their belief that most future parents lacked awareness of the benefits of

PCC[32].

361
Patient-led conversations

Despite believing that PCC is a part of their role, HPs also perceive that the onus of

responsibility to seek PCH information and PCC is shared with the patient and that

improved public education on PCH risks and health behaviours and the benefits of

PCC would support this to occur more frequently in the absence of 'routine' PCH risk

screening and PCH promotion[23, 33, 34, 36]. For example, in an Australian cross-

sectional study[23], approximately half of GPs' (n=304) stated that they discuss

fertility with male patients 'opportunistically' when consulted about a reproductive

health matter when a 'patient specifically asks for advice' and when consulting with a

couple who 'plan to have children;' very few said that they raise the subject with men

'routinely'[23]. These findings were similar to a qualitative study[36] from the US

where the frequency with which HPs (n=92) engaged in conversations about

reproductive goals ranged from a 'routine' component of each visit to 'episodically' or

only in response to patients' 'question' or 'request.' In a qualitative study from Africa,

all the HPs (n=20) felt that women also have a role in demanding PCC services and

seeking the services[34].

Discussion

Findings from this systematic review provide up-to-date insights into the areas that

provide opportunities to improve PCC delivery within healthcare settings based on

HP's beliefs and attitudes. These include professional and public health education to

increase PCH knowledge and awareness, clinical resources to support HPs providing

PCC, and HP referral networks to support effective PCC delivery to prospective

parents.

386
One of the main findings of this review is that, overall, HPs report low PCH literacy

yet would like to receive training, suggesting that they may benefit from PCH

education. Our findings add to those reported in previous systematic reviews (e.g.,

HPs wanting more training to improve confidence to provide PCC and lack in

knowledge of PCH is a barrier to PCC)[6, 7] indicating little has changed and the

PCH education needs of HPs remains to be addressed. There is a high variability of

the content and recommendations across existing PCC-CPGs and a need for

guidelines that are yet to address all clinical PCC areas[11]. Comprehensive PCC-

CPGs that address all evidence-based PCC areas are needed to assist HPs in

delivering PCC[11]. However, more than simply developing CPGs to manage the PCH

knowledge of HPs is required. As supported by this review's findings, clinicians often

lack awareness of existing guidelines. Interventions that support the knowledge of

HPs and effect changes to practice behaviour are needed. Training interventions

would benefit from nuanced consideration of the skills, knowledge, time, and

funding for targeted HP groups. PCH knowledge may also need to be tiered so

foundational training can be included in clinical degrees, with more advanced

training available for qualified clinicians. To tailor and co-develop such interventions

to specific HPs, further understanding of their knowledge gaps and their current and

potential scope of practice concerning PCC is needed.

406
In this review, some HPs reported believing that the onus of responsibility should be

with the patient to seek PCH information and PCC without routine PCH risk

screening and PCH promotion. This belief assumes patient awareness of PCH risk

factors and the benefits of PCC. With the current level of public health education and

understanding of PCH risk factors, this belief may need to be challenged. Similar

findings have been reported by HPs in previous systematic reviews (e.g., client's lack

of awareness of PCC, the benefits of PCC and initiation for PCC)[6, 7]. A focus on

public education of PCH is also needed to increase the public's PCH literacy.

Adherence to guidelines during preconception is an issue previously described in a

systematic review reporting on dietary guideline adherence during preconception

and pregnancy[39]. This review also highlighted the importance of acknowledging

the influence of demographic and social factors on guideline adherence[39].

Implementation interventions are needed that support knowledge translation and

changes to health behaviour through strategies that improve whole population health

through the life course (e.g., PCH school education) as well as targeted interventions

to those life course phases where becoming pregnant is more likely or possible (e.g.,

PCH information resources and support tools) or intended (e.g., PCH education and

counselling programmes)[40]. As such, Hall and colleagues' proposed model for PCC

integrates education, digital health interventions, campaigns and social media for

raising PCH awareness among HPs and the public and includes the individualised

and specialised provision of PCC by a range of HPs able to provide clinic-based

counselling, motivational interviewing, provision of supplements and

interconception interventions[5].

430
Some HPs report that the need for clinical PCH resources and time are persistent

barriers to providing effective PCC. HPs would like access to PCH information

resources to support them in delivering PCC. Specifically, HPs identified a range of

resources they would find helpful in clinics, from clinician checklists and clinician

websites to patient information factsheets and patient websites. Clinicians report

that lack of consultation time to provide effective PCC is a barrier, and, in some

instances, clinicians would like to be better funded for PCC by receiving third-party

reimbursement. Similar findings have been reported by HPs in previous systematic

reviews (e.g., a lack of clinical time for PCC and reimbursement for that time,

needing physical space or clinically relevant information such as PCC tools checklists

and PCC guidelines)[6, 7]. Clinical resources are available, providing organisations

and clinicians with information pertinent to implementing PCC programmes[10] and

PCC-CPGs (e.g., PCH risk screening checklist or tool and PCC intervention

guidelines)[9], yet high-quality guidelines on PCC are lacking[11]. Future focus is

needed on assessing the PCH behaviour outcomes associated with healthcare

services that implement PCC programmes and when clinicians can access clinically

relevant PCH information resources and PCC-CPGs informed by high-quality

evidence.

449
450
This review also found that many HPs believe they have a role in PCC. However, they

also attest that primary care HPs (primarily GPs) should be the leading providers.

There are two issues needing further consideration. First, a precedent exists for PCC

to be provided by a broader range of HPs, with the opportunity to improve referral

networks. Second, these HPs may be under the assumption that the patient's GP has

already provided PCC. These findings are barriers to delivering PCC and were

reported by HPs in previous systematic reviews (e.g., a lack of clarity regarding who

is responsible for providing PCC[6, 7] and poor communication between HPs)[7]. A

PCC model that supports collaboration between HPs and utilisation of all capable

HPs[41-43] provides an opportunity to deliver PCC that improves PCH outcomes

throughout the community, including at-risk populations and across the life

course[12]. Meaningful involvement of all key stakeholders is required to co-design

PCC healthcare services that support a coordinated healthcare workforce[42, 43].

Further research is needed to understand the outcomes of an integrated PCC

healthcare services delivery model on PCH risks, PCH behaviours and maternal and

child health outcomes[5].

467
468

Limitations

Although the studies in this review represent a wide range of HPs from varied countries or regions, not all HPs who may have a role in PCC were represented in the study populations. The methodological quality of the qualitative studies was good. However, the quantitative study quality was varied. The heterogeneous nature of the study aims and data collection methods precluded the pooling of data for meta-analysis. We should also acknowledge that three studies[19, 30, 32]were included in a previous review[7]. The research question driving our study had a broader focus than the review by Goossens and colleagues and identified an additional 18 papers; however, to avoid overlapping between the two reviews, the findings of the three[19, 30, 32] articles were reported only when other papers in our review shared similar results.

Conclusions

HPs report insufficient knowledge about PCH, lack PCH training, and want education on PCH risks and interventions to improve their confidence in providing PCC and enhanced access to clinical PCH information resources. One of the barriers they experience to providing effective PCC is an insufficient amount of consultation time. Most HPs believe it is their role to provide PCC, and referral networks could improve the delivery of coordinated preconception interventions to identify and support patients with preconception risks. HPs want increased public health education that improves patient awareness of PCH risks and benefits of PCC. These barriers and facilitators to the provision of PCC reported by HPs in studies conducted in the past five years have remained very similar when compared to previous reviews of earlier studies on the topic. To improve the provision of PCC

going forward, there is a need to co-develop professional and public PCH education,

PCC clinical resources, and an integrated PCC healthcare service model.

List of abbreviations

AXIS - Appraisal tool for Cross-Sectional Studies

CASP - Critical Appraisal Skills Programme

CPG – Clinical practice guideline

GP – General Practitioner

HP – Health professional

MCaFHN – Maternal child and family health nurses

PCC – Preconception Care

PCH – Preconception health

PROSPERO - Prospective Register of Systematic Reviews

UK – United Kingdom

US – United States

Declarations

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Availability of data and materials

All data generated during this study are included in this published article [and its

supplementary information files].

Competing interests

The authors declare they have no competing interests.

Funding

The lead reviewer, CC, received an Australian Government Research Training

Program Scholarship. DS is supported by the National Institute for Health and Care

Research (NIHR) through an NIHR Advanced Fellowship (NIHR302955) and the

NIHR Southampton Biomedical Research Centre (NIHR203319). The views

expressed are those of the author(s) and not necessarily those of the NIHR or the

Department of Health and Social Care. AS is supported by an Australian Research

Council Future Fellowship (FT220100610). Funding sources have not had any role in

conducting the systematic review.

Authors' contributions

CC designed the protocol, performed the searches, screening, quality appraisal, data

extraction, and data analysis, and drafted, reviewed, and edited the manuscript. DS

was the second reviewer for article screening at the title and abstract stage; AS was

the second reviewer for article screening at the full-text stage, quality appraisal and

data extraction; and AS, DS and EM reviewed and edited the manuscript.

Acknowledgements
534
535   Not applicable.

537   **Supplementary File 1. Search Strategy**

538   **Supplementary File 2. Full-text articles excluded with reason**

REFERENCES

1. Caut C, Schoenaker D, McIntyre E, Vilcins D, Gavine A, Steel A. Relationships between Women's and Men's Modifiable Preconception Risks and Health Behaviors and Maternal and Offspring Health Outcomes: An Umbrella Review. Semin Reprod Med. 2022;40(03/04):170-83. doi: 10.1055/s-0042-1744257.
2. Daly M, Kipping RR, Tinner LE, Sanders J, White James W. Preconception exposures and adverse pregnancy, birth and postpartum outcomes: Umbrella review of systematic reviews. Paediatric and Perinatal Epidemiology. 2022;36(2):288-99. doi: https://doi.org/10.1111/ppe.12855.
3. Carter T, Schoenaker D, Adams J, Steel A. Paternal preconception modifiable risk factors for adverse pregnancy and offspring outcomes: a review of contemporary evidence from observational studies. BMC Public Health. 2023;23(1):509.
4. Christiansen C, Chandra-Mouli V, Ogbaselassie L, Willumsen J, Mason E. Meeting to develop a global consensus on preconception care to reduce maternal and childhood mortality and morbidity: World Health Organization Headquarters, Geneva, 6–7 February 2012: meeting report. 2013.
5. Hall J, Chawla M, Watson D, Jacob CM, Schoenaker D, Connolly A, et al. Addressing reproductive health needs across the life course: an integrated, community-based model combining contraception and preconception care. The Lancet Public Health. 2023;8(1):e76-e84. doi: https://doi.org/10.1016/S2468-2667(22)00254-7.
6. Steel A, Lucke J, Reid R, Adams J. A systematic review of women's and health professional's attitudes and experience of preconception care service delivery. Family Practice. 2016;33(6):588-95.
7. Goossens J, De Roose M, Van Hecke A, Goemaes R, Verhaeghe S, Beeckman D. Barriers and facilitators to the provision of preconception care by healthcare providers: a systematic review. International journal of nursing studies. 2018;87:113-30.
8. Atrash H, Jack B. Preconception care to improve pregnancy outcomes: The science. J Hum Growth Dev. 2020;30(3):334-41.
9. Atrash H, Jack B. Preconception care to improve pregnancy outcomes: clinical practice guidelines. Journal of Human Growth and Development. 2020;30(3):407-16.
10. Atrash H, Jack B. Preconception care: developing and implementing regional and national programs. J Hum Growth Dev. 2020;30(3):363-71.
11. Dorney E, Boyle JA, Walker R, Hammarberg K, Musgrave L, Schoenaker D, et al. A Systematic Review of Clinical Guidelines for Preconception Care. Semin Reprod Med. 2022;40(03/04):157-69. doi: 10.1055/s-0042-1748190.
12. Boyle JA, Black K, Dorney E, Amor DJ, Brown L, Callander E, et al. Setting Preconception Care Priorities in Australia Using a Delphi Technique. Semin Reprod Med. 2022;40(03/04):214-26. doi: 10.1055/s-0042-1749683.
13. Page MJ, McKenzie JE, Bossuyt PM, Boutron I, Hoffmann TC, Mulrow CD, et al. Updating guidance for reporting systematic reviews: development of the PRISMA 2020 statement. Journal of Clinical Epidemiology. 2021;134:103-12. doi: https://doi.org/10.1016/j.jclinepi.2021.02.003.
14. Veritas Health Innovation. Covidence systematic review software. Melbourne, Australia 2022.

15. Critical Appraisal Skills Programme: CASP Qualitative Checklist. https://casp-
uk.net/images/checklist/documents/CASP-Qualitative-Studies-Checklist/CASP-
Qualitative-Checklist-2018_fillable_form.pdf (2018). Accessed.
16. Downes MJ, Brennan ML, Williams HC, Dean RS. Development of a critical
appraisal tool to assess the quality of cross-sectional studies (AXIS). BMJ Open.
2016;6(12):e011458. doi: 10.1136/bmjopen-2016-011458.
17. Pope C, Ziebland S, Mays N. Analysing qualitative data. Bmj.
2000;320(7227):114-6.
18. Briggs A, Nouri PK, Galloway M, O'Leary K, Pereira N, Lindheim SR. Expanded
carrier screening: a current survey of physician utilization and attitudes. J Assist
Reprod Genet. 2018;35(9):1631-40. doi: https://dx.doi.org/10.1007/s10815-018-
1272-8.
19. Fieldwick D, Smith A, Paterson H. General practitioners and preconception
weight management in New Zealand. Australian & New Zealand Journal of
Obstetrics & Gynaecology. 2017;57(4):420-5. doi:
https://dx.doi.org/10.1111/ajo.12609.
20. Ganganna A, Devishree G, Devishree G. Opinion of dentists and gynecologists on
the link between oral health and preterm low birth weight: "Preconception care -
treat beyond the box". Journal of the Indian Society of Pedodontics & Preventive
Dentistry. 2017;35(1):47-50. doi: 10.4103/0970-4388.199231.
21. Guess K, Malek L, Anderson A, Makrides M, Zhou SJ. Knowledge and practices
regarding iodine supplementation: A national survey of healthcare providers.
Women Birth. 2017;30(1):e56-e60. doi:
https://dx.doi.org/10.1016/j.wombi.2016.08.007.
22. Hammarberg K, Taylor L. Survey of Maternal, Child and Family Health Nurses'
attitudes and practice relating to preconception health promotion. Aust J Prim
Health. 2019;25(1):43-8. doi: https://dx.doi.org/10.1071/PY18078.
23. Hogg K, Rizio T, Manocha R, McLachlan RI, Hammarberg K. Men's
preconception health care in Australian general practice: GPs' knowledge, attitudes
and behaviours. Aust J Prim Health. 2019;25(4):353-8. doi: 10.1071/py19069.
24. Kizirian NV, Black KI, Musgrave L, Hespe C, Gordon A. Understanding and
provision of preconception care by general practitioners. Australian & New Zealand
Journal of Obstetrics & Gynaecology. 2019;59(6):799-804. doi: 10.1111/ajo.12962.
25. Manze MG, Calixte C, Romero DR, Roberts L, Perlman M, Langston A, et al.
Physician perspectives on routine pregnancy intention screening and counseling in
primary care. Contraception. 2020;101(2):91-6. doi:
https://dx.doi.org/10.1016/j.contraception.2019.11.004.
26. Mida LA, della Zazzera V, Fontaine-Bisson B. Knowledge, attitude and practice of
physicians regarding periconceptional folic acid for women at low risk of a neural
tube defect affected pregnancy. Preventive Medicine Reports. 2021;22 (no
pagination). doi: http://dx.doi.org/10.1016/j.pmedr.2021.101327.
27. Ukoha WC, Dube M. Primary health care nursing students' knowledge of and
attitude towards the provision of preconception care in KwaZulu-Natal. Afr.
2019;11(1):e1-e8. doi: https://dx.doi.org/10.4102/phcfm.v11i1.1916.
28. Wang C, Luo S, Yang D, Yan J, Ding Y, Wang S, et al. Current practice and
perspectives of healthcare providers regarding preconception care for women with
type 1 diabetes in China. Diabetes Metab Res Rev. 2021;37(4):e3454. doi:
https://dx.doi.org/10.1002/dmrr.3454.
29. Kurniawati W, Afiyanti Y, Prasetyo S, Achadi EL, Kumboyono K. The perspective
of healthcare practitioners on preconception care at primary healthcare in Jakarta: A

qualitative study. International Journal of Africa Nursing Sciences. 2021;15 (no pagination). doi: http://dx.doi.org/10.1016/j.ijans.2021.100351.

30. Poels M, Koster MP, Franx A, van Stel HF. Healthcare providers' views on the delivery of preconception care in a local community setting in the Netherlands. BMC Health Serv Res. 2017;17(1):92. doi: https://dx.doi.org/10.1186/s12913-017-2051-4.

31. Janssens S, Chokoshvili D, Vears DF, De Paepe A, Borry P. Pre- and post-testing counseling considerations for the provision of expanded carrier screening: exploration of European geneticists' views. BMC Med Ethics. 2017;18(1):46. doi: https://dx.doi.org/10.1186/s12910-017-0206-9.

32. M'Hamdi H I, van Voorst SF, Pinxten W, Hilhorst MT, Steegers EA. Barriers in the Uptake and Delivery of Preconception Care: Exploring the Views of Care Providers. Matern Child Health J. 2017;21(1):21-8. doi: https://dx.doi.org/10.1007/s10995-016-2089-7.

33. Molinaro ML, Evans M, Regnault TRH, de Vrijer B. Translating developmental origins of health and disease in practice: health care providers' perspectives. J Dev Orig Health Dis. 2021;12(3):404-10. doi: https://dx.doi.org/10.1017/S2040174420000483.

34. Munthali M, Chiumia IK, Mandiwa C, Mwale S. Knowledge and perceptions of preconception care among health workers and women of reproductive age in Mzuzu City, Malawi: a cross-sectional study. Reprod Health. 2021;18(1):229. doi: https://dx.doi.org/10.1186/s12978-021-01282-w.

35. Ojifinni OO, Ibisomi L. Health care providers perceptions about preconception care in ibadan, southwest nigeria: A qualitative study. Maternal and Child Health Journal. 2022. doi: 10.1007/s10995-021-03305-1.

36. Simone J, Hoyt MJ, Storm DS, Finocchario-Kessler S. Models of HIV Preconception Care and Key Elements Influencing These Services: Findings from Healthcare Providers in Seven US Cities. AIDS Patient Care STDS. 2018;32(7):272-81. doi: https://dx.doi.org/10.1089/apc.2017.0299.

37. Devido JA, Doswell WM, Braxter BJ, et al. Experiences of Parish Nurses in Providing Diabetes Education and Preconception Counseling to Women With Diabetes. JOGNN: Journal of Obstetric, Gynecologic and Neonatal Nursing. 2017;46(2):248-57.

38. Maas VYF, Poels M, Hölscher IM, van Vliet-Lachotzki EH, Franx A, Koster MPH. How to improve preconception care in a local setting? Views from Dutch multidisciplinary healthcare providers. Midwifery. 2022:103274. doi: https://doi.org/10.1016/j.midw.2022.103274.

39. Caut C, Leach M, Steel A. Dietary guideline adherence during preconception and pregnancy: A systematic review. Maternal & Child Nutrition. 2020;16(2):e12916. doi: https://doi.org/10.1111/mcn.12916.

40. Barker M, Dombrowski SU, Colbourn T, Fall CHD, Kriznik NM, Lawrence WT, et al. Intervention strategies to improve nutrition and health behaviours before conception. Lancet. 2018;391(10132):1853-64. doi: 10.1016/s0140-6736(18)30313-1.

41. Steel A, Schloss J, Leach M, Adams J. The naturopathic profession in Australia: A secondary analysis of the Practitioner Research and Collaboration Initiative (PRACI). Complementary Therapies in Clinical Practice. 2020;40:101220. doi: https://doi.org/10.1016/j.ctcp.2020.101220.

42. Walker R, Morris H, Lang S, Hampton K, Boyle J, Skouteris H. Co-designing preconception and pregnancy care for healthy maternal lifestyles and obesity prevention. Women and Birth. 2020;33(5):473-8. doi: https://doi.org/10.1016/j.wombi.2019.11.005.

693   43. Steel A. Creating a Vision for Family-Centered Health Services for the Five
694   'Trimesters': Co-Design Meeting Summary Report. University of Technology Sydney
695   2021.
696