Health professionals' beliefs and attitudes towards preconception care: A systematic review

Authors

Cherie Caut[1], Danielle Schoenaker[2,3,4], Erica McIntyre[5,6], Amie Steel[1]

1. Australian Research Centre in Complementary and Integrative Medicine, School of Public Health, Faculty of Health, University of Technology Sydney, Building 10, Level 8, 235-253 Jones Street, Ultimo NSW 2007 Australia.
2. School of Human Development and Health, Faculty of Medicine, University of Southampton, Southampton, UK.
3. MRC Lifecourse Epidemiology Centre, University of Southampton, Southampton, UK.
4. NIHR Southampton Biomedical Research Centre, University of Southampton and University Hospital Southampton NHS Foundation Trust, Southampton, UK.
5. Institute for Sustainable Futures, University of Technology Sydney, Building 10, Level 10, 235-253 Jones Street, Ultimo NSW 2007 Australia. Visiting Fellow
6. Research Institute for Innovative Solutions for Well-being and Health, University of Technology Sydney, Ultimo NSW, Australia.

Corresponding author: Cherie Caut

Corresponding author email: cherie.d.caut@student.uts.edu.au

Abstract

Background

Health professionals have previously identified various barriers and factors that would help facilitate preconception care services in healthcare settings. Clinically relevant preconception information and clinical practice guidelines have since been developed to aid health professionals in preconception care delivery. This systematic review aimed to 1) synthesise recent literature (past 8 years) describing health professionals' beliefs and attitudes towards preconception care services or programmes and 2) determine if the experience of health professionals providing preconception care has changed compared to literature reviews conducted more than 8 years ago.

Methods

Five databases were searched between 27/01/2016 and 20/11/2024. Primary quantitative and qualitative research studies were included if they examined health professionals' beliefs and attitudes towards delivering preconception care services or programmes. Study quality was assessed using the CASP Checklist (qualitative studies) and AXIS tool (quantitative studies). Data synthesis used thematic categorisation adapted from the framework approach.

Results

Twenty-seven studies were included (n=11 qualitative, n=14 quantitative, n=2 mixed-

methods studies). Methodological quality was generally good for qualitative studies

but varied for quantitative studies. The results covered three categories: 1)

addressing preconception care health literacy (i.e. lack of knowledge, awareness,

training and resources), 2) *clinical practicalities of preconception care* (i.e. need for

coordination of care and clarity on role responsibility), and 3) *the role of the patient*

(i.e. need for public health education to support patient-led conversations).

Conclusions

Little has changed regarding the barriers and facilitators to providing preconception

care reported by health professionals. To improve the provision of preconception

care, there is a need to co-develop professional and public preconception health

education, clinical resources, and a coordinated preconception healthcare service

model.

Keywords

preconception care, health professionals, beliefs and attitudes, systematic review

60
61

Introduction

People's modifiable preconception health (PCH) and health behaviours—such as body composition, lifestyle behaviours, nutrition, environmental exposures, and birth spacing—can affect maternal and child health[1-3]. The need to address such risk factors has led to clinical and public health measures that aim to screen for PCH risks and implement intervention strategies to optimise PCH and the health behaviours of prospective parents before conception, known as preconception care (PCC)[4]. The importance of PCC is highlighted by health policies and strategies produced by a number of countries around the world[5]. It has received dedicated attention from the World Health Organization[4] as a critically important component of healthcare that can impact multiple generations.

While PCC is valuable and important for prospective parents, healthcare providers experience a range of barriers to implementing PCC services or programmes[6, 7]. Previous systematic reviews that included studies conducted more than five years ago have reported health professionals' (HPs) experiences providing PCC and describe the types of barriers they experience, including but not limited to poor interprofessional communication, insufficient clinical time, funding, clinically relevant information, and public and HP awareness of the benefits of PCC[6, 7]. Conversely, HPs reported that adequate knowledge of PCC, prospective parents discussing their intention to become pregnant or requesting PCC, and clinical PCC resources would enable them to facilitate PCC[7]. To address some of these barriers and promote PCC services, clinician-focussed PCC information and guidelines have been developed[8, 9], and relevant information for implementing PCC programmes in organisations[10]. A recent systematic review, however, identified that existing

clinical practice guidelines (CPGs) on PCC supported by high-quality evidence are

lacking[11]. Emerging research efforts aim to establish population PCH priorities and

co-develop strategies to address them at a healthcare services level[12].

While these guidelines aim to assist HPs providing PCC, an up-to-date

understanding of HPs' views towards PCC delivery is needed to identify if the

increasing availability of guidelines has reduced or changed previously reported

barriers and if HPs also require other types of support to implement PCC

meaningfully. Therefore, this systematic review aimed to 1) synthesise recent

literature describing health professionals' beliefs and attitudes towards

preconception care services or programmes and 2) compare these experiences to

previous systematic review findings[6, 7].

Methods

This systematic review is reported following the Preferred Reporting Items for

Systematic Reviews and Meta-analyses 2020[13] and was prospectively registered on

the International Prospective Register of Systematic Reviews (PROSPERO;

CRD42021249386).

Eligibility criteria

Original primary research (quantitative and qualitative studies) published between

27/01/2016 and 20/11/2024 that sampled HPs (including, but not limited to, general

practitioners, midwives, and obstetricians) and examined their beliefs and/or

attitudes towards delivering PCC services or programmes were eligible for inclusion.

Information sources

Keyword and MeSH terms were employed in the databases MEDLINE (OVID), EMBASE (OVID), Maternity and Infant Care (OVID), CINAHL (EBSCO), and PsycINFO (EBSCO), with the following limits: title and abstract, studies in humans, published between 27/01/2016 to 27/01/2022. A second search was conducted for articles published between 28/01/2022 to 20/11/2024. No limits to language were applied. The search strategy is presented in Supplementary File 1.

Selection process

The search was conducted on 27 January 2022 and 20/11/2024. CC completed electronic database searches and downloaded citations and abstracts into EndNoteX9 citation management software. Duplication screening occurred before citations were exported into Covidence systematic review software[14]. Articles were screened by title and abstract initially by CC, and by DS and AS to establish reliability. Full-text articles were downloaded and screened initially by CC, followed by AS, to establish reliability before final inclusion for review. There were no disagreements between reviewers. References lists of the included studies were searched for additional eligible studies. Article exclusion reasons from the full-text screening stage were recorded.

Data collection process

CC initially extracted data from eligible studies into a customised form, followed by AS to establish reliability.

Data items

Data extracted included the study reference, title, aims, type of PCC service, location,

population, study design, data collection method(s), sample size, and findings on HP

beliefs and attitudes towards PCC.

Critical appraisal

The Critical Appraisal Skills Programme (CASP) Qualitative Checklist[15] evaluated

study reporting and methodological quality. Quantitative studies were assessed via

the Appraisal tool for Cross-Sectional Studies (AXIS)[16]. CC critically appraised

included studies, and AS assessed 10% of included studies for reliability.

Disagreements were discussed until a consensus was reached. A third reviewer (EM

or DS) was invited to adjudicate if unresolved.

Synthesis methods

A framework approach for applied and policy-relevant research was employed to

analyse the data, identify common themes across the studies, and then categorise

these findings to determine the key themes[17]. Findings on beliefs and attitudes

towards PCC from the included studies were extracted by CC reading and re-reading

each paper, and reviewed by AS, DS and EM. Common themes were then identified

across the studies and further defined into categories. Each study was assigned as

many categories as relevant to their reported findings.

Results

Study selection

Searches retrieved 604 articles. After removing duplicates, 376 titles and abstracts

were screened (Figure 1). Thirty-seven full-text articles were checked for eligibility;

27 met the inclusion criteria and were included in this review. Reasons for exclusion

at full-text screening are presented in Supplementary File 2.

163
164 **[Insert Figure 1. Prisma flow diagram]**

165
Study characteristics

All articles (n=27) were published in English between 2017 and 2023 (Table 1).

Fourteen studies[18-31] employed a cross-sectional study design and survey method

for data collection. Eleven qualitative studies were identified utilising focus groups

(n=4)[32-35] or interviews (n=7)[36-42]. Two studies[43, 44] collected data using

mixed methods; Devido et al. [43] employed a focus group and survey, and Mass et

al. [44] a survey and interactive workshop. The included studies were conducted in

Australia (n=6)[21-24, 29, 30]; the United States (US; n=5)[18, 25, 41-43]; the

Netherlands (n=4)[31, 33, 37, 44]; United Kingdom (UK) (n=2)[35, 38]; India

(n=2)[20, 34]; and one study each in New Zealand[19], Europe[36], Indonesia[32],

Canada[26], Malawi[39], South Africa[27], Nigeria[40], and China[28]. A range of

HPs were represented, including but not limited to pharmacists, nurses, midwives,

general practitioners, obstetricians and gynaecologists. Sample size varied among the

qualitative (n=11 to 92), quantitative (n=77 to 992) and mixed-methods (n=48 to

299) studies.

181
182 **[Insert Table 1. Characteristics of included studies]**

183
Critical appraisal of included studies

Quality assessment details are presented in Table 2 and Table 3 for qualitative and

quantitative studies, respectively.

Most of the qualitative studies fulfilled all the CASP checklist[15] criteria for

assessing methodological quality. One study[39] needed to be more transparent on

the appropriateness of the recruitment strategy, consideration of researcher bias and

rigour of data analysis. The methodological quality of the quantitative studies was

varied[16]. Sample size justification, non-response bias and funding sources or

conflicts of interest were the most poorly reported criteria, followed by the use of

instruments or measurements that had been trialled, piloted, or published

previously.

[Item Table 2. CASP Quality Assessment]

[Item Table 3. AXIS Quality Assessment]

199
200 Results of syntheses

201 Three common thematic categories across the findings from the reviewed studies

202 were identified and defined: *addressing preconception care health literacy, clinical*

203 *practicalities of preconception care*, and *the role of the patient*.

204
205 **Addressing preconception care health literacy**

206 Twenty-three studies reported findings related to addressing PCC health literacy[18-

207 26, 28-36, 38, 39, 41-43]. These studies addressed PCH knowledge, clinical training

208 needs and access to resources.

209
Preconception health knowledge gaps

Insufficient knowledge about PCH as a barrier to HPs' PCC practices was commonly

reported. For example, an Australian cross-sectional study of GPs (n=110) found

their lack of knowledge of PCC guidelines to be one of the most common barriers to

their delivery of PCC[24]; this view was shared by 40% of midwives in a second

Australian study[21] and 95% of nurses in another Australian study[30], a broader

group of HPs from a UK qualitative study.[38] The lack of knowledge as a barrier to

PCC provision also applied to GPs (n=304) in Australia providing care to men[23], of

whom almost all respondents (90%) indicated that they did not feel confident about

their knowledge of the factors affecting male fertility[23]. A cross-sectional study on

midwives (n=338) from Australia[29] found that most (85%) rated their overall

knowledge about pre and interconception health for women as excellent or above

average, although for men or partners, a higher proportion of participants (40%)

reported their overall knowledge as below average, poor or none. Participants with

more than 11 years of experience were more likely to report above average to

excellent knowledge (OR 3.11; 95% CI 1.09, 8.85), although, for men or partners, this

association was limited to midwives with more than 21 years of experience (OR 2.20;

95% CI 1.18, 4.11). Another cross-sectional study from Australia[30], on nurses

(n=152) found that only one in 20 respondents agreed that they had excellent

knowledge about PCC for women and men. When asked to rate their knowledge of

the importance of women's health in the preconception period, over half (54%)

reported average to below-average levels. Regarding the importance of men's health

in the preconception period, this increased to two-thirds of respondents (67%)

reporting average to below average knowledge. In contrast, a study[28] of HPs in

China found that 26.9% of those providing care to women with type 1 diabetes felt

they lacked the knowledge to deliver PCC to this population. At the same time, a US

study reported a significant positive association between teaching self-efficacy and

preconception counselling knowledge among parish nurses (n=48) providing

diabetes education to women with diabetes[43]. Only one cross-sectional study

sampling dentists and gynaecologists (n=300) from India investigated clinician

characteristics associated with PCH knowledge[20]. It was found in logistic

regression analyses that older age doctors (Adjusted odds ratio [AOR]=1.055 (95%

Confidence interval [CI]: 1.055-1.092, *p*=0.002), the field of specialisation - dentists

(AOR=1.635, 95% CI: 1.064-2.511, *p*=0.002), and years of practicing experience > 5

years (AOR=2.254, 95% CI: 1.46-3.45, *p*=0.001) were significantly associated with

higher knowledge levels of periodontal health and adverse pregnancy outcomes[20].

One qualitative study from India on healthcare workers (n=45)[34] found that most

had some knowledge about preconception care, limited to adolescent health and

family planning services. A qualitative study from the UK on pharmacists (n=11)[35]

found that lower knowledge scores were related to a lack of training and a lack of

confidence in oneself.

252
Awareness of preconception risks and guidelines

HPs reported needing to be made aware of PCH risks and of existing CPGs designed

to assist them in screening for PCH risks and delivering PCH interventions[19-21, 23,

24, 26, 43]. This issue was highlighted in a study involving dentists and

gynaecologists (n=300), where only 79% were aware of the association between

periodontal health and preterm low birth weight[20]. In Australia, a study of GPs

(n=110) found that only 53% were aware of PCC-CPGs[24]. Similarly, a qualitative

study on healthcare providers (n=20) from the US[42] showed a lack of awareness of

published PCC guidelines. In contrast, healthcare workers (n=45) in a qualitative

study from India were aware that there is no formal PCC program in the country[34].

Even where HPs reported being aware of guidelines, familiarity with the

recommendations may pose another barrier, as indicated in an Australian cross-

sectional study where fewer HPs (including GPs, obstetricians and gynaecologists,

midwives and dietitians) were aware of the recommended dose (38%) or duration

(44%) of preconception iodine supplementation[21]. This was despite 71% of those

HPs being aware that the National Health and Medical Research Council

recommends this intervention[21]. In this Australian study, awareness of the

recommendation was positively associated with recommending iodine

supplements[21]. In contrast, in a cross-sectional study of GPs (n=200) in New

Zealand (NZ), most GPs reported being aware of the risks of obesity in pregnancy,

over 50% of these GPs reported practice that was inconsistent with guidelines, and

only 12% of these same GPs were aware of the New Zealand Ministry of Health

Guidance for Healthy Weight in Pregnancy[19]. A similar finding was found in a

Canadian cross-sectional study assessing periconceptional folic acid

recommendations of GPs (n=77), where only half knew the correct dose and duration

of folic acid for low-risk women[26]. In the Netherlands, most (64.2 %) but not all

the respondents in a cross-sectional study on midwives and obstetricians(n=83)[31]

believe the scientific proof of the effectiveness of PCC to be sufficient.

282
The desire for further training

HPs reported wanting more information and education on PCH risks and

interventions[21-23, 33, 39] to improve their confidence in delivering PCC[22, 23,

30]. For example, in a cross-sectional study of Australian health professionals

(n=396), HPs (including GPs, obstetricians and gynaecologists, midwives and

dietitians) indicated they would be interested in receiving new information relating

to iodine[21], and in another Australian cross-sectional study almost all maternal

child and family health nurses (MCaFHN) (n=192) agreed that more information and

formal education opportunities on the topic would increase their confidence to

discuss PCH[22]. Most (87%) nurses (n=152) in another cross-sectional study from

Australia[30] indicated that more education would increase their confidence in

delivering PCC. Most midwives (n=338) in a cross-sectional study from Australia[29]

desired further education on managing pre-existing health conditions (69%) and on

optimising reproductive health (66%). Healthcare workers (n=45) suggested the

need for program-specific guidelines and training in a qualitative study from

India[34]. There was a consensus among pharmacists (n=11) in a qualitative study

from the UK[35] that insufficient training opportunities exist. A majority (54.6%) of

Australian GPs (n=304) suggested that more information and education about

factors that affect male fertility would make them more confident to talk to male

patients about fertility[23]. Some studies specifically identified lack of training as the

barrier to providing PCC and an increased likelihood of HPs providing PCH

information and PCC to their patients if they had received further training[18, 25, 28,

32, 36, 41]. For example, in a cross-sectional study from the US, 15% of GPs (n=443)

reported they needed training before implementing PCC[25]. In another cross-

sectional study from the US, obstetricians and gynaecologists (n=297) reported

offering expanded carrier screening more commonly if they were fellowship-trained

(80%) compared to those who were not (70%)[18].

310
Access to resources

Ten studies reported findings related to clinician access to resources supporting PCC

delivery[21-24, 30, 38, 41, 43]. The types of resources participants suggested include

sample meal plans[21], screening guidelines, relevant research findings[21], online

tools[22, 30], factsheets, trustworthy websites[22, 23, 30], waiting room posters[23],

checklists, brochures[24], expert clinician champions, and medical record

prompts[41]. Two other qualitative studies reported structural resource

requirements such as PCC policy and procedures used to support them to implement

PCC programs into clinical practice settings, as reported in a study of HPs (n=92)

from the US[41] and a funding model through third-party reimbursement to improve

the delivery of PCC as described by HPs (n=30) in a study from the Netherlands[33].

322
Clinical practicalities of preconception care

Twenty studies reported findings related to the clinical practicalities of PCC[20-25,

27-30, 32-35, 37, 39-42, 44].

Coordination of care

In some studies, HPs reported that a lack of interprofessional referral and care

coordination was a barrier to providing PCC[20, 28, 29, 32-34, 37, 41, 44]. One

cross-sectional study of dentists and gynaecologists (n=300) from India, for

example, found that gynaecologists rarely refer to dentists despite being aware of the

link between periodontal health and preterm low birth weight, and only 12% of

gynaecologists referred patients to the dentists in the preconception period.*[20]*

Several studies[20, 28, 32, 33] reported HPs describing a desire for an integrated,

multi-disciplinary approach to providing PCC services, greater coordination and

referral networks between HPs[29, 33, 35, 37, 41, 44] and emphasised the

importance of patient follow-up[41]. In a study from the Netherlands, HPs (n=299)

suggested that while the responsibility for providing PCC consultations is best suited

to primary care, many other HPs involved may act as referrers towards PCC[44].

Role responsibility

Most HPs in the included studies believed it to be part of their role to provide

preconception risk screening, PCH promotion and provision or referral for PCH

interventions. These include GPs[23, 24, 30, 44], MCaFHNs[22], midwives[29, 44],

nurses[27], pharmacists[35], obstetricians and gynaecologists[42] and specialist

physicians[40]. In a qualitative study from the Netherlands, HPs (including

midwives, obstetricians and gynaecologists, fertility specialists, GPs, preventive child

health care workers, maternity health care providers, physiotherapists, pharmacists

and dieticians) (n=30) expressed that the provision of PCC is challenging due to

unclear allocation of responsibilities[33]. Although HPs (including midwives, nurses,

obstetricians and gynaecologists and GPs) acknowledged that the responsibility to

provide PCC consultations lies with all professions[39] in an Australian cross-

sectional study, 84% of GPs (n=110) reported that they should be the primary

providers of PCC[24]. In a mixed-methods study from the Netherlands, HPs

(including nurses, midwives, GPs, physiotherapists, preventive child healthcare

professionals, dieticians, policy officers, maternity care assistants, and

gynaecologists) (n=299) suggested that the responsibility for providing PCC

consultations lies within primary care, mainly GPs (95.6%) and midwives

(94.4%)[44]. In this same study, HPs found it significantly more challenging to start

a conversation about a wish to conceive than midwives (26.8% versus 20.2%,

p=0.006)[44]; they felt less competent to provide preconception information (32.3%

versus 15.1%, p=<0.001)[44]. Similar findings were reported in a cross-sectional

study from the US (n=443), where most GPs (88%) felt pregnancy intention

screening should be routinely included in primary care[25]. In contrast, in a

qualitative study from Nigeria, HPs (including nurses and specialist physicians)

(n=26) stated that PCC services should be offered at all levels of health care with

referral when needed[40]. Specialist physicians from this Nigerian study also

identified the relevance of PCC to their practice, stating that those with chronic

diseases would benefit more[40]. In a cross-sectional study from Australia on

midwives' (n=338), most (88%) reported that they often encounter health states that

could be managed before pregnancy (88%)[29]. Pharmacists (n=11) in a qualitative

study from the UK[35] discussed how they are frequently asked for conception

advice, particularly by those experiencing difficulties in conceiving or those wanting

advice on optimising health to conceive.

375
Clinician time

HPs report that one of the barriers to providing effective PCC is an insufficient

amount of consultation time[21, 22, 24, 29, 30, 32, 34, 35]. For example, in an

Australian cross-sectional study[21], the main reason for not discussing dietary

sources of iodine with women was insufficient time[21]. Time constraints were also

reported by MCaFHNs (n=192)[22], GPs (n=110)[24], pharmacists[35], and a

broader range of HPs (n=32)[32, 34], as the most frequently endorsed barrier to

promoting PCH. Several of the MCaFHNs (n=192) suggested that adding a scheduled

visit dedicated to interconception health advice to those who want it would be

helpful[22].

The role of the patient

Twelve studies reported findings related to the role of the patient[18, 23, 31, 34, 35,

37-39, 41, 42, 44].

390
Community education and health promotion

Four studies indicated that HPs would like to see an increase in public health

education that improves their patients' knowledge of PCH risk factors and the

benefits of PCC[18, 37, 44] and perceive the absence of such community awareness

as a barrier to PCC[34, 37]. Most (56%) obstetricians and gynaecologists (n=297)

believed that offering expanded carrier screening to their patients should be

restricted to those diseases important to the couple, and 52% believed that screening

should only occur when the clinical significance of each disease being screened for is

understood by the couple[18]. In a qualitative study from the Netherlands, HPs

(including midwives, GPs, obstetricians and gynaecologists, cardiologists and

gastroenterologists) (n=20) stated that barriers affecting the uptake and delivery of

PCC included their belief that most future parents lacked awareness of the benefits of

PCC[37]. These findings were similar to another qualitative study from India[34],

where healthcare workers (n=45) found that preconception was not viewed as a

critical phase in the woman's reproductive cycle, and women would rarely think of

eating a balanced diet before pregnancy.

407
Patient-led conversations

Despite believing that PCC is a part of their role, HPs also perceive that the onus of

responsibility to seek PCH information and PCC is shared with the patient and that

improved public education on PCH risks and health behaviours and the benefits of

PCC would support this to occur more frequently in the absence of 'routine' PCH risk

screening and PCH promotion[23, 29-31, 34, 38, 39, 41, 42]. For example, in an

Australian cross-sectional study[23], approximately half of GPs' (n=304) stated that

they discuss fertility with male patients 'opportunistically' when consulted about a

reproductive health matter when a 'patient specifically asks for advice' and when

consulting with a couple who 'plan to have children;' very few said that they raise the

subject with men 'routinely'[23]. In another cross-sectional study (n=152) from

Australia[30], 74% of nurses stated they discuss PCC in their practice, although only

13% do so 'routinely', and of these, more preconception discussions are held with

women than with men. These findings were similar to a qualitative study[41] from

the US where the frequency with which HPs (n=92) engaged in conversations about

reproductive goals ranged from a 'routine' component of each visit to 'episodically' or

only in response to patients' 'question' or 'request.' In a qualitative study from Africa,

all the HPs (n=20) felt that women also have a role in demanding PCC services and

seeking the services[39]. Conversely, in a cross-sectional study from Australia,

midwives (n=338)[29] with a registered nursing qualification were less likely to

agree that planning to conceive is a personal decision that should only be discussed

when initiated by the woman (OR 0.55; 95% CI 0.35, 0.87).

Discussion

Findings from this systematic review provide up-to-date insights into the areas that

provide opportunities to improve PCC delivery within healthcare settings based on

HP's beliefs and attitudes. These include professional and public health education to

increase PCH knowledge and awareness, clinical resources to support HPs providing

PCC, and HP referral networks to support effective PCC delivery to prospective

parents.

438
One of the main findings of this review is that, overall, HPs report low PCH literacy

yet would like to receive training, suggesting that they may benefit from PCH

education. Our findings add to those reported in previous systematic reviews (e.g.,

HPs wanting more training to improve confidence to provide PCC and lack in

knowledge of PCH is a barrier to PCC)[6, 7] indicating little has changed and the

PCH education needs of HPs remains to be addressed. There is a high variability of

the content and recommendations across existing PCC-CPGs and a need for

guidelines that are yet to address all clinical PCC areas[11]. Comprehensive PCC-

CPGs that address all evidence-based PCC areas are needed to assist HPs in

delivering PCC[11]. However, more than simply developing CPGs to manage the PCH

knowledge of HPs is required. As supported by this review's findings, clinicians often

lack awareness of existing guidelines. While some studies have investigated

interventions that support the knowledge of HPs and effect changes to practice

behaviour[45, 46], further research is needed. Training interventions would benefit

from nuanced consideration of the skills, knowledge, time, and funding for targeted

HP groups. PCH knowledge may also need to be tiered so foundational training can

be included in clinical degrees, with more advanced training available for qualified

clinicians. To tailor and co-develop such interventions to specific HPs, further

understanding of their knowledge gaps and their current and potential scope of

practice concerning PCC is needed.

459
In this review, some HPs reported believing that the onus of responsibility should be

with the patient to seek PCH information and PCC without routine PCH risk

screening and PCH promotion. This belief assumes patient awareness of PCH risk

factors and the benefits of PCC. With the current level of public health education and

understanding of PCH risk factors, this belief may need to be challenged. Similar

findings have been reported by HPs in previous systematic reviews (e.g., client's lack

of awareness of PCC, the benefits of PCC and initiation for PCC)[6, 7]. A focus on

public education of PCH is also needed to increase the public's PCH literacy.

Adherence to guidelines during preconception is an issue previously described in a

systematic review reporting on dietary guideline adherence during preconception

and pregnancy[47]. This review also highlighted the importance of acknowledging

the influence of demographic and social factors on guideline adherence[47].

Implementation interventions are needed that support knowledge translation and

changes to health behaviour through strategies that improve whole population health

through the life course (e.g., PCH school education) as well as targeted interventions

to those life course phases where becoming pregnant is more likely or possible (e.g.,

PCH information resources and support tools) or intended (e.g., PCH education and

counselling programmes)[48]. As such, Hall and colleagues' proposed model for PCC

integrates education, digital health interventions, campaigns and social media for

raising PCH awareness among HPs and the public and includes the individualised

and specialised provision of PCC by a range of HPs able to provide clinic-based

counselling, motivational interviewing, provision of supplements and

interconception interventions[5].

483
Some HPs report that the need for clinical PCH resources and time are persistent

barriers to providing effective PCC. HPs would like access to PCH information

resources to support them in delivering PCC. Specifically, HPs identified a range of

resources they would find helpful in clinics, from clinician checklists and clinician

websites to patient information factsheets and patient websites. Clinicians report

that lack of consultation time to provide effective PCC is a barrier, and, in some

instances, clinicians would like to be better funded for PCC by receiving third-party

reimbursement. Similar findings have been reported by HPs in previous systematic

reviews (e.g., a lack of clinical time for PCC and reimbursement for that time,

needing physical space or clinically relevant information such as PCC tools checklists

and PCC guidelines)[6, 7]. Clinical resources are available, providing organisations

and clinicians with information pertinent to implementing PCC programmes[10] and

PCC-CPGs (e.g., PCH risk screening checklist or tool and PCC intervention

guidelines)[9], yet high-quality guidelines on PCC are lacking[11]. Future focus is

needed on assessing the PCH behaviour outcomes associated with healthcare

services that implement PCC programmes and when clinicians can access clinically

relevant PCH information resources and PCC-CPGs informed by high-quality

evidence.

502
503
This review also found that many HPs believe they have a role in PCC. However, they

also attest that primary care HPs (primarily GPs) should be the leading providers.

There are two issues needing further consideration. First, a precedent exists for PCC

to be provided by a broader range of HPs, with the opportunity to improve referral

networks. Second, these HPs may be under the assumption that the patient's GP has

already provided PCC. These findings are barriers to delivering PCC and were

reported by HPs in previous systematic reviews (e.g., a lack of clarity regarding who

is responsible for providing PCC[6, 7] and poor communication between HPs)[7]. A

PCC model that supports collaboration between HPs and utilisation of all capable

HPs[49-51] provides an opportunity to deliver PCC that improves PCH outcomes

throughout the community, including at-risk populations and across the life

course[12]. Meaningful involvement of all key stakeholders is required to co-design

PCC healthcare services that support a coordinated healthcare workforce[50, 51].

Further research is needed to understand the outcomes of an integrated PCC

healthcare services delivery model on PCH risks, PCH behaviours and maternal and

child health outcomes[5].

520
521
Limitations

This review should be viewed within the context of its limitations. The authors

complied with an accepted review methodology[52], although only 10% of the data

extraction was verified by a second author. As the data being extracted was not

complex and there were no differences identified in the 10% of papers that were

checked, the impact of this difference is expected to be minimal. Although the studies

in this review represent a wide range of HPs from varied countries or regions, not all

HPs who may have a role in PCC were represented in the study populations. The

methodological quality of the qualitative studies was good. However, the quantitative

study quality was varied. The heterogeneous nature of the study aims and data

collection methods precluded the pooling of data for meta-analysis. We should also

acknowledge that three studies[19, 33, 37]were included in a previous review[7]. The

research question driving our study had a broader focus than the review by Goossens

and colleagues and identified an additional 24 papers; however, to avoid overlapping

between the two reviews, the findings of the three[19, 33, 37] articles were reported

only when other papers in our review shared similar results.

Conclusions

HPs report insufficient knowledge about PCH, lack PCH training, and want

education on PCH risks and interventions to improve their confidence in providing

PCC and enhanced access to clinical PCH information resources. One of the barriers

they experience to providing effective PCC is an insufficient amount of consultation

time. Most HPs believe it is their role to provide PCC, and referral networks could

improve the delivery of coordinated preconception interventions to identify and

support patients with preconception risks. HPs want increased public health

education that improves patient awareness of PCH risks and benefits of PCC. These

barriers and facilitators to the provision of PCC reported by HPs in studies

conducted in the past five years have remained very similar when compared to

previous reviews of earlier studies on the topic. To improve the provision of PCC

going forward, there is a need to co-develop professional and public PCH education,

PCC clinical resources, and an integrated PCC healthcare service model.

List of abbreviations

AXIS - Appraisal tool for Cross-Sectional Studies

CASP - Critical Appraisal Skills Programme

CPG – Clinical practice guideline

GP – General Practitioner

HP – Health professional

MCaFHN – Maternal child and family health nurses

PCC – Preconception Care

PCH – Preconception health

PROSPERO - Prospective Register of Systematic Reviews

UK – United Kingdom

US – United States

Declarations

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Availability of data and materials

All data generated during this study are included in this published article [and its

supplementary information files].

Competing interests

The authors declare they have no competing interests.

Funding

The lead reviewer, CC, received an Australian Government Research Training

Program Scholarship. DS is supported by the National Institute for Health and Care

Research (NIHR) through an NIHR Advanced Fellowship (NIHR302955) and the

NIHR Southampton Biomedical Research Centre (NIHR203319). The views

expressed are those of the author(s) and not necessarily those of the NIHR or the

Department of Health and Social Care. AS is supported by an Australian Research

Council Future Fellowship (FT220100610). Funding sources have not had any role in

conducting the systematic review.

Authors' contributions

CC designed the protocol, performed the searches, screening, quality appraisal, data

extraction, and data analysis, and drafted, reviewed, and edited the manuscript. DS

was the second reviewer for article screening at the title and abstract stage; AS was

the second reviewer for article screening at the full-text stage, quality appraisal and

data extraction; and AS, DS and EM reviewed and edited the manuscript.

Acknowledgements

591
592   Not applicable.

594   **Supplementary File 1. Search Strategy**

595   **Supplementary File 2. Full-text articles excluded with reason**

REFERENCES

1. Caut C, Schoenaker D, McIntyre E, Vilcins D, Gavine A, Steel A. Relationships between Women's and Men's Modifiable Preconception Risks and Health Behaviors and Maternal and Offspring Health Outcomes: An Umbrella Review. Semin Reprod Med. 2022;40(03/04):170-83. doi: 10.1055/s-0042-1744257.
2. Daly M, Kipping RR, Tinner LE, Sanders J, White James W. Preconception exposures and adverse pregnancy, birth and postpartum outcomes: Umbrella review of systematic reviews. Paediatric and Perinatal Epidemiology. 2022;36(2):288-99. doi: https://doi.org/10.1111/ppe.12855.
3. Carter T, Schoenaker D, Adams J, Steel A. Paternal preconception modifiable risk factors for adverse pregnancy and offspring outcomes: a review of contemporary evidence from observational studies. BMC Public Health. 2023;23(1):509.
4. Christiansen C, Chandra-Mouli V, Ogbaselassie L, Willumsen J, Mason E. Meeting to develop a global consensus on preconception care to reduce maternal and childhood mortality and morbidity: World Health Organization Headquarters, Geneva, 6–7 February 2012: meeting report. 2013.
5. Hall J, Chawla M, Watson D, Jacob CM, Schoenaker D, Connolly A, et al. Addressing reproductive health needs across the life course: an integrated, community-based model combining contraception and preconception care. The Lancet Public Health. 2023;8(1):e76-e84. doi: https://doi.org/10.1016/S2468-2667(22)00254-7.
6. Steel A, Lucke J, Reid R, Adams J. A systematic review of women's and health professional's attitudes and experience of preconception care service delivery. Family Practice. 2016;33(6):588-95.
7. Goossens J, De Roose M, Van Hecke A, Goemaes R, Verhaeghe S, Beeckman D. Barriers and facilitators to the provision of preconception care by healthcare providers: a systematic review. International journal of nursing studies. 2018;87:113-30.
8. Atrash H, Jack B. Preconception care to improve pregnancy outcomes: The science. J Hum Growth Dev. 2020;30(3):334-41.
9. Atrash H, Jack B. Preconception care to improve pregnancy outcomes: clinical practice guidelines. Journal of Human Growth and Development. 2020;30(3):407-16.
10. Atrash H, Jack B. Preconception care: developing and implementing regional and national programs. J Hum Growth Dev. 2020;30(3):363-71.
11. Dorney E, Boyle JA, Walker R, Hammarberg K, Musgrave L, Schoenaker D, et al. A Systematic Review of Clinical Guidelines for Preconception Care. Semin Reprod Med. 2022;40(03/04):157-69. doi: 10.1055/s-0042-1748190.
12. Boyle JA, Black K, Dorney E, Amor DJ, Brown L, Callander E, et al. Setting Preconception Care Priorities in Australia Using a Delphi Technique. Semin Reprod Med. 2022;40(03/04):214-26. doi: 10.1055/s-0042-1749683.
13. Page MJ, McKenzie JE, Bossuyt PM, Boutron I, Hoffmann TC, Mulrow CD, et al. Updating guidance for reporting systematic reviews: development of the PRISMA 2020 statement. Journal of Clinical Epidemiology. 2021;134:103-12. doi: https://doi.org/10.1016/j.jclinepi.2021.02.003.
14. Veritas Health Innovation. Covidence systematic review software. Melbourne, Australia 2022.

15. Critical Appraisal Skills Programme: CASP Qualitative Checklist. https://casp-
uk.net/images/checklist/documents/CASP-Qualitative-Studies-Checklist/CASP-
Qualitative-Checklist-2018_fillable_form.pdf (2018). Accessed.

16. Downes MJ, Brennan ML, Williams HC, Dean RS. Development of a critical
appraisal tool to assess the quality of cross-sectional studies (AXIS). BMJ Open.
2016;6(12):e011458. doi: 10.1136/bmjopen-2016-011458.

17. Pope C, Ziebland S, Mays N. Analysing qualitative data. Bmj.
2000;320(7227):114-6.

18. Briggs A, Nouri PK, Galloway M, O'Leary K, Pereira N, Lindheim SR. Expanded
carrier screening: a current survey of physician utilization and attitudes. J Assist
Reprod Genet. 2018;35(9):1631-40. doi: https://dx.doi.org/10.1007/s10815-018-
1272-8.

19. Fieldwick D, Smith A, Paterson H. General practitioners and preconception
weight management in New Zealand. Australian & New Zealand Journal of
Obstetrics & Gynaecology. 2017;57(4):420-5. doi:
https://dx.doi.org/10.1111/ajo.12609.

20. Ganganna A, Devishree G, Devishree G. Opinion of dentists and gynecologists on
the link between oral health and preterm low birth weight: "Preconception care -
treat beyond the box". Journal of the Indian Society of Pedodontics & Preventive
Dentistry. 2017;35(1):47-50. doi: 10.4103/0970-4388.199231.

21. Guess K, Malek L, Anderson A, Makrides M, Zhou SJ. Knowledge and practices
regarding iodine supplementation: A national survey of healthcare providers.
Women Birth. 2017;30(1):e56-e60. doi:
https://dx.doi.org/10.1016/j.wombi.2016.08.007.

22. Hammarberg K, Taylor L. Survey of Maternal, Child and Family Health Nurses'
attitudes and practice relating to preconception health promotion. Aust J Prim
Health. 2019;25(1):43-8. doi: https://dx.doi.org/10.1071/PY18078.

23. Hogg K, Rizio T, Manocha R, McLachlan RI, Hammarberg K. Men's
preconception health care in Australian general practice: GPs' knowledge, attitudes
and behaviours. Aust J Prim Health. 2019;25(4):353-8. doi: 10.1071/py19069.

24. Kizirian NV, Black KI, Musgrave L, Hespe C, Gordon A. Understanding and
provision of preconception care by general practitioners. Australian & New Zealand
Journal of Obstetrics & Gynaecology. 2019;59(6):799-804. doi: 10.1111/ajo.12962.

25. Manze MG, Calixte C, Romero DR, Roberts L, Perlman M, Langston A, et al.
Physician perspectives on routine pregnancy intention screening and counseling in
primary care. Contraception. 2020;101(2):91-6. doi:
https://dx.doi.org/10.1016/j.contraception.2019.11.004.

26. Mida LA, della Zazzera V, Fontaine-Bisson B. Knowledge, attitude and practice of
physicians regarding periconceptional folic acid for women at low risk of a neural
tube defect affected pregnancy. Preventive Medicine Reports. 2021;22 (no
pagination). doi: http://dx.doi.org/10.1016/j.pmedr.2021.101327.

27. Ukoha WC, Dube M. Primary health care nursing students' knowledge of and
attitude towards the provision of preconception care in KwaZulu-Natal. Afr.
2019;11(1):e1-e8. doi: https://dx.doi.org/10.4102/phcfm.v11i1.1916.

28. Wang C, Luo S, Yang D, Yan J, Ding Y, Wang S, et al. Current practice and
perspectives of healthcare providers regarding preconception care for women with
type 1 diabetes in China. Diabetes Metab Res Rev. 2021;37(4):e3454. doi:
https://dx.doi.org/10.1002/dmrr.3454.

29. Bradfield Z, Leefhelm E, Soh SE, et al. The MidPIC study: Midwives' knowledge,
perspectives and learning needs regarding preconception and interconception care.
PLoS ONE. 2023;18(11).

30. Dorney E, Millard J, Hammarberg K, Griffin K, Gordon A, McGeechan K, et al.
Australian primary health care nurses' knowledge, practice and attitudes relating to
preconception care: learnings for service implementation. Aust J Prim Health.
2022;28(1):63-8. doi: 10.1071/PY21104.
31. Scheele J, Smith SM, Wahab RJ, Bais B, Steegers–Theunissen RPM, Gaillard R,
et al. Current preconception care practice in the Netherlands — An evaluation study
among birth care professionals. Midwifery. 2023;127:N.PAG-N.PAG. doi:
10.1016/j.midw.2023.103855.
32. Kurniawati W, Afiyanti Y, Prasetyo S, Achadi EL, Kumboyono K. The perspective
of healthcare practitioners on preconception care at primary healthcare in Jakarta: A
qualitative study. International Journal of Africa Nursing Sciences. 2021;15 (no
pagination). doi: http://dx.doi.org/10.1016/j.ijans.2021.100351.
33. Poels M, Koster MP, Franx A, van Stel HF. Healthcare providers' views on the
delivery of preconception care in a local community setting in the Netherlands. BMC
Health Serv Res. 2017;17(1):92. doi: https://dx.doi.org/10.1186/s12913-017-2051-4.
34. Chutke AP, Doke PP, Gothankar JS, Pore PD, Palkar SH, Patil AV, et al.
Perceptions of and challenges faced by primary healthcare workers about
preconception services in rural India: A qualitative study using focus group
discussion. Frontiers in public health. 2022;10:888708. doi:
https://dx.doi.org/10.3389/fpubh.2022.888708.
35. Silverio SA, Karki P, Lakhani S, Alter M, Weinman J, Flynn AC. 'People still want
a face and that's where we can fill in': a qualitative study of community pharmacists'
experiences of providing healthcare advice about preconception and pregnancy. The
International journal of pharmacy practice. 2023;31(1):38-45. doi:
https://dx.doi.org/10.1093/ijpp/riac097.
36. Janssens S, Chokoshvili D, Vears DF, De Paepe A, Borry P. Pre- and post-testing
counseling considerations for the provision of expanded carrier screening:
exploration of European geneticists' views. BMC Med Ethics. 2017;18(1):46. doi:
https://dx.doi.org/10.1186/s12910-017-0206-9.
37. M'Hamdi H I, van Voorst SF, Pinxten W, Hilhorst MT, Steegers EA. Barriers in
the Uptake and Delivery of Preconception Care: Exploring the Views of Care
Providers. Matern Child Health J. 2017;21(1):21-8. doi:
https://dx.doi.org/10.1007/s10995-016-2089-7.
38. Molinaro ML, Evans M, Regnault TRH, de Vrijer B. Translating developmental
origins of health and disease in practice: health care providers' perspectives. J Dev
Orig Health Dis. 2021;12(3):404-10. doi:
https://dx.doi.org/10.1017/S2040174420000483.
39. Munthali M, Chiumia IK, Mandiwa C, Mwale S. Knowledge and perceptions of
preconception care among health workers and women of reproductive age in Mzuzu
City, Malawi: a cross-sectional study. Reprod Health. 2021;18(1):229. doi:
https://dx.doi.org/10.1186/s12978-021-01282-w.
40. Ojifinni OO, Ibisomi L. Health care providers perceptions about preconception
care in ibadan, southwest nigeria: A qualitative study. Maternal and Child Health
Journal. 2022. doi: 10.1007/s10995-021-03305-1.
41. Simone J, Hoyt MJ, Storm DS, Finocchario-Kessler S. Models of HIV
Preconception Care and Key Elements Influencing These Services: Findings from
Healthcare Providers in Seven US Cities. AIDS Patient Care STDS. 2018;32(7):272-
81. doi: https://dx.doi.org/10.1089/apc.2017.0299.
42. Nacev EC, Greene MZ, Taboada MP, Ehrenthal DB. Factors Influencing Provider
Behavior Around Delivery of Preconception Care. Matern Child Health J.
2022;26(7):1567-75. doi: 10.1007/s10995-022-03411-8.

43. Devido JA, Doswell WM, Braxter BJ, et al. Experiences of Parish Nurses in
Providing Diabetes Education and Preconception Counseling to Women With
Diabetes. JOGNN: Journal of Obstetric, Gynecologic and Neonatal Nursing.
2017;46(2):248-57.
44. Maas VYF, Poels M, Hölscher IM, van Vliet-Lachotzki EH, Franx A, Koster MPH.
How to improve preconception care in a local setting? Views from Dutch
multidisciplinary healthcare providers. Midwifery. 2022:103274. doi:
https://doi.org/10.1016/j.midw.2022.103274.
45. de Castro Nascimento N, Borges ALV, Fujimori E, Reis-Muleva B. Training in
preconception care focused on primary health care providers: Effects on
preconception care knowledge and provision. Heliyon. 2024;10(9).
46. Hauser KW, Lilly CM, Frías JL. Florida health care providers' knowledge of folic
acid for the prevention of neural tube defects. SOUTHERN MEDICAL JOURNAL-
BIRMINGHAM ALABAMA-. 2004;97:437-9.
47. Caut C, Leach M, Steel A. Dietary guideline adherence during preconception and
pregnancy: A systematic review. Maternal & Child Nutrition. 2020;16(2):e12916. doi:
https://doi.org/10.1111/mcn.12916.
48. Barker M, Dombrowski SU, Colbourn T, Fall CHD, Kriznik NM, Lawrence WT, et
al. Intervention strategies to improve nutrition and health behaviours before
conception. Lancet. 2018;391(10132):1853-64. doi: 10.1016/s0140-6736(18)30313-1.
49. Steel A, Schloss J, Leach M, Adams J. The naturopathic profession in Australia: A
secondary analysis of the Practitioner Research and Collaboration Initiative (PRACI).
Complementary Therapies in Clinical Practice. 2020;40:101220. doi:
https://doi.org/10.1016/j.ctcp.2020.101220.
50. Walker R, Morris H, Lang S, Hampton K, Boyle J, Skouteris H. Co-designing
preconception and pregnancy care for healthy maternal lifestyles and obesity
prevention. Women and Birth. 2020;33(5):473-8. doi:
https://doi.org/10.1016/j.wombi.2019.11.005.
51. Steel A. Creating a Vision for Family-Centered Health Services for the Five
'Trimesters': Co-Design Meeting Summary Report. University of Technology Sydney
2021.
52. Shea BJ, Reeves BC, Wells G, Thuku M, Hamel C, Moran J, et al. AMSTAR 2: a
critical appraisal tool for systematic reviews that include randomised or non-
randomised studies of healthcare interventions, or both. bmj. 2017;358.
793

Identification of studies via databases and registers

Identification

Records identified from Databases (n=602)
MEDLINE(OVID) (n=166);
EMBASE(OVID) (n=253);
MATERNITY AND INFANT CARE(OVID) (n=35);
CINAHL(EBSCO) (n=119);
PSYCHINFO(EBSCO) (n=29)
Records identified from hand searches (n=2)

Records removed *before* screening:
Duplicate records removed by human (n=223); by automation tool (n=5)

Screening

Records screened (n=376) → Records excluded by human (n=339)

Reports sought for retrieval (n=37) → Reports not retrieved (n=0)

Reports assessed for eligibility (n=37) → Reports excluded:
(n=5) Not full text publication (e.g., conference abstract or poster)
(n=2) Wrong outcomes (e.g., not beliefs and/or attitudes)
(n=2) Wrong population (e.g., not health professional, not human)
(n=1) Wrong exposure (e.g., does not include preconception care within scope of service)

Included

Studies included in review (n=27)