1 **A Qualitative Study of People with Alzheimer's Disease in a Memory Clinic Considering**
2 **Lecanemab Treatment**
3

4 Anna L Parks, MD (1), Ayush Thacker, BS (2), Daniel Dohan, PHD (3) Liliana A Ramirez Gomez,
5 MD (4), Christine S Ritchie, MD, MSPH (2, 5) Joanna Paladino, MD* (2, 5) and Sachin J Shah,
6 MD, MPH (2, 6)*
7
(1) University of Utah, Division of Hematology and Hematologic Malignancies, Salt Lake City,
UT, USA
(2) Massachusetts General Hospital, Mongan Institute Center for Aging and Serious Illness,
Boston, MA, USA
(3) University of California San Francisco, Institute for Health Policy Studies, San Francisco,
CA, USA,
(4) Massachusetts General Hospital, Memory Disorders Division, Department of Neurology,
Boston, MA, USA
(5) Massachusetts General Hospital, Division of Palliative Care and Geriatric Medicine, Boston,
MA, USA
(6) Massachusetts General Hospital, Division of General Internal Medicine, Boston, MA, USA
19
20 *Co-senior authors
21
22 Word count: 3680
23
Contact
Anna L. Parks, MD
2000 Circle of Hope Drive
Salt Lake City, Utah 84112
Anna.parks@hsc.utah.edu
801-213-2648
30
31 Running title: Qualitative Study of Lecanemab Treatment

32
33

Abstract

Background

People with Alzheimer's disease (AD) now have access to disease-modifying treatment with

anti-amyloid monoclonal antibodies (mAbs). Their perception of risks and benefits and approach

to treatment decisions remain unknown.

Objective

We aimed to understand how people with AD weigh the benefits and costs of anti-amyloid

mAbs and incorporate these into decisions about treatment.

Methods

We conducted semi-structured interviews with people with biomarker- or imaging-confirmed AD

and mild or moderate cognitive impairment who were seen at memory care clinics and

discussed lecanemab with a clinician. Interviews were recorded, transcribed, and deidentified.

Thematic analysis identified themes and subthemes.

Results

Among 22 participants (mean age 70, 8 [36%] women, 22 [100%] White), analysis revealed 3

major themes and associated subthemes: 1) People with AD sought and obtained information

from different sources—advocacy organizations, the Internet, and clinicians; 2) Hopes, expected

benefits, and the existential threat of dementia drove willingness and readiness to start

lecanemab; 3) Individual traits, family factors, and degree of trust in expertise influenced how

people balanced risks and benefits. Some would accept treatment at any cost; others carefully

weighed risks and burdens, but were motivated by supportive families, insurance coverage, and

trust in expertise; for a few, costs decidedly outweighed their personal benefits. People with AD

desired more individualized information and to hear more from patients who took the

medication.

Conclusion

Results from this first qualitative study of people with AD considering treatment with anti-amyloid

mAbs can inform clinician, health system and policy efforts to individualize decisions.

Key words

Alzheimer's disease, anti-amyloid monoclonal antibodies, qualitative methods, shared decision

making

Introduction

Approval of anti-amyloid monoclonal antibodies (mAbs) is a major advance in the field of Alzheimer's disease (AD). This medication class is the first potentially disease-modifying therapy for people with AD. Lecanemab, approved by the Food and Drug Administration (FDA) in 2023, can slow cognitive decline in early AD by six months.[1] Donanemab was FDA-approved in July 2024, and clinical trials of additional mAbs are in process.[2]

Alongside therapeutic optimism is growing recognition of potential tradeoffs that complicate treatment decisions. Side effects include intracranial swelling (amyloid-related imaging abnormalities-edema), and intracranial bleeding (amyloid-related imaging abnormalities-hemorrhage [ARIA-H]).[3] Certain comorbidities and medications, like anticoagulants, and genotypes, like apolipoprotein E ε4 (ApoE4), increase the ARIA-H risk.[4] Although most ARIA is asymptomatic, it can result in death and disability.[5] How patients fare long-term remains unknown. Treatment requires time expenditure, including subspecialty visits, diagnostic testing, monitoring, and infusions every two-to-four weeks. And anti-amyloid mAbs are financially costly for individuals and society.[6]

Many have called for shared decision-making to address the complexity of treatment decisions.[4,5,7–9] To be effective, shared decision-making requires understanding how patients perceive the risks and benefits of treatment and how they approach decisions.[10] In the new era of anti-amyloid mAbs, our aim was to understand how people with AD weigh potential benefits of treatment against costs and how they make treatment decisions.

Methods

Study Design and Data Collection

We conducted an exploratory qualitative study using semi-structured interviews of people seen

in memory clinics with AD between February-March 2024. A multidisciplinary team developed an

interview guide using the Ottawa Decision Support Framework.[11] The guide included questions

about 1) information received, 2) hopes, motivations, and concerns about treatment, 3) level of

certainty about starting treatment, 4) factors influencing decision-making, and 5) the decision-

making process. Interviews were conducted remotely by Zoom or telephone by trained study

personnel (A.P., A.T.). The institutional review boards of participating institutions approved this

study as exempt. We adhered to Consolidated Criteria for Reporting Qualitative Studies

(COREQ) guidance.[12]

Setting and Participants

Participants were recruited by mail and telephone from memory clinic referral centers at two

academic medical centers, the University of Utah and Massachusetts General Hospital. **Figure**

1 shows a flowchart of recruitment. We screened memory clinic referrals and included subjects

who:1) were English-speakers, 2) between ages 50-90 years, 3) had at least one memory clinic

visit where lecanemab was discussed, 4) underwent initial diagnostic testing to confirm AD and

assess for lecanemab eligibility (e.g., amyloid on cerebrospinal fluid and/or positron-emission

tomography, magnetic resonance imaging and *ApoE4 testing*). To reflect the full breadth of

patients with whom clinicians have shared decision-making discussions, neither consent for nor

subsequent receipt of lecanemab were inclusion criteria. We included subjects with Montreal

Cognitive Assessment (MOCA) scores $\geq$16 (mild or moderate cognitive impairment), consistent

with current appropriate use criteria for lecanemab.[5] We included two participants with

biomarker or imaging-confirmed AD who had normal MOCA score ($\geq$26) but clinical diagnosis of

mild cognitive impairment based on Quick Dementia Rating System or Clinical Dementia

Rating-Sum of Boxes score. These two participants were treated with lecanemab and were

included to accurately represent real-world clinical practice. Participants completed verbal

informed consent and an online demographic questionnaire before participation and received a

$50 gift card.

Data Analysis

Interviews were digitally recorded, transcribed, and de-identified. Qualitative thematic analysis

was performed in March-April 2024 using a stepwise approach and hybrid deductive-inductive

coding methods.[13,14] The primary analytic team was comprised of AP (female physician-

researcher with qualitative research expertise), AT (male clinical research coordinator), and JP

(female researcher with serious illness communication and qualitative research expertise).

Coders independently coded the first five transcripts deductively using the interview guide as a

framework; additional codes were added inductively to create the preliminary codebook. AT, AP,

and JP coded the remaining transcripts and met every five interviews to discuss discrepancies,

resolve differences by consensus, revise codes, ensure iterative content building, and examine

and address how personal experiences or perspectives might influence the analysis. The final

codebook was applied to all interviews. Using meetings with two additional experts (DD, SS),

codes were grouped to develop preliminary themes and sub-themes. Associations were

confirmed by the entire group to identify final themes. AP, AT, and JP reviewed transcripts

iteratively to assess data saturation, or when interviews were not gleaning new information, at

which point recruitment was stopped.[15–17]

Results

Characteristics for the 22 participants are provided in **Table 1**. The study cohort included 12

(55%) people from Massachusetts General Hospital and 10 (45%) from the University of Utah.

Mean age was 70 years, 8 (36%) were women, and 22 (100%) were non-Hispanic White. Eight

(36%) had less than a college degree, 8 (50%) had annual income less than $100,000 (6 [27%]

were unsure or had no income), and 6 (27%) lived rurally. Mean Montreal Cognitive Assessment

(MOCA) was 20 (standard deviation 3); 2 (9%) had normal cognitive performance (MOCA ≥26),

15 (68%) had mild cognitive impairment (MOCA 18-25), and 5 (23%) had moderate cognitive

impairment (MOCA 10-17). A majority (12 [55%]) were ready to decide on mAb treatment, and

the remainder were evenly split between ready but wanting more information (5 [23%]) and not

ready (5 [23%]). A small proportion would decline (3 [14%]), a similar proportion needed more

information (4 [18%]), and most would pursue treatment (15 [68%]).

154
155   **Table 1 Patient Characteristics**
156

Characteristic	N=22
Age mean (SD)	70 (10)
Institution n (%)	
Massachusetts General Hospital	12 (55%)
University of Utah	10 (45%)
Gender n (%)	
Male	14 (64%)
Female	8 (36%)
Orientation n (%)	
Heterosexual or straight	22 (100%)
Race n (%)	
White	22 (100%)
Ethnicity n (%)	
Non-Hispanic	22 (100%)
Geographic Location n (%)	
Suburban	14 (64%)
Rural	6 (27%)

Characteristic	N=22
Urban	2 (9.1%)
Religious Affiliation n (%)	
Christian	16 (73%)
Agnostic	3 (14%)
Atheist	1 (4.5%)
Buddhist	1 (4.5%)
Prefer not to answer	1 (4.5%)
Highest level of education obtained n (%)	
High School	1 (4.5%)
Some college	5 (23%)
Associate's Degree	2 (9.1%)
Bachelor's Degree	7 (32%)
Master's Degree	4 (18%)
Doctorate Degree	3 (14%)
Household Income n (%)	
$30,000-59,999	1 (4.5%)
$60,000-99,999	7 (32%)
$100,000 or more	8 (36%)
Unsure/No Income	6 (27%)
Montreal Cognitive Assessment (MOCA) score mean (SD)	20 (2)
Normal cognitive performance (MOCA ≥26) n (%)*	2 (9)
Mild cognitive impairment (MOCA 18-25) n (%)	15 (68)
Moderate cognitive impairment (MOCA 10-17) n (%)**	5 (23)
How ready do you feel to decide about treatment n (%)	

Characteristic	N=22
Ready to make a decision	12 (55%)
Ready to make a decision but has more questions	5 (23%)
Not ready to make a decision	5 (23%)
If you had to decide today about treatment n (%)	
Yes to treatment	15 (68%)
Not sure, need more information	4 (18%)
No to treatment	3 (14%)

158　Through analysis, we identified three primary themes and associated subthemes (**Figure 2**),

159　with representative quotations in the manuscript text. **Table 2** provides additional supportive

160　quotations.

162　**Table 2 Additional Illustrative Quotations from People with AD by Theme**

Theme	Illustrative quotation
People with AD sought and obtained information from different sources	
Advocacy organizations, the Internet and the media as resources for self-study	*"I think we went to the conference in [city] the year before. And that was when they were going through the approval process and that was a big focus of the conference, so that pretty much made me more aware… it was kind of emphasizing how this was a new avenue to explore. And so that's what got me interested." (ID 12)*
Variably detailed information from clinicians	*"It wasn't until our local doctor here gave us a whole fistful, this little stack of articles that described what it was and how it worked and what are the complications, and what are some of the things that would make it difficult for me, like the distance. It was like an article from The New England Journal of Medicine and one from the American Medical Association. He's one of these people that really keeps on top of things. I mean they were legitimate articles. Because he has access to all that kind of information through his— he's a neurologist." (ID 15)*
Motivating interactions with treated patients and informed family members	*"Our son is… a [medical professional]. He runs the tests and all this, exactly this, but with younger people. And he's been very helpful with phone calls and his network that he's got fairly wide,*

Theme	Illustrative quotation
	going to conferences and things, talking with people. And has been instrumental." (ID 25)

Hopes, expected benefits, and the existential threat of dementia driving willingness and readiness to start anti-amyloid mAbs.

Theme	Illustrative quotation
Stalling, reversal or cure of cognitive decline	*"Obviously the hope is that it can stall this disease. At least I'd feel better if they could give us some kind of parameters around...You know, for somebody with your subtype, you know, what we're seeing thus far, it can stall it two years, three years." (ID 5)*
Optimism drawn from mechanism of action and novelty compared to non-pharmacologic options	*"I think the biggest selling point was this is the best there is. Nothing any better and this is the way we have to go, or should go, and I was all for that." (ID 24)*
Stopgap until more effective therapy	*"I just made the decision that, for me, not only because of the immediacy of the potential help from lecanemab, but I don't know what's gonna happen in the future... hopefully, this gives me more time to see if something else happens as well." (ID 2)*
Motivated by desire to buy time for enjoyable activities, preserve independence, and do "something" vs nothing	*"I feel like I could manage going forward with the capabilities that I've retained thus far to function fairly independently if needed. So, I'm in a good spot, so if we can stop it in its tracks, that would be wonderful." (ID 5)*

Individual traits and preferences, family factors, degree of trust in expertise influence, and sense of pressure or urgency influenced how people with AD balance risks and benefits in decision-making.

Theme	Illustrative quotation
Impact of personality	*"I have the ability to pass over the negativity a bit. So my answer to that is I—when the negatives came up, I didn't listen very well...I think I kind of passed over it. I was really centered on the idea that I was going to get better. I let the rest—if there was any rest, I let it fly by. You know, being an [80-85]-year-old man that is cognitively challenged is not all bad. You're kind of a funny old bastard wandering through life. And I kind of like being that. It's not that bad." (ID 16)*
Perceived inevitability of decline	*"I mean, the writing's on the wall, so I might as well, you know, I'm not just gonna sit back and accept all this if something that can be done." (ID 12)*
Weighing side effects, costs and logistical burdens	*"Well, the biggest drawback is how much it's gonna cost. I don't know if I can afford it. My understanding was that there would be*

Theme	Illustrative quotation
	some, but I don't know what the price is and the cost—out of pocket. That will make—that's-that's a stop right there, if it's too high…That to me is the only thing that will possibly stop.” (ID 10)
Importance of supportive families	*“How I rationalize that in my mind and then turn around and had a conversation with my husband, I said, ‘If I had breast cancer, nobody would question if I was going for chemotherapy tomorrow, but technically chemotherapy is a poison.’ And at some point long before we became conditioned to hearing about chemotherapy, it sounded big, bad, and scary too… once I kind of laid that scenario out about the chemotherapy, I think it kind of put everything into perspective and nobody's really said anything negative since.” (ID 5)*
Trust in expertise	*“I feel, I gotta think the FDA has done its job. Like I think that you guys are, you know, have one of the-the best hospitals in the world and the best doctors. So if you're gonna have this disease, I think I have it at the right place with the right hospital, with the right doctors, with the right partner.” (ID 12)*
Family, physicians and sense of personal urgency to act now as sources of pressure	*“I kinda got the impression that he [the physician] was recommending it or that it would give me more years. Pushing it's too strong a word, but that he was encouraging it in some way… I got the impression he was definitely pro doing the drugs. I don't know if that's based on good results he's seen in patients or whether it's just kinda the latest thing to try. The latest thing to do.” (ID 15)*
Making calculated risks and conscious tradeoffs	*I'm ready to do it, willing to do it…Is it a perfect world? Not always. But I'm fine with that. Calculated risk.” (ID 8)*
Sense of urgency to decide	*“I'm ready. 'Cause I feel like the longer it takes, I feel like I lose more cognitive abilities. And so I'm anxious to get started in the hopes that this will slow everything down. Because once you lose what you lose, you don't get it back.” (ID 6)*
Autonomy	*“I think it's a combination of two things. I'd like to talk to the doctor to understand what he or she knows and take that information with the family and say listen, there's no guarantees. But I honestly think that my family would say, ‘Go for it.’ Got nothin' to lose, really, in a sense.” (ID 8)*
Reasons for declining	*“The other red flag was that it talked in there was, that it wouldn't be good for people with autoimmune diseases. And I have lupus, but it hasn't been active for a number of years. So that still was a red flag for me. (ID 15)*

Theme	Illustrative quotation
	"My first reaction was you have got to be kidding me. But that's because I went right away to the side effects. Because I've got other things going on and that I'm takin' meds for." (ID 25)
	"If I had to decide right now, I'd probably be hesitant because I don't know enough about the medication...The over-dosage on the medication that made me throw up so much—that would cause some hesitancy." (ID 22)

People with AD sought and obtained information from different sources

Nearly all respondents reported self-directed information gathering. In addition to the Internet

and media, advocacy organizations played a prominent role in disseminating information and

empowering patients.

"I've actually gone out and sought out information on my own…I was able to connect

from the Alzheimer's page to an article on it. And I basically printed that and then kind of

read it in its entirety so that I could highlight areas of concern for myself." (ID5)

Most learned about anti-amyloid mAbs from their clinicians. Some learned from primary care

providers—often the physician to whom they first presented with cognitive complaints—while

others obtained information from subspecialty providers. Level of detail conveyed by clinicians

varied, ranging from comprehensive to only a broad overview.

"You can do it over at our local hospital. I don't really know much more than that...It just

was, like, a couple of minutes or something. He just said it'd be an infusion, and so I

asked him if that was—I don't know. He didn't really say much." (ID 18)

A few participants experienced motivating narratives from other people with AD who had

received anti-amyloid mAbs. In other instances, information from family members with

healthcare expertise played a role.

"…I met a gal who was a little younger than me who's been getting the treatment and

she really feels like it's helped her quite a bit. She's done very well on it, so that was

another exciting thing." (ID12)

Hopes, expected benefits, and the existential threat of dementia drive willingness and readiness

to start lecanemab

Delaying progression of cognitive decline dominated the hopes for treatment. Some participants'

expectations exceeded the benefits seen in clinical trials. Several hoped for a cure, while others

hoped that lecanemab would reverse memory loss or completely remove amyloid from the

brain.

"[I hope] that they'll come in in a year and say, 'We can't find any amyloid plaque—- in

your brain. And that the tau tangles, or whatever they are, seem not to be there.'" (ID 9)

Patient motivation to accept treatment came from the opportunities to a) gain more time feeling

like themselves and doing enjoyable activities; b) have a plan to fight dementia rather than

succumbing to the status quo; or c) preserve independence.

"I would like to be able to live a life and be able to speak and not get wound up in my

mind.... This is the anxiety.... I'm hoping it can help not only just the hope, but it's good to

have a plan and, 'This is what I'm gonna do.' And 'I've got good people around me.' I feel

good when that happens, I have to say." (ID 8)

Some people felt motivated knowing that they were receiving the "latest and greatest" compared

to existing non-pharmacologic alternatives and that they were still cognitively able to assess the

risks and benefits.

"I'm, you know, cognitively up with what people are saying to me and what I can read

about or have my wife read to me about. So, I'm still pretty confident and —how do I put

this? I'm worth saving." (ID 22)

Respondents also described lecanemab as a stopgap until the development of more effective

therapies. This was coupled with anxiety about whether receiving lecanemab would preclude

them from future treatments.

"The other thing that comes in is if you get on something, does it preclude the something

better coming along? In the next two years you know, or whatever it is, that's better. But

because you've been on this or on this, you can't do that. I don't know how likely that is."

(ID 25)

Some described feeling conflicted because they worried about tradeoffs (e.g., diminished quality

of life from complications). Several raised concerns that lecanemab would have side effects like

chemotherapy (e.g., nausea, hair loss) because both are infusions.

"It's because, you know, I've had so much trouble with drug reactions in my life that I've

tried as best I can to stay off drugs and stay away from drugs. So, I'm real—would be

really concerned about, I'd rather have like two good years with as clear a mind as I can

have, rather than having three years, but being so affected by the drugs that I'm taking

that it wouldn't be worth the extra year." (ID 11)

Individual traits and preferences, family factors, degree of trust in expertise influence, and sense

of pressure or urgency influenced how people with AD balance risks and benefits in decision-

making.

When asked to consider risks and benefits, some noted that their approach was rooted in their

personal tendency to "look past the negatives", rather than the specifics of anti-amyloid mAbs.

For others, the perceived inevitability of decline without intervention helped them decide to

accept treatment at any cost.

"What do I have to lose? Is basically the way I look at it. I mean, even if it does cause

problems for me, I'm headed that way already. I'm not—might be a different kind of

problem. I don't care." (ID 10)

The complication of ARIA-H weighed on most participants' minds. Although people with AD

wanted more information on their individual risk, they felt that bleeding risk was not high and

surmountable. Very few mentioned death as a potential outcome.

"I just know that there's things that could go bad, and I'm ready for that. I'd rather be able

to get onto the drug and… I'd rather be able⬚ on it. At least try it, and… if I die, I die."

(ID 17)

Costs, insurance coverage, and logistical hurdles also influenced willingness to pursue

treatment. Some felt that infusions every two weeks would not affect their lifestyle, while others

found this disruptive. People with AD likened the logistics of anti-amyloid mAbs to receiving

chemotherapy for cancer.

"It also goes with a medication that caused you to lose what you already have as far as

your physicalness, your aliveness, those kinds of things. So if it's a medication that's

going to maybe give you longer memory, but less life, that would make a difference." (ID

25)

Family also influenced how people with AD weighed side effects and costs. Some described

family members as being very supportive, while others described hesitation. Those whose

desire for treatment conflicted with family members reminded them of their autonomy despite

their declining cognition and again drew comparisons to cancer to address concerns. A few

equated forgoing treatment with unnecessarily burdening family.

"My husband has been wonderful about it. I think he has more worries about the risks

than I do, partly because he's like, 'I'd rather have you even if you don't know who I am

anymore. I'd rather have you as long as I can.'" (ID2)

People with AD voiced trust in expertise and felt that the medical team took their individualized

risk and benefit into account. Faith in the drug approval process and their healthcare system

also influenced decision-making.

"I know that no doctor is perfect. I do honestly believe that they know a lot more about

this than I do—in many ways. And I trust that they're very, very good. And that's the only

way I know how to answer...I don't even think about it because I trust it. Unless

something went wrong, then it might change... It's like I say, I feel like something

happens that makes me lose that trust. That's hard to get back." (ID 10)

People with AD described a sense of pressure to accept treatment. Pressure came from family

members, clinicians, or a sense of personal urgency to act now (versus waiting until they may

not have a choice).

"I suppose I could wait later, but I also feel like the damage that is happening, I know at

some point—I don't wanna get to the point where I don't have the choice anymore,

right?" (ID 2)

When making a final treatment decision, a small proportion indicated that they would decide

alone, and a small proportion would make a group decision with family. The majority described

gathering input from physicians and family but ultimately felt that the decision was theirs to

make.

"I will decide on my own—with agreement from the doctor. I'm certainly willing to listen to

any concerns from my family member, but ultimately it is my body, my decision." (ID 5)

Ideal future state of knowledge and decision-making

We also synthesized information that people with AD wanted to improve their knowledge and

decision-making (**Table 3**).

Table 3: Desired information and support for anti-amyloid mAb treatment decisions

Desired information and support	Representative quotation
Personalized information on benefits	*"I need to know what the results of other patients are. And preferably other patients that fit my same general condition, age, that kind of stuff..." (ID 26)*
Personalized information on risks	*"Obviously, I wanna know how prevalent the bad side effects are. If it's a little discomfort on getting the infusion, that's not a problem. The things that I'm concerned about are the things that would either make it far worse or sooner than letting it run its course." (ID 3)*

Desired information and support	Representative quotation
Detailed and personalized information on ARIA-H and its treatments	*"That sounds pretty drastic. It could be. I'd wanna know more about why... And if there can be blood in the brain, you know, is that—do I need an operation to get that fixed? Am I gonna die, or will that—is that normal? (ID 18)*
Updates on long-term outcomes from patients in the trials	*"I read about the studies and the results of their initial trials, but I guess I don't know if there's been any documentation or anywhere I should look for follow up information on what's been learned now to date. Not just from the initial study but what's been learned from circling back to those original patients. Like is it still stalled or did they have to do another course?" (ID 5)*
Testimonials from people with AD to give a "human face" to statistics	*"But in terms of, just kind of on a daily basis how it would help you out or not help you out. Not just the—'cause the-the stuff we read was all like statistics on, you know, improved this for so many months or weeks or a year or whatever. There was nothing personal about how people who've taken it actually reacted to it and how they felt about it." (ID 15)*
Handout	*"Is there-are there papers? I don't wanna read all the big papers. But like is there a one or two side paper or something? It would be nice to have something like that from the doctors not the drug company." (ID 4)*
Website	*"A website? You mean, like, that'll give me information? Yeah. That would be helpful. Maybe to reassure☐—the information, I guess, is the best." (ID 17)*

Discussion

To our knowledge, this is the first study evaluating how people with AD understand and

deliberate about anti-amyloid mAb treatment (**Figure 2**). People with AD sought out information

to supplement knowledge from their physicians. They hoped that state-of-the-art treatment

would forestall inexorable cognitive decline and bridge to more effective treatments. They were

motivated by taking action that could buy more time, rather than passively letting nature take its

course. Diminished quality of life due to side effects, costs and burdens of treatment were

concerns. How people with AD balanced these tradeoffs was determined by personality, family,

and confidence in expertise.

Our findings help clarify a new stage in the dementia disease trajectory: decisions about

disease-modifying therapy. Before effective therapies were developed, qualitative studies in

people with AD focused on experiences and end-of-life care. Our findings have some

similarities, including a shared desire for greater guidance from clinicians on expected disease

course. Distinct from our findings, participants in prior studies focused on behavioral (versus

pharmacologic) interventions and the need for more palliative care.[18–21] Research examining

decision-making for people with dementia focuses on moving to a nursing home and advance

care planning. Some key facets overlap with our findings, like variation in perceptions of risks

and benefits, emotional complexity, and desire for autonomy. [22,23] In a recent systematic review

of 13 qualitative studies, true decisional autonomy was rare and confined to minor decisions,

whereas it was nearly universal in our population with mild symptoms.[24]

Our study participants were predominantly White, although they exhibited more educational,

income and geographic diversity. During the recruitment period, 97% of patients seen at the two

study sites' memory care clinics were White, which is in line with the population characteristics

at memory care clinics nationally.[25] It is also consistent with a recent analysis of real-world rates

of lecanemab prescription between January 2023 and February 2024, which revealed that 80%

of patients receiving lecanemab were White.[26] Decades of social science research have

empirically demonstrated that medical breakthroughs, even when beneficial to health,

exacerbate health inequalities.[27–29] Treatments that are expensive, complicated or time-

consuming—like anti-amyloid mAbs—are more likely to create disparities. This is because those

with socioeconomic advantages (educational, financial, social) will employ them to overcome

these barriers and obtain access to new treatments. The lack of diversity in patients seen in

these two memory care clinics and among participants in our study, therefore, suggests a

recapitulation of this phenomenon and an imperative to intervene. Subsequent studies will

examine whether our findings apply in settings with greater diversity that reflect all people with

AD. As anti-amyloid mAb science continues to advance, concerted efforts to reduce cost and

complexity of care should promote broad access and reduce disparities for people with AD.[29]

These results also reveal the sensitivity and skill with which clinicians caring for people with

dementia must approach conversations about treatment. Our results inform how communication

strategies could be adapted for this need. Some participants had misconceptions about

lecanemab's benefits, and potential side effects were considered minor. Given the gravity of a

dementia diagnosis, expressions of "unrealistic hope" may reflect normal coping or a lack of

cognitive understanding.[30,31] Some clinicians may be prone to overstating benefits and

minimizing harms, while others focus on potential adverse sequelae.[32] Our results also

underscore the salience and complexity of emotions in treatment decisions for people with AD,

such as grief over loss of identity and worries about burdening family. Evidence-based

communication training could address these needs, including sharing personalized risks and

benefits, naming uncertainty, using pictorials, creating space for questions, partnering in

decision-making, and using empathy.[33–37]

Our findings highlight that promoting informed decisions about anti-amyloid mAbs would benefit

from incorporating messaging channels outside the clinic. External resources were important for

people with AD. Heightened media attention to anti-amyloid mAbs may have led to passive

exposure.[38–40] Alternative resources may also have filled a vacuum while awaiting subspecialty

appointments or when incomplete information was provided by clinicians, a phenomenon seen

previously in dementia.[18] Interestingly, no people with AD reported hearing about lecanemab

through direct-to-consumer marketing, which is both common and can distort decision-

making.[41,42] Given patients' multiple streams of information, interventions to provide education

and decision-making support should use various modalities and start before and continue after

interaction with the healthcare system.

Patient-drawn parallels between lecanemab and chemotherapy raise potential opportunities to

learn from both the successes and failures of oncology care models. This comparison is apt

given the rapid transformation of AD care into a landscape not unlike oncology care. People with

dementia who might previously have felt stigmatized and hopeless about their diagnosis may

now identify with cancer patients who often have access to effective therapies and are valorized

for their bravery.[43] Clinicians caring for people with memory loss must now add responsibilities

that are common among oncologists like in-depth side effect monitoring.[44] Healthcare systems

must now provide services common in oncology care, including helping people with dementia

navigate complicated visit schedules. Lessons learned about the limitations of oncology care

models can be applied, such as the potential for financial considerations to affect treatment

decisions and the importance of eliciting patients' values and aligning care to meet them.[45,46]

Successful aspects ofoncology models could be adapted and their pitfalls avoided to support

people with dementia, clinicians and healthcare systems in making informed, cost-conscious

and goal-concordant decisions about anti-amyloid mAbs.[47,48]

Limitations

Although we reached data saturation, our study may not capture the full spectrum of patient

perspectives. Aspects of decision-making overlapped with those in previous studies in

dementia, suggesting broader relevance.[18–21]Our participants had some of the earliest access to

anti-amyloid mAbs. These early adopters may not reflect all people with AD. For example, we

identified consistent themes of self-efficacy, autonomy, and eagerness to start. Our findings

could simply be interpreted as natural enthusiasm in response to relief from a devastating—and

previously unmodifiable—diagnosis. Additionally, participants were recruited from memory care

clinics at academic medical centers who may differ from those in community settings or with

limited access to specialized care. As noted above, our sample lacked racial and ethnic

diversity so may not be generalizable, and future studies targeting underrepresented

populations will advance our understanding of shared decision-making. Future studies should

also examine how patients' cognitive status affects their perception of risks and benefits and

decision-making. Patients' views on treatment can evolve with time; longitudinal studies

examining changes in decision-making would also offer important insights.

Conclusions

In this first qualitative study of front-line people with dementia considering treatment with anti-

amyloid mAbs, we found variation in hopes for treatment, information sought and obtained, and

contextual factors in decision-making, supporting the need for an individualized approach.

Identifying these features can help identify targets for interventions to improve individualized

decision-making for anti-amyloid mAb treatment.

Acknowledgements

Author Contributions

Anna L Parks, MD (Conceptualization; Data curation; Formal analysis; Funding acquisition;

Investigation; Methodology; Project administration; Resources; Supervision; Validation; Writing –

original draft)

Ayush Thacker, BS (Data curation; Formal analysis; Methodology; Project administration;

Writing – review & editing)

Daniel Dohan, PHD (Formal analysis; Investigation; Methodology; Resources; Supervision;

Writing – review & editing)

Liliana A Ramirez Gomez, MD (Conceptualization; Investigation; Writing – review & editing)

Christine S Ritchie, MD, MSPH (Resources; Supervision; Writing – review & editing)

Joanna Paladino, MD (Conceptualization; Data curation; Formal analysis; Investigation;

Methodology; Writing – original draft)

Sachin J Shah, MD, MPH (Conceptualization; Formal analysis; Funding acquisition;

Methodology; Writing – original draft)

Statements and Declarations

Ethical considerations

The institutional review boards of participating institutions approved this study as exempt.

Consent to participate

Study participants completed verbal informed consent for participation.

Consent for publication

Not applicable

Declaration of conflicting interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship,

and/or publication of this article.

Funding statement

Dr. Shah reports research funding from NIA (K76AG074919). Dr. Parks reports research funding
from NIA (K76AG083304).
431

432 *Data availability*

433 The data supporting the findings of this study are available within the article and/or its

434 supplemental material

Figures

Figure 1 Timeline of Recruitment and Interviews in Relation to Clinical Evaluation for People with AD Considering Anti-Amyloid Monoclonal Antibody Treatment

Figure 2 Major Themes, Subthemes and Implications Identified in People with AD Considering Anti-Amyloid Monoclonal Antibody Treatment

Supplement

Appendix 1 COREQ checklist
Appendix 2 Master coded thematic statements

References

1. van Dyck CH, Swanson CJ, Aisen P, et al. Lecanemab in Early Alzheimer's Disease. *New England Journal of Medicine* 2023; 388: 9–21.

2. Yadollahikhales G, Rojas JC. Anti-Amyloid Immunotherapies for Alzheimer's Disease: A 2023 Clinical Update. *Neurotherapeutics* 2023; 20: 914–931.

3. Sperling RA, Jack CR, Black SE, et al. Amyloid-related imaging abnormalities in amyloid-modifying therapeutic trials: recommendations from the Alzheimer's Association Research Roundtable Workgroup. *Alzheimers Dement* 2011; 7: 367–385.

4. Ko D, Pascual-Leone A, Shah SJ. Use of Lecanemab for Patients With Cardiovascular Disease: The Challenge of Uncertainty. *JAMA*. Epub ahead of print 15 March 2024. DOI: 10.1001/jama.2024.2991.

5. Cummings J, Apostolova L, Rabinovici GD, et al. Lecanemab: Appropriate Use Recommendations. *J Prev Alzheimers Dis* 2023; 10: 362–377.

6. Arbanas JC, Damberg CL, Leng M, et al. Estimated Annual Spending on Lecanemab and Its Ancillary Costs in the US Medicare Program. *JAMA Intern Med* 2023; 183: 885–889.

7. Widera EW, Brangman SA, Chin NA. Ushering in a New Era of Alzheimer Disease Therapy. *JAMA* 2023; 330: 503–504.

8. Sarkisian CA, Romanov A, Mafi JN. Talking With Patients About the New Anti-amyloid Alzheimer Disease Medications. *Ann Intern Med* 2024; 177: 246–248.

9. Chin NA, Widera E, Brangman SA, et al. Monoclonal anti-amyloid antibodies for the treatment of Alzheimer's disease and the hesitant geriatrician. *Journal of the American Geriatrics Society* 2024; 72: 643–645.

10. Shay LA, Lafata JE. Where is the evidence? A systematic review of shared decision making and patient outcomes. *Med Decis Making* 2015; 35: 114–131.

11. Stacey D, Légaré F, Boland L, et al. 20th Anniversary Ottawa Decision Support
Framework: Part 3 Overview of Systematic Reviews and Updated Framework. *Med*
Decis Making 2020; 40: 379–398.

12. Tong A, Sainsbury P, Craig J. Consolidated criteria for reporting qualitative research
(COREQ): a 32-item checklist for interviews and focus groups. *International*
Journal for Quality in Health Care 2007; 19: 349–357.

13. Vaismoradi M, Turunen H, Bondas T. Content analysis and thematic analysis:
Implications for conducting a qualitative descriptive study. *Nurs Health Sci* 2013;
15: 398–405.

14. Braun V, Clarke V. Using thematic analysis in psychology. *Qualitative Research in*
Psychology 2006; 3: 77–101.

15. Malterud K, Siersma VD, Guassora AD. Sample Size in Qualitative Interview
Studies: Guided by Information Power. *Qual Health Res* 2016; 26: 1753–1760.

16. Hennink MM, Kaiser BN, Marconi VC. Code Saturation Versus Meaning Saturation:
How Many Interviews Are Enough? *Qual Health Res* 2017; 27: 591–608.

17. What is an adequate sample size? Operationalising data saturation for theory-
based interview studies - PubMed, https://pubmed.ncbi.nlm.nih.gov/20204937/
(accessed 11 July 2024).

18. Shafir A, Ritchie CS, Garrett SB, et al. "Captive by the uncertainty" – Experiences
with anticipatory guidance for people living with dementia and their caregivers at a
specialty dementia clinic. *J Alzheimers Dis* 2022; 86: 787–800.

19. Harrison KL, Garrett SB, Halim M, et al. "I didn't sign up for this": perspectives from
persons living with dementia and care partners on challenges, supports, and
opportunities to add geriatric neuropalliative care to dementia specialty care. *J*
Alzheimers Dis 2022; 90: 1301–1320.

20. Bunn F, Goodman C, Sworn K, et al. Psychosocial Factors That Shape Patient and
Carer Experiences of Dementia Diagnosis and Treatment: A Systematic Review of
Qualitative Studies. *PLoS Med* 2012; 9: e1001331.

21. Prorok JC, Horgan S, Seitz DP. Health care experiences of people with dementia
and their caregivers: a meta-ethnographic analysis of qualitative studies. *CMAJ*
2013; 185: E669–E680.

22. Young JA, Lind C, Orange J. A qualitative systematic review of experiences of
persons with dementia regarding transition to long-term care. *Dementia* 2021; 20:
5–27.

23. Sellars M, Chung O, Nolte L, et al. Perspectives of people with dementia and
carers on advance care planning and end-of-life care: A systematic review and
thematic synthesis of qualitative studies. *Palliat Med* 2019; 33: 274–290.

24. Bhatt J, Walton H, Stoner CR, et al. The nature of decision-making in people living
with dementia: a systematic review. *Aging & Mental Health* 2020; 24: 363–373.

25. Boyd ND, Naasan G, Harrison KL, et al. Characteristics of people with dementia
lost to follow-up from a dementia care center. *Int J Geriatr Psychiatry*; 37. Epub
ahead of print January 2022. DOI: 10.1002/gps.5628.

26. Research T. Real-world prescribing of lecanemab in the United States. *Truveta*,
https://www.truveta.com/blog/research/ispor-2024-lecanemab/ (2024, accessed 9
October 2024).

27. P GD, N LD. A Theory of Health Disparities and Medical Technology. *The BE
Journal of Economic Analysis & Policy* 2005; 4: 1–30.

28. Lutfey K, Freese J. Toward Some Fundamentals of Fundamental Causality:
Socioeconomic Status and Health in the Routine Clinic Visit for Diabetes. *American
Journal of Sociology* 2005; 110: 1326–1372.

29. Social Conditions as Fundamental Causes of Health Inequalities: Theory,
Evidence, and Policy Implications - Jo C. Phelan, Bruce G. Link, Parisa Tehranifar,
2010, https://journals-sagepub-
com.ezproxy.lib.utah.edu/doi/10.1177/0022146510383498 (accessed 9 October
2024).

30. Patients' Expectations of the Benefits and Harms of Treatments, Screening, and
Tests: A Systematic Review | Nursing | JAMA Internal Medicine | JAMA Network,
https://jamanetwork.com/journals/jamainternalmedicine/article-abstract/2038981
(accessed 12 June 2024).

31. Brenner K, Greer JA, Jackson V, et al. Development of a Prognostic Awareness
Impact Scale for Patients with Advanced Cancer. *J Palliat Med* 2022; 25: 445–454.

32. Morgan DJ, Pineles L, Owczarzak J, et al. Clinician Conceptualization of the
Benefits of Treatments for Individual Patients. *JAMA Network Open* 2021; 4:
e2119747.

33. Barry MJ, Edgman-Levitan S. Shared decision making--pinnacle of patient-
centered care. *N Engl J Med* 2012; 366: 780–781.

34. Dahm MR, Cattanach W, Williams M, et al. Communication of Diagnostic
Uncertainty in Primary Care and Its Impact on Patient Experience: an Integrative
Systematic Review. *J Gen Intern Med* 2023; 38: 738–754.

35. Paladino J, Bernacki R, Neville BA, et al. Evaluating an Intervention to Improve
Communication Between Oncology Clinicians and Patients With Life-Limiting
Cancer: A Cluster Randomized Clinical Trial of the Serious Illness Care Program.
JAMA Oncol 2019; 5: 801–809.

36. Paladino J, Koritsanszky L, Nisotel L, et al. Patient and clinician experience of a
serious illness conversation guide in oncology: A descriptive analysis. *Cancer Med*
2020; 9: 4550–4560.

37. Braddock CH, Edwards KA, Hasenberg NM, et al. Informed decision making in
outpatient practice: time to get back to basics. *JAMA* 1999; 282: 2313–2320.

38. Allen A. The real costs of the new Alzheimer's drug, Leqembi — and why taxpayers
will foot much of the bill - CBS News, https://www.cbsnews.com/news/alzheimers-
drug-leqembi-cost-patients-taxpayers/ (2023, accessed 12 June 2024).

39. Leqembi and Alzheimer's: What to Know About the New Drug, Treatment and
Benefits - The New York Times,
https://www.nytimes.com/2024/06/10/well/mind/alzheimers-treatment-leqembi.html
(accessed 12 June 2024).

40. J, o, n, et al. An experimental Alzheimer's drug outperforms one just approved by
the FDA. *NPR*, 17 July 2023, https://www.npr.org/sections/health-
shots/2023/07/17/1188075646/donanemab-experimental-alzheimers-drug-
outperforms-lecanemab-leqembi (17 July 2023, accessed 12 June 2024).

41. Schwartz LM, Woloshin S. Medical Marketing in the United States, 1997-2016.
JAMA 2019; 321: 80–96.

42. O'Donoghue AC, Johnson M, Sullivan HW, et al. Aging and Direct-to-Consumer
Prescription Drug Television Ads: The Effects of Individual Differences and Risk
Presentation. *Journal of Health Communication* 2019; 24: 368–376.

43. Robb KA, Simon AE, Miles A, et al. Public perceptions of cancer: a qualitative study
of the balance of positive and negative beliefs. *BMJ Open* 2014; 4: e005434.

44. Garrett SB, Matthews TM, Abramson CM, et al. Before Consent: Qualitative
Analysis of Deliberations of Patients With Advanced Cancer About Early-Phase
Clinical Trials. *JCO Oncol Pract* 2020; 16: e56–e63.

45. Mitchell AP, Rotter JS, Patel E, et al. Association Between Reimbursement
Incentives and Physician Practice in Oncology: A Systematic Review. *JAMA Oncol*
2019; 5: 893–899.

46. Wright AA, Zhang B, Ray A, et al. Associations between end-of-life discussions,
patient mental health, medical care near death, and caregiver bereavement
adjustment. *JAMA* 2008; 300: 1665–1673.

47. Shickh S, Leventakos K, Lewis MA, et al. Shared Decision Making in the Care of
Patients With Cancer. *Am Soc Clin Oncol Educ Book* 2023; e389516.

48. Henrikson NB, Shankaran V. Improving Price Transparency in Cancer Care. *J
Oncol Pract* 2016; 12: 44–47.

584
585

Major Themes	Subthemes	Implications
People with dementia sought and obtained information from different sources	• Advocacy organizations, the Internet and the media as resources for self-study • Variably detailed information from clinicians • Motivating interactions with treated patients and informed family members	Incorporate messaging approaches and channels outside the clinical encounter and healthcare system.
Hopes, expected benefits, and the existential threat of dementia driving willingness and readiness to start anti-amyloid mAbs.	• Stalling, reversal or cure of cognitive decline • Optimism drawn from mechanism of action and novelty compared to non-pharmacologic options • Stopgap until more effective therapy • Motivated by desire to buy time for enjoyable activities, preserve independence, and do "something" vs nothing	Apply communication skills that balance validating the importance of hope with accurate portrayal of personalized risks and benefits.
Individual traits and preferences, family factors, and degree of trust in expertise influence how people with dementia balance risks and benefits.	• Impact of personality • Perceived inevitability of decline • Weighing side effects, costs and logistical burdens • Importance of supportive families • Trust in expertise • Family and physicians as sources of pressure • Making calculated risks and conscious tradeoffs • Sense of urgency to decide • Autonomy	Adapt communication training that acknowledges and builds upon strong emotions. Capitalize on patient-identified parallels with oncology and consider adapting oncology models of care.