Running title: High Seroprevalence to *Aedes*-borne arboviruses in Ethiopia

Keywords: dengue, Zika, chikungunya, *Aedes aegypti*, Ethiopia, arbovirus, seroprevalence

Title: High Seroprevalence to *Aedes*-borne arboviruses in Ethiopia: a Cross-sectional Survey in 2024

Authors: *[1,2]Daniel M. Parker; [3]Werissaw Haileselassie; [4]Temesgen Sisay Hailemariam; [5]Arsema

Workenh; [3]Salle Workineh; [1]Xiaoming Wang; [1]Ming-Chieh Lee; [1,2]Guiyun Yan

Affiliations:

1. Department of Population Health and Disease Prevention; University of California, Irvine; U.S.A

2. Department of Epidemiology & Biostatistics; University of California, Irvine; U.S.A

3. School of Public Health, College of Health Sciences; Addis Ababa University; Addis Ababa,

Ethiopia.

4. Medical Diagnostic Laboratory, Tikur Anbessa Specialized Hospital; College of Health

Sciences, Addis Ababa University; Addis Ababa Ethiopia

5. Dil chora General Hospital, Dire Dawa Health Bureau, Dire Dawa City, Ethiopia

Author email addresses:

DMP: dparker1@hs.uci.edu ; ORCiD: https://orcid.org/0000-0002-5352-7338

WH: werissawhaileselassie@yahoo.com

TSH: temesegen.sisay@aau.edu.et

AW: Arsea2004@hotmail.com

SW: saleabera@yahoo.com

XW: xiaomiw1@uci.edu

ML: mingchil@uci.edu

GY: guiyuny@hs.uci.edu

*corresponding author: Daniel M. Parker; dparker1@hs.uci.edu ; ORCiD: https://orcid.org/0000-0002-

5352-7338

ABSTRACT

Aedes-borne diseases infect millions of people each year. In the last decade several arbovirus outbreaks

have been reported in Ethiopia. Arbovirus diagnosis and surveillance is lacking and the true burden is

unknown. In this study we conducted a seroprevalence survey using a commercially available test kit

that that tests for immunological responses to IgM and IgG for dengue (DENV), Zika (ZIKV), and

chikungunya (CHIKV) viruses in Dire Dawa city, eastern Ethiopia. We found a high IgG

seroprevalence for DENV (76%), CHIKV (44%), and ZIKV (38%), and <20% IgM seropositivity

across all viruses. As a comparison, we conducted serosurveillance in Addis Ababa, the national capital

with no reported history of arbovirus outbreaks. The highest seropositivity we found was to IgM for

DENGV at approximately 3%. Our results suggest both past and recent widespread exposure to these

arboviruses, underscoring the need for improved surveillance and public health interventions in

Ethiopia.

BACKGROUND

Aedes borne diseases contribute significantly to the global burden of disease, and are especially prevalent in tropical and subtropical settings. *Ae. aegypti*, and to a lesser extent *Ae. albopictus*, are the major vectors of disease and three viruses of primary concern are are dengue virus (DENV, from genus *Orthoflavivirus*), Zika (ZIKV, from genus *Orthoflavivirus*), chikungunya (CHIKV, from genus *Alphavirus*).

DENV, CHIKV, and ZIKV viruses each represent significant public health challenges with distinct impacts and distributions. DENV, endemic in tropical and subtropical regions, is responsible for an estimated 400 million infections annually, with severe cases leading to dengue hemorrhagic fever and dengue shock syndrome[1]. CHIKV was first detected in Tanzania in 1952 and in recent years has caused epidemics in Africa, Asia, Europe, and the Americas[2,3]. The disease is characterized by debilitating joint pain and arthritis that can last weeks to months, with significant economic implications on top of health impacts[4]. ZIKV was first detected in Uganda in 1947 and in the last decades there have been epidemics in Africa, Asia, the Americas, and the Pacific[5]. While most ZIKV infections are mild, the disease has been associated with congenital abnormalities (e.g. microcephaly) when transmitted from mother to fetus[6–8]. Collectively, these mosquito-borne viruses cause substantial morbidity, impact quality of life, and strain healthcare systems globally.

While these viruses cause considerable disease globally, infections are presumptively diagnosed in much of the world, based on symptoms and without laboratory confirmation of the cause of disease. Dengue and chikungunya infections result in similar symptoms and may be confused in some settings in the absence of laboratory confirmation. Zika viruses are often asymptomatic. All three may be drastically under-diagnosed in some settings and surveillance is lacking in much of the African continent so that the true burden of these diseases is unknown.

Seroprevalence surveys offer the possibility to evaluate both the long- and short-term

epidemiology of these diseases. Antibodies to IgM typically emerge shortly after infection (normally

within days for each virus) and can persist for months. Antibodies to IgG typically emerge weeks after

infection and can persist for years (in the case of dengue, may be lifelong). Seroprevalence estimates

from IgM are therefore normally indicative of recent epidemiology of these diseases whereas

seroprevalence estimates of IgG are indicative of long-term dynamics. In some high transmission

settings, most of the population will be seropositive before reaching adulthood[9,10]. Cross-reactivity can

complicate seroprevalence surveys that attempt to distinguish exposure to these three viruses[11].

Several dengue fever outbreaks have been documented in Ethiopia since 2013, with recurrent

outbreaks occurring in the Eastern part of the nation[12–15]. Chikungunya outbreaks have also been

reported more recently in the Eastern part of the nation[16,17], and seropositivity among symptomatic

individuals has been reported from the Northwest[18] and more recently in the Northeast part of the

nation in patients with suspected malaria infections[19]. Dire Dawa, a major city in the Eastern part of

Ethiopia has been repeatedly impacted by dengue infections and more recently by a major chikungunya

outbreak that infected over 40,000 individuals[16]. Entomological surveys have confirmed the presence of

Ae. aegypti in this city as well[20]. Research on Zika infections in this setting is scarce[21]. Surveillance is

lacking for all three viruses.

The primary objective of this research were to assess the seropositivity to DENGV, CHIKV, and

ZIKV in Dire Dawa, Ethiopia. Since age patterns in seropositivity can reflect both recent and historical

transmission intensity, our goal was also to examine potential associations between age and

seropositivity. We conducted secondary analyses of cross-reactivity.

DATA AND METHODS

Study location

Dire Dawa is a major trade city in Eastern Ethiopia. The city has a hot, arid climate with a hot

season from February through June and a two rainy seasons: a major one from July through September

and a minor rainy season In March – May. Temperatures normally range from 58F to 91F (14.4C to

32.7C). The wettest month is normally August, with a typical month experiencing a 5 inches (12.7cm)

of cumulative precipitation. The estimated population of the city in 2024 is 485,558 and the city is at

approximately 1,276 meters elevation.

We conducted a seroprevalence survey in Dire Dawa in March 2024. A total of 340 individuals

were recruited for the survey in Dire Dawa. Individuals were selected using a household registry and a

table of random numbers, with households being the unit of sampling. In a selected household, all

individuals were invited to participate in the survey regardless of age. We specifically targeted kebeles

(administrative units) with known dengue outbreaks in previous years for this survey.

For comparison, we conducted a secondary seroprevalence survey on the outskirts of Addis

Ababa using the same approach (selecting individuals by household). Addis Ababa is the capital and

largest city in Ethiopia, with an estimated population in 2024 of 5,703,628. In contrast to Dire Dawa,

Addis Ababa has a mild and temperate climate, rarely experiencing temperatures above 80 F (26.7C)

and typically ranging from 48F to 75F (8.9C to 23.9C). The rainy season follows the same general

pattern as Dire Dawa, but with August typically experiencing 8 inches (20.3cm) of cumulative

precipitation. Addis Ababa is at an elevation of approximately 2,355 meters.

For this secondary survey we recruited a total of 180 individuals to the seroprevalence survey.

We hypothesized that exposure to *Ae. aegypti* and associated viruses would be lower in Addis Ababa

than in Dire Dawa and wanted to conduct this secondary survey to compare overall seroprevalence

results to Dire Dawa, in part because we lacked a diagnostic control. We targeted 2 kebeles in Addis

Ababa for this secondary survey.

Seropositivity test

We used the Chembio DPP® ZCD IgM/IgG system for the seroprevalence survey[22]. This system is an immunochromatographic test, meant for detection and differentiation of IgG and IgM antibodies to DENV, CHIKV, and ZIKV (i.e. each individual potentially testing positive for 6 different outcomes). The test kit comes with a reader, which is used 10 to 15 minutes after adding buffer to the test device.

The reader for the rapid diagnostic test also outputs a numeric value of the intensity of the immunological response. The numeric value corresponds to the intensity of the colored lines on the test device, which are proportional to the presence of antibodies. The reader uses photometry to measure light reflectance from the lines on the test strip, with the intensity of reflected light being inversely proportional to the relevant antibody responses. This quantitative measure provides an opportunity to look at patterns of cross-reactivity across viruses and immunological measures.

Questionnaire

The survey team visited each household with a tablet that was loaded with basic demographic questions (age and gender) and test responses, and test kits.

Generalized Estimating Equations

We used a generalized estimating equations (GEE) model to estimate the overall seroprevalence to IgG and IgM for all three viruses, accounting for the household clustering design of the study by specifying an exchangeable correlation structure. We used bootstrapping to obtain robust confidence intervals around the seroprevalence estimates. We then used another GEE model to estimate age-specific seropositivity to IgG and IgM to each of the three viruses (DENGV, CHIKV, and ZIKV). To model the relationship between age and seropositivity, we incorporated natural spline functions,

allowing us to capture potential non-linear patterns in seropositivity across age. The model estimated

the probability of being seropositive for each virus as a function of age, adjusted for household

clustering. We then used these model estimates to generate predicted probabilities of seropositivity

across age groups. Confidence intervals for these predictions were calculated by deriving the standard

errors of the linear predictor, which were then transformed to the probability scale using the logistic

function. These predicted probabilities, along with their 95% confidence intervals, were plotted to

visualize how IgG and IgM seropositivity changes with age for each virus, while accounting for

clustering at the household level.

Assessment of cross-reactivity

We used linear mixed models, with the quantitative immunological measures for each of the

viruses and for both IgG and IgM as outcome variables, and with the other IgG or IgM measures for

each virus as predictors, while accounting for household clustering using a random intercept for

household. For models that used IgM measures as the outcome we also included IgG measures as

covariates because we hypothesized that previous exposure to other viruses could potential impact

immunological reactions to a more recent infection.

Assessment of household clustering

The use of linear mixed models in our assessment of cross-reactivity also allowed us to

investigate household clustering in immunological responses. To quantify household clustering we used

the intraclass correlation coefficient (ICC). The ICC quantifies the proportion of variance attributable

to between-group differences (in this case, households) compared to within-group differences

(individuals). The mixed models were fitted with random intercepts for households, allowing us to

estimate the variance in responses across households. By comparing the random intercept variance to

the total variance (random intercept variance plus residual variance), we calculated the ICC for each

model. This provided insight into how much of the variation in IgG and IgM responses could be

attributed to clustering within households versus individual-level variability.

Ethics review

Ethics approval was obtained from the Institutional Review Board of the College of Health Sciences,

Addis Ababa University. Verbal and written informed consent were required from individuals before

participating in the study. Assent/consent for participation for those under 18 years old was granted by

their parent or legal guardian.

RESULTS

Overall seroprevalence

A total of 340 individuals were recruited for the seroprevalence survey in Dire Dawa.

Seropositivity to dengue virus IgG was 76% (95% Cis: 0.71 – 0.80). Chikungunya and Zika IgG

seropositivity were 44% (0.38 – 0.50) and 38% (0.31 – 0.44), respectively. Seropositivity to IgM

antibodies, typically indicating recent infections, was much lower (**Table 1**). Dengue and chikungunya

IgM was approximately the same (0.17 (0.13 – 0.21) versus 0.18 (0.13 – 0.22), respectively). Zika IgM

was much lower, at approximately 0.03 (0.02 – 0.05).

In comparison, few individuals from the secondary seroprevalence survey in Addis Ababa tested

positive (not enough for accurate seroprevalence estimates). A total of 180 individuals were recruited in

this survey and out of them 5 tested positive for DENV IgG, 2 for ZIKV IgG, and 1 for CHIKV IgG.

Six tested positive for DENV IgM, 1 tested positive for ZIKV IgM, and 1 tested positive for CHIKV

IgM. Again, each of the ZIKV IgG positives were also DENV IgG positive. A total of 11 individuals

were found seropositive (combining IgG and IgM, and for all three viruses), these individuals ranged in

age from 25 to 36 (mean = 30.2; median = 28), and each reported travel to other regions in Ethiopia

(five specifically mentioned travel to Dire Dawa).

Age patterns in seropositivity

Age patterns in predicted probability of seropositivity (**Figure 1**) largely followed the age

patterns of the age-specific seropositivity calculations (**Supplemental Figure 1**). By age 10, 52% of all

sampled individuals were seropositive to dengue IgG. By age 40, approximately 86% of all sampled

individuals were seropositive to dengue IgG. By age 30, over half (51%) of all individuals were

seropositive to chikungunya IgG. Finally, by age 20 over 37% of all individuals were seropositive to

Zika IgG. The potential cross-over between IgM for dengue and chikungunya was likewise evident in

these estimates, though the confidence bands around the estimates were sufficiently large to warrant

caution in interpreting this result.

Potential for cross reactivity

All Zika IgG positive individuals were also dengue IgG positive, suggesting a potential for

cross-reactivity. Indeed, both are flaviviruses (i.e. genus *Orthoflavivirus*), and so cross-reactivity would

not be surprising. 253 of the tested individuals were dengue IgG seropositive and 126 were Zika IgG

positive.

An analysis of the quantified immunological response to IgG antibodies for Zika and dengue

showed that individuals who were categorized as Zika IgG negative but dengue IgG positive had very

low Zika IgG responses. Those who were both Zika IgG and dengue IgG positive had much higher

responses than those who were dengue IgG positive but not Zika IgG positive (**Supplemental Figure**

2).

Individuals who were dengue IgG positive and Zika IgG positive had significantly higher

quantitative dengue IgG values when compared to those who were dengue IgG reactive but Zika IgG

non-reactive (**Supplemental Table 2C**). In general, cross reactivity was high with regard to

quantitative dengue IgG outcomes for those who have had a previous Zika exposure. The inverse (cross

reactivity where dengue virus exposure increases Zika IgG) appears minimal in comparison. We

therefore hypothesize that the Zika IgG positives are not solely the results of cross-reactivity with

dengue virus but rather indication of true previous infections with Zika virus. Furthermore, not all Zika

IgM positives were also dengue IgM positive.

We generated a Spearman's correlation matrix from all possible immunological outcomes

(**Figure 2**). The strongest associations were between Zika IgG and dengue IgG, chikungunya IgG and

chikungunya IgM, followed by Zika IgG and dengue IgM, and dengue IgG and IgM.

Our linear mixed models analyses (**Supplemental Tables 1 and 2**) showed that Zika IgG was a

strong predictor of dengue IgG values (**Supplemental Table 1**). Chikungunya IgG was also associated

but the association was smaller than with Zika IgG. Dengue IgG was a strong predictor of Zika IgG.

Chikungunya IgG was negatively associated with Zika IgG. Finally, Zika IgG was positively associated

with chikungunya IgG values while dengue IgG was negatively associated with chikungunya IgG.

Zika IgM was positively associated with dengue IgM, as was chikungunya IgM (**Supplemental**

Table 2). Likewise, dengue and chikungunya IgM values were positively associated with Zika IgM.

Finally, dengue and Zika IgM were both positively associated with chikungunya IgM. IgG values were

statistically significant predictors of each IgM outcome, but their effect sizes were small.

Household clustering

The ICC values revealed varying degrees of household clustering for the different viruses and

immune responses. For DENGV, we observed moderate clustering in IgG responses (ICC = 10.3%) and

a low degree of clustering for IgM responses (ICC = 1.8%). CHIKV showed a similar pattern, with

clustering in IgG (ICC = 7.1%) and a stronger clustering effect in IgM responses (ICC = 14.0%). Zika,

on the other hand, displayed no household-level clustering for either IgG or IgM responses (ICC = 0

for both). These results suggest that household membership affected the variability in DENGV and

CHIKV responses to a moderate extent, while it had no impact on ZIKV responses.

DISCUSSION

Using a cross-sectional seroprevalence survey from Dire Dawa, Ethiopia we found and here

report a high seroprevalence to three arboviruses of concern: DENV, CHIKV, and ZIKV. These results

existed for both IgG and IgM suggesting both a history of widespread exposure to these viruses, and

recent exposure. The age patterns for DENV and CHIKV IgG seroprevalence in particular suggest that

most of the population in this setting has been exposed to these viruses by age 30 (over 50%

seroprevalence to CHIKV IgG and over 80% to DENGV IgG). We also found evidence of household

clustering with regard to DENGV and CHIKV exposure, but not ZIKV exposure.

While approximately 25% of all children ages 0 – 4 were seropositive to DENGV IgG, none

were seropositive to CHIKV IgG. In 2019 there was a major CHIKV outbreak in this setting, with over

40,000 individuals diagnosed with the disease[16]. It is therefore possible that much of the seroprevalence

to CHIKV IgG that we found was the result of this previous epidemic, or to the 2019 epidemic and

others. We did, however, find individuals in most ages (but not the youngest age group) who were

seropositive to CHIKV IgM, suggesting that there has potentially been some recent transmission of this

disease or a related alphavirus, among individuals from the study location. IgM to Chikungunya has

been shown to persist in some individuals over long periods of time (28 months post infection[23]) but we

believe this is unlikely to explain the IgM seropositivity to CHIKV from our survey (now 5 years after

the major 2019 epidemic).

While dengue and chikungunya epidemics have previously been reported in Ethiopia, data on

exposure to Zika is scarce. One survey from Gambella (in far West Ethiopia) estimated 2.9, 15.6, and

27.3% seroprevalence for Yellow Fever, chikungunya, and Zika virus[24]. Another report of Zika

seropositivity in Ethiopia[21]. Zika virus disease has been reported in traveler to Europe coming from

Kenya[25]. Zika virus was originally discovered on the African continent, but surveillance remains low[5].

One limitation of our work is the use of a single screening kit. It might have been preferable to

have another test (e.g. ELISA) for comparison on at least a subsample. A few other studies have used

the same test kit in research settings. One study suggested general low sensitivity of this test in

comparison to ELISA. However, specificity was high (sensitivity/specificity: Dengue 43.92/100%;

Zika 25.86/98.81%; CHIKV 37.78/99.35%). If the same pattern holds true in this setting it would

indicate that the estimated seroprevalence is an underestimate[26]. Another study focusing on IgM alone

found sensitivity/specificity of the ZCD test to Zika IgM was 79.0/97.1%; dengue was 90.0/89.2%; and

for chikungunya it was 90.6/97.2%[27]. We take the results of these previous studies to suggest that for

ZIKV and CHIKV in particular, our results may be underestimates.

Cross reactivity may have influenced some of our results. DENGV and ZIKV are both

flaviviruses, and cross-reactivity between the two is well-known but not completely understood[11,28,29]. If

other alphaviruses are circulating in this setting, they could potentially lead to false-positive CHIKV

seropositivity. Cross-reactivity can have clinical implications, as is already well-known within DENGV

infections across serotypes and has been hypothesized between DENGV and ZIKV infections. While

DENGV infections are known to provide long-term immunity, the immunity is sero-type specific and

infection by another serotype can lead to more severe disease outcomes (including death) through

antibody dependent enhancement (ADE)[30–32]. ADE may also occur across flaviviruses – with the

potential for previous ZIKV infection leading to severe dengue disease even in the primary infection by

DENGV[33]. Current dengue vaccines are only provided to populations that already have previous

exposure to DENGV (i.e. the vaccines should only be provided to seropositive individuals) as

vaccinating individuals who are seronegative can increase their risk of severe disease when they are

infected by DENGV[34].

Cross-reactivity is further complicated by the fact that these three viruses are all vectored by the

same mosquito (e.g. *Ae. aegypti*), meaning that individuals who have been infected with dengue in this

setting are likely to also have been at risk of chikungunya (which has been confirmed in this setting)

and Zika virus infection, if the respective viruses are present in, or introduced to, this location. That is,

individuals in this setting who acquire dengue infections are likely to also be at risk of CHIKV or

ZIKV infections, if those viruses are present, given the shared mosquito vector that is present in this

setting.

This seroprevalence survey was meant to look for age-specific seroprevalence in kebeles in Dire

Dawa that are known to have previously had dengue outbreaks. While we used a clustered random

sampling approach (randomly selecting households within the kebeles), our results may not be

representative of the entire city of Dire Dawa. It would be valuable to conduct further surveys that

would be representative of the city, as well as in other parts of Ethiopia.

In summary, we found evidence of widespread exposure to DENGV, CHIKV, and ZIKV in Dire

Dawa, Ethiopia. This aligns with reported outbreaks of dengue fever in this setting, and of the massive

chikungunya fever outbreak in 2019. We also found evidence of exposure to ZIKV, which has rarely

been reported from this nation. Not only did we find evidence of previous exposure to ZIKV, we also

found evidence of recent exposure (i.e. IgM seropositivity) to all three arboviruses. While we cannot

rule out the potential for cross-reactivity influencing our results, they are suggestive of a large burden

of *Ae. aegypti*-borne diseases in a setting where this important vector of disease is known to be present.

Repeated infections can in some cases lead to more severe disease outcomes, for example through

ADE. More research into arboviral diseases and their mosquito vectors in this setting is warranted both

to confirm our results and to develop a better understanding of the burden of different arboviral

diseases, any potential age and gender associated risk factors, and all so that public health interventions

can be formulated to address these diseases. As the world has experienced pandemic levels of *Aedes*-

borne disease several times since 2019, it is increasingly important to have increased surveillance for

these diseases and for public health agencies to plan accordingly.

ACKNOWLEDGMENTS

We are thankful for the community members and community health workers in Dire Dawa and Addis

Ababa who participated in this research.

FINANCIAL SUPPORT

This study was partially funded through a Council on Research, Computing and Libraries (CORCL)

pilot grant to DMP and through NIH grants D43 TW001505 and U19 AI129326 to GY.

334 **REFERENCES**

1. Bhatt S, Gething PW, Brady OJ, et al. The global distribution and burden of dengue. *Nature*. 2013;496(7446):504-507. doi:10.1038/nature12060

2. Chikungunya fact sheet. Accessed August 26, 2024. https://www.who.int/news-room/fact-sheets/detail/chikungunya

3. Staples JE, Breiman RF, Powers AM. Chikungunya Fever: An Epidemiological Review of a Re-Emerging Infectious Disease. *Clinical Infectious Diseases*. 2009;49(6):942-948. doi:10.1086/605496

4. Mora-Salamanca AF, Porras-Ramírez A, De la Hoz Restrepo FP. Estimating the burden of arboviral diseases in Colombia between 2013 and 2016. *International Journal of Infectious Diseases*. 2020;97:81-89. doi:10.1016/j.ijid.2020.05.051

5. Zika virus. Accessed August 26, 2024. https://www.who.int/news-room/fact-sheets/detail/zika-virus

6. Araújo TVB de, Ximenes RA de A, Miranda-Filho D de B, et al. Association between microcephaly, Zika virus infection, and other risk factors in Brazil: final report of a case-control study. *The Lancet Infectious Diseases*. 2018;18(3):328-336. doi:10.1016/S1473-3099(17)30727-2

7. Oliveira WK de, França GVA de, Carmo EH, Duncan BB, Kuchenbecker R de S, Schmidt MI. Infection-related microcephaly after the 2015 and 2016 Zika virus outbreaks in Brazil: a surveillance-based analysis. *The Lancet*. 2017;390(10097):861-870. doi:10.1016/S0140-6736(17)31368-5

8. Musso D, Ko AI, Baud D. Zika Virus Infection — After the Pandemic. *New England Journal of Medicine*. 2019;381(15):1444-1457. doi:10.1056/NEJMra1808246

9. Rodríguez-Barraquer I, Buathong R, Iamsirithaworn S, et al. Revisiting Rayong: Shifting Seroprofiles of Dengue in Thailand and Their Implications for Transmission and Control. *American Journal of Epidemiology*. 2014;179(3):353-360. doi:10.1093/aje/kwt256

10. Vongpunsawad S, Intharasongkroh D, Thongmee T, Poovorawan Y. Seroprevalence of antibodies to dengue and chikungunya viruses in Thailand. *PLOS ONE*. 2017;12(6):e0180560. doi:10.1371/journal.pone.0180560

11. Priyamvada L, Suthar MS, Ahmed R, Wrammert J. Humoral Immune Responses Against Zika Virus Infection and the Importance of Preexisting Flavivirus Immunity. *The Journal of Infectious Diseases*. 2017;216(suppl_10):S906-S911. doi:10.1093/infdis/jix513

12. Shimelis T, Mulu A, Mengesha M, et al. Detection of dengue virus infection in children presenting with fever in Hawassa, southern Ethiopia. *Sci Rep*. 2023;13(1):7997. doi:10.1038/s41598-023-35143-2

13. Gutu MA, Bekele A, Seid Y, et al. Another dengue fever outbreak in Eastern Ethiopia—An emerging public health threat. *PLoS Negl Trop Dis*. 2021;15(1):e0008992. doi:10.1371/journal.pntd.0008992

14. Mesfin Z, Ali A, Abagero A, Asefa Z. Dengue Fever Outbreak Investigation in Werder Town, Dollo Zone, Somali Region, Ethiopia. *Infect Drug Resist*. 2022;15:7207-7217. doi:10.2147/IDR.S368562

15. Degife LH, Worku Y, Belay D, Bekele A, Hailemariam Z. Factors associated with dengue fever outbreak in Dire Dawa administration city, October, 2015, Ethiopia - case control study. *BMC Public Health*. 2019;19(1):650. doi:10.1186/s12889-019-7015-7

16. Geleta D, Tesfaye N, Ayigegn H, Waldetensai A, Gemechu F, Amare H. Epidemiological Description of Chikungunya Virus Outbreak in Dire Dawa Administrative City, Eastern Ethiopia, 2019. *Int J Clin Exp Med Sci*. 2020;6(3):41-45. doi:10.11648/j.ijcems.20200603.13

17. Chikungunya disease outbreak affects 20,000 people in Ethiopia - Xinhua | English.news.cn.http://www.xinhuanet.com/english/2019-09/03/c_138362347.htm. Accessed August 26, 2024.

18. Ferede G, Tiruneh M, Abate E, et al. Evidence of chikungunya virus infection among febrile patients in northwest Ethiopia. *International Journal of Infectious Diseases*. 2021;104:183-188. doi:10.1016/j.ijid.2020.12.057

19. Zerfu B, Kassa T, Mamo G, Larrick JW, Legesse M. High seroprevalence of IgM antibodies against chikungunya among patients with acute febrile illness seeking healthcare in a malaria-endemic area in the Afar Region, Northeast Ethiopia. *SAGE Open Medicine*. 2024;12:20503121241276557. doi:10.1177/20503121241276557

20. Yared S, Gebressilasie A, Worku A, et al. Breeding habitats, bionomics and phylogenetic analysis of Aedes aegypti and first detection of Culiseta longiareolata, and Ae. hirsutus in Somali Region, eastern Ethiopia. *PLOS ONE*. 2024;19(1):e0296406. doi:10.1371/journal.pone.0296406

21. Mengesha Tsegaye M, Beyene B, Ayele W, et al. Sero-prevalence of yellow fever and related Flavi viruses in Ethiopia: a public health perspective. *BMC Public Health*. 2018;18(1):1011. doi:10.1186/s12889-018-5726-9

22. *Chembio DPP® ZCD IgM/IgG System*. Accessed September 18, 2024. https://chembio.com/wp-content/uploads/2021/12/MSDS-018-DPP-ZCD-IgM-IgG-System-65-9565-0-Rev-1.pdf

23. Costa DM do N, Coêlho MRCD, Gouveia PA da C, et al. Long-Term Persistence of Serum-Specific Anti-Chikungunya IgM Antibody - A Case Series of Brazilian Patients. *Rev Soc Bras Med Trop*. 54:e0855-2020. doi:10.1590/0037-8682-0855-2020

24. Asebe G, Michlmayr D, Mamo G, et al. Seroprevalence of Yellow fever, Chikungunya, and Zika virus at a community level in the Gambella Region, South West Ethiopia. *PLOS ONE*. 2021;16(7):e0253953. doi:10.1371/journal.pone.0253953

25. Gossner CM, Hallmaier-Wacker L, Briet O, et al. Arthropod-borne diseases among travellers arriving in Europe from Africa, 2015 to 2019. *Eurosurveillance*. 2023;28(7):2200270. doi:10.2807/1560-7917.ES.2023.28.7.2200270

26. Chakma R, Sriburin P, Sittikul P, et al. Arbovirus Seroprevalence Study in Bangphae District, Ratchaburi Province, Thailand: Comparison between ELISA and a Multiplex Rapid Diagnostic Test (Chembio DPP® ZCD IgG). *Tropical Medicine and Infectious Disease*. 2022;7(11):378. doi:10.3390/tropicalmed7110378

27. Boeras D, Diagne CT, Pelegrino JL, et al. Evaluation of Zika rapid tests as aids for clinical diagnosis and epidemic preparedness. *EClinicalMedicine*. 2022;49:101478. doi:10.1016/j.eclinm.2022.101478

28. Priyamvada L, Hudson W, Ahmed R, Wrammert J. Humoral cross-reactivity between Zika and dengue viruses: implications for protection and pathology. *Emerging Microbes & Infections*. 2017;6(1):1-6. doi:10.1038/emi.2017.42

29. Wen J, Shresta S. Antigenic cross-reactivity between Zika and dengue viruses: is it time to develop a universal vaccine? *Current Opinion in Immunology*. 2019;59:1-8. doi:10.1016/j.coi.2019.02.001

30. Guzman MG, Alvarez M, Halstead SB. Secondary infection as a risk factor for dengue hemorrhagic fever/dengue shock syndrome: an historical perspective and role of antibody-dependent enhancement of infection. *Arch Virol*. 2013;158(7):1445-1459. doi:10.1007/s00705-013-1645-3

31. Halstead SB. Neutralization and antibody-dependent enhancement of dengue viruses. *Adv Virus Res*. 2003;60:421-467. doi:10.1016/s0065-3527(03)60011-4

32. Pang T, Cardosa MJ, Guzman MG. Of cascades and perfect storms: the immunopathogenesis of dengue haemorrhagic fever-dengue shock syndrome (DHF/DSS). *Immunology & Cell Biology*. 2007;85(1):43-45. doi:10.1038/sj.icb.7100008

33. Bonheur AN, Thomas S, Soshnick SH, et al. A fatal case report of antibody-dependent enhancement of dengue virus type 1 following remote Zika virus infection. *BMC Infectious Diseases*. 2021;21(1):749. doi:10.1186/s12879-021-06482-0

34. Robinson ML, Durbin AP. Dengue vaccines: implications for dengue control. *Curr Opin Infect Dis*. 2017;30(5):449-454. doi:10.1097/QCO.0000000000000394

342 **FIGURES:**

344 **Figure 1**: Age specific probability of IgG and IgM seropositivity from the GEE models

346 **Figure 2**: Spearman correlation matrix for correlations between IgG and IgM for each arbovirus

 TABLES

367   **Table 1**: Seropositivity estimates and confidence intervals, from GEE models using an exchangeable

368   correlation structure to account for potential clustering in households.

		Serporevalence	
	Disease	**Estimate**	**95% Cis**
IgG	Dengue	0.76	(0.71 – 0.80)
	Chikungunya	0.44	(0.38 – 0.50)
	Zika	0.38	(0.31 – 0.44)
IgM	Dengue	0.17	(0.13 – 0.21)
	Chikungunya	0.18	(0.13 – 0.22)
	Zika	0.03	(0.02 – 0.05)

Spearman Correlation Matrix of IgG and IgM Values