Associations of Genetic Variants in the Dopamine Transporter with Problematic Sexual

Behavior and Reward Deficiency Syndrome

Authors

Shui Jiang, PhD

Jerome C. Foo, PhD

Xiuying Hu, BM, MSc

Bradley Green, PhD

Randolph Arnau, PhD

Pennie J. Carnes, MA

Behavioral Addictions Studies and Insights Consortium*

Rick Isenberg, MD

David Wishart, PhD

Patrick J. Carnes, PhD

Esther Fujiwara, PhD

Katherine J. Aitchison, BM BCh, PhD, FRCPsych

Affiliations

Department of Psychiatry, Faculty of Medicine and Dentistry, College of Health Sciences,

University of Alberta, Edmonton, AB, Canada (SJ, JCF, EF, KJA); Department of Genetic

Epidemiology in Psychiatry, Central Institute of Mental Health, Mannheim, Germany (JCF);

Neuroscience and Mental Health Institute, University of Alberta, Edmonton, AB, Canada (JCF,

KJA); Department of Medical Genetics, Faculty of Medicine and Dentistry, College of Health

Sciences, University of Alberta, Edmonton, AB, Canada (XH, KJA); Department of Psychology

and Counseling, University of Texas at Tyler, Tyler, TX, USA (BG); School of Psychology,

University of Southern Mississippi, Hattiesburg, MS, USA (RA); Gentle Path at the Meadows,

Wickenburg, AZ, USA (PJC, PC); Psychological Counselling Services, Ltd., Scottsdale, AZ,

USA (RI); Department of Biological Sciences, Faculty of Science, College of Natural and

Applied Sciences, University of Alberta, Edmonton, AB, Canada (DW); Women and Children's

Health Research Institute, University of Alberta, Edmonton, AB, Canada (KJA); and the

Psychiatry Section, Division of Clinical Sciences, Northern Ontario School of Medicine,

Thunder Bay, ON, Canada (KJA)

Correspondence to:

Shui Jiang, PhD

Tel number: 780-492-4018

Email: shui2@ualberta.ca

5-020, Katz Group Centre for Pharmacy and Health Research, Edmonton, AB, Canada

*Consortium authors in supplementary material

Sources of support/funding for the work

The work herein was supported by an Alberta Centennial Addiction and Mental Health Research

Chair and transitional funding (to KJA), Canada Foundation for Innovation (CFI), John R. Evans

Leaders Fund (JELF) grant (32147 - Pharmacogenetic translational biomarker discovery),

Alberta Innovation and Advanced Education Small Equipment Grants Program (to KJA), and a

research grant and philanthropic support from the American Foundation for Addiction Research

(to KJA). A Fulbright-Canada-Palix Foundation grant (to PC) assisted with his contributions to

study design, project management, and collaborative working. SJ was supported by an Alberta

Innovates Postdoctoral Recruitment Fellowship.

Conflicts of interest:

While Dr. Carnes was previously a Board member of the American Foundation for Addiction

Research, he is no longer a member. Moreover, neither the Foundation nor any other of the

funders played any role in study design, or in data analysis or interpretation thereof. All other

authors declare no conflicts of interest.

Abstract word count: < 250 (247)

Manuscript word count (not including references, tables, or figure legends): <3500 (3461)

Number of references: <40 (39)

ABSTRACT

Objectives

Problematic sexual behavior (PSB) is defined by recurrent sexual behaviors that are difficult to

control, causing social and functional impairments. PSB can co-occur with reward deficiency

syndrome (RDS), but this relationship remains unclear. RDS has been associated with the 10/10

genotype of 3' variable number tandem repeat (VNTR) in the dopamine transporter gene

SLC6A3, which is implicated in the reward pathway. This study investigates the genetic

relationship between PSB and RDS, testing their association with *SLC6A3* 3' VNTR genotype.

Methods

PSB patients from addiction treatment facilities (n=454), and comparison participants (with

PSB=82; without PSB, n=888) were recruited. PSB was measured by the Sexual Addiction

Screening Test-Revised (SAST-R) Core and RDS was measured using a composite variable from

a custom test battery. DNA was collected from saliva and buccal swabs. Genotyping was

performed using polymerase chain reaction (PCR), and regression analyses were conducted to

investigate the association of *SLC6A3* 3' VNTR genotype with PSB and RDS.

Results

The 10/10 genotype of *SLC6A3* 3' VNTR was associated with RDS in a combined analysis of all

groups, and with PSB only in comparison participants. In patients, rare *SLC6A3* 3' VNTR

genotypes (3-, 6-, 8-, 11-repeat alleles) were associated with PSB. No genotype showed

relationships to RDS in only PSB patients.

Conclusions

82 The link of the 10/10 genotype to RDS indicates that PSB could be a manifestation of RDS in

83 non-clinical populations; rare genotypes might be associated with clinical forms of PSB, together

84 with comorbid psychopathology.

86 **Keywords:** addiction; behavioral addiction; reward deficiency syndromes; genetics; the

87 dopamine transporter gene

1. INTRODUCTION

Problematic sexual behavior (PSB) is defined as difficult to control recurrent sexual behaviors that cause social and functional impairments.[1] PSB can be referred to as sex addiction,[2] or compulsive sexual behavior.[3] The scientific debate on the nosology and nature of PSB continues, but it is recognized that PSB often co-occurs with neuropsychiatric conditions such as reward deficiency syndrome,[4,5] possibly suggesting common underlying biology.

Reward deficiency syndrome (RDS), often described as an "octopus of behavioral dysfunction," is not a single mental health condition, but a collection of conditions including addictive, compulsive, and impulsive behaviors, and personality disorders, which share endophenotypes characterized by hypodopaminergia in the reward circuit.[6-8] Although the relationship between RDS and PSB is not well characterized, there is emerging evidence that they may overlap.[1] Jiang et al. (2023) reported that compared to those without PSB, university students with PSB showed significantly higher rates of conditions with known alterations in the dopaminergic reward pathway which would fall under the umbrella of RDS (e.g., Attention Deficit/Hyperactivity Disorder [ADHD], obsessive-compulsive disorder [OCD], and personality disorders).[1]

Addictive behaviors and symptoms are core features of RDS,[9] and there are several genetic variants associated with vulnerability, maintenance, and relapse in addiction.[10-12] It is possible that at least part of the phenotypic overlap between PSB and RDS may be driven by genetic variants common to the two. For example, in one study, 81% of sexual addiction patients reported that at least one parent had an addictive disorder, with 36% indicating that a parent had a sexual addiction.[13] One of the potential genetic variants for addictive disorders is the dopamine transporter gene *SLC6A3 (solute carrier family 6 member 3)*, located at chromosome 5p15.33.

There is a 40-base pair variable number tandem repeat (VNTR) in the 3' untranslated region of

SLC6A3 with 9- and 10- repeat alleles being the most common.[14] Studies have shown that

individuals with the *SLC6A3* 10/10 genotype express higher levels of dopamine transporter than

9-repeat allele carriers, and this is reflected in lower striatal dopamine among 10-repeat

carriers.[15]

The dopamine transporter mediates the reuptake of dopamine from the synaptic cleft,[16]

and the 10/10 genotype of the *SLC6A3* 3' VNTR has been implicated in various conditions

associated with an altered reward system, including ADHD and addictive disorders.[12,17,18] For

example, Cornish et al. (2005) reported that the 10/10 genotype was associated with higher

ADHD symptom scores in boys.[19] A meta-analysis by Ma et al., (2016) showed that individuals

carrying the 10/10 genotype were 17% less likely to quit smoking compared to those who was

the 9-repeat alleles.[10] Furthermore, a previous study indicated that the 10-repeat allele of the

SLC6A3 3' VNTR was associated with phenotypes directly related to sexual behavior. In a study

of men, Guo et al. (2007) reported that individuals with at least one 10-repeat allele of the

SLC6A3 3' VNTR showed an 80%-100% increase in the number of sexual partners compared to

those homozygous for the 9-repeat allele.[20,21] While the 10/10 variant seems to be more

consistently identified as the risk genotype, some studies probing gene-environment (G×E)

interactions pointed to a greater sensitivity of 9-repeat carriers to environmental influences. For

example, Erblich and colleagues (2004) reported that after stress, carriers of the 9-repeat allele

showed a cigarette craving four times greater than non-carriers.[22] This contrasts with findings of

potentially protective effects of the 10-repeat variant under adverse environmental influences

(stress or childhood trauma) as seen in ADHD,[19] smoking,[10] and number of sexual partners.[20,21]

However, these complex interactions have rarely been explored and contradictory findings exist

as well.[23]

Apart from common alleles such as 9- and 10-repeat, rare genetic variants, such as 3-, 5-,

7- , 8-, and 11-repeats are also observed.[14] Despite their low frequency and smaller contribution

to population-wide heritability compared to common variants, these rare variants can still

significantly increase individual risk of illness due to their pronounced impact on protein

function or expression.[24] A previous study including 471 children with or without ADHD found

that *SLC6A3* rare genotypes (8/10, 7/10 and 10/11) were associated with ADHD.[25] However,

research on these rare genotypes in candidate gene studies is limited, and these variants have

often been excluded due to their low frequency and the resulting small sample sizes.

The aim of the present study was to investigate the relationship between PSB and RDS at

the candidate gene level in PSB patients and in a non-clinical sample, with a focus on *SLC6A3*

3'VNTR genotype. We expected to extend the association between PSB and RDS that we had

previously reported in the non-clinical comparison participants to a clinical sample. We

hypothesized that the 10/10 genotype might be associated with PSB and/or with RDS.

2. METHODS

2.1.Patients and comparison participants

The clinical cohort comprised inpatients and outpatients receiving treatment for PSB at eight

addiction treatment facilities in the United States (n=512). The recruitment of comparison

participants (n=3530) was undertaken at a Canadian university as previously described.[1] Since

the allele frequencies of the *SLC6A3* 3' VNTR are observed to vary by ethnicity,[26] genotypic and

phenotypic analyses were restricted to individuals of European ancestry (n=1434, Supplemental

Digital Content, Figure 1).

All participants provided written informed consent. The Quorum Review Institutional

Review Board (Seattle, Washington; Protocol Number: 2016-001) and the University of Alberta

Research Ethics Board (Protocol: Pro00066552) approved the study.

2.2.Assessments

We screened for PSB with the 20-item Sexual Addiction Screening Test-Revised (SAST-R)

Core.[27] We measured four addictive dimensions of PSB: relationship disturbance caused by

sexual behaviors (SAST-R Core items 6, 8, and 16, and we also included item 26 from the

additional SAST-R subscales[27]), loss of control despite problems (SAST-R Core items 10, 12,

15, and 17), preoccupation about sex (SAST-R Core items 3, 18, 19 and 20), and affect

disturbance (such as anxiety and depression related to sexual behaviors, SAST-R Core items 4,

5, 11, 13 and 14). Each question is responded to with a binary choice (yes/no=1/0), and the

threshold for screening positive for PSB in the general population is 6.[27] The PSB variable we

used in all analyses (referred to as the "adjusted SAST-R Core") was the scale total minus the

first two questions, which inquire about a history of childhood sexual trauma and parental

trouble with sexual behavior, as previously described.[1] These items were excluded due to their

overlap with other variables in our analysis, such as "self-reported childhood trauma" and

"family history of PSB." This approach was taken to avoid redundancy and increase the

specificity of the PSB regression, even though there are a part of SAST-R Core.

To derive our RDS measure, several relevant conditions were assessed, including

nicotine dependence (Fagerström Test for Nicotine Dependence [FTND]), pathological gambling

(DSM-5 Pathological Gambling Diagnostic Form [DPGDF]), personality disorders (8-item

Standardized Assessment of Personality-Abbreviated Scale as a Self-Administered Screening

Test [SA-SAPAS]), ADHD (Adult ADHD Self-Report Scale [ASRS] version 1.1), compulsive

buying (Richmond Compulsive Buying Scale [RCBS]), and internet addiction (Internet

Addiction Test [IAT]), as previously described.[1] To represent the overall RDS construct, a

dummy coded binary variable was created (1=positive screening in any of the RDS components,

0=no). In addition, we collected self-reported information about demographic details, childhood

trauma (self-reported childhood physical, emotional and sexual trauma, and family history of

domestic violence, combined with item 1 from the SAST-R Core "sexually abused as a child",

1=yes, 0=no, as previously described[1]), past or current mental health diagnoses (1=yes, 0=no),

number of past or current mental health diagnoses, and number of mental health condition(s) in

the family history. A positive family history of PSB (1=yes, 0=no) was ascertained through item

2 of the SAST-R Core ("parental problems with sexual behavior") and self-report data, as

previously described.[1] Since OCD in particular is reported to be associated with PSB[1] and RDS,[4]

we also used available data from MINI-International Neuropsychiatric Interview (MINI) (n=715;

379 comparison participants without PSB, 72 comparison participants with PSB, 264 patients),

in combination with self-reported past/current OCD diagnoses (n=967, 448 comparison

participants without PSB, 89 comparison participants with PSB, 430 patients) to estimate rates of

OCD.

2.3. Genotyping

DNA was extracted from saliva using Oragene OG-500 kits (DNAGenotek, Kanata, Canada)

and/or from buccal swabs (Isohelix via D-Mark Biosciences, Toronto, Canada). Genotyping of

the 40 bp *SLC6A3* 3' VNTR was performed by a polymerase chain reaction (PCR). DNA was

amplified with an initial denaturation step at 95°C for 2 minutes, followed by 35 cycles of

denaturation at 95°C for 15 seconds, annealing at 64°C for 30 seconds, extension at 72°C for 1

minute, and a terminal extension step of 72°C for 5 minutes (modified from Vandenbergh et al.,

1992).[14] The primer sequences were: forward 5'-TGTGGTGTAGGGAACGGCCTGAG-3', and

reverse 5-CTTCCTGGAGGTCACGGCTCAAGG-3'.[14] PCRs were resolved on a 2.25% agarose

gel. The 9-repeat number was at 440 bp and the 10-repeat number was at 480 bp on the gel;

genotypes other than 9/9, 9/10, or 10/10 are considered rare genotypes.[14] Since the allele

frequencies of the *SLC6A3* 3' VNTR vary by ethnicity,[26] genotypic and phenotypic analyses

were conducted in only those of European ancestry.

2.4.Data analysis

Data were analyzed using STATA.[28] To examine the association of PSB, participants were

categorized into three groups: (1) PSB patients (n = 405; patients who scored ≥11 on the

unadjusted SAST-R Core); (2) Comparison participants with PSB (n=85) scoring ≥6 on the

unadjusted SAST-R Core,[27] and (3) Comparison participants without PSB (n=439) scoring <6 on

the unadjusted SAST-R Core and also screening negative on other relevant measurements.[27]

In the final analysis of PSB, a total of 929 participants (524 from the comparison group

and 405 from the patient group) were included, after the removal of 59 individuals who were not

of European ancestry or did not provide DNA. The SAST-R Core, developed based on a large

clinical cohort (665 men and 274 women) of inpatients and outpatients in treatment for PSB,

classifies individuals at a cut-point of 6, achieving a sensitivity of 0.817 and specificity of

0.778.[27,29] However, to enhance the specificity of the genetic association study, we selected a

cut-off score of 11 on the SAST-R Core to focus on individuals with more pronounced PSB

symptoms. This stricter threshold was implemented to ensure clearer signals for the genetic

analysis by minimizing ambiguous cases, thereby prioritizing the identification of true positives

over the risk of false negatives. Thus, we excluded 19 patients who scored <11 on the unadjusted

SAST-R Core.[28,29] In addition, to enhance the specificity of testing for associations with PSB, we

excluded a total of 432 participants who screened negative for PSB, but positive for RDS or

mental health conditions(s). This was done due to the consideration of potentially shared genetic

etiologies between RDS, PSB, and mental health conditions. For the validation of RDS, all

available data from patients (n=376) and comparison participants (n=668) were included.

2.4.1.   Descriptive statistics

To investigate if those with or without PSB have different clinical and demographic features,

group differences were examined with Pearson's χ^2 test or a Mann-Whitney U test or a Kruskal-

Wallis test, followed by Dunn's test. To test if the distribution of $SLC6A3$ genotype differed

between different groups, Pearson's χ^2 was conducted. Tetrachoric (for binary variables) and

Kendall's τ (for ordinal variables) correlations were used to examine the correlation of clinical

and demographic variables with severity of PSB.

2.4.2.   Regression analyses

To investigate the association of demographic and clinical variables with PSB and RDS,

backward stepwise logistic regressions were performed. Regarding PSB, two regression models

were performed: the first model included all comparison participants (with and without PSB), the

second model included PSB patients and comparison participants without PSB. Initial

independent variables in both models were: age, gender, $SLC6A3$ genotype, childhood trauma,

self-reported mental health condition(s) (a binary variable), number of mental health condition(s)

in the family history (a continuous variable), and family history of PSB or of OCD. We also

included an interaction term of childhood trauma and $SLC6A3$ genotype to test for potential G×E

interactions.[23,30,31] To validate the association of the 10/10 genotype with RDS, one regression

model was applied using the entire cohort. Initial independent variables were identical but

additionally included "group" (comparison participants vs. patients). Predictors were retained in

the models if the model passed the link test for correct specification and showed non-significant

Hosmer-Lemeshow χ^2, their inclusion did not introduce multicollinearity, and they remained

significant in the final model. A *P*-value of <0.05 was considered significant.

3. RESULTS

A total of 8.5% of comparison participants (82 out of 970) scored ≥ 6 on the unadjusted SAST-R

Core, meeting the threshold for PSB in the general population.[23] Patients showed the highest

scores in adjusted SAST-R Core and addictive dimensions of PSB (*P*<0.001, Table 1).

Compared to comparison participants, patients were significantly older (*P*<0.001), had a higher

proportion of male participants, and higher rates of RDS (*P*<0.001, Table 1). Furthermore,

family history of mental health condition(s) and experiencing childhood trauma increased in

frequency from comparison without PSB to comparison with PSB to PSB patients (*P*<0.001,

Table 1).

The distributions of *SLC6A3* genotypes (9/9, 9/10, and 10/10) were in Hardy-Weinberg

Equilibrium in both comparison participants and in patients (Supplemental Digital Content,

Table 1). Results showed that comparison participants with the 10/10 genotype had trend level

elevated rates of PSB compared to 9-repeat carriers (*P*=0.050, Table 2). They also displayed

significantly higher RDS total scores (*P*=0.0042), especially regarding ADHD symptoms

(ASRS, *P*=0.021) and internet addiction (IAT, *P*=0.0073), followed by a trend level elevation in

compulsive buying (RCBS, *P*=0.050).

In comparison participants, the 10/10 genotype showed trend level correlations with RDS

(τ=0.077, *P*=0.088), with self-reported mental health condition(s) (τ=0.081, *P*=0.073), as well as

with PSB (τ=0.082, *P*=0.067; Supplemental Digital Content, Figure 2). In patients, however, the

10/10 genotype was not significantly correlated with PSB or RDS. Our results higher rates of

OCD in *SLC6A3* 3' VNTR rare genotype carriers (*P*=0.044, Table 2). Additional analyses on

correlations between genotype and SA-SAPAS items are shown in Supplemental Digital

Content, Table 2.

3.1.Regression analyses

In both regression models for PSB, *SLC6A3* genotype, childhood trauma, family history of PSB,

and gender were retained as independent variables (Model 1 in comparison participants: pseudo

R^2=0.12, n=509, Figure 1; Model 2 in PSB patients and comparison participants without PSB:

pseudo R^2=0.58, n=739, Figure 2). In Model 1, the 10/10 genotype of *SLC6A3* 3' VNTR (OR:

1.97, 95% confidence interval (CI): 1.15-3.38, *P*=0.013), childhood trauma (OR: 2.31, 95% CI:

1.21-4.41, *P*=0.011), family history of PSB (OR: 4.49, 95% CI: 1.85-10.90, *P*<0.001) and male

gender (OR: 4.54, 95% CI: 2.64-7.83, *P*<0.001) were associated with PSB. In Model 2, *SLC6A3*

3' VNTR rare genotypes (OR: 5.69, 95% CI: 1.08-30.07, *P*=0.041), childhood trauma (OR: 9.78,

95% CI: 4.91-19.49, *P*<0.001), family history of PSB (OR:45.65, 95% CI: 15.56-133.90,

P<0.001), and male gender (OR: 142.77, 95% CI: 52.11-391.15, *P*<0.001) were associated with

PSB. The 10/10 genotype was not significantly associated with PSB in model 2 (*P*=0.72). The

interaction term of childhood trauma and *SLC6A3* genotype was not significant in either

comparison participants (*P*=0.36) or in patients (*P*=0.26).

For RDS, the final model retained the interaction term of *SLC6A3* genotype and

childhood trauma, self-reported mental health condition(s), *SLC6A3* genotype, childhood trauma,

and group status on the entire cohort (pseudo R^2=0.13, n=1044, Figure 3). Significant parameters

included the 10/10 genotype of *SLC6A3* 3' VNTR (OR:2.39, 95% CI:1.52-3.77, *P*<0.001), self-

reported mental health condition(s) (OR:2.64, 95% CI: 1.96-3.56, *P*<0.001), childhood trauma

(OR:2.81, 95% CI:1.72-4.60, *P*<0.001), and being a PSB patient (OR:4.10, 95% CI:2.96-5.68,

P<0.001). The interaction term of the 10/10 genotype and childhood trauma was significantly

associated with decreased odds of RDS (OR:0.32, 95% CI:0.17-0.60, *P*=0.011, the detailed

interaction terms and predicted margins are provided in Supplemental Digital Content, Table 3).

4.   DISCUSSION

In our study, the 10/10 genotype of the SLC6A3 3' VNTR was associated with PSB in healthy

young adults, and with RDS across the entire cohort, suggesting that this genotype may serve as

a shared genetic risk factor for both PSB and RDS. This indicated that PSB could be a

manifestation of RDS in this cohort of healthy young adults. However, in the clinical cohort, it

was rare genotypes, rather than the 10/10 genotype, that showed the association to PSB. This

observation aligns with the understanding that both common and rare genetic variants can be

associated with a psychiatric phenotype, with common genotypes typically exerting a smaller

impact, while rare genotypes tend to have a larger impact and are associated with more severe

symptoms.[24]

308   Whether PSB should be regarded as a behavioral addiction or compulsive disorder

remains an open question.[32,33] Of note, Koob and colleagues (2016) conceptualized addiction as

a cycle with three stages, linked to disruptions in dopaminergic and serotoninergic pathways.[34]

They further indicated that impulsivity is often seen in the early stages of addiction, and

compulsivity evolves as the condition progresses.[34] Impulsive, reward-seeking tendencies of

PSB[31,32] could be more prominent in milder forms of PSB, such as in the comparison group, and

appear to be associated with the relatively common 10/10 VNTR genotype. In contrast, the

compulsive features may prevail in clinical PSB, accompanied by multiple mental health

comorbidities (e.g., OCD), and then be associated with rare VNTR genotypes.

Through our convenience sampling we were not able to match comparison participants

and PSB patients on age and other demographic variables. These group differences could have

driven different manifestations of PSB, which may be associated with different genetic risk

factors. Riemersma et, al. (2013) proposed a "new generation of sex addiction," or

"comtemporary sex addiction." While both "classical sex addiction" and "contemporary sex

addiction" could be assaociated with alterations in the reward pathway, Riemersma et, al. (2013)

suggested that "contemporary sex addiction" is primarily linked to reward seeking, whereas

"classical sex addiction," is predominantly associated with trauma, emotional dysregulation, and

insecure attachment, and may involve alterations in the prefrontal and limbic regions.[35,36] In our

study, comparison participants with PSB were significantly younger and a lower ratio of them

reporting childhood trauma than PSB patients. Hence, the relevance of the 10/10 genotype to

RDS in the entire cohort might have been influenced by the larger proportion of younger

participants who could have had such "contemporary" PSB. In contrast, rare genotypes of the

SLC6A3 3' VNTR, higher rates of childhood trauma and mental health comorbidities such as

OCD could potentially be associated with more "classical" sex addiction in the clinical PSB

group. Interestingly, our results suggest that in patients, this VNTR may have a potential link to

OCD, with those who carried rare genotypes having higher rates of OCD. Some previous

research has suggested a potential association between this VNTR and OCD;[37,38] however,

further studies are needed to confirm these associations.

We also investigated specific G×E interactions (*SLC6A3* 3' VNTR genotype and

childhood trauma) in the context of hypodopaminergic phenotypes of PSB and RDS. Although

such interactions have been explored in some relevant psychiatric conditions, particularly in

substance use, the results have been inconsistent or inconclusive.[23] A systematic review

examining G×E between dopaminergic genes (*DRD2*, *DRD4*, and *SLC6A3*) and psychological,

family, and socio-demographic factors in relation to substance abuse reported that five out of

nine studies identified at least one significant interaction.[23] In the present study, we observed that

the 9-repeat allele could be associated with RDS when childhood trauma was present. It is

possible that the 9-repeat allele carriers may be more sensitive to environmental influences. For

example, other research has found that children with conduct disorder and the 9/9 genotype

showed stronger reductions in symptom severity in response to both negative or positive

parenting practices, compared to 10-repeat allele carriers.[31]

This study had several limitations and the sociodemographic differences between the

comparison participants and patients in these convenience samples are important to

acknowledge. As mentioned above, age influences the phenotype and genetic susceptibility to

many psychiatric symptoms, presumably including PSB and RDS.[35,39] Furthermore, the analysis

was limited to those of European ancestry and included relatively few women. Also, the sample

size for rare genotypes was limited, making it challenging to investigate the differential

phenotypes between common and rare genotypes. Additionally, only one genetic variant in the

SLC6A3 gene and only one gene in the dopaminergic pathway were investigated in this study. It

is possible that other genes, gene-gene interactions and G×E interactions may be associated with

PSB and could have impacted the results. Thus, the findings should be interpreted with caution,

and sample recruitment and matching should be optimized in future studies. These should also

incorporate a broader range of common genetic variants or use a genome-wide approach, which

would offer deeper insights into the genetic foundations of PSB.

5. CONCLUSION

Our results suggest that both common and rare variants of *SLC6A3* may be associated with PSB.

The link of the 10/10 genotype to RDS indicates that PSB could be a manifestation of RDS in

comparison participants, whereas rare *SLC6A3* variants might be associated with clinical PSB,

together with comorbid mental health issues.

6. Acknowledgement

The authors would like to express their gratitude to all those who supported data collection at the university: Keanna Wallace, Esther Yang, Leslie Roper, Garima Aryal, Dawon Lee, and Rohit J Lodhi.

Lemeshow χ2 test was insignificant (P=0.75), suggesting good model fit. The mean VIF and the

condition number were 1.06 and 3.38, respectively. The model accuracy, area under the ROC

curve, sensitivity, specificity, and F score were 85.07%, 73.87%, 3.90%, 99.54%, and 0.073,

respectively. The exact P-values were as follows: family history of PSB (P=9.19×10^{-4}), gender

(men: P=5.17×10^{-8}). Power analysis demonstrated that the regression model had a power of over

0.90. *Abbreviation: PSB: problematic sexual behavior; SLC6A3: solute carrier family 6 member*

3; VIF: variance inflation factor.

~~the link test with a significant P (hat) (P=<0.001) and a nonsignificant P (hatsq) (P=0.36). The~~

probability of the Hosmer-Lemeshow $\chi2$ test was insignificant (*P*=0.98), suggesting a good

model fit. The mean VIF and the condition number were 1.60 and 4.18, respectively. The model

accuracy, area under the ROC curve, sensitivity, specificity, and F score were 85.39%, 94.03%,

70.63%, 95.64%, and 0.80, respectively. The exact *P*-values were as follows: childhood trauma

(*P*=8.72×10^{-11}), family history of PSB (*P*=3.43×10^{-12}), and gender (men, *P*<2.00×10^{-16}). Power

analysis demonstrated that the regression model had a power of over 0.90. *Abbreviation: PSB:*

problematic sexual behavior; SLC6A3: solute carrier family 6 member 3; VIF: variance inflation

factor.

Figure 3. Forest Plot of Reward Deficiency Syndrome (RDS) Logistic Regression Analysis

outliers, with 91 of them being RDS positive, a pseudo R^2=0.13, P<0.001). The model passed the

link test with a significant P(hat) (P<0.001) and a nonsignificant P (hatsq) (P=0.81). The

probability of the Hosmer-Lemeshow χ2 test was insignificant (P=0.99), suggesting a good

model fit. The mean VIF and the condition number were 2.49 and 10.03, respectively. The model

accuracy, area under the ROC curve, sensitivity, specificity, and F score were 68.29%, 72.00%,

67.46%, 69.16%, and 0.71, respectively. The exact P-values were as follows: the interaction term

of 10/10 genotype and childhood trauma (P=3.68×10^{-4}), the 10/10 genotype of *SLC6A3* 3'

VNTR (P=1.73×10^{-4}), self-reported mental health condition(s) (P=1.46×10^{-10}), childhood

trauma (P=3.88×10^{-5}) and being PSB patients (P<2.00×10^{-16}). Power analysis demonstrated that

the regression model had a power of over 0.90. *Abbreviation: PSB: problematic sexual behavior;*

SLC6A3: solute carrier family 6 member 3; VIF: variance inflation factor.

417   **Table 1. Demographic and Clinical Variables across Subgroups: Scores Indicate Median (Ranges) or Frequencies**

418   **(Percentages)**

Variables		Comparison participants without PSB (n=266-448)	Comparison participants with PSB (n=55-89)	PSB Patients (n=418-434)
Age *		23 (18-55) [a]	22 (18-66) [a]	43 (19-80) [b]
Gender *	Women	309 (69.8%)	35 (42.2%)	23 (6.3%)
	Men	134 (30.2%)	50 (58.8%)	408 (94.7%)
SLC6A3 genotype, *P*=0.100	9/9 and 9/10	210 (46.9%)	35 (39.3%)	209 (48.2%)
	10/10	228 (50.9%)	54 (60.7%)	209 (48.2%)
	Rare	10 (2.2%)	0 (0%)	16 (3.6%)
Problematic Sexual Behavior - adjusted SAST-R core *		0 (0-5) [a]	7 (4-18) [b]	15 (10-18) [c]
Relationship disturbance *		0 (0-2) [a]	1 (0-4) [b]	4 (0-4) [c]
Loss of control *		0 (0-2) [a]	2 (0-4) [b]	4 (0-4) [c]
Preoccupation *		0 (0-3) [a]	1 (0-4) [b]	2 (0-4) [c]

Variables	Comparison participants without PSB (n=266-448)	Comparison participants with PSB (n=55-89)	PSB Patients (n=418-434)
Affect disturbance *	0 (0-3) [a]	3 (1-5) [b]	4 (2-5) [c]
Reward deficiency syndrome *	0 (0%)	60 (67.4%)	318(84.1%)
FTND *	0 (0-1) [a]	0 (0-7) [b]	0 (0-12) [b]
DPGDF (P=0.051)	0 (0-3) [a]	0 (0-9) [b]	0 (0-9) [b]
SA-SAPAS *	1 (0-3) [a]	1 (0-8) [b]	2 (0-8) [c]
ASRS *	1 (0-3) [a]	3 (0-6) [b]	3 (0-6) [b]
RCBS *	10 (6-24) [a]	11 (6-40) [b]	11 (6-42) [b]
IAT *	22 (0-30) [a]	30 (0-62) [b]	38 (0-97) [c]
Childhood trauma *	67 (15.1%)	33 (37.1%)	290 (70.9%)
Self-reported mental health condition(s) *	0 (0%)	24 (27.0%)	277 (65.2%)
Number of self-reported mental health condition(s) *	0 [a]	0 (0-12) [b]	1 (0-9) [c]

Variables	Comparison participants without PSB (n=266-448)	Comparison participants with PSB (n=55-89)	PSB Patients (n=418-434)
Number of mental health condition(s) in family history *	3 (0-17) [a]	4 (0-17) [b]	6 (0-24) [c]
Family history of PSB *	17 (3.8%)	19 (21.6)	273 (62.8%)
OCD *	0 (0%)	8 (9.0%)	61 (14.1%)

*: significant omnibus effect ($P<0.001$)

[a,b,c]: different superscripts denote significant differences ($P < 0.05$) between groups

Kruskal-Wallis H test and Dunn's post-hoc tests for continuous variables; Pearson's or Fisher's χ^2 test (cells < 5) for categorical

variables; Abbreviations: *PSB: problematic sexual behavior; SAST-R: Sexual Addiction Screening Test-Revised; ASRS: Adult ADHD*

Self-Report Scale version 1.1; FTND: Fagerström Test for Nicotine Dependence; IAT: Internet Addiction Test; OCD: obsessive-

compulsive disorder; RCBS: Richmond Compulsive Buying Scale

 Table 2. Distribution of Demographic and Clinical Variables by *Solute Carrier Family 6 Member 3 (SLC6A3)* Genotypic

 Group: Scores Indicate Median (Ranges) or Frequencies (Percentages)

Variables	Comparison participants (n= 55-89 with PSB; n=266-438 without PSB)		PSB patients (391-418) and comparison participants without PSB (n=266-438)		
	9/9 and 9/10 genotypes)	10/10 genotype	9/9 and 9/10 genotypes	10/10 genotype	Rare genotypes
PSB	*30 (12.2%)[†]*	*52 (18.4%)[†]*	200 (48.1%)	202 (46.1%)	16 (61.5%)
Adjusted SAST-R core	1 (0-14)	1 (0-18)	4.5 (0-18)	3 (0-18)	12.5 (0-18)
Relationship disturbance	0 (0-3)	0 (0-4)	3 (0-4)	3 (0-4)	3 (0-4)
Loss of control	0 (0-4)	0 (0-4)	2 (0-4)	1 (0-4)	3 (0-4)
Preoccupation	0 (0-4)	0 (0-4)	1 (0-4)	1 (0-4)	1.5 (0-4)
Affect disturbance	0 (0-4)	0 (0-5)	2 (0-5)	2 (0-5)	4 (0-5)
Reward deficiency syndrome	*21 (8.6%)[†]*	*39 (13.8%)[†]*	155 (38.3%)	152 (36.4%)	11(45.8%)
FTND	0 (0-4)	0 (0-7)	0 (0-12)	0 (0-11)	0 (0-4)
DPGDF	0 (0-9)	0 (0-9)	0 (0-8)	0 (0-9)	0 (0-3)

Variables	Comparison participants (n= 55-89 with PSB; n=266-438 without PSB)		PSB patients (391-418) and comparison participants without PSB (n=266-438)		
	9/9 and 9/10 genotypes)	10/10 genotype	9/9 and 9/10 genotypes	10/10 genotype	Rare genotypes
SA-SAPAS	1 (0-7)	1 (0-8)	1 (0-8)	1 (0-8)	1 (0-6)
ASRS	**1 (0-6)**[*]	**1 (0-6)**[*]	2 (0-6)	2 (0-6)	1 (0-6)
RCBS	**10 (6-30)**[*]	**10 (6-40)**[*]	10 (6-32)	10 (6-32)	10 (6-42)
IAT	**22 (0-55)**[*]	**24 (2-62)**[*]	25 (0-89)	26 (1-97)	28 (4-88)
Childhood trauma	45 (18.4%)	55 (19.7%)	169 (41.7%)	176 (41.7%)	12 (46.1%)
Self-reported mental health condition(s)	*7 (2.9%)*[†]	*17 (6.0%)*[†]	131 (31.7%)	134 (30.9%)	12 (48.0%)
Number of self-reported mental health condition(s)	*0 (0-10)*[†]	*0 (0-12)*[†]	0 (0-6)	0 (0-9)	0 (0-9)
Number of mental health condition(s) in family history	3 (0-17)	3 (0-17)	4 (0-24)	4 (0-19)	4.5 (0-14)
Family history of PSB	15 (6.1%)	21 (7.5%)	142 (33.9%)	138 (31.6%)	10 (38.5%)

Variables	Comparison participants (n= 55-89 with PSB; n=266-438 without PSB)		PSB patients (391-418) and comparison participants without PSB (n=266-438)		
	9/9 and 9/10 genotypes)	10/10 genotype	9/9 and 9/10 genotypes	10/10 genotype	Rare genotypes
OCD	3 (1.2%)	5 (1.8%)	27 (6.5%)[*]	29 (6.6%)[*]	5 (19.2%)[*]

[*]: **$P<0.05$**; [†]: $P<0.1$

Mann-Whitney U test for two-group comparisons of continuous variables; Kruskal-Wallis H test and Dunn's post-hoc tests for three-

group comparisons of continuous variables; Pearson's or Fisher's χ^2 test (cells < 5) for categorical variables; Abbreviations: *PSB:*

problematic sexual behavior; SAST-R: Sexual Addiction Screening Test-Revised; ASRS: Adult ADHD Self-Report Scale version 1.1;

FTND: Fagerström Test for Nicotine Dependence; IAT: Internet Addiction Test; OCD: obsessive-compulsive disorder; RCBS:

Richmond Compulsive Buying Scale

References

1.   Jiang S, Wallace K, Yang E, et al. Logistic Regression With Machine Learning Sheds
Light on the Problematic Sexual Behavior Phenotype. *J Addict Med*. Mar-Apr 01
2023;17(2):174-181. doi:https://doi.org/10.1097/ADM.0000000000001078

2.   Carnes P. *Don't call it love: Recovery from sexual addiction*. Bantam Books; 1991:439 p.

3.   Kuzma JM, Black DW. Epidemiology, prevalence, and natural history of compulsive
sexual behavior. *Psychiatr Clin North Am*. Dec *2008*;*31*(*4*):603-11.
doi:https://10.1016/j.psc.2008.06.005

4.   Blum K, Braverman ER, Holder JM, et al. The Reward deficiency syndrome: a
biogenetic model for the diagnosis and treatment of impulsive, addictive, and compulsive
behaviors. *J Psychoactive Drugs*. Nov 2000;32:1-112.
doi:http://doi.org/10.1080/02791072.2000.10736099

5.   Comings DE, Blum K. Reward deficiency syndrome: genetic aspects of behavioral
disorders. *Prog Brain Res*. 2000/01/01 2000;126:325-341. doi:https://doi.org/10.1016/S0079-
6123(00)26022-6

6.   Blum K, Gold M, Demetrovics Z, et al. Substance use disorder a bio-directional subset of
reward deficiency syndrome. *Front Biosci (Landmark Ed)*. Jun 1 2017;22(9):1534-1548.
doi:http://doi.org/10.2741/4557

7.   Gondre-Lewis MC, Bassey R, Blum K. Pre-clinical models of reward deficiency
syndrome: A behavioral octopus. *Neurosci Biobehav Rev*. Aug 2020;115:164-188.
doi:https://doi.org/10.1016/j.neubiorev.2020.04.021

8.   Blum K, Dennen CA, Elman I, et al. Should Reward Deficiency Syndrome (RDS) Be
Considered an Umbrella Disorder for Mental Illness and Associated Genetic and Epigenetic
Induced Dysregulation of Brain Reward Circuitry? *J Pers Med*. 2022;12(10):1719.
doi:http://doi.org/10.3390/jpm12101719

9.   Blum K, Baron D, McLaughlin TP, Giordano J, Ceccanti MD, Badgaiyan R. Summary
document research on RDS anti-addiction modeling: annotated bibliography. *J Addict
Psychiatry*. 2024;8(1):1-33.

10.   Ma Y, Yuan W, Cui W, Li MD. Meta-analysis reveals significant association of 3'-UTR
VNTR in SLC6A3 with smoking cessation in Caucasian populations. *Pharmacogenomics J*. Feb
2016;16(1):10-7. doi:http://doi.org/10.1038/tpj.2015.44

11.   Franklin TR, Lohoff FW, Wang Z, et al. DAT genotype modulates brain and behavioral
responses elicited by cigarette cues. *Neuropsychopharmacology*. Feb 2009;34(3):717-28.
doi:https://doi.org/10.1038/npp.2008.124

12.   Reith MEA, Kortagere S, Wiers CE, et al. The dopamine transporter gene SLC6A3:
multidisease risks. *Mol Psychiatry*. Feb 2022;27(2):1031-1046.
doi:http://doi.org/10.1038/s41380-021-01341-5

13. Schneider JP, M.A. M. Couple recovery from sexual addiction / co addiction: Results of a
survey of 88 marriages. *Sexual Addiction and Compulsivity: The Journal of Treatment and*
Prevention. 1996;3:111-126.

14. Vandenbergh DJ, Persico AM, Hawkins AL, et al. Human dopamine transporter gene
(DAT1) maps to chromosome 5p15.3 and displays a VNTR. *Genomics*. *Dec* 1992;14(4):1104-6.
doi:https://10.1016/s0888-7543(05)80138-7

15. Dreher JC, Kohn P, Kolachana B, Weinberger DR, Berman KF. Variation in dopamine
genes influences responsivity of the human reward system. *Proc Natl Acad Sci U S A*. Jan 13
2009;*106*(2):617-22. doi:https://doi.org/10.1073/pnas.0805517106

16. Vandenbergh DJ, Persico AM, Uhl GR. A human dopamine transporter cDNA predicts
reduced glycosylation, displays a novel repetitive element and provides racially-dimorphic TaqI
RFLPs. *Brain Res Mol Brain Res*. Sep 1992;15(1-2):161-6. doi:https://10.1016/0169-
328x(92)90165-8

17. Salatino-Oliveira A, Rohde LA, Hutz MH. The dopamine transporter role in psychiatric
phenotypes. *Am J Med Genet B Neuropsychiatr Genet*. Mar 2018;177(2):211-231.
doi:http://doi.org/10.1002/ajmg.b.32578

18. Volkow ND, Wang G-J, Newcorn JH, et al. Motivation deficit in ADHD is associated
with dysfunction of the dopamine reward pathway. *Mol Psychiatry*. 2011;16(11):1147-1154.
doi:http://doi.org/10.1038/mp.2010.97

19. Cornish KM, Manly T, Savage R, et al. Association of the dopamine transporter (DAT1)
10/10-repeat genotype with ADHD symptoms and response inhibition in a general population
sample. *Mol Psychiatry*. Jul 2005;10(7):686-98. doi:https://10.1038/sj.mp.4001641

20. Guo G, Tong Y, Xie C-W, Lange LA. Dopamine transporter, gender, and number of
sexual partners among young adults. *Eur J Hum Genet*. 2007;15(3):279-287.
doi:http://doi.org/10.1038/sj.ejhg.5201763

21. Guo G, Tong Y, Cai T. Gene by social context interactions for number of sexual partners
among white male youths: Genetics-informed sociology. *Am J Sociol*. 2008;114(S1):S36-S66.
doi:https://doi.org/10.1086/592207

22. Erblich J, Lerman C, Self DW, Diaz GA, Bovbjerg DH. Stress-induced cigarette craving:
effects of the DRD2 TaqI RFLP and SLC6A3 VNTR polymorphisms. *Pharmacogenomics J*.
2004/04/01 2004;4(2):102-9. doi:https://doi.org/10.1038/sj.tpj.6500227

23. Jiang Z, Chen Z, Chen X. Candidate gene-environment interactions in substance abuse: A
systematic review. *Plos one*. 2023;18(10):e0287446.
doi:http://doi.org/10.1371/journal.pone.0287446.

24. Andreassen OA, Hindley GFL, Frei O, Smeland OB. New insights from the last decade
of research in psychiatric genetics: discoveries, challenges and clinical implications. *World*
Psychiatry. Feb 2023;22(1):4-24. doi:http://doi.org/10.1002/wps.21034

25. Sery O, Paclt I, Drtilkova I, et al. A 40-bp VNTR polymorphism in the 3'-untranslated
region of DAT1/SLC6A3 is associated with ADHD but not with alcoholism. *Behav Brain Funct*.
Jun 11 2015;11:21. doi:https://doi.org/10.1186%2Fs12993-015-0066-8

26.   Doucette-Stamm LA, Blakely DJ, Tian J, Mockus S, Mao JI. Population genetic study of
the human dopamine transporter gene (DAT1). *Genet Epidemiol*. 1995;12(3):303-8.
doi:https://doi.org/10.1002/gepi.1370120307

27.   Carnes P, Green B, Carnes S. The same yet different: Refocusing the Sexual Addiction
Screening Test (SAST) to reflect orientation and gender. *Sexual Addiction & Compulsivity*.
2010;17(1):7-30. doi:https://doi.org/10.1080/10720161003604087

28.   *Stata Statistical Software: Release 17*. StataCorp LLC.; 2023.

29.   Giordano AL, Hedden LE, Kim SW, Lundeen LA, Lu Z. Childhood trauma and sex
addiction among adult men. *J Addict Offender Couns*.
2024;doi:https://doi.org/10.1002/jaoc.12134

30.   Franke B, Buitelaar JK. Gene-environment interactions. *Oxford Textbook of Attention
Deficit Hyperactivity Disorder*. 2018:35.

31.   Lahey BB, Rathouz PJ, Lee SS, et al. Interactions Between Early Parenting and a
Polymorphism of the Child's Dopamine Transporter Gene in Predicting Future Child Conduct
Disorder Symptoms. *J Abnorm Psychol*. Feb 2011;120(1):33-45.
doi:http://doi.org/10.1037/a0021133

32.   Kraus SW, Voon V, Potenza MN. Should compulsive sexual behavior be considered an
addiction? https://doi.org/10.1111/add.13297. *Addiction*. Dec 2016;111(12):2097-2106.
doi:10.1111/add.13297

33.   Walton MT, Bhullar N. Compulsive Sexual Behavior as an Impulse Control Disorder:
Awaiting Field Studies Data. *Arch Sex Behav*. 2018/07/01 2018;47(5):1327-1331.
doi:http://doi.org/10.1007/s10508-018-1200-0

34.   Koob GF, Volkow ND. Neurobiology of addiction: a neurocircuitry analysis. *Lancet
Psychiatry*. Aug 2016;3(8):760-773. doi:https://10.1016/S2215-0366(16)00104-8

35.   Riemersma J, Sytsma M. A new generation of sexual addiction. *Sexual Addiction &
Compulsivity*. 2013;20(4):306-322.

36.   Schmidt C, Morris LS, Kvamme TL, Hall P, Birchard T, Voon V. Compulsive sexual
behavior: Prefrontal and limbic volume and interactions. *Hum Brain Mapp*. Mar
2017;38(3):1182-1190. doi:https://doi.org/10.1002/hbm.23447

37.   Billett E, Richter M, Sam F, et al. Investigation of dopamine system genes in obsessive-
compulsive disorder. *Psychiatric genetics*. 1998;8(3):163-170.

38.   Hemmingsa SMJ, Kinnear CJ, Niehaus DJH, et al. Investigating the role of dopaminergic
and serotonergic candidate genes in obsessive-compulsive disorder. *Eur Neuropsychopharmacol*.
2003/03/01 2003;13(2):93-98. doi:https://doi.org/10.1016/S0924-977X(02)00129-3

39.   Grunblatt E, Werling AM, Roth A, Romanos M, Walitza S. Association study and a
systematic meta-analysis of the VNTR polymorphism in the 3'-UTR of dopamine transporter
gene and attention-deficit hyperactivity disorder. *J Neural Transm (Vienna)*. Apr
2019;126(4):517-529. doi:http://doi.org/10.1007/s00702-019-01998-x