Impact of 10-valent pneumococcal conjugate vaccine on the epidemiology of otitis media

with otorrhea among Bangladeshi children

Hakka Naziat[1,2], Md Hafizur Rahman[1], Rajib C Das[1], Maksuda Islam[1], Shampa Saha[1], Md.

Manzoor Hussain[3], Asif Sattar[3], Mohammad J Uddin[1], Md. Jahangir Alam[4], Cynthia G.

Whitney[5], Senjuti Saha[1], Yogesh Hooda[1, *], Samir K Saha[1,2, *]

1. Child Health Research Foundation, Dhaka, Bangladesh

2. Department of Microbiology, Bangladesh Shishu Hospital and Institute, Dhaka, Bangladesh

3. Department of Ear, Nose and Throat, Bangladesh Shishu Hospital and Institute, Dhaka,

Bangladesh

4. Department of Pediatrics, Bangladesh Shishu Hospital and Institute, Dhaka, Bangladesh

5. Rollins School of Public Health, Emory University, Atlanta, USA

* Corresponding author:

Samir K. Saha PhD FRCPath FAAM

samir@chrfbd.org

Child Health Research Foundation, SEL Huq Skypark, 23/2 Khilji Road, Mohammadpur, Dhaka,

1207

AND

Yogesh Hooda PhD

yhooda@chrfbd.org

Child Health Research Foundation, SEL Huq Skypark, 23/2 Khilji Road, Mohammadpur, Dhaka,

1207

Keywords: *Streptococcus pneumoniae*, pneumococcal conjugate vaccines, otitis media,

Bangladesh

ABSTRACT

In 2015, Bangladesh introduced the ten-valent pneumococcal conjugate vaccine (PCV10). This study evaluates the impact of PCV10 on pneumococcal otitis media (OM), a prevalent middle ear infection amongst children in low-and-middle income countries, including Bangladesh. We analyzed ear swabs from OM cases with otorrhea collected at the largest pediatric hospital in Dhaka, Bangladesh from April 2014 to March 2019. The study involved identifying pathogens and conducting pneumococcal serotyping using the Quellung reaction.

Four years post PCV10 introduction, the rate of detection of pneumococcus among otorrhea cases decreased from 18.4% (164/892) in 2014-15 to 15.6% (581/3735) in 2018-19, reflecting a 15.4% reduction. Notably, vaccine serotypes (VTs) demonstrated a significant decline throughout the post-PCV years, with a 44.1% reduction compared to the pre-PCV period. Specific VTs like 14, 6B and 19F exhibited significant reductions of 60.2%, 57.5% and 42.9% respectively. Conversely, non-vaccine serotypes (NVTs) showed non-significant increase of 8.6%; specifically serotype 35B showed a 70.1% increase. Administration of two or three doses of PCV10 provided 84.4% protection against OM cases caused by serotypes covered by PCV10.

This is the first report from South Asia assessing PCV's impact on pneumococcal OM. It demonstrates a 15.4% reduction in pneumococcal isolation over four years and over 80% efficacy of PCV10 in preventing vaccine type pneumococcal OM cases. While some vaccine serotypes decreased significantly, the impact of PCV10 on overall pneumococcal OM cases was dampened by increasing isolation of non-vaccine serotypes.

INTRODUCTION

Streptococcus pneumoniae, or pneumococcus, is a significant bacterial pathogen responsible for a range of pediatric infections worldwide, including meningitis, sepsis, pneumonia and otorrhea [1,2]. Given the considerable global burden of pneumococcus-associated morbidity and mortality, the World Health Organization has recommended the inclusion of pneumococcal conjugate vaccine (PCV) in the routine immunization programs of all countries and assessment of its impact on pneumonia, other invasive diseases, and otitis media (OM) [3,4]. In March 2015, Bangladesh introduced the 10-valent pneumococcal conjugate vaccine (PCV10), targeting 10 serotypes: 1, 4, 5, 6B, 7F, 9V, 14, 18C, 19F, and 23F [5]. In addition, PCV10 was also anticipated to provide cross-protection against 6A due to immunologic similarities with vaccine serotype 6B [6–8]. Bangladesh has successfully achieved high coverage (99.2% coverage for ≥ 2 doses) of PCV10 across the country [9], and recent studies have reported a decline in severe radiologically confirmed pneumonia amongst children under the age of five years at two sites in rural Bangladesh – Mirzapur [5] and Sylhet [10]. Nevertheless, the broader impact of PCV10 on other pneumococcal diseases remain to be assessed.

Otitis media is among the most common bacterial infections diagnosed in children globally, with peak incidences typically within the first two years of life [11,12]. OM can range from asymptomatic cases, which usually resolve or are treated without complications, to more severe forms such as recurrent, nonresponsive, dry perforation, spontaneously draining, and chronic suppurative OM [13]. In Bangladesh, pneumococcus is a principal cause of OM, accounting for 18% of all OM cases [14]. PCV10 and PCV10□+□6A serotypes accounted for 8% and 9% of all OM cases and 46% and 49% of pneumococcus-positive cases, respectively, and these serotypes

exhibited a higher likelihood to be non-susceptible to at least one antibiotic (erythromycin and/or

trimethoprim-sulfamethoxazole) than nonvaccine serotypes (91% vs. 77%). A recent Cochrane

review including 15 publications from 11 trials reported that administration of the licensed

seven-valent pneumococcal conjugate vaccine (PCV7) and ten-valent PCV (PCV10) during early

infancy substantially reduced the relative risk of pneumococcal OM [15].

To assess the impact of PCV10 on pneumococcal OM, our research group conducted a study to

identify and characterize bacterial pathogens of OM in the ear, nose, and throat (ENT) outpatient

department of Bangladesh Shishu Hospital and Institute from 2014 to 2019, spanning one year

prior to PCV10 introduction and four years following the PCV10 introduction. The study aimed

to determine a) the impact of PCV10 on percentage of OM cases that were pneumococcus-

positive, b) changes in serotype distribution amongst pneumococcal OM cases, c) variations in

the age of infection and duration of ear discharge, and d) the antimicrobial susceptibility of

pneumococcus isolated from OM cases.

METHODS

The methodology employed in this study adhered strictly to those outlined in our previously

published baseline study [14], with no modifications. This consistency ensured the comparability

of results across both study periods. Details are provided below.

Study site and population

The study was conducted at Bangladesh Shishu Hospital and Institute (BSHI, formerly known as

Dhaka Shishu Hospital or DSH), the largest pediatric hospital in Bangladesh, which serves

children from 0 to 18 years. This hospital has a dedicated out-patient department (OPD) for the

management of all cases seeking care for ear, nose, and throat (ENT) conditions.

Case identification and enrollment criteria

Children presenting with symptoms suggestive of otitis media (OM) were identified by a study

physician based on World Health Organization-recommended clinical criteria and otoscopic

examinations [16,17]. The inclusion criteria for OM cases were: (1) acute onset of one or more

sign/symptoms of infection such as fever, earache, ear tugging or irritability, and (2) middle ear

inflammation with/without middle ear effusion or spontaneous discharge (otorrhea). From

enrolled children with/without otorrhea (not caused by otitis externa), data on sex, age, duration

of discharge, and antibiotic use within the previous seven days were recorded. Informed consent

was obtained from caregivers for children with otorrhea to collect specimens.

Specimen collection

Specimens were collected from children presenting with otorrhea by trained research assistants.

The external auditory canal was swabbed after the removal of any debris, using a flocked swab,

and placed into a tube containing STGG medium (2% skim milk, 3% tryptone, 10% glycerol and

0.5% glucose). Each specimen was cultured within three hours in the laboratory and examined

for the presence of bacterial pathogens.

Laboratory methods

All specimens were cultured on blood agar with and without gentamicin (in 5% CO_2

environment), chocolate agar with bacitracin and MacConkey agar [17]. Growth on these

selective media were examined for the presence of *Haemophilus influenzae*, *S. pneumoniae*, and

other bacterial pathogens, such as *Staphylococcus aureus*, *Moraxella catarrhalis*, *Pseudomonas*

spp.

Pneumococcal isolates were serotyped using the Quellung reaction [18]. Antibiotic susceptibility

testing was conducted using the disc diffusion method adhering to the latest Clinical and

Laboratory Standards Institute (CLSI) guidelines [19]. Antibiotics tested included penicillin (1

μg oxacillin disc), trimethoprim-sulfamethoxazole (1/19, 25 μg disc), chloramphenicol (30 μg

disc), erythromycin (15 μg disc) and ciprofloxacin (5 μg disc). Non-susceptible pneumococcal

isolates were further tested using the E-test to determine minimum inhibitory concentrations.

Indirect cohort analysis

Vaccine effectiveness was assessed by comparing the vaccination status of children with

vaccine-type OM (VT-OM) and non-vaccine-type OM (NVT-OM). This analysis assumes that

vaccination does not influence the risk for non-vaccine-type disease among vaccinated

individuals. A recent study has shown that results derived from the indirect cohort method were

comparable to that of a standard case-control study [20,5], which is complex and difficult to

implement. VT-OM was defined as OM cases in which PCV10 serotypes or 6A were identified,

all other serotypes were considered NVT. All pneumococcal OM cases from April 2015 through

March 2019 were used for this analysis. Based on the immunization status verified by the

vaccination card or the caregiver's verbal confirmation (in cases where the card was not available

or lost), children were classified into two groups: vaccinated and unvaccinated. The unvaccinated

cohort included verbally confirmed non-vaccinated children and 2 - <24-month-old children who

were not age-eligible for PCV at the time of the initial pneumococcal vaccination program (21st

March 2015). As Bangladesh's EPI did not have catch-up campaigns and only children aged 6

weeks were eligible for vaccine introduction, this strategy led to an increase of vaccine eligible

children gradually with time. To identify vaccine eligible children, we back calculated the

birthdate of children and considered those born on 7th February 2015 or later as eligible to

receive the 1st dose of pneumococcal vaccine at 6 weeks of age. Children who sought medical

care two weeks or more after receiving an initial dose were considered vaccinated. Any child

who came to the hospital within 14 days of receiving the 1st dose of PCV10 vaccine was

considered non-vaccinated. Children who received two or three doses of PCV10 were considered

individually for calculating vaccine impact, using children with no doses were used the

comparison group. We calculated the odds ratio of receipt of PCV10 dose among children who

had vaccine-type (PCV10) OM versus non-vaccine-type OM and used logistic regression to

estimate vaccine effectiveness, calculated as: Vaccine effectiveness = (1-odds ratio for PCV10

vaccination) x 100%.

Statistical data analysis

Data analyses were conducted using STATA 14.0 (Stata Corp LP, College Station, TX).

Epidemiological data were analyzed to elucidate the serotype distribution amongst

pneumococcal OM cases, duration of ear discharge, and antibiotic susceptibility profiles of *S.*

pneumoniae isolates. Common contaminants (e.g., *Bacillus spp.*, *Staphylococcus spp.* etc.) were

excluded.

Ethical clearance

This surveillance protocol was approved by the Institutional Review Board of Bangladesh

Institute of Child Health at BSHI. Specimens were collected as part of routine patient care.

RESULTS

From April 2014 to March 2019, a total of 5,040 OM cases were identified, with 4,279 (93.8%)

classified as otorrhea cases (Figure 1, Table 1). Ear swabs were successfully collected from

4,626 (97.8%) of these cases. A pathogen could be identified in 2,181 of the 4,626 samples

(47.1%; Supplemental Table 1). *H. influenzae* was the most frequently detected pathogen

(782/2,181; 35.8%), followed by *S. pneumoniae* (581/2,181; 26.6%), *S. aureus* (514/2,181;

23.6%) and *Pseudomonas spp.* (365/2,181; 16.7%). The trends in isolation of all detected causes

of OM cases are shown in Figure 2. In 554 out of the 2,181 culture-positive cases (25.4%),

multiple organisms were detected.

Table 1: Total, PCV10-type and non-PCV10-type *S. pneumoniae* obtained from OM cases in

this study.

	April 2014-March 2015	April 2015-March 2016	April 2016-March 2017	April 2017-March 2018	April 2018-March 2019	Total post-PCV
OM episodes recorded	1111	1089	979	909	952	3929
Episodes with otorrhea	981	1061	963	840	884	3748
Specimens collected	891	1054	960	839	882	3735
Total *S. pneumoniae* detected	164	167	148	153	113	581
PCV10-type *S. pneumoniae* detected	76	72	50	28	28	178
Non-PCV10 type *S.*	88	95	98	125	85	403

Figure 1: Study design highlighting the number of yearly *S. pneumoniae* isolates collected

during the baseline (Pre-PCV, April 2014-March 2015) and present study period (Post-PCV,

April 2015- March 2019).

Trends in number and serotype distribution of pneumococcal positive otorrhea cases

From April 2014 to March 2015, 164 pneumococcal OM cases were identified. This number

reduced to 113 by April 2018 to March 2019. Over the four years following the introduction of

PCV10, the pneumococcal positivity among otorrhea cases decreased from 18.4% (164/892) to

15.6% (581/3,735) [15.4% reduction (95% CI: -1.2-28.9%); p=0.04] (Figure 3). The detection

rate for *Haemophilus influenzae, Staphylococcus aureus* and *Pseudomonas spp* was not affected

(Figure 2).

Figure 2: Trends in isolation of all cause OM, OM caused by *Haemophilus influenzae*,

Streptococcus pneumoniae, *Staphylococcus aureus, Pseduomonas spp*.

Vaccine serotypes displayed a declining trend over the post-PCV year, resulting in a 44.1% (95%

CI: 25.8 - 57.5%, p<0.001) reduction compared to the pre-PCV period (Figure 3). Notably VTs

like 14, 6B and 19F showed significant reductions of 60.2% (95% CI: 10.7% - 81.7%; p=0.009),

57.5% (95% CI: -9.01-82.3%; p=0.026) and 42.9% (95% CI: 2.6-65.5%; p=0.015), respectively

(Figure 4A). No protection was seen for Serotype 1 (35.6%, 95% CI: -85.7%-83.7%; p=0.215).

While the percentage of OM cases attributable to 6A declined, this change was not statistically

significant (10.4% reduction, 95% CI: -270.7% - 71.5%; p=0.405).

Non-PCV10 serotypes showed a slight increase although not significant (8.6% increase, 95% CI:

-15.4-28.2%, p=0.225), particularly 35B, which exhibited a 70.1% increase (95% CI: -18.6 % -

96.6%; p=0.04) (Figure 3, Sup Figure 1). Certain serotypes absent in the pre-PCV period were

notably present in substantial numbers during the post-PCV period, including serotype 17F

(N=13), 12F (N=9), and 18A (N=7) (Sup Figure 1).

Figure 3: Percent of OM cases attributable to *S. pneumoniae* at Bangladesh Shishu Hospital and Institute in Dhaka, Bangladesh from April 2014 to March 2019. Trends of all *S. pneumoniae* (green), PCV10-type *S. pneumoniae* (VT, blue) and non-PCV10-type *S. pnuemoniae* (NVT, red) isolated from OM cases is shown.

Figure 4: Trends in the proportion of OM cases attributable to vaccine-type *S. pneumoniae* from April 2014 to March 2019. A) The yearly trends of percentage of VT-OM cases caused by different VT pneumococcal serotypes included in PCV7 (pink), PCV10-GSK (dark blue), PCV10-SII (purple), PCV13 (sky blue), PCV15 (yellow) and PCV20 (green) are shown. B) The estimated coverage of different PCV vaccines in the pre-PCV and post-PCV period.

Based on our data, the estimated serotype coverage of different PCVs for OM cases amongst Bangladeshi infants during the pre-PCV period (2014-2015) was 37.8% for PCV7, 46.3% for $PCV10_{GSK}$ 62.2% for $PCV10_{SII}$, 72.6% for PCV13, 72.6 % for PCV15 and 79.3% for PCV20. Four years after the introduction of PCV10 (2018-2019), the estimated serotype coverage changed to 23.8% for PCV7, 30.6% for $PCV10_{GSK}$, 41.7% for $PCV10_{SII}$, 53.5% for PCV13, 54.4% for PCV15 and 64.5% for PCV20 (Figure 4B).

Age and ear discharge duration

The median age of a child with pneumococcal OM was 10.0 months (95%CI: 9.0-13.0) in the pre-PCV period and 13.0 months (95%CI, 11.0-14.0) in the post-PCV period (Sup Figure 2). The median ear discharge duration was 5 days (95% CI: 5-6) in pre-PCV period and 5 days (95% CI: 4-5) in the post-PCV period (Sup Figure 3). Trend analysis found no statistical difference in the median age or ear discharge duration throughout the study period.

Trends of antibiotic susceptibility of pneumococcus

A large percentage of pneumococcal isolates were resistant to cotrimoxazole (58%) and erythromycin (72%), while low rates of resistance were observed for penicillin (0.95%), chloramphenicol (5.5%) and ciprofloxacin (6.9%). No statistical difference in trends was observed for the five antibiotics tested over the study period (Figure 5).

Figure 5: Trends in antimicrobial susceptibility of *S. pneumoniae* isolates against five antibiotics (penicillin, chloramphenicol, cotrimoxazole, erythromycin and ciprofloxacin the during the study period. The proportion of isolates that were resistant to each drug is shown.

PCV10 effectiveness calculation

Between March 21, 2015, and March 31, 2019, 581 cases of pneumococcal OM were identified, with 323 children being eligible for PCV10 vaccination based on their age. Among these, 178 (30.6%) were of vaccine-type (VT) and 403 (69.4%) were non-vaccine-type (NVT). Of the vaccinated children (n=307), 55 had VT OM, and 252 had NVT OM. Among the unvaccinated (n=274), 123 had VT OM, and 151 had NVT OM. The median age of vaccinated children and unvaccinated children were 7.0 and 33.5 months, respectively. The effectiveness of the 1st PCV10 dose was 83.5% (95% CI: 29.4% - 96.2%; p-value =0.011), 2nd PCV10 dose was 82.1% (95% CI: 42.2% - 94.5%; p-value =0.003) and 3rd PCV10 dose was 84.3% (95% CI: 57.1% -

257   94.5%; p-value<0.001) (Table 2). Two or three PCV10 doses provided 84.4% (95% CI: 57.1% –

258   94.3%; p-value <0.001) protection against VT OM (Table 2).

260   **Table 2:** Indirect cohort analysis for PCV10 effectiveness calculation.

PCV doses	VT vaccinated case/Total	NVT vaccinated case/Total	vaccine effectiveness (95% CI)	p-value
1ˢᵗ dose	4/14 (28.6%)	17/24 (70.8%)	83.5% (29.4%, 96.2%)	0.0106
2ⁿᵈ dose	11/21 (52.4%)	43/50 (86.%)	82.1% (42.2%, 94.5%)	0.0034
3ʳᵈ dose	46/56 (82.1%)	212/219 (96.8%)	84.8% (57.9, 94.5%)	0.0003
1ˢᵗ+2ⁿᵈ dose	15/25 (60%)	60/67 (89.6%)	82.5% (46.4%, 94.3%)	0.0020
2ⁿᵈ+3ʳᵈ dose	57/67 (85.1%)	255/262 (97.3%)	84.4% (57.1%, 94.3%)	0.0003

DISCUSSION

Otitis media continues to be one of the most prevalent pediatric infections in Bangladesh, and a

common cause of antibiotic prescriptions. This study, the largest case series of OM from

Bangladesh - a country with a population of 170 million, examined the impact of PCV10 on

pneumococcal OM. Specifically, it assessed changes in serotype distribution, duration of ear

discharge, age distribution, and antibiotic susceptibility, allowing for the calculation of PCV10's

effectiveness in preventing OM cases cause by vaccine serotypes. To our knowledge, this

represents the first report from South Asia investigating the impact of PCVs on OM.

A key finding from our study is the 15.4% reduction in pneumococcal positivity among otorrhea

cases over four years following the introduction of PCV10. This reduction is consistent with

systematic analyses of studies evaluating the impact of PCV on OM, which have reported

reductions ranging from 8% to 42.7% for PCV7, 5.6% to 84% for PCV10, and 2.2% to 68% for

PCV13 [15]. Similar results were seen from studies in Vietnam, a southeast Asian country that

introduced PCV10 in March 2017 and observed a ~40% reduction in pneumococcal OM cases

[21].

Our data indicated a 44.1% reduction in VT OM cases, with significant decreases observed in

most PCV10 serotypes, including 19F, 6B, and 14. Limited cross-protection was noted for

serotype 6A, as reported in several studies [15,22]. In contrast, no significant difference was

noted for serotype 1 and low case numbers precluded effective estimates for serotypes 4, 5, and

7F. As expected, PCV10 had no effect on non-vaccine-type (NVT) OM cases. An increase in

certain NVT serotype's such as 35B was observed, along with a rise in the number of serotypes

post-vaccination, frequency of NVT OM cases did not change through the study period. These

changes in serotype distribution did not result in differences in median age, ear discharge

duration, or the antimicrobial susceptibility against five commonly used antibiotics.

An interesting trend was observed with serotype 19A, which became an increasingly common

cause of invasive pneumococcal disease in some countries following the introduction of PCVs

that did not include the 19A antigen [23,24]. Some studies suggested that the 19F antigen

included in PCV10 could offer cross-protection for 19A [8]. Initially, we observed a reduction

in the proportion of 19A cases for two years post-vaccination. Genomic sequencing of isolates

could offer an understanding of the changes in the observed phenotype [25].

Using indirect cohort analysis, the vaccine efficacy of PCV10 against VT pneumococcal OM

cases was calculated at 84.4%, aligning with efficacy observed in other settings, in both

developed and developing countries [15,21]. We also evaluated the coverage of other

pneumococcal vaccines, noting coverage rates of 41.7% for Serum Institute PCV10, 53.5% for

PCV13, 54.4% for PCV15, and 64.5% for PCV20. Incorporating PCV15 or PCV20 could

potentially increase the proportion of OM cases preventable by vaccination, suggesting a benefit

from higher valent pneumococcal vaccines in enhancing disease coverage.

Several limitations should be considered when interpreting our results. First, tympanocentesis

was not performed on children without otorrhea, which may have led to lack of detection of

certain pathogens. However, since 94% of the suspected OM cases presented with otorrhea, this

limitation likely had minimal, if any, impact on our analysis. Plus, the possibility of missing

bacterial pathogens in this study was low as we used selective and enriched (nonselective) media

specifically for pneumococcus and *H. influenzae*. Second, we did not test for viral pathogen in

the specimens, which may also play a role as co-pathogens in OM cases. Third, the study was

conducted at the outpatient department of a single hospital in Dhaka, the capital city of

Bangladesh. It is likely that severe OM cases, especially those with draining ears, are more

inclined to seek care at facilities like Bangladesh Shishu Hospital and Institute, which is

equipped to handle more complex conditions. Conversely, milder cases of OM are often

managed in private clinics or pharmacies. Therefore, the impact of PCV10 observed in this study

might not fully represent its effect across different healthcare settings and regions within

Bangladesh. Moreover, having only one year of pre-PCV data limited our ability to discern

longer-term trends in serotype distribution pre- and post-vaccination. Finally, the lack of a

population-based denominator in our surveillance data precludes incidence calculations.

Despite these limitations, our study provides robust evidence of the significant impact of PCV10

on pneumococcal OM cases due to vaccine serotypes in Bangladesh. Studies that detail OM

outcomes by specific pneumococcal serotypes are exceedingly rare, making our data particularly

valuable to the existing literature. This serotype-specific analysis offers a critical perspective on

vaccine effects, especially given the high burden of OM. In the absence of population-based

studies, the large number of cases identified through our hospital-based surveillance enabled a

detailed examination of PCV10's impact in Bangladesh. This study underscores the value of

serotype-specific investigations in understanding and enhancing vaccine effectiveness in

preventing serious health outcomes. Collectively, these findings underscore the importance of

continued support for PCV vaccination, highlighting its positive impact on the overall well-being

of children not only in Bangladesh but across the region.

AUTHOR CONTRIBUTION

SKS conceived the study, provided supervision, and secured funding. HN, MHR, ShS, MI

conducted the laboratory tests. SKS, MH and MJA supervised the sample collection. HN, RCD

and YH performed the data analysis and generated the figures. YH, SS, HN and SKS wrote the

manuscript. YH, SS, CJW and SKS reviewed the manuscript.

FUNDING STATEMENT

This work was supported by Pfizer Inc (Grant number: WI209075). The funding organization had

no role in the design, implementation, data analysis, manuscript writing or decision to publish

this study.

CONFLICT OF INTEREST

Authors declare no conflict of interest.

REFERENCES

[1] Wahl B, O'Brien KL, Greenbaum A, Majumder A, Liu L, Chu Y, et al. Burden of Streptococcus pneumoniae and Haemophilus influenzae type b disease in children in the era of conjugate vaccines: global, regional, and national estimates for 2000–15. The Lancet Global Health 2018;6:e744–57. https://doi.org/10.1016/S2214-109X(18)30247-X.

[2] Troeger C, Blacker B, Khalil IA, Rao PC, Cao J, Zimsen SRM, et al. Estimates of the global, regional, and national morbidity, mortality, and aetiologies of lower respiratory infections in 195 countries, 1990–2016: a systematic analysis for the Global Burden of Disease Study 2016. The Lancet Infectious Diseases 2018;18:1191–210. https://doi.org/10.1016/S1473-3099(18)30310-4.

[3] Pneumococcal conjugate vaccine for childhood immunization--WHO position paper. Wkly Epidemiol Rec 2007;82:93–104.

[4] Pneumococcal conjugate vaccines: WHO position paper n.d. https://www.who.int/publications-detail-redirect/10665-310968 (accessed April 1, 2024).

[5] Malaker R, Hasanuzzaman M, Hooda Y, Rahman H, Chandra Das R, Kanon N, et al. Effectiveness of ten-valent pneumococcal conjugate vaccine on invasive pneumococcal disease among children <2 years old: A prospective population-based study in rural Bangladesh. Vaccine 2024;42:255–62. https://doi.org/10.1016/j.vaccine.2023.12.007.

[6] Saha S, Hasan M, Kim L, Farrar JL, Hossain B, Islam M, et al. Epidemiology and risk factors for pneumonia severity and mortality in Bangladeshi children <5 years of age before 10-valent pneumococcal conjugate vaccine introduction. BMC Public Health 2016;16:1233. https://doi.org/10.1186/s12889-016-3897-9.

[7] Whitney CG, Farley MM, Hadler J, Harrison LH, Bennett NM, Lynfield R, et al. Decline in invasive pneumococcal disease after the introduction of protein-polysaccharide conjugate vaccine. N Engl J Med 2003;348:1737–46. https://doi.org/10.1056/NEJMoa022823.

[8] Hausdorff WP, Hoet B, Schuerman L. Do pneumococcal conjugate vaccines provide any cross-protection against serotype 19A? BMC Pediatr 2010;10:4. https://doi.org/10.1186/1471-2431-10-4.

[9] DGHS Health Bulletin n.d. https://old.dghs.gov.bd/index.php/en/publications/health-bulletin/dghs-health-bulletin (accessed April 1, 2024).

[10] McCollum ED, Ahmed S, Roy AD, Chowdhury NH, Schuh HB, Rizvi SJR, et al. Effectiveness of the 10-valent pneumococcal conjugate vaccine against radiographic pneumonia among children in rural Bangladesh: A case-control study. Vaccine 2020;38:6508–16. https://doi.org/10.1016/j.vaccine.2020.08.035.

[11] Monasta L, Ronfani L, Marchetti F, Montico M, Brumatti LV, Bavcar A, et al. Burden of Disease Caused by Otitis Media: Systematic Review and Global Estimates. PLOS ONE 2012;7:e36226. https://doi.org/10.1371/journal.pone.0036226.

[12] Jin Y, Yang X, Sun H, Zhang J, Yang S, Jiang S, et al. Global, Regional, and National Burdens of Otitis Media From 1990 to 2019: A Population Based Study. Ear and Hearing n.d.:10.1097/AUD.0000000000001453. https://doi.org/10.1097/AUD.0000000000001453.

[13] O'Neill P, Roberts T, Bradley Stevenson C. Otitis media in children (acute). BMJ Clin Evid 2007;2007:0301.

[14] Naziat H, Saha S, Islam M, Saha S, Uddin MJ, Hussain M, et al. Epidemiology of Otitis Media With Otorrhea Among Bangladeshi Children: Baseline Study for Future Assessment

of Pneumococcal Conjugate Vaccine Impact. The Pediatric Infectious Disease Journal
2018;37:715. https://doi.org/10.1097/INF.0000000000002077.

[15] Sévaux JL de, Venekamp RP, Lutje V, Hak E, Schilder AG, Sanders EA, et al.
Pneumococcal conjugate vaccines for preventing acute otitis media in children. Cochrane
Database of Systematic Reviews 2020. https://doi.org/10.1002/14651858.CD001480.pub6.

[16] Rosario DC, Mendez MD. Chronic Suppurative Otitis. StatPearls, Treasure Island (FL):
StatPearls Publishing; 2024.

[17] Grevers G, Wiedemann S, Bohn J-C, Blasius R-W, Harder T, Kroeniger W, et al.
Identification and characterization of the bacterial etiology of clinically problematic acute
otitis media after tympanocentesis or spontaneous otorrhea in German children. BMC Infect
Dis 2012;12:312. https://doi.org/10.1186/1471-2334-12-312.

[18] Saha SK, Baqui AH, Darmstadt GL, Ruhulamin M, Hanif M, El Arifeen S, et al.
Comparison of Antibiotic Resistance and Serotype Composition of Carriage and Invasive
Pneumococci among Bangladeshi Children: Implications for Treatment Policy and Vaccine
Formulation. Journal of Clinical Microbiology 2003;41:5582–7.
https://doi.org/10.1128/JCM.41.12.5582-5587.2003.

[19] Jorgensen JH, Howell AW, Maher LA. Quantitative antimicrobial susceptibility testing of
Haemophilus influenzae and Streptococcus pneumoniae by using the E-test. J Clin
Microbiol 1991;29:109–14. https://doi.org/10.1128/jcm.29.1.109-114.1991.

[20] Bar-Zeev N, Swarthout TD, Everett DB, Alaerts M, Msefula J, Brown C, et al. Impact and
effectiveness of 13-valent pneumococcal conjugate vaccine on population incidence of
vaccine and non-vaccine serotype invasive pneumococcal disease in Blantyre, Malawi,
2006–18: prospective observational time-series and case-control studies. The Lancet Global
Health 2021;9:e989–98. https://doi.org/10.1016/S2214-109X(21)00165-0.

[21] Toizumi M, Satoh C, Quilty BJ, Nguyen HAT, Madaniyazi L, Le LT, et al. Effect of
pneumococcal conjugate vaccine on prevalence of otitis media with effusion among
children in Vietnam. Vaccine 2022;40:5366–75.
https://doi.org/10.1016/j.vaccine.2022.07.047.

[22] Eskola Juhani, Kilpi Terhi, Palmu Arto, Jokinen Jukka, Eerola Mervi, Haapakoski Jaason,
et al. Efficacy of a Pneumococcal Conjugate Vaccine against Acute Otitis Media. New
England Journal of Medicine 2001;344:403–9.
https://doi.org/10.1056/NEJM200102083440602.

[23] Ardanuy C, Rolo D, Fenoll A, Tarrago D, Calatayud L, Liñares J. Emergence of a
multidrug-resistant clone (ST320) among invasive serotype 19A pneumococci in Spain. J
Antimicrob Chemother 2009;64:507–10. https://doi.org/10.1093/jac/dkp210.

[24] Richter SS, Heilmann KP, Dohrn CL, Riahi F, Diekema DJ, Doern GV. Pneumococcal
Serotypes before and after Introduction of Conjugate Vaccines, United States, 1999–20111.
Emerg Infect Dis 2013;19:1074–83. https://doi.org/10.3201/eid1907.121830.

[25] Nakano S, Fujisawa T, Chang B, Ito Y, Akeda H, Fujita J, et al. Whole-Genome Analysis-
Based Phylogeographic Investigation of Streptococcus pneumoniae Serotype 19A Sequence
Type 320 Isolates in Japan. Antimicrob Agents Chemother 2022;66:e0139521.
https://doi.org/10.1128/AAC.01395-21.

SUPPLEMENTARY FIGURES

Sup Figure 1: Yearly distribution of OM cases attributable to *S. pneumoniae* serotypes not covered by PCVs from April 2014 to March 2019.

Sup Figure 2: Age distribution of OM cases attributable to *S. pneumoniae* during the study period.

442 **Sup Figure 3:** Ear discharge of OM cases attributable to *S. pneumoniae* during the study period.

445 **Sup Table 1:** List of all bacterial pathogens isolated from otitis media episodes detected at

446 Bangladesh Shishu Hospital and Institute during the study period: April 2015 – March 2019

447 (N=3735).

Bacterial pathogen detected	Number
Haemophilus influenzae	418
Streptococcus pneumoniae	292
Staphylococcus aureus	308
Pseudomonas spp	286
Acinetobacter spp	43
Proteus mirabilis	16
Klebsiella pneumoniae	47
Aeromonas hydrophila	1
Escherichia coli	38
Streptococcus pyogenes	101
Enterobacter spp	27
Serratia spp	6
Candida	15
Citrobacter spp	4

Morganella Morganii	4
Streptococcus agalactiae	6
Klebsiella oxytoca	3
Proteus vulgaris	8
Providencia spp	4
Streptococcus pneumoniae + Haemophilus influenzae	171
Streptococcus pneumoniae + Streptococcus pyogenes	1
Streptococcus pneumoniae + Staphylococcus aureus	32
Streptococcus pneumoniae + Pseudomonas spp	14
Streptococcus pneumoniae + Acinitobacter spp	3
Streptococcus pneumoniae + Klebsiella pneumoniae	7
Streptococcus pneumoniae + Escherichia coli	5
Streptococcus pneumoniae + Enterobacter spp	1
Streptococcus pneumoniae + Providencia spp	3
Streptococcus pneumoniae + Candida	1
Streptococcus pneumoniae + Proteus mirabilis	4
Streptococcus pneumoniae + Moraxella catarrhalis	1
Haemophilus influenzae + Staphylococcus aureus	78
Haemophilus influenzae + Streptococcus agalactiae	1
Haemophilus influenzae + Moraxella catarrhalis	1
Haemophilus influenzae + Pseudomonas spp	12
Haemophilus influenzae + Streptococcus pyogenes	20
Haemophilus influenzae + Enterobacter spp	6
Haemophilus influenzae + Acinitobacter spp	3
Haemophilus influenzae + Klebsiella pneumoniae	7
Haemophilus influenzae + Escherichia coli	6
Haemophilus influenzae + Providencia spp	1
Haemophilus influenzae + Proteus mirabilis	1
Haemophilus influenzae + Streptococcus spp	1
Haemophilus influenzae + Candida	1
Streptococcus pyogenes + Acinitobacter spp	1
Streptococcus pyogenes + Klebsiella pneumoniae	2
Streptococcus pyogenes + Enterobacter spp	2
Streptococcus pyogenes + Staphylococcus aureus	15
Streptococcus pyogenes + Pseudomonas spp	7
Moraxella catarrhalis + Acinitobacter spp	1
Moraxella catarrhalis + Pseudomonas spp	1
Staphylococcus aureus + Pseudomonas spp	12
Staphylococcus aureus + Escherichia coli	9

Staphylococcus aureus + Acinetobacter spp	6
Staphylococcus aureus + Enterobacter spp	10
Staphylococcus. aureus + Klebsiella pneumoniae	8
Staphylococcus aureus + Providencia spp	1
Staphylococcus aureus + Serratia spp	1
Staphylococcus aureus + Klebsiela oxytoca	1
Klebsiella pneumoniae + Escherichia coli	5
Klebsiella pneumoniae + Enterobacter spp	2
Klebsiella pneumoniae + Pseudomonas spp	10
Klebsiela oxytoca + Pseudomonas spp	1
E. coli + Pseudomonas spp	2
Enterobacter spp + Pseudomonas spp	2
Proteus mirabilis + Pseudomonas spp	4
Proteus mirabilis + Serratia spp	1
Acinitibacter sp+ Pseudomonas spp	1
Acinitibacter sp+ Citobacter spp	1
Streptococcus pneumoniae + Haemophilus influenzae + Acinitobacter spp	1
Streptococcus pneumoniae + Haemophilus influenzae + Klebsiella pneumoniae	1
Streptococcus pneumoniae + Haemophilus influenzae + Escherichia coli	2
Streptococcus pneumoniae + Haemophilus influenzae + Enterobacter spp	2
Streptococcus pneumoniae + Haemophilus influenzae + Providencia spp	1
Streptococcus pneumoniae + Haemophilus influenzae + Staphylococcus aureus	23
Streptococcus pneumoniae + Haemophilus influenzae + Pseudomonas spp	3
Streptococcus pneumoniae + Haemophilus influenzae + Streptococcus pyogenes	3
Streptococcus pneumoniae + Streptococcus pyogenes + Pseudomonas spp	2
Streptococcus pneumoniae + Proteus vulgaris + Pseudomonas spp	1
Streptococcus pneumoniae + Klebsiella pneumoniae + Pseudomonas spp	1
Streptococcus pneumoniae + Staphylococcus aureus + Acinitobacter spp	2
Streptococcus pneumoniae + Staphylococcus aureus + Klebsiella pneumoniae	1
Streptococcus pneumoniae + Staphylococcus aureus + Enterobacter spp	1
Streptococcus pneumoniae + Staphylococcus aureus + Proteus mirabilis	1
Streptococcus pneumoniae + Staphylococcus aureus +Pseudomonas spp	1
Haemophilus influenzae + Staphylococcus aureus + Escherichia coli	1
Haemophilus influenzae + Staphylococcus aureus + Acinitobacter spp	2
Haemophilus influenzae + Pseudomonas spp + Acinitobacter spp	1
Haemophilus influenzae + Pseudomonas spp + Escherichia coli	1
Haemophilus influenzae + Klebsiella pneumoniae + Acinitobacter spp	2
Haemophilus influenzae + Klebsiella pneumoniae + Enterobacter spp	1
Haemophilus influenzae + Klebsiella pneumoniae + Escherichia coli	1

Haemophilus influenzae + Streptococcus pyogenes + Acinitobacter spp	1
Haemophilus influenzae + Streptococcus pyogenes + Klebsiella pneumoniae	1
Haemophilus influenzae + Streptococcus pyogenes + Staphylococcus aureus	5
Haemophilus influenzae + Streptococcus pyogenes + Streptococcus agalactiae	1
Haemophilus influenzae + Staphylococcus aureus + Pseudomonas spp	2
Streptococcus pyogenes + Staphylococcus aureus + Pseudomonas spp	1
Streptococcus pyogenes + Staphylococcus aureus + Acinitobacter spp	1
No bacterial pathogen detected	1554
Total	3735