Digital phenotyping from wearables using AI characterizes psychiatric disorders and identifies genetic associations

Jason J. Liu[1,2,*], Beatrice Borsari[1,2,*], Yunyang Li[3], Susanna Liu[1,2], Yuan Gao[4], Xin Xin[1,2], Shaoke Lou[1,2], Matthew Jensen[1,2], Diego Garrido-Martín[5], Terril Verplaetse[6], Garrett Ash[7,8,9], Jing Zhang[10], Matthew J. Girgenti[6], Walter Roberts[6,9,#], Mark Gerstein[1,2,3,9,11,#]

[1] Program in Computational Biology and Bioinformatics, Yale University, New Haven, CT, 06511, USA
[2] Department of Molecular Biophysics and Biochemistry, Yale University, New Haven, CT, 06511, USA
[3] Department of Computer Science, Yale University, New Haven, CT, 06511, USA
[4] Yale School of Public Health, Division of Health Informatics, Yale University, New Haven, CT, 06511, USA
[5] Department of Genetics, Microbiology and Statistics, Universitat de Barcelona (UB), Barcelona, 08028, Spain
[6] Department of Psychiatry, Yale University School of Medicine, New Haven, CT, 06511, USA
[7] Section of General Internal Medicine, Yale University School of Medicine, New Haven, CT, 06511, USA
[8] Center for Pain, Research, Informatics, Medical Comorbidities and Education Center (PRIME), VA Connecticut Healthcare System, West Haven, CT, 06516, USA
[9] Department of Biomedical Informatics and Data Science, Yale University, New Haven, CT, 06511, USA
[10] Department of Computer Science, University of California, Irvine, CA, 92697, USA
[11] Department of Statistics and Data Science, Yale University, New Haven, CT, 06511, USA

* Co-first authors
# Co-corresponding authors (pi@gersteinlab.org)

Abstract

Psychiatric disorders are complex and influenced by both genetic and environmental factors. However, studying the full spectrum of these disorders is hindered by practical limitations on measuring human behavior. This highlights the need for novel technologies that can measure behavioral changes at an intermediate level between diagnosis and genotype. Wearable devices are a promising tool in precision medicine, since they can record physiological measurements over time in response to environmental stimuli and do so at low cost and minimal invasiveness. Here we analyzed wearable and genetic data from a cohort of the Adolescent Brain Cognitive Development study. We generated >250 wearable-derived features and used them as intermediate phenotypes in an interpretable AI modeling framework to assign risk scores and classify adolescents with psychiatric disorders. Our model identifies key physiological processes and leverages their temporal patterns to achieve a higher performance than has been previously possible. To investigate how these physiological processes relate to the underlying genetic architecture of psychiatric disorders, we also utilized these intermediate phenotypes in univariate and multivariate GWAS. We identified a total of 29 significant genetic loci and 52 psychiatric-associated genes, including *ELFN1* and *ADORA3*. These results show that wearable-derived continuous features enable a more precise representation of psychiatric disorders and exhibit greater detection power compared to categorical diagnostic labels. In summary, we demonstrate how consumer wearable technology can facilitate dimensional approaches in precision psychiatry and uncover etiological linkages between behavior and genetics.

Introduction

Psychiatric disorders of childhood and adolescence currently affect 1 in 7 youths in the United States and globally[1,2]. Externalizing disorders such as attention-deficit/hyperactivity disorder, and internalizing disorders such as anxiety, are among the most prevalent and represent a wide spectrum of dysfunctional behavior patterns[3]. Treatment barriers are complex and multifaceted but major contributors include our limited understanding of psychiatric phenotypes and difficulty identifying youth individuals that experience these disorders.

Traditionally, psychiatric disorders have been conceptualized as categorical macrophenotypes, based on clinical manifestations of a disease which are defined according to the number and type of symptoms, and the presence of distress or impairment[4-6]. While this has practical benefits in terms of reliability and ease of diagnosis, it poses several challenges to the research of these disorders, and consequently to the development of treatments. In fact, psychiatric disorders are complex and often comorbid, and this high degree of heterogeneity is not always accurately translated into categorical diagnosis labels, which may be defined by arbitrary cut-offs. Instead, intermediate phenotypes (i.e., quantitative traits that are positioned between genotype and macrophenotype) may better capture the heterogeneity potentially missed by existing diagnostic categories[7-9]. Additionally, genetic penetrance is expected to be higher for these intermediate phenotypes compared to macrophenotypes, enabling improved dissection of the genetic architecture underlying psychiatric disorders[10]. Nevertheless, many genome-wide association studies (GWAS) aimed at identifying genetic variants or biomarkers for psychiatric disorders do not consider these intermediate phenotypes and instead rely on dichotomised (i.e., binary) traits. In fact, identifying intermediate phenotypes with clinical and biological relevance remains a challenge[11].

Therefore, to improve our understanding of psychiatric disorders it is important that we identify intermediate phenotypes that not only offer a more comprehensive representation of an individual's behavior with respect to their environment, but also relate well with existing clinical definitions and aid in diagnosis. Once identified, these intermediate phenotypes can then be also used to guide more comprehensive studies to identify genetic associations and biomarkers that may ultimately improve precision treatments.

To achieve this goal, it is important to leverage new emerging technologies that can quantitatively assess an individual's behavioral patterns[12]. Wearable sensors such as smartwatches collect data that reflect physical and physiological processes (e.g., movement, pulse, metabolic intake), and can be used to infer higher-order behavioral events (e.g., sleep, exercise) and their temporal dynamics. Because of the documented relationship between such higher-order behavioral events and mental health, and given their low cost and minimal invasiveness, wearable devices have emerged as promising tools for mental health monitoring and psychiatric evaluation[13-15].

Therefore, wearable sensors have the potential for capturing intermediate phenotypes relevant to behavior and psychiatric disorders, ultimately enabling improved GWAS. However, significant computational challenges remain in generating intermediate phenotypes from wearable-derived data that describe the full spectrum of a given psychiatric disorder. Moreover, further curation of these intermediate phenotypes is necessary to identify genetic associations that have clinical and biological relevance.

To address these limitations, we developed an AI modeling framework that flexibly leverages data
from wearable devices to generate intermediate phenotypes in the form of static and dynamic
digital features. We establish the validity of these digital features as intermediate phenotypes by
classifying externalizing and internalizing disorders with an accuracy beyond baseline expectation,
and even surpassing the performance of some other gold-standard intermediate phenotypes such
as fMRI measurements[16-19]. Interpretability modules in our AI framework enable us to identify
key temporal and physiological insights between clinical diagnosis and digital features, further
supporting the validity of using these wearable-derived features as intermediate phenotypes. We
further curate these intermediate phenotypes and employ them in GWAS models to identify
genetic associations and biomarkers that capture the continuous spectrum of psychiatric disorders
and behavioral patterns. Finally, we identify 29 significant loci, several of which overlap
previously reported genetic variants associated with behavioral traits and mental illnesses and are
proximal to genes with a documented role in neurodevelopmental and psychiatric disorders.
104
In sum, this work shows how wearable devices can advance our understanding of psychiatric
disorders by establishing a more objective and dimensional approach that can ultimately lead to
improved treatments in precision psychiatry.

Results

Leveraging the Adolescent Brain Cognitive Development cohort

To improve our understanding of psychiatric disorders, we leveraged and analyzed a dataset from
a cohort of US adolescents recruited by the NIH Adolescent Brain Cognitive Development
Consortium (ABCD) project, consisting of clinical, wearable, and genetic data (**Fig. 1**). The ABCD
cohort consists of a total of 11,878 adolescents (5682 males and 6196 females), of age between
nine and fourteen years and belonging to four different ethnicities (**Suppl. Fig. S4.1**). We
identified nine categories of psychiatric phenotypes (**Suppl. Table S1.1**), which were established
using a gold standard parent diagnostic semi-structured interview (Kiddie Schedule for Affective
Disorders and Schizophrenia-5)[20]. The healthy controls represented adolescents who did not meet
the criteria for any of those nine psychiatric disorders. We defined these clinical labels as the
categorical macrophenotypes in the study (**Fig. 1A-B**). Our modeling framework also utilized data
from cognitive tests (e.g., NIH Toolbox) and behavioral checklists (**Fig. 2A, Suppl. Fig. S2.1**).
121
Generating intermediate phenotypes from wearable-derived data

We processed data obtained from FitBit smartwatches, which comprise measurements of heart
rate, calories, activity intensity, steps, metabolic equivalents, sleep level and sleep intensity (**Fig.**
1C, Suppl. Table S1.2)[21]. These measurements quantify an individual's physiological processes
and their real-time changes in response to environmental stimuli, and can thus provide key
information about an individual's behavior.
128
129 To reconstruct the full spectrum of an individual's behavioral functioning from these data, we
130 applied two different feature engineering techniques, allowing us to generate wearable-derived
131 dynamic and static features, which we consider as intermediate phenotypes. The dynamic features
132 preserve the time-varying nature of the original data as a time series, enabling sequential and
133 temporal patterns of the data to be retained. In contrast, the static features summarize patterns of

Figure 1. Leveraging clinical, digital, and genetic data of the ABCD cohort to improve characterization of psychiatric disorders.
A) Framework schematic describing how intermediate phenotypes from wearable-derived data are leveraged to better understand the association between macrophenotype and genotype. The link between intermediate phenotype and macrophenotype serves as construct validity and aid in diagnostics. Wearable GWAS is performed through genotype-to-intermediate-phenotype association studies. **B)** The Adolescent Brain Cognitive Development (ABCD) cohort contains 11,878 individuals spanning nine different categorical macrophenotypes based on clinical diagnosis from the Kiddie Schedule for Affective Disorders and Schizophrenia-5. A breakdown of the counts of each disorder is shown in the bottom bar graph, with anxiety disorder and ADHD being the most prevalent. "Bipolar" refers to bipolar or psychotic disorders. **C)** Digital data from FitBit biosensors are collected for 5,339 individuals. The collected time series data are then processed into dynamic and static features, with information spanning various physiological and higher order processes. **D)** Genetic data are collected by the ABCD consortium through Smokescreen genotyping array. Imputed genotypes are used for downstream GWAS analyses. The genotype arrays are subjected to best-practice processing and QC to ensure included individuals and SNPs are of high quality. PCA performed on 8,791 individuals and 157,556 genotyped SNPs reveals distinct ancestral clusters across the cohort and the inferred genotype principal components (PCs) are used as covariates in downstream analyses.

134 the digital data to produce time-invariant, quantitative features that are commonly used in
135 downstream modeling[14,22].
136
137 To generate dynamic features, we performed signal imputation and processing after filtering the
138 individuals with sparse data, and obtained 48 channels of time series (**Figure 2B**, **Suppl. Fig.**
139 **S2.3**). Compared to the static features, this further processing allowed us to preserve both local
140 and global temporal patterns potentially relevant to characterizing behavior and neurological
141 response to stimuli[8,23].
142
To generate static features, we first collected a total of 49 FitBit summary-based features (**Suppl.**
Table S2.1a-b). We next applied descriptive statistics (e.g. mean, median, etc.) to each of these
features and generated a total of 258 static features for each individual (**Fig. 2C, Suppl. Table**
S2.2)[14,24]. We then grouped these static features into seven main clusters, each of which
summarizes different aspects of physiological and behavioral processes, such as heart rate, sleep
duration and quality, metabolic intake or physical activity (**Fig. 2D**).
149
150 Altogether, static and dynamic features represent the physiological and behavioral profiles of the
151 adolescents, and can be leveraged as intermediate phenotypes in a wide range of analyses to better
152 characterize psychiatric phenotypes, such as generating disorder-specific probability risk scores,
153 macrophenotype classification, model interpretability, and biomarker identification via wearable
154 GWAS.
155
Predicting psychiatric phenotypes from wearable-derived intermediate phenotypes
To demonstrate the validity of static and dynamic features as clinically relevant intermediate
phenotypes and to evaluate their utility as a diagnostic tool, we employed these features in an array
of classification tasks to identify individuals with either an externalizing (ADHD) or internalizing
(anxiety) disorder from their typically developing peers. We selected ADHD and anxiety due to
their high prevalence in adolescents, which is mirrored in the cohort (**Fig. 1B**)[25].
162
We applied a gradient boosting machine learning algorithm, XGBoost, for classification tasks
using static features (**Fig. 2C-D**)[26]. On the other hand, to fully leverage the time series nature of
the dynamic features, we used a convolutional neural network for time series, featuring depthwise
separable convolution, called Xception (**Fig. 2E**)[27]. Variable convolutional filters and residual
(skip) connections, coupled with efficient parametrization, allow information encoded in both
small and large receptive fields to be more optimally leveraged. Practically, this framework takes
into account local and global patterns of physiology and behavior when performing downstream
classification of psychiatric disorders. In both modeling approaches we included covariates that
accounted for demographic features, family history of disorders, and other clinical information
(**Fig. 2A, Suppl. Table S1.2**). To assess the benefit, in terms of model performance, of including
wearable-derived data, we also trained a baseline model using just the covariates, which served as
a comparison to the models including static or dynamic features. In practice, this comparison
allowed us to determine whether wearable-derived features can improve diagnostic accuracy
relative to that achievable using only a widely used broadband behavior rating scale.
177
178 After data filtering, we first used static features to classify 216 individuals with ADHD (an
179 externalizing disorder) versus 1,737 of their typically developing peers (healthy controls) (**Fig. 3A**

Figure 2. Workflow for data processing, feature engineering, and model architecture.
A) ABCD cohort metadata including various demographic features, cognitive test scores, and clinical characteristics are used as covariates and represent the input features used in our baseline comparison model. Features shown in this plot correspond to the filtered set of individuals with wearable data. **B)** Digital data collected by wearable biosensors are used to generate dynamic features after signal processing and imputation steps. Together with the processed covariates, these time series features represent the input features for the dynamic model. **C)** Summary statistics applied to digital data collected by wearables are used to generate a total of 258 static features. In addition to the covariates, these are the input features used in the static model. The static model leverages the machine learning framework, XGBoost, for downstream tasks such as risk score generation, classification, model interpretability, and wearable GWAS. **D)** Hierarchical clustering of the static features yields seven distinct physiological clusters of wearable data. **E)** The dynamic model is based on the Xception deep learning framework, and uses the generated 48 channels from the dynamic features and covariates as input into a convolution-like model. The architecture consists of six inception layers and residual connections. Global average pooling and a fully connected layer allow for similar downstream tasks as mentioned in C).

and Suppl. Fig. 3.1). Using static features with XGBoost, we achieved an average area under the receiver operating characteristic curve (AUROC) of 0.87 and precision of 0.79. When using the dynamic features and Xception, we were able to achieve an average AUROC of 0.89 and precision of 0.83. The baseline model consisting of only the covariates achieved an average AUROC of 0.83, suggesting that the inclusion of wearable-derived features facilitates a clinically meaningful improvement in diagnostic accuracy. This improvement between the baseline and dynamic features model demonstrates statistical significance (one-sided t-test between baseline and dynamic features model, *p* value = 0.0022).

Second, we evaluated the performance of our model using static or dynamic features in the classification of 666 individuals diagnosed with anxiety disorder (internalizing disorder) versus 1,737 of their typically developing peers (healthy controls) (**Fig. 3B and Suppl. Fig. 3.2)**. Here, we again repeated the use of the same modeling framework, i.e., static features with XGBoost and dynamic features with Xception, and compared it to the baseline covariate model. We found that static and dynamic features achieve an average AUROC of 0.69 and 0.71 and precision of 0.64 and 0.68, respectively. In both models, the performance was greater than that of the baseline model (average AUROC of 0.67), with the dynamic features model showing the most significant performance improvement (one-sided t-test between baseline and dynamic feature model, *p* value = 0.00016). Overall, the fact that the models using dynamic features achieved the highest performance suggests the usefulness of the temporal patterns intrinsic to wearable-derived data towards understanding behavior.

Interpreting wearable features prioritized by the deep learning model

Deep learning methods are typically characterized by complex internal structures that cannot be easily interpreted by humans. While maximizing the classification accuracy is one crucial aspect for characterizing complex phenotypes, understanding which features are most important in terms of their individual contribution to performance is also critical. To this end, we utilized ablation techniques to determine the relative contribution of each individual feature to model performance. For the ADHD classification task, heart rate was the most important feature (largest change in AUROC), followed by other dynamic features (i.e., sleep, steps, METs) as well as covariates such as demographics, family history, and cognitive scores from picture memory and stop-signal reaction time tests (**Fig. 3C, Suppl. Fig. S3.3**).

On the other hand, the ablation study for the anxiety classification task revealed a different set of important features. In this case, sleep quality and stage, calories, and step count were the most important dynamic features, whereas heart rate features, which were extremely important for classifying ADHD, were not prioritized in the anxiety model (**Fig. 3D, Suppl. Fig. S3.5**). Additionally, while the anxiety model prioritized some covariates that were relevant also for the ADHD model (e.g., sex, family history, and family divorce), cognitive scores from tests such as picture memory did not appear to be important for the identification of individuals diagnosed with anxiety, consistent with theory-driven accounts of neurocognitive aspects of anxiety disorders[28].

Additionally, we assessed the importance for model performance of dynamic features at various times throughout the day by adapting gradient-weighted class activation mapping (Grad-CAM) strategies[29]. We calculated the relative importance of each time point in our dynamic features. For ADHD, we observed enriched significance of the heart rate dynamic feature around the early

Figure 3. Performance and interpretability of psychiatric phenotype classification models.
A-B) Model performance for baseline, static, and dynamic models employed for classifying individuals with ADHD (blue, top) or individuals with anxiety disorder (green, bottom) versus healthy controls. *P* values were calculated using one-sided t-test. **C-D)** Feature importance based on ablation studies for the dynamic model for ADHD (blue, top) and anxiety disorder (green, bottom) classification. Wearable-derived dynamic features are shown in red font and clinical features (covariates) are shown in black font. Feature importance is equivalent to the decrease in model performance (AUROC) after removal of the given feature. **E-F)** Temporal importance during a 48-hour period for dynamic features in ADHD (blue, top) or anxiety disorder (green, bottom) classification based on the GRAD-CAM interpretability module. Importance is represented as the GRAD-CAM score, based on each time points contribution towards model performance.

afternoon, potentially suggesting stronger behavioral differences between adolescents with ADHD
and their typically developing peers (healthy controls) during this time of day (**Fig. 3E, Suppl.**
Fig. S3.4). This is consistent with clinical research demonstrating time-of-day effects on ADHD
symptom expression[30]. In contrast, sleep-related dynamic features during the night are much more
informative in classifying anxiety, consistent with clinical expectations (**Fig. 3F, Suppl. Fig.**
S3.6)[31]. Together, the ablation studies suggest a role for wearable-derived features to not only
serve as quantitative intermediate phenotypes, but also to more closely reveal insights into the
behavioral and physiological temporal patterns related to categorical macrophenotypes.
234
Using wearable-derived features as intermediate phenotypes for wearable GWAS
Our accurate classification of both internalizing and externalizing psychiatric phenotypes based
on wearable-derived features suggests that these features can serve as useful intermediate
phenotypes and may be leveraged to identify genetic associations with psychiatric conditions (**Fig.**
4, Suppl. Fig. S4.1-S4.6). To this end, we first focused specifically on ADHD, given the higher
predictive power observed with our models (**Fig. 3A-B**) and its higher estimated heritability
compared to anxiety (77-88% vs. 30-60%)[32,33]. We selected 1,191 individuals (137 individuals
with ADHD and 1,054 healthy control individuals) with genetic and wearable data available, and
performed a GWAS using the continuous prediction scores obtained from our wearable modeling
framework. In practice, these scores represent risk probabilities for ADHD[34,35]. We identified 10
genome-wide (p value $< 5 \cdot 10^{-8}$) significant loci and 21 psychiatric or brain-related genes (**Fig. 4B,**
Table 1, Suppl. Fig. S4.7-S4.9, Suppl. Table S4.1). Three of the identified genes (*ADORA3*,
PSMD11 and *DLG4*) have been previously associated with ADHD, bolstering the overall
functional significance of the results[36-38]. Furthermore, several of these loci overlap with
previously reported GWAS SNPs related to ADHD, neuroticism, sleep disruption and other
clinically relevant traits (**Fig. 4B**, **Suppl. Fig. S4.10, Suppl. Table S4.2**). Note that here we used
a continuous risk score as a univariate response variable for the GWAS. In comparison, when
performing a traditional case–control GWAS for ADHD on the same set of individuals using the
binary diagnostic label (presence/absence of disorder) as response variable, we did not identify
any significant loci (**Fig. 1A, Fig. 4A, Suppl. Fig. S4.7-S4.9**). This result is consistent with the
higher statistical power of continuous measurements over dichotomized (i.e., binary) traits (**Suppl.**
Fig. S4.11), and with the findings that intermediate phenotypes show higher genetic penetrance
compared to macrophenotypes[7,39-41].
258
While the analysis above collapses wearable-derived features into a single continuous variable that
summarizes the risk score for a particular disorder (i.e., GWAS with a univariate response), it is
also possible to directly use the full set of wearable-derived features as a more comprehensive and
richer phenotype to represent the continuum of psychiatric disorders and their latent manifestations
(i.e., GWAS with a multivariate response). In fact, these features can collectively capture
behavioral patterns by measuring physiological processes and their real-time changes in response
to environmental stimuli, and unlike disease risk scores, are not restricted to a specific cohort of
individuals[42,43]. In what follows, we employed a multivariate nonparametric test of association to
regress the vector of wearable-derived features on the genotype of each genetic variant, employing
a larger cohort that spans healthy controls and individuals with any psychiatric disorder (n =
2,410)[44]. From this novel type of GWAS we identified 19 significant loci and 31 genes with a
documented role in neurodevelopmental and psychiatric disorders (**Fig. 4C, Table 1, Suppl. Fig.**
S4.12-S4.14, and Suppl. Table S4.3). Many of these loci overlap previously identified GWAS

Figure 4. Manhattan plots summarizing the results of the univariate and multivariate wearable GWAS.
A) Left panel: Case-control GWAS on 1,191 individuals from the ABCD cohort. We employed the clinical diagnosis label as the univariate binary response variable for the GWAS (1 = "individual with ADHD", $n = 137$; 0 = "healthy control individual", $n = 1,054$). Right panel: Manhattan plot showing the $-\log_{10} p$ value of association between the genetic variants and the univariate binary response variable. No genetic variants passed the genome-wide significance threshold (p value $< 5 \cdot 10^{-8}$; blue line). Genetic variants with a suggestive p value ($< 10^{-5}$) are represented as green dots.
B) Analogous representation to panel A) using the wearable-derived risk scores for ADHD as univariate continuous response variable. The GWAS was performed on the same set of 1,191 individuals and using the same set of covariates as in panel A). 10 and 427 loci passed the p value thresholds of $5 \cdot 10^{-8}$ and $1 \cdot 10^{-5}$, respectively. A detailed list of genome-wide significant loci is provided in **Table 1** and **Suppl. Table S4.1**. Loci chr1:111,372,165-111,482,359, chr17:7,101,607-7,101,608, and chr17:32,256,997-32,283,356 are proximal to genes *ADORA3* (72 Kb), *DLG4* (86 Kb) and *PSMD11* (174 Kb) (highlighted in dark blue) respectively, which have been previously associated with ADHD. Other proximal genes related to other psychiatric disorders are highlighted in pink (evidence obtained from OpenTargets). Brain-related traits associated with genetic variants overlapping the ten genome-wide significant loci are highlighted in orange. GWAS associations were obtained from the EBI-NHGRI GWAS catalog. **C)** Analogous representation to panel A) using the 258 wearable-derived static features as multivariate continuous response variable. The GWAS was performed on a set of 2,410 individuals (both healthy controls and individuals with any disorder). 19 and 314 loci passed the p value thresholds of $5 \cdot 10^{-8}$ and $1 \cdot 10^{-5}$, respectively. A detailed list of genome-wide significant loci is provided in **Table 1** and **Suppl. Table S4.3**. Neuropsychiatric-related genes proximal to the identified loci are shown in pink. Brain-, heart-, and sleep-related traits with associated variants overlapping the 19 loci are highlighted in orange.

GWAS Method	Locus	Chr	Start	End	Lead Variant	Position	p value	Genes
Univariate continuous	1	1	111,372,165	111,482,359	rs114081965	111,372,166	1.11E-08	*TMIGD3, ADORA3, RAP1A, CHI3L2*
Univariate continuous	2	3	161,873,055	161,927,820	rs79203233	161,909,261	3.89E-08	-
Univariate continuous	3	4	184,417,766	184,424,056	rs1425551	184,421,904	2.15E-08	*IRF2, CASP3, PRIMPOL*
Univariate continuous	4	10	121,524,611	121,582,200	rs140794722	121,524,612	2.31E-09	*FGFR2*
Univariate continuous	5	11	38,982,793	39,384,610	rs151239852	39,273,497	3.80E-08	-
Univariate continuous	6	14	26,214,683	26,216,532	rs149074469	26,216,532	2.95E-08	*NOVA1*
Univariate continuous	7	14	62,903,677	63,014,798	rs143225169	63,014,798	3.89E-08	*KCNH5, RHOJ*
Univariate continuous	8	17	7,101,607	7,101,608	rs11653054	7,101,608	8.51E-09	*CLEC10A, DLG4*
Univariate continuous	9	17	32,256,997	32,283,356	rs6505293	32,270,863	1.12E-08	*RHBDL3, RHOT1, C17orf75, ZNF207, PSMD11, LRRC37B, CDK5R1, MYO1D*
Univariate continuous	10	19	4,495,610	4,495,611	rs150855276	rs150855276	2.75E-08	-
Multivariate continuous	11	2	65,047,524	65,158,021	rs147959551	65,140,366	4.47E-08	*RAB1A, ACTR2, SLC1A4, SPRED2*
Multivariate continuous	12	7	1,789,321	1,791,353	rs113525298	1,791,353	5.10E-09	*MAD1L1, ELFN1, PSMG3, MAFK*
Multivariate continuous	13	14	23,392,601	23,418,974	rs365990	23,392,602	5.33E-09	*MYH6, CMTM5, IL25, BCL2L2-PABPN1, BCL2L2*
Multivariate continuous	14	16	79,283,253	79,302,474	rs8051625	79,288,217	5.98E-09	*WWOX*
Multivariate continuous	15	X	10,444,818	10,481,837	rs73492938	10,467,098	1.03E-08	*MID1, CLCN4*
Multivariate continuous	16	X	16,350,347	16,647,557	rs149504239	16,454,383	1.09E-09	-
Multivariate continuous	17	X	22,606,616	22,606,617	-	22,606,617	7.23E-12	*PHEX*
Multivariate continuous	18	X	27,164,449	28,601,484	rs200021485	27,782,503	7.89E-10	-
Multivariate continuous	19	X	30,757,116	30,896,124	rs150685307	30,831,630	1.70E-08	*TAB3, GK*
Multivariate continuous	20	X	40,530,738	40,530,739	rs149441354	40,530,739	3.17E-08	*ATP6AP2*
Multivariate continuous	21	X	45,687,487	45,858,851	-	45,819,973	8.04E-09	-
Multivariate continuous	22	X	62,901,143	64,518,530	rs7883352	63,935,445	1.09E-08	*ARHGEF9*
Multivariate continuous	23	X	67,188,492	68,693,105	rs180773472	68,693,105	9.82E-11	*OPHN1, AR*
Multivariate continuous	24	X	96,759,291	96,961,667	-	96,759,292	4.40E-10	-
Multivariate continuous	25	X	102,017,055	103,738,764	-	102,017,056	1.99E-10	*RAB40AL, GPRASP2*
Multivariate continuous	26	X	124,254,095	124,260,611	rs12012355	124,254,420	2.08E-10	*TEX13D, SH2D1A, TENM1, STAG2*
Multivariate continuous	27	X	143,176,627	143,212,922	rs12011450	143,190,440	1.69E-08	*SPANX4*
Multivariate continuous	28	X	146,722,378	146,739,375	rs5966302	146,729,545	6.26E-09	-
Multivariate continuous	29	X	148,716,235	148,915,910	rs57168704	148,907,585	5.71E-09	*AFF2*

Table 1. Results for the 29 genetic loci identified by the univariate and multivariate continuous wearable GWAS.
For each locus we report the GWAS Method (univariate or multivariate continuous) that identifies the locus, the genomic coordinates in human assembly GRCh38, and the lead variant rsID with corresponding genomic position and p value. Brain-related or neuropsychiatric genes proximal to the locus are also listed (**Suppl. Tables S4.1** and **S4.3**).

272 SNPs related to heart, sleep, metabolism, and brain traits (**Fig. 4C**, **Suppl. Fig. S4.15**). This aligns
273 with the close association between physiological functions, the central nervous system, and
274 individual behavior.
275
Functionally dissecting and interpreting novel wearable GWAS loci

To further investigate the loci identified by the behavioral GWAS, we dissected the variants using
a battery of publicly available genomic resources[45,46]. Many of these loci overlap either GTEx
expression quantitative trait loci (eQTLs) or ENCODE candidate cis-regulatory elements (cCREs),
suggesting a link between the biochemical activity of these variants and their functional impact on
macrophenotype (**Suppl. Table S4.3**). We also explored the impact of these loci beyond
behavioral traits and their relationship with clinical psychopathology. For example, behavioral
traits significantly associated with a specific genetic variant may correlate with clinical symptoms
of a specific psychiatric cohort. Indeed, in some cases we show that the genetic variant in question
is also differentially enriched between that specific psychiatric cohort and healthy individuals.
286
For instance, we found the minor allele (G) at rs365990 to be significantly associated with an
increase in mean heart rate and a decrease in interday heart rate variation (**Fig. 5A, left**). The
variant, missense for *MYH6*, had been previously linked to atrial fibrillation, ventricular
tachycardia and resting heart rate, and the entire locus shows a significant enrichment of chromatin
features in heart samples compared to other tissues and organs (**Suppl. Fig. S5.1**)[47-50]. We also
found the same allele to be enriched in the bipolar/psychotic disorder cohort compared to healthy
controls (**Fig. 5A, right**). This cohort included youth meeting criteria for bipolar or unspecified
psychotic spectrum disorder, and such severe pathology is known to be associated with
characteristic irregularities in heart activity[51-53]. SNP rs365990 is also a GTEx eQTL for the
CMTM5 gene (**Suppl. Fig. S5.2**), which is highly expressed in brain subregions and has been
implicated in stress response and childhood adversity, further supporting the relevance of this locus
for psychiatric conditions in addition to heart pathophysiology[46,54].
299
In a similar fashion, we explored variants rs113525298 and rs147959551. The minor allele at
rs113525298 is associated with prolonged periods in bed and shorter vigorously active time during
the day, and appears at a lower frequency in the ADHD cohort compared to healthy individuals
(**Fig. 5B**). This suggests a potential protective role of the allele against hyperactivity disorders,
further supported by the proximity of the SNP to *ELFN1*, previously implicated in the
pathophysiology of ADHD[55,56]. On the other hand, we found the minor allele at rs147959551
associated with shorter sedentary time at night and a shorter period of time identified as sleep based
on heart rate, two features suggestive of sleep disruption (**Fig. 5C**). The same allele is also enriched
in individuals with depression disorder, consistent with growing evidence implicating sleep
impairment as a transdiagnostic feature of many forms of adolescent psychopathology (**Fig. 5C,**
right)[57,58].
311
312 Overall, these results highlight how wearable-derived features can be leveraged as intermediate
313 phenotypes in GWAS and enable the identification of genetic variants relevant to clinical
314 psychiatry with significant effects on exhibited behavior in adolescents.
315

Figure 5. Exploring the genetic-physiological-psychiatric axis with wearable GWAS.
A) Left panel: rs365990 (chr14:23,392,602, A/G) is located in exon 25 of *MYH6*, and is associated with changes in wearable-derived heart rate features (multivariate GWAS *p* value = 5.33·10⁻⁹). The boxplots show distributions of covariate-adjusted mean and interday coefficient of variation (CV) for heart rate across genotype groups at rs365990 (AA *n* individuals = 1,228; AG *n* individuals = 1,509; GG *n* individuals = 519). *P* values (two-sided Wilcoxon Rank-Sum test) for each pairwise comparison are also displayed, encoded as follows: *** (*p* ≤ 0.001), ** (0.001 < *p* ≤ 0.01), * (0.01 < *p* ≤ 0.05), n.s. (*p* > 0.05). For visualization purposes, outliers are not shown. Right panel: enrichment, displayed as odds-ratio (log2 OR; y axis) of the minor allele (G) in individuals with different psychiatric disorders (x axis) compared to healthy controls. OR estimates and 95% confidence interval (error bar) are displayed. The red horizontal dashed line indicates no enrichment. The G allele is significantly more enriched in individuals with bipolar/psychotic disorder compared to healthy controls (two-sided Fisher test p value: 8.00·10⁻³; FDR-adjusted *p* value: 7.00·10⁻²). **B)** Similar representation for rs113525298 (chr7:1,791,353; AA *n* individuals = 2,294; AG *n* individuals = 101; GG *n* individuals = 15). rs113525298 is located 125 Kb from *ELFN1*, a gene that encodes a postsynaptic protein involved in the temporal dynamics of interneuron recruitment[65,66]. *Elfn1* mutant mice exhibit hyperactivity that is treatable by psychostimulant medication[55,56]. The G allele at rs113525298 is associated with increased minimum number of first-out-of-bed minutes and decreased minimum number of total-vigorously-active minutes (multivariate GWAS *p* value = 5.10·10⁻⁹), and is significantly more enriched in healthy controls compared to individuals with ADHD (two-sided Fisher test *p* value: 9.00·10⁻⁴; FDR-adjusted *p* value: 6.00·10⁻³). **C)** Similar representation for rs147959551 (chr2:65,140,366; AA *n* individuals = 2,279; AG *n* individuals = 117; GG *n* individuals = 14), located near a cluster of genes relevant for several psychiatric disorders, such as *ACTR2* (schizophrenia; 87 Kb), *SLC1A4* (schizophrenia, bipolar disorder, major depressive disorder; 117 Kb) and *SPRED2* (schizophrenia, OCD; 170 Kb)[67-77]. The G allele of rs147959551 is associated with decreased mean number of sedentary-time-at-night minutes and decreased maximum number of sleep-based-on-heart-rate minutes (multivariate GWAS *p* value = 4.47·10⁻⁸), and is significantly more enriched in individuals with depressive disorder compared to healthy controls (two-sided Fisher test *p* value: 9.74·10⁻³; FDR-adjusted *p* value: 7.80·10⁻²).

Discussion

Psychiatric disorders have been traditionally described with diagnostic categories based on retrospective self-report of symptom sets. However, current efforts in the field are increasingly leveraging novel technologies to transition from retrospective self-reporting and fixed symptom sets to more dimensional conceptualizations, aiming to capture the complex and heterogeneous nature of psychiatric disorders for more accurate research into their underlying structure[6]. One approach to enhancing dimensional models is the use of intermediate phenotypes—quantitative traits linked more closely to a disorder's underlying molecular pathways. Although intermediate phenotypes have been derived from cellular, tissue, and organ levels of information, computational strategies that generate useful intermediate phenotypes in the behavioral domain are currently limited. Wearable biosensors such as smartwatches offer a unique opportunity to objectively study psychiatric disorders in a non-invasive way by measuring their underlying physiological foundations of behavior over time.

Towards this end, we used wearable data to generate static and dynamic features that were employed by our AI modeling framework as intermediate phenotypes to distinguish between adolescents with and without psychiatric disorders. Models utilizing these wearable-derived intermediate phenotypes performed comparably to those based on more expensive data sources such as fMRI measurements[18,59]. To gain critical theoretical insights and inform treatment development efforts, we augmented the modeling framework with interpretability modules, allowing us to pinpoint temporal and functional regions of the time series that were highly correlated with overall disease state[60]. These interpretability modules have the potential to facilitate mechanistic studies that offer deeper insight into the underlying complexities of these disorders. For example, our interpretability modules revealed that heart rate time series held high importance in predicting ADHD. This finding aligns with the clinical manifestation of ADHD – affected children are characterized by episodes of heightened arousal that are often incongruent with environmental demands[61]. Conversely, the interpretability modules identified sleep intensity and quality as key predictors in our anxiety disorder models, in line with known disruptions in sleep patterns and circadian rhythms commonly seen in youth with anxiety disorders[62].

Wearable-derived intermediate phenotypes are not just effective for detecting the presence of psychiatric disorders in individuals; they also serve as a valuable research tool for understanding the correspondence between behavior patterns and molecular attributes. This comprehensive approach helps to uncover the foundational elements of pathological behavior patterns. In this context, we focused on establishing links with genetics. Specifically, we showed that these intermediate phenotypes can serve as response variables in GWAS models. Their continuous nature enhances statistical power compared to categorical diagnostic labels. Furthermore, we took advantage of the features' correlated structure to create multivariate response variables for GWAS. This strategy is statistically advantageous because it mitigates the multiple testing burden associated with evaluating the numerous (>250) independent features. Conversely, from a biological standpoint, these wearable GWAS allowed us to explore triaxial associations encompassing genetic, physiological, and psychiatric factors. Utilizing our framework, we successfully identified a significant association between a missense variant of the *MYH6* gene, which encodes the cardiac muscle myosin, and heart rate patterns. Heart activity receives complex inputs from the CNS, which implies behavioral influence and, in combination with our GWAS, supports the notion of a gene-behavior-disorder pathway[63]. Building on this finding, we discovered

enrichment of the same genetic variant among individuals with bipolar/psychotic disorders, psychiatric conditions known to be associated with characteristic irregularities in heart activity[51]. While additional research is needed to confirm such associations, our findings resonate with the objectives of the RDoC initiative[6]. Specifically, wearable-derived intermediate phenotypes serve as objective markers of behavior, bridging lower-level biological systems like genetics to broader psychiatric disorders.

While we have employed these wearable-derived intermediate phenotypes in a targeted research context (i.e., to enhance a psychiatric GWAS), their broad applicability make them promising for other domains of health research. For example, the risk scores generated by our AI-modeling framework could be used to assess disorder severity, and the genetic variants identified by our wearable GWAS could be employed to construct more comprehensive polygenic risk scores for behavioral and psychiatric disorders. Unlike other diseases (e.g., cancer) where objective biomarkers are more common, psychiatry faces a significant barrier in treatment due to the lack of objective and sensitive screening methods[64]. Therefore, these physiological and genetic features could be leveraged as objective biomarkers to more accurately subtype patients within diagnostic categories, which in turn could help move towards precision treatment delivery in psychiatry.

Although the results presented in this study require further experimental validation, they illuminate the transformative potential of wearable devices combined with AI modeling frameworks for deepening our understanding of complex behavioral and psychiatric traits. We anticipate that further development of our AI modeling framework, coupled with an expanded array of wearable devices, could fundamentally transform how psychiatric disorders are measured and understood in both research and clinical settings. This could lead to more nuanced digital intermediate phenotypes and open new avenues for the study of human behavior.

Acknowledgements

This work was funded by support from the National Institute of Mental Health (RF1MH123245) and the National Institute on Alcohol Abuse and Alcoholism (U54AA027989).

Data used in the preparation of this article were obtained from the Adolescent Brain Cognitive DevelopmentSM (ABCD) Study (https://abcdstudy.org), held in the NIMH Data Archive (NDA). This is a multisite, longitudinal study designed to recruit more than 10,000 children age 9-10 and follow them over 10 years into early adulthood. The ABCD Study® is supported by the National Institutes of Health and additional federal partners under award numbers U01DA041048, U01DA050989, U01DA051016, U01DA041022, U01DA051018, U01DA051037, U01DA050987, U01DA041174, U01DA041106, U01DA041117, U01DA041028, U01DA041134, U01DA050988, U01DA051039, U01DA041156, U01DA041025, U01DA041120, U01DA051038, U01DA041148, U01DA041093, U01DA041089, U24DA041123, U24DA041147. A full list of supporters is available at https://abcdstudy.org/federal-partners.html. A listing of participating sites and a complete listing of the study investigators can be found at https://abcdstudy.org/consortium_members/. ABCD consortium investigators designed and implemented the study and/or provided data but did not necessarily participate in the analysis or writing of this report. This manuscript reflects the views of the authors and may not reflect the opinions or views of the NIH or ABCD consortium investigators.

Author Contributions

Conceptualization: JL, BB, WR, MG

Methodology: JL, BB, YL, SL, YG, XX, SKL, MJ, DGM, TV, GA, JZ, MJG, WR, MG

Investigation: JL, BB, YL, SL, YG

Visualization: JL, BB, YL, SL, YG

Funding acquisition: MG

Data curation: TV, WR, MG

Supervision: WR, MG

Writing – original draft: JL, BB

Writing – review & editing: JL, BB, YL, SL, YG, XX, SKL, MJ, DGM, TV, GA, JZ, MJG, WR, MG

Competing interests

The authors declare they have no competing interests.

Methods

Dataset Description

ABCD Dataset

The Adolescent Brain Cognitive Development (ABCD) study is a comprehensive longitudinal project initiated in 2015 with the purpose of characterizing the neural, cognitive, and behavioral aspects of adolescent development. Commissioned by a consortium of U.S. federal agencies, ABCD investigators deeply phenotyped a large and representative sample of children aged 9-14 with plans to track their development into early adulthood. The ABCD dataset incorporated multimodal brain imaging data, substance use history, behavioral and psychological measures, genetic data, and an all-encompassing collection of demographic, physical health and activity, mental health, and environmental information, including data derived from wearable devices.

Clinical Diagnoses

Clinical diagnoses were operationalized using the parent report version of the Kiddie Schedule for Affective Disorders and Schizophrenia (KSADS). The KSADS is a gold standard semi-structured diagnostic interview that is used to establish a broad range of clinical diagnoses in children and adolescents. It has been used previously to define clinical groups in case/control studies conducted with data from the ABCD study[59].

Cohort Definitions

We identified several clinical groups of interest in order to evaluate our framework across different forms of psychopathology. The *nonclinical comparison* cohort was composed of youth who did not meet any current diagnostic criteria for any disorder on their most recent administration of the parent report KSADS. Similar diagnostic categories, based on ICD10 diagnostic codes, were combined to create cohorts with sufficient sample sizes for our modeling framework. Each clinical cohort was composed of the following diagnostic groups, based on the current reported symptom sets (**Suppl. Table S1.1**). Some codes were included in multiple categories to balance the need for sufficient sample size and homogenized cohort definitions.

Preprocessing and Quality Control of Wearable Device Data

We commenced by combining data from seven distinct wearable-derived modalities (heart rate, calories, intensity, steps, METs, sleep level, and sleep intensity) for 5,339 individuals into a single dataframe, resulting in highly sparse data structures. We excluded individuals with at least one missing wearable modality, leaving us with 3,538 participants. To address the impact of missing values on further analysis, we implemented a rigorous quality control procedure. In the initial phase, we examined all potential time windows for two selected days each week per data modality. Our objective was to balance the maximization of data inclusion with the assurance of its quality. We pinpointed the time window that offered the best alignment - that is, the period which had the highest number of valid measurements across all modalities. This procedure enabled us to determine the most suitable time window for downstream analysis, taking into account both the richness of the data and the necessity for top-quality inputs. In the next stage, we established a criterion that each day must have at least 60% valid measurements within the identified optimal window for an individual. Participants who did not meet this standard were removed from our

464   dataset. We provided a visual representation of the data processing and QC steps in **Suppl. Fig.**
465   **S2.3.**
466

Imputation
Missing values are still existent in the resulting QC-controlled time windows. To approach the
data missingness, we devised an imputation strategy for categorical and quantitative data
modalities, respectively. For the categorical data, we introduced a *'Not Recorded'* category into the
frame for imputation and subsequently applied label encoding. For the quantitative data, we used
the '*drift*' method from the sktime package (v0.19.1) with its default settings[78]. Recognizing that
these imputation strategies may not be adept at capturing non-polynomial dynamics, we further
included an indicator time series for each data modality:

476
$$T_{\text{indicator}}(i) = \mathbf{1}(T(i) = \text{Na})$$

where $\mathbf{1}(\cdot)$ is the indicator function and $T(i) = \text{Na}$ indicates the data at time step i is missed.
We concatenated the indicator time series with the imputed time series along the channel
dimension. The indicator time series serves as a mask that shows where imputations have been
made, while the imputed time series contains both the actual and imputed data. By including this
additional indicator time series, we are effectively providing the model with the flexibility to learn
an adaptive imputation strategy, where the model can learn how to treat imputed data points based
on the surrounding, non-imputed data.
484

485   **Machine Learning Classifier**
486

487   **Problem Formulation**
488   We first formulated the phenotype classification as a canonical machine learning task with
489   *manually engineered features*, which is outlined as follows. Given an input for a set of features,
490   X, *machine learning classification* (MLC) targets an output value y which represents the
491   macrophenotype of the subject:
492

493
$$X \in \mathcal{R}^{N \times d} \mapsto y$$

494   Here, N is the number of individuals and d is the number of features. Specifically, we chose the
495   curated *XGBoostRegessor* model implemented in xgboost package (v1.7.5) as our backbone ML
496   models, i.e.:

497
$$\text{XGBoostRegressor}(X) \mapsto y$$

XGBoost (eXtreme Gradient Boosting) has emerged as an effective machine learning framework,
noted for its optimized speed, scalability, and robustness[26]. As a variant of gradient boosted
decision trees, XGBoost is tailored for efficiency and demonstrates consistent performance across
diverse machine learning applications. Central to XGBoost is its adeptness at engineering trees
which pinpoint and rectify residuals from prior iterations, continually refining model accuracy. In
this work, we take advantage of the strengths that XGBoost offers, guided by a carefully crafted
set of features.
506

Feature Engineering
Our feature engineering for the XGBoost model is elaborated below. Specifically, the time-
invariant wearable features X_w were primarily derived from summary statistics of the time-series

wearable. We identified seven clusters of time-invariant wearable features from a total of 258
features. We further included curated covariates X_{cov} as additional features to supplement the time-
invariant wearables features. Covariates used for machine learning model include demographic
background (sex, race, age at second-year follow-up, divorced parents, parents' level of education,
parent income, adoption) family history of psychiatric illness (bipolar disorder, schizophrenia,
antisocial behavior, nervous breakdown, psychiatric treatment, hospital admission, suicide),
cognitive scores (flanker test, picmemory, process speed, reading score, stop reaction time, etc.),
and child behavioral checklist (CBCL internalizing and externalizing scores). Our complete
features set encompasses both static wearable features and covariates:

$$ X = \text{Concat}\left(\left\{ X_{\text{cov}}, X_w^1, X_w^2, \cdots, X_w^7 \right\}\right) $$

where $\text{Concat}(\cdot)$ denotes concatenation on the feature dimension. This enabled us to characterize
a nuanced interplay of wearable features with individual covariates, substantially accentuating the
power of our model.

Clustering of Wearable Static Features
We considered the 258 wearable static features in a subset of 2,410 ABCD individuals with
complete genetic, wearable and covariate information (see Methods section "Multivariate
wearable GWAS"). We computed Pearson's *r* correlation coefficients between all possible pairs
of features, and used these correlation values as distance measures to perform hierarchical
clustering (R function "hclust" & clustering method "Complete"). We also performed k-means
clustering of the correlation matrix by varying the number of clusters from two to twenty (R
function "kmeans", with nstart = 10), and chose an optimal number of seven clusters based on the
elbow curve of the total within-cluster sum of squares. A heatmap representation of the seven
clusters is shown in **Figure 2D**, and the list of static features for each cluster is provided in **Suppl.**
Tables S2.3-S2.4.

Class Balancing
Imbalanced training labels, where one class substantially outnumbered the other (e.g. 1,737 control
individuals versus 216 ADHD individuals), pose a substantial impact on the model performance.
To address this issue and ensure a more robust model, we implemented stochastic downsampling
techniques on classes with higher representation in each run of the model. To formalize this, we
assume two classes, A and B, where $|A|$ and $|B|$ represent the number of instances in each class.
Assuming $|A| \geq |B|$, we calculate the ratio r :

$$ r = \frac{|B|}{|A|} $$

546 We then randomly select a subset A' from A such that:

$$ |A'| = r \times |A| $$

549 The downsampled dataset will then consist of A' and B:

$$ \text{Downsampled Dataset} = A' \cup B $$

Multichannel Time Series Classifier

Problem Formulation

We formulate the phenotype classification as a multichannel time series classification problem which is described as follows. Given an input multichannel time series X:

$$X = \text{CONCAT}\{X_W, X_W^{\text{indicator}}, X_{\text{cov}}\},$$

where $X_W \in \mathcal{R}^{N \times c_w \times L}$, $X_W^{\text{indicator}} \in \{0, 1\}^{N \times c_w \times L}$, $X_{\text{cov}} \in \mathcal{R}^{N \times c_{\text{cov}} \times L}$. Here, N is the number of samples, , c_w the number of wearable modalities, L the number of measurements, X_W the multichannel wearables data, $X_W^{\text{indicator}}$ the multichannel indicator data (See section Imputation), and X_{cov} covariates (detail in next section). The *multi-channel time series classification* (MCTSC) targets an output value y which represents the macrophenotype of the subject:

$$X \in \mathcal{R}^{N \times C \times L} \mapsto y$$

$C = 2 \times c_w + c_{\text{cov}}$ is the number of input time-series channels. We further define a parameterized model which maps X to the output y:

$$f_\theta(X) \rightarrow y$$

f represents the mapping function, which is parameterized by θ. To optimize θ, we employed *cross-entropy loss with label smoothing* as the objective function, which is defined as:

$$\text{CE}(y, \hat{y}) := -\sum_{k=1}^{K} y_k \log(\hat{y_k})$$

where K denotes the number of classes. We employed a label smoothing regularizer to the ground truth label:

$$y_k^{\text{ls}} = y_k(1 - \alpha) + \frac{\alpha}{K}$$

Here, α is a smoothing parameter (we chose 0.1). This label smoothing technique helps to prevent the model from becoming too confident about the class labels, which could potentially bolster its generalization ability.

Covariate Integration

In order to integrate both covariates and time-series data for classification, we adapted the same covariates described in XGBoost feature engineering into a time-series format. Essentially, we transformed these variables into time-invariant sequences, where the value for each covariate remains the same at every time step. The transformed time-series covariates were then merged with wearable sensor data along the channel dimension. This approach allows the model to capture potential interactions between covariates and wearable measures, wherein the model can adjust its weights accordingly if a certain covariate influences the interpretation of the wearable data.

Xception Encoder

The XceptionTime encoder harnesses the power of one-dimensional convolutional neural networks (1d-CNNs) as its underlying architecture[27]. The model is structured with convolutional

597 filters of various sizes, which are sequentially followed by MaxPooling, Batch Normalization, and
598 ReLU activation functions, which form residual connections. Formally:

$$H_{\text{bottleneck}}^l = \text{Conv1D}(H^{l-1})$$
$$H_{\text{MaxConvPool}}^l = \text{Conv1D}(\text{MaxPool}(H^{l-1}))$$
$$\Delta H^l = \oplus \left\{ \oplus_k \left\{ \text{XceptionConv}_k(H_{\text{bottleneck}}^l) \right\}, H_{\text{MaxConvPool}}^l \right\}$$
$$\Delta H^l = \text{BatchNorm}(\Delta H^l)$$
$$\Delta H^l = \text{ReLU}(\Delta H^l)$$
$$H^l = H^{l-1} + \Delta H^l$$

Here, $\text{Conv1D}(\cdot)$ denotes 1d convolution, $\text{XceptionConv}(\cdot)$ represents depthwise
separable convolution, $\text{BatchNorm}(\cdot)$ represents Batch Normalization, $\text{ReLU}(\cdot)$ represents
ReLU activation function, and $\oplus$ aggregates feature maps from convolution filters of different
sizes. A visual representation of the model could be found in **Suppl. Fig. S2.4**. In summary, the
input feature maps are first projected to a bottleneck features map where the number of input
channels is much larger than the number of output channels. A sequential operation of max pooling
and 1d convolution is then performed on the input features maps to increase the expressivity of the
model. The variation in the size of Xception convolutional filters gives rise to multi-level receptive
fields, allowing the model to aggregate and process information at different levels of granularity
or resolution. Such a property is particularly advantageous when dealing with data from wearable
devices, as wearable data often exhibits both local patterns (i.e., minute-by-minute changes) and
global trends (i.e., hourly or daily rhythms).

The XceptionTime encoder introduces a modification to the vanilla 1d convolution model by
substituting the 1d convolution with a 1d depth-wise separable convolution. The operation can be
broken down into two steps:

$$\text{XceptionConv}(H^{l-1}) := \text{PointwiseConv}(\text{DepthwiseConv}(H^{l-1}))$$

In contrast to the traditional convolution operation, the depth-wise separable convolution first
applies a convolutional filter to each channel individually. This is followed by a 1x1 pointwise
convolution module, which performs a linear combination of the outputs across channels. This
process reduces the computational complexity of the model while still allowing for complex
feature extraction. These steps are described in detail below:

Depthwise Convolution: This applies a single filter to each input channel which can be expressed
as:

$$F_c^d = H_c * K_c^d$$

635 where F_c^d is the output feature map for channel c after the depthwise convolution, H_c is the input
636 feature map for channel c, and K_c^d is the depthwise filter (or kernel) for channel c. $*$ denotes the
637 convolution operation.

639 *Pointwise Convolution*: This operation combines the outputs from the depthwise convolution
640 across channels:

$$F^p = \oplus_c Y_c^d * K^p$$

642 where F^p is the output feature map after the pointwise convolution, Y_c^d is the input feature map
643 for channel c, K^p is the pointwise filter, which has a spatial dimension of 1x1 and operates across
644 all channels, and $\oplus$ is used to denote the aggregation of feature maps from all channels.
645
646 **Model Training and Evaluation**
647
Training Details
We split the dataset into 70% training set and 30% test set. We ran different experiments with 10
random seeds, and the final results were calculated as the mean of the 10 runs. This helps to
mitigate the risk of overfitting on a specific split, and provides a more robust estimate of the model
performance. We used ADAM as the optimizer for training, with $1\text{x}10^{-3}$ as the initial learning rate.
The neural network model is trained on an NVIDIA V100 32GB graphical processing unit using
the PyTorch and tsai deep learning libraries[79,80].
655
656 **Risk Scores**
657 In our study, we computed risk scores $\text{RS} \in \mathcal{R}^{N \times |\mathcal{K}|}$ by extracting the final layer of the deep
658 learning model, specifically the softmax probability. For the XGBoost model, we leverage the
659 pred_prob method implemented in the XGBoost library. Specifically:
660

$$\text{RS}(j) := P(y = j | x) = \frac{e^{f_j}}{\sum_{k \in \mathcal{K}} e^{f_k}}$$

661
where $f \in \mathcal{R}^{N \times K}$ is either the XceptionTime logits in the XceptionTime model, or the sum of
outputs from all trees in the XGBoostRegressor model. The softmax function, used in the final
layer of the deep learning model, returns probabilities for each category in a multi-class problem
that sum up to 1. Similarly, XGBoost's predict_proba method generates class probabilities as
output. These scores can serve as a measure of the 'risk' or likelihood associated with each class or
outcome. We utilized these risk scores as the response variable in our subsequent GWAS study
(see Methods section "Univariate Wearable GWAS"). This approach not only bridged the gap
between deep learning modeling and GWAS but also significantly enhanced the power of our
GWAS studies.
671
672 **Model Interpretability**
673
Ablation Method for Step and Feature Importance
The ablation method we present was used to measure the importance of features in a dataset.
Ablation methods are based on randomly rearranging the values of a feature or a group of features
across all subjects in the dataset, and then calculating an importance score based on the decrease
in a chosen metric. In our case, we utilized the *Area Under the Receiver Operating Characteristic*
curve (AUROC) as the metric to calculate this score. The rationale behind this method is that if a
feature is important for model predictions, shuffling the values of that feature should disrupt the
model's ability to make accurate predictions, leading to a decrease in the chosen performance
metric. The larger the decrease, the more important the feature is considered to be.
683
The ablation importance score can be applied to calculate both feature importance and step
importance. For step importance, the implementation is slightly different. Instead of shuffling
individual features, we shuffled the values within selected windows of the time series. The time

series was divided into windows of a chosen length (in our case, 1 hour), and these windows were
then shuffled across all subjects, allowing us to assess the importance of information at different
time steps or periods. If the model performance significantly decreases when the values within a
certain time window are shuffled, the information within that time window is important for the
model predictions.
692
693 **Grad-CAM**
694 The weighted class activation mapping (CAM) method is a well-established technique for
695 examining how a trained model makes its predictions[29]. In the context of time-series data, it can
696 highlight which time steps are particularly influential in the model's decision-making process.
697
We first computed the gradient of the score for the predicted class y with respect to the feature
map of the first layer activations $A_0^{c,i}$ of a convolutional layer. This gradient, denoted as $\frac{\partial y}{\partial A_0^{c,i}}$,
provides a measure of how a small change in the activation of the convolutional layer could affect
the final prediction of the model. To convert these gradients into a measure of importance for each
channel (indexed by c), we employed a global average pooling, which calculates the average of all
gradients across the sequence length (indexed by i). This resulted in a set of channel-wise gradient
averages, denoted as α_c. Mathematically, this is expressed as:
705

$$\alpha_c := \frac{1}{Z} \sum_{i=1}^{L} \frac{\partial y}{\partial A_0^{c,i}}$$

706
707 where Z is a normalization constant, typically the total number of elements in the layer, and L is
708 the length of the sequence.
709
710 We next generated the Gradient-weighted Class Activation Mapping (Grad-CAM). This is a visual
711 representation of the importance of each time step for the model's predictions. The Grad-CAM,
712 denoted as L_{GM}^i, is defined as:
713

$$L_{GM}^i = \mathrm{RELU}\left(\sum_c \alpha_c A_0^{c,i}\right)$$

714
715 where the ReLU (Rectified Linear Unit) function is used to ensure that only features with a positive
716 influence on the class of interest result in high activation. Essentially, this means that only the time
717 steps that positively contribute to the model's decision will have high importance scores.
718
719 Finally, for each time step, we computed the average Grad-CAM scores across the entire test set.
720 This allowed us to determine which time steps in the input data were most influential in the model
721 predictions.
722
723 **Genome-wide Association Studies (GWAS)**
724
Quality Control of Genetic Data
We obtained genotyped and imputed genetic data for 11,099 individuals as part of the ABCD Data
Release 3 (https://abcdstudy.org/scientists/data-sharing/). We used the genotyped data to infer

population stratification and the imputed data to perform the different GWAS described below
(see Methods sections "Univariate wearable GWAS" and "Multivariate wearable GWAS").
We applied a quality control (QC) protocol on the genotyped data[81]. Specifically, we performed
the QC steps described in
https://github.com/MareesAT/GWA_tutorial/blob/master/1_QC_GWAS.zip (file
"1_Main_script_QC_GWAS.txt") using PLINK v1.90b6.21[82]. Briefly, of the initial set of 516,598
variants, we kept those with a missingness rate across individuals < 0.02 (n = 481,920). Of the
initial set of 11,099 individuals, we kept those with a missingness rate across variants < 0.02 (n =
10,660). Next, we considered only variants located on autosomal chromosomes (n = 470,076),
those with Minor Allele Frequency (MAF) > 0.01 (n = 427,704), and those that did not deviate
from Hardy-Weinberg equilibrium (p value $\geq 10^{-10}$; n = 370,002). These variants were pruned to
a final set of 156,556 variants (window size = 50; number of variants to shift the window at each
step = 5; multiple correlation coefficient 0.2). We computed the heterozygosity rate for each
individual using the pruned set of variants, and kept individuals with a heterozygosity rate
deviating less than 3 standard deviations from the mean (n = 10,467). We also used pruned variants
to assess cryptic relatedness by identifying groups of individuals with Proportion Identity-By-
Descent (pi_hat) > 0.2. For every group of related individuals, we then selected the individual with
the lowest variant missingness rate, leaving a total of 8,816 individuals. We used PLINK2 to
perform a Principal Component Analysis (PCA) on the 156,556 pruned genotyped variants from
the 8,816 selected individuals. We integrated the PCA results with the ethnicity score group
information provided by the ABCD metadata, which was available for 8,791 individuals (**Fig. 1D**
and **Suppl. Fig. S4.1-S4.4**).
750
751 We filtered the imputed genetic variants for MAF > 0.01 and estimated imputation accuracy (R^2) >
752 0.3, and obtained a final set of 11,954,686 variants for the GWAS analysis (**Suppl. Fig. S4.5**). We
753 also computed distributions of R^2 for all (genotyped and imputed) variants, and of empirical leave-
754 one-out R^2 (ER^2) for genotyped variants (**Suppl. Fig. S4.6**).
755
Covariates included in the GWAS
We considered five different groups of covariates: basic (sex, age at second-year follow-up, first
five genotype PCs), behavioral (CBCL internalizing and externalizing scores, DSM internalizing
and externalizing scores), family history of psychiatric illness (bipolar disorder, schizophrenia,
antisocial behavior, nervous breakdown, psychiatric treatment, hospital admission, suicide),
family situation (divorced parents, parents' level of education, family income, adoption), and other
(ACS raked propensity score, DNA extraction batch). 3,579 of the previously selected 8,791
individuals reported complete information for these 24 covariates.
764
Univariate Wearable GWAS
For this analysis, we focused on a subset of 1,191 individuals that were either diagnosed with
ADHD (n = 137) or belonged to the non-clinical control group (n = 1,054). We performed a GWAS
testing for association between genetic variants and ADHD diagnosis, encoded as a binary
outcome (ADHD = 1, control = 0; univariate binary GWAS; **Figure 4A**). We also obtained, for
each individual, ten different ADHD risk scores based on the XGBoost and Xception predictive
models (see Methods section "Risk Scores"). Specifically, we used risk scores from the following
six models: baseline model using CBCL externalizing score ("CBCL ext."); baseline model using
CBCL internalizing score ("CBCL int."); XGBoost model using wearable features ("XGB");

XGBoost model using wearable features and CBCL scores ("XGB + CBCL"); Xception model
using wearable features ("Xception"); and Xception model using wearable features and CBCL
scores ("Xception + CBCL"). For models "XGB", "XGB + CBCL", "Xception" and "Xception +
CBCL", we also implemented the "liability-CC" trait methodology[35]. This methodology consists
of converting the predictive modeling risk score of the cases (individuals with ADHD) to a value
of 1, while keeping the original risk scores for the controls. These four additional types of scores
are labeled as "XGB v2", "XGB + CBCL v2", "Xception v2" and "Xception + CBCL v2".

We performed ten different GWAS to test for associations between genetic variants and each of
these ten scores (univariate continuous GWAS; **Figure 4B**). We used PLINK2 to perform both
binary and continuous univariate GWAS, and the FUMA platform for loci definition (reference
panel population: "1000G Phase3 ALL")[83,84]. We first ran all GWASs using only the set of basic
covariates (sex, age, first five population structure PCs), as these were also used in previous GWAS
for ADHD[85,86]. These results and the corresponding p value quantile-quantile plots are shown in
Figure 4A-B, **Table 1** (GWAS Method: "univariate continuous"), **Suppl. Fig. S4.7-S4.8** ("Basic
covariates"), and **Suppl. Table S4.1**. We also repeated both continuous and binary GWAS to
include all covariates described in Methods section "Covariates included in the GWAS", apart
from the CBCL and DSM behavioral scores, which were employed as features for the predictive
models that generated the risk scores (**Suppl. Fig. S4.7** and **S4.9** - "All covariates").

Statistical Power of Binary *vs.* Continuous Traits
To compare the statistical power of genetic association testing using binary and continuous traits,
we simulated a cohort of 1,500 individuals. In each individual i, we generated biallelic SNPs with
a binomial model (i.e., the genotype at each SNP followed a binomial distribution, with the number
of trials equal to 2 and probability of success on each trial equal to a given MAF). We chose the
cohort size to approximate the number of individuals ($n = 1{,}191$) in the univariate GWAS for
ADHD described above (Methods section "Univariate wearable GWAS"). For each individual i,
we then simulated a continuous trait (C_i) as the sum of the genotype effect (b) at a given SNP with
genotype x_i (0, 1, or 2) and random noise (e_i):

$$C_i = x_i \cdot b + e_i$$
$$\text{where}$$
$$\mathbf{b} \sim U(0,1)$$
$$\mathbf{e} \sim N(0,1)$$

810   We also simulated a binary trait for each individual i (B_i), following

$$B_i = 1 \text{ if } C_i > \text{median}(C), \text{ otherwise } 0$$

where C is the vector of simulated continuous traits for the entire cohort.

For a particular genotype effect b, we ran 10,000 simulations. Under this scenario, we estimated
the power of the simulated continuous and binary traits as the fraction of significant (i.e.
Benjamini-Hochberg adjusted p value < 0.05) linear and logistic regression tests, respectively. We
employed linear and logistic regression as implemented in the R functions "lm" (library "stats")

and "glm" (family = binomial; library "MASS"), respectively. Overall, we simulated 50 different
values of b across nine different MAFs (**Suppl. Fig. S4.11**).
822
Multivariate Wearable GWAS
This second type of GWAS consists in testing the association between genetic variants and a multi-
trait (multivariate) phenotype. In this case, we define the multivariate phenotype as the vector of
static summary features obtained from an individual's wearable device (see Methods section
"Clustering of Wearable Static Features"). For this analysis, we considered all individuals with
complete genetic, wearable, and covariate data independently of their diagnosis.
829
We first conducted a wearable GWAS using the 14 static features related to heart rate as the
multivariate response variable, which were available for 3,256 individuals (features: InterdayCV,
InterdaySD, IntradayCV_mean, IntradayCV_median, IntradayCV_sd, IntradayMean_mean,
IntradayMean_median, IntradayMean_sd, IntradaySD_mean, IntradaySD_median,
IntradaySD_sd, Mean, Median, STD). We next aimed to include all 258 static features, which were
available for 2,410 individuals, and applied two different strategies to reduce the dimensionality
of the multivariate response. In one case, we performed a PCA of the individuals based on their
values for the 258 features, and used the first five PCs as the multivariate response. In the second
case, we considered each of the seven clusters of static features as a separate multivariate response,
and performed a GWAS for each cluster (see also Methods section "Clusters of static summary
features" below). Therefore, we ran a total of nine multivariate GWAS (one for heart rate features,
one for the first five PCs of all features, and one for each of the seven clusters of features). For all
multivariate GWAS, we defined a model that included the genotype and 24 covariates as
independent variables (see Methods section "Covariates included in the GWAS").
844
We used the Multivariate Asymptotic Non-parametric Test of Association R package (MANTA,
https://github.com/dgarrimar/manta) to test for association between genetic variants and the
multivariate wearable trait, and performed all the analyses within a containerized Nextflow
pipeline, available at https://github.com/dgarrimar/mvgwas-nf[44,87]. Since MANTA is a non-
parametric method, normalization of the GWAS traits was not required. After performing the
different GWAS runs, we used FUMA for loci definition (reference panel population: "1000G
Phase3 ALL")[84]. These results and the corresponding p value quantile-quantile plots are shown in
Figure 4C, **Table 1** (GWAS Method: "multivariate continuous"), **Suppl. Fig. S4.12**, and **Suppl.**
Table S4.3. As MANTA p values do not come from a normal distribution, we employed λ_X
(instead of the commonly used λ_G) to estimate the genomic inflation factor[88].
855
Genome-wide *vs.* Study-wide Significance
We selected the conventional genome-wide significant p value threshold of $5 \cdot 10^{-8}$ to identify
significant loci from all GWAS runs. However, in line with previous GWAS studies, we also
considered a study-wide significance threshold to account for the fact that multiple GWAS were
performed[89]. In our case, the study-wide significant thresholds are $5 \cdot 10^{-9}$ ($5 \cdot 10^{-8}$ / 10 GWAS runs)
for the univariate continuous GWAS for ADHD, and $5.56 \cdot 10^{-9}$ ($5 \cdot 10^{-8}$ / 9 GWAS runs) for the
multivariate wearable GWAS. Based on these thresholds, one locus from the ADHD GWAS and
nine loci from the wearable GWAS would pass the study-wide significance threshold. Similar to
other GWAS, we also considered a suggestive p value threshold of $1 \cdot 10^{-5}$ (**Figure 4** and **Suppl.**
Fig. S4.7)[89].

Chromosome X
Because the multivariate GWAS (specifically in runs corresponding to feature clusters 2 and 7) reported a large number of significant loci on chromosome X (**Table 1** and **Suppl. Table S4.3**), we implemented additional quality controls to account for potential bias in chromosome X variants. First, we tested whether chromosome X variants showed systematically lower p values compared to variants located on autosomal chromosomes. To do this, we computed the proportion of variants with p value $< 10^{-4}$ from each autosomal chromosome, and performed a one-sided Fisher's exact test to evaluate whether this proportion was significantly lower compared to variants on chromosome X for the same GWAS run. We found that this was the case only in GWAS runs for clusters 2, 4, and 7 (**Suppl. Fig. S4.13**). We reasoned that if an unknown systematic bias related to chromosome X was truly present (e.g., genotyping issues), we would observe the same situation for every cluster of features. Given that we did not identify significant loci on chromosome X for cluster 4, this analysis ruled out any unaccounted systematic bias related to chromosome X, and confirms that wearable features in clusters 2 and 7 indeed show stronger association with variants located on chromosome X. We also checked for imputation bias in chromosome X variants that could systematically differentiate female and male individuals. To do this, we performed a PCA of the 2,410 individuals based on their imputed genotypes at chromosome X variants, and did not observe a separation between female and male individuals (**Suppl. Fig. S4.14**).

Neuropsychiatry-related Proximal Genes and eGenes
For each genome-wide significant locus, we retrieved the ten closest genes when considering a window of $\pm$ 250 Kb from the center of the locus, using the GENCODE human genome annotation version 41[90]. Next, we labeled as "neuropsychiatric-related" those proximal genes that are associated with psychiatric disorders according to OpenTargets (https://platform.opentargets.org/) (**Suppl. Tables S4.1** and **S4.3**)[91]. We further intersected our catalog of genome-wide significant loci with previous eQTL catalogs using BEDTools intersect (v2.30.0), and identified a subset of proximal neuropsychiatric-related genes with eQTLs overlapping our list of loci[46,92-95]. We labeled these genes as "neuropsychiatric-related proximal eGenes" (**Suppl. Tables S4.1** and **S4.3**).

Chromatin Dissection of locus chr14:23392601-23418974
We first performed an exploratory analysis by intersecting our two lists of significant loci with the ENCODE4 registry of candidate cis-regulatory elements (cCREs)(https://www.encodeproject.org/search/?type=Annotation&encyclopedia_version=current&annotation_type=candidate+Cis-Regulatory+Elements&annotation_type=chromatin+state&annotation_type=representative+DNase+hypersensitivity+sites&status=released&encyclopedia_version=ENCODE+v4) (**Suppl. Tables S4.1 and S4.3**)[45]. Given the documented role of locus chr14:23392601-23418974 in heart-related traits and diseases, we next evaluated the enrichment of heart-specific epigenetic features (nucleosome positioning, histone modifications, and transcription factor (TF) binding) at this locus. We downloaded peak calling files for DNase-seq, ATAC-seq, ChIP-seq (histone marks & TFs) and Mint-ChIP-seq for histone marks available for human biosamples from the ENCODE portal (https://www.encodeproject.org/metadata/?control_type%21=%2A&status=released&perturbed=false&assay_title=Histone+ChIP-seq&assay_title=TF+ChIP-seq&assay_title=DNase-seq&assay_title=ATAC-seq&assay_title=Mint-ChIP-seq&files.file_type=bigBed+narrowPeak&replicates.library.biosample.donor.organism.scientific

[_name=Homo+sapiens&type=Experiment&files.analyses.status=released&files.preferred_defaul](https://www.encodeproject.org/)
[t=true](https://www.encodeproject.org/); access date: 09/27/2022)[45,96]. We then grouped human biosamples based on their
"biosample   ontology   organ   slim"
(https://www.encodeproject.org/report/?type=Experiment&control_type!=*&status=released&pe
[rturbed=false&assay_title=TF+ChIP-seq&assay_title=Histone+ChIP-seq&assay_title=DNase-](https://www.encodeproject.org/)
[seq&assay_title=ATAC-seq&assay_title=Mint-ChIP-](https://www.encodeproject.org/)
[seq&replicates.library.biosample.donor.organism.scientific_name=Homo+sapiens&field=biosam](https://www.encodeproject.org/)
[ple_ontology.organ_slims&field=biosample_ontology.cell_slims&field=biosample_ontology.sys](https://www.encodeproject.org/)
[tem_slims&field=%40id&field=biosample_ontology.term_name](https://www.encodeproject.org/)). To test the tissue-specific
enrichment of chromatin features in a particular organ, we computed the number of times any of
the five significant variants at the locus overlapped a peak from experiments in that organ
compared to all other organs (two-sided Fisher's exact test, Benjamini-Hochberg adjusted p value
< 0.1). For this analysis, we counted only once those overlaps involving variants that are < 100 bp
apart.
926
Exploring the Genetic-behavioral-psychiatric Axis
The multivariate wearable GWAS allowed us to first perform an exploratory analysis to identify
genetic variants associated with any of the seven clusters of wearable-derived features (genome-
wide significant p value < $5 \cdot 10^{-8}$). To identify the specific features that are driving the significant
association between the cluster and the variant, we next performed unpaired Wilcoxon rank tests
between all three groups of genotype individuals (i.e., AA $vs.$ AG, AA $vs.$ GG, and AG $vs.$ GG)
for each feature within a particular cluster. We then selected features where at least one of the three
Wilcoxon tests reported a Benjamini-Hochberg-adjusted p value < 0.1, and showed examples for
three SNPs in **Figure 5A-C** (left panel). For each of these three examples, we next evaluated the
enrichment of the minor allele (in all three cases the G allele) in individuals within a specific
psychiatric cohort $vs.$ non-clinical control individuals (two-sided Fisher exact test; **Figure 5A-C**
right panel). Given the reduced number of individuals with GG genotype for SNPs rs113525298
and rs147959551 (15 and 14, respectively), in these two cases the enrichment of the minor allele
was tested by merging individuals with AG and GG genotypes. For SNP rs365990, the enrichment
was computed only on individuals with GG genotype. For all tests, we required at least one
individual to be present in every cell of the 2×2 matrix employed for the Fisher's exact test ($a =$
n individuals with minor allele AND part of the psychiatric cohort; $b = n$ individuals without minor
allele AND part of the psychiatric cohort; $c = n$ individuals with minor allele AND part of the
healthy controls; $d = n$ individuals without minor allele AND not part of the healthy controls).
946
Intersection of genome-wide significant loci with the GWAS Catalog
To assess the clinical relevance of our GWAS loci, we intersected them with variants from the
NHGRI-EBI GWAS catalog (https://www.ebi.ac.uk/gwas/; access date: 05/16/2023). For the loci
identified by our ADHD GWAS, we considered overlaps with brain- or neuropsychiatric-related
GWAS hits (**Figure 4B**). Because our wearable-derived features are mostly related to heart, sleep,
metabolism and physical activity, for the wearable GWAS loci we considered any overlaps with
heart-, sleep-, metabolism- and physical activity-related GWAS hits. Additionally, given the
presence of individuals with psychiatric disorders in the wearable GWAS cohort, we also
considered intersections with brain- or neuropsychiatric-related GWAS hits (**Figure 4C**). We
acknowledge that colocalization analysis would be the most appropriate way to compute these
intersections, and we performed this analysis for ADHD GWAS loci (see Methods section

"Colocalization analysis"). However, MANTA does not provide estimates of variant effect sizes
that can be directly employed in co-localization analysis. For this reason, we evaluated the strength
of these intersections against a null distribution. Specifically, we computed the proportion of
GWAS variants associated with a particular trait that overlap our significant loci, and compared it
to the proportions observed across 10,000 random sets of genomic loci with the same size and
chromosome location. We report the percentile of our GWAS enrichments compared to the null
distribution in **Suppl. Fig. S4.10** (univariate GWAS for ADHD) and **S4.15** (multivariate GWAS).
965
Colocalization Analysis
We performed colocalization analysis using the R package *coloc* (function coloc.abf, default
parameters) on the results obtained from the univariate continuous GWAS[97]. Specifically, we
focused on two of the seven overlapping brain-related traits with available GWAS summary
statistics (**Figure 4B** and **Suppl. Table S4.2**), and tested the hypothesis of signal co-localization
between our ADHD risk scores at the intersecting loci. Locus chr17:32256997:32283356 reported
a posterior probability of 0.99 for signal co-localization with a locus previously associated with
chronotype measurement[98]. We also tested locus chr7:68219282:68338849 (suggestive
association at p value $< 10^{-5}$) for co-localization with a previously reported locus for ADHD[85]. In
this case, given that the two traits being tested are the same, we set all three parameters p1, p2 and
p12 equal to $1 \cdot 10^{-5}$, and reported a posterior probability of 0.25.
977
978 **Code Availability**
979 The code for the paper is publicly available at https://github.com/gersteinlab/ABCD.
980
Data and Materials Availability

The data used in this study is available through the NIMH ABCD NDA portal
(https://nda.nih.gov/general-query.html?q=query=featured-
datasets:Adolescent%20Brain%20Cognitive%20Development%20Study%20(ABCD)). Data
used in the preparation of this article were obtained from the Adolescent Brain Cognitive
DevelopmentSM (ABCD) Study (https://abcdstudy.org), held in the NIMH Data Archive (NDA).
This is a multisite, longitudinal study designed to recruit more than 10,000 children age 9-10 and
follow them over 10 years into early adulthood. The ABCD Study® is supported by the National
Institutes of Health and additional federal partners under award numbers U01DA041048,
U01DA050989, U01DA051016, U01DA041022, U01DA051018, U01DA051037,
U01DA050987, U01DA041174, U01DA041106, U01DA041117, U01DA041028,
U01DA041134, U01DA050988, U01DA051039, U01DA041156, U01DA041025,
U01DA041120, U01DA051038, U01DA041148, U01DA041093, U01DA041089,
U24DA041123, U24DA041147. A full list of supporters is available at
https://abcdstudy.org/federal-partners.html. A listing of participating sites and a complete listing
of the study investigators can be found at https://abcdstudy.org/consortium_members/. ABCD
consortium investigators designed and implemented the study and/or provided data but did not
necessarily participate in the analysis or writing of this report. This manuscript reflects the views
of the authors and may not reflect the opinions or views of the NIH or ABCD consortium
investigators.

References

Zablotsky, B., Terlizzi, E. P. & National Center for Health Statistics (U.S.). in *NCHS data brief,*   1 online resource (7 pages, 1 unnumbered page).

UNICEF. *Impact of COVID-19 on poor mental health in children and young people 'tip of the iceberg' – UNICEF*, <https://www.unicef.org/philippines/press-releases/impact-covid-19-poor-mental-health-children-and-young-people-tip-iceberg-unicef#:~:text=According%20to%20the%20latest%20available,needs%20and%20mental%20health%20funding> (2021).

CDC. *Data and Statistics on Children's Mental Health*, <https://www.cdc.gov/childrensmentalhealth/data.html> (2023).

McGorry, P. D. & Nelson, B. Transdiagnostic psychiatry: premature closure on a crucial pathway to clinical utility for psychiatric diagnosis. *World Psychiatry* **18**, 359-360 (2019). https://doi.org/10.1002/wps.20679

Hartmann, J. A. *et al.* Pluripotential Risk and Clinical Staging: Theoretical Considerations and Preliminary Data From a Transdiagnostic Risk Identification Approach. *Front Psychiatry* **11**, 553578 (2020). https://doi.org/10.3389/fpsyt.2020.553578

Insel, T. *et al.* Research Domain Criteria (RDoC): Toward a New Classification Framework for Research on Mental Disorders. *Am J Psychiat* **167**, 748-751 (2010). https://doi.org/10.1176/appi.ajp.2010.09091379

Flint, J., Timpson, N. & Munafo, M. Assessing the utility of intermediate phenotypes for genetic mapping of psychiatric disease. *Trends Neurosci* **37**, 733-741 (2014). https://doi.org/10.1016/j.tins.2014.08.007

Leuchter, A. F., Hunter, A. M., Krantz, D. E. & Cook, I. A. Intermediate phenotypes and biomarkers of treatment outcome in major depressive disorder. *Dialogues Clin Neurosci* **16**, 525-537 (2014). https://doi.org/10.31887/DCNS.2014.16.4/aleuchter

Sanchez-Roige, S. & Palmer, A. A. Emerging phenotyping strategies will advance our understanding of psychiatric genetics. *Nat Neurosci* **23**, 475-480 (2020). https://doi.org/10.1038/s41593-020-0609-7

Meyer-Lindenberg, A. & Weinberger, D. R. Intermediate phenotypes and genetic mechanisms of psychiatric disorders. *Nat Rev Neurosci* **7**, 818-827 (2006). https://doi.org/10.1038/nrn1993

Meyer-Lindenberg, A. Intermediate or brainless phenotypes for psychiatric research? *Psychol Med* **40**, 1057-1062 (2010). https://doi.org/10.1017/s0033291709991929

Koppe, G., Guloksuz, S., Reininghaus, U. & Durstewitz, D. Recurrent Neural Networks in Mobile Sampling and Intervention. *Schizophr Bull* **45**, 272-276 (2019). https://doi.org/10.1093/schbul/sby171

Gomes, N., Pato, M., Lourenco, A. R. & Datia, N. A Survey on Wearable Sensors for Mental Health Monitoring. *Sensors (Basel)* **23** (2023). https://doi.org/10.3390/s23031330

Dunn, J. *et al.* Wearable sensors enable personalized predictions of clinical laboratory measurements. *Nat Med* **27**, 1105-1112 (2021). https://doi.org/10.1038/s41591-021-01339-0

Insel, T. R. Digital Phenotyping: Technology for a New Science of Behavior. *JAMA* **318**, 1215-1216 (2017). https://doi.org/10.1001/jama.2017.11295

Cronbach, L. J. & Meehl, P. E. Construct validity in psychological tests. *Psychol Bull* **52**, 281-302 (1955). https://doi.org/10.1037/h0040957

Jablensky, A. Psychiatric classifications: validity and utility. *World Psychiatry* **15**, 26-31 (2016). https://doi.org/10.1002/wps.20284

Sen, B., Borle, N. C., Greiner, R. & Brown, M. R. G. A general prediction model for the detection of ADHD and Autism using structural and functional MRI. *PLoS One* **13**, e0194856 (2018). https://doi.org/10.1371/journal.pone.0194856

Kim, W. P. *et al.* Machine Learning-Based Prediction of Attention-Deficit/Hyperactivity Disorder and Sleep Problems With Wearable Data in Children. *JAMA Netw Open* **6**, e233502 (2023). https://doi.org/10.1001/jamanetworkopen.2023.3502

van Dijk, M. T., Murphy, E., Posner, J. E., Talati, A. & Weissman, M. M. Association of Multigenerational Family History of Depression With Lifetime Depressive and Other Psychiatric Disorders in Children: Results from the Adolescent Brain Cognitive Development (ABCD) Study. *JAMA Psychiatry* **78**, 778-787 (2021). https://doi.org/10.1001/jamapsychiatry.2021.0350

Anttila, V. *et al.* Analysis of shared heritability in common disorders of the brain. *Science* **360**, eaap8757 (2018). https://doi.org/10.1126/science.aap8757

Mishra, T. *et al.* Pre-symptomatic detection of COVID-19 from smartwatch data. *Nat Biomed Eng* **4**, 1208-1220 (2020). https://doi.org/10.1038/s41551-020-00640-6

Liu, J. *et al.* Bayesian structural time series for biomedical sensor data: A flexible modeling framework for evaluating interventions. *PLoS Comput Biol* **17**, e1009303 (2021). https://doi.org/10.1371/journal.pcbi.1009303

Bent, B. *et al.* The digital biomarker discovery pipeline: An open-source software platform for the development of digital biomarkers using mHealth and wearables data. *J Clin Transl Sci* **5**, e19 (2020). https://doi.org/10.1017/cts.2020.511

Salari, N. *et al.* The global prevalence of ADHD in children and adolescents: a systematic review and meta-analysis. *Ital J Pediatr* **49**, 48 (2023). https://doi.org/10.1186/s13052-023-01456-1

Chen, T. Q. & Guestrin, C. XGBoost: A Scalable Tree Boosting System. *Kdd'16: Proceedings of the 22nd Acm Sigkdd International Conference on Knowledge Discovery and Data Mining*, 785-794 (2016). https://doi.org/10.1145/2939672.2939785

Chollet, F. Xception: Deep Learning with Depthwise Separable Convolutions. *Proc Cvpr Ieee*, 1800-1807 (2017). https://doi.org/10.1109/Cvpr.2017.195

Kofler, M. J. *et al.* Working memory and information processing in ADHD: Evidence for directionality of effects. *Neuropsychology* **34**, 127-143 (2020). https://doi.org/10.1037/neu0000598

Selvaraju, R. R. *et al.* Grad-CAM: Visual Explanations from Deep Networks via Gradient-based Localization. *Ieee I Conf Comp Vis*, 618-626 (2017). https://doi.org/10.1109/Iccv.2017.74

Imeraj, L. *et al.* Time-of-day effects in arousal: disrupted diurnal cortisol profiles in children with ADHD. *J Child Psychol Psychiatry* **53**, 782-789 (2012). https://doi.org/10.1111/j.1469-7610.2012.02526.x

Alvaro, P. K., Roberts, R. M. & Harris, J. K. A Systematic Review Assessing Bidirectionality between Sleep Disturbances, Anxiety, and Depression. *Sleep* **36**, 1059-1068 (2013). https://doi.org/10.5665/sleep.2810

Grimm, O., Kranz, T. M. & Reif, A. Genetics of ADHD: What Should the Clinician Know? *Curr Psychiatry Rep* **22**, 18 (2020). https://doi.org/10.1007/s11920-020-1141-x

Purves, K. L. *et al.* A major role for common genetic variation in anxiety disorders. *Mol Psychiatry* **25**, 3292-3303 (2020). https://doi.org/10.1038/s41380-019-0559-1

Yang, L., Wang, S. & Altman, R. B. POPDx: an automated framework for patient phenotyping across 392 246 individuals in the UK Biobank study. *J Am Med Inform Assoc* **30**, 245-255 (2023). https://doi.org/10.1093/jamia/ocac226

Yang, L., Sadler, M. C. & Altman, R. B. Genetic association studies using disease liabilities from deep neural networks. *medRxiv* (2023). https://doi.org/10.1101/2023.01.18.23284383

Chen, Y. C. *et al.* Neuroanatomic, epigenetic and genetic differences in monozygotic twins discordant for attention deficit hyperactivity disorder. *Mol Psychiatry* **23**, 683-690 (2018). https://doi.org/10.1038/mp.2017.45

Karlsson Linner, R. *et al.* Multivariate analysis of 1.5 million people identifies genetic associations with traits related to self-regulation and addiction. *Nat Neurosci* **24**, 1367-1376 (2021). https://doi.org/10.1038/s41593-021-00908-3

Fan, Z. *et al.* DLGAP1 and NMDA receptor-associated postsynaptic density protein genes influence executive function in attention deficit hyperactivity disorder. *Brain Behav* **8**, e00914 (2018). https://doi.org/10.1002/brb3.914

Royston, P., Altman, D. G. & Sauerbrei, W. Dichotomizing continuous predictors in multiple regression: a bad idea. *Stat Med* **25**, 127-141 (2006). https://doi.org/10.1002/sim.2331

Breitling, L. P. & Brenner, H. Odd odds interactions introduced through dichotomisation of continuous outcomes. *J Epidemiol Community Health* **64**, 300-303 (2010). https://doi.org/10.1136/jech.2009.089458

Schmitz, S., Adams, R. & Walsh, C. The use of continuous data versus binary data in MTC models: a case study in rheumatoid arthritis. *BMC Med Res Methodol* **12**, 167 (2012). https://doi.org/10.1186/1471-2288-12-167

Schandry, R. Heart beat perception and emotional experience. *Psychophysiology* **18**, 483-488 (1981). https://doi.org/10.1111/j.1469-8986.1981.tb02486.x

Hsueh, B. *et al.* Cardiogenic control of affective behavioural state. *Nature* **615**, 292-299 (2023). https://doi.org/10.1038/s41586-023-05748-8

Garrido-Martin, D., Calvo, M., Reverter, F. & Guigo, R. A fast non-parametric test of association for multiple traits. *Genome Biol* **24**, 230 (2023). https://doi.org/10.1186/s13059-023-03076-8

Consortium, E. P. *et al.* Expanded encyclopaedias of DNA elements in the human and mouse genomes. *Nature* **583**, 699-710 (2020). https://doi.org/10.1038/s41586-020-2493-4

Consortium, G. T. The GTEx Consortium atlas of genetic regulatory effects across human tissues. *Science* **369**, 1318-1330 (2020). https://doi.org/10.1126/science.aaz1776

Landrum, M. J. *et al.* ClinVar: improvements to accessing data. *Nucleic Acids Res* **48**, D835-D844 (2020). https://doi.org/10.1093/nar/gkz972

Fu, T. *et al.* Association of the MYH6 Gene Polymorphism with the Risk of Atrial Fibrillation and Warfarin Anticoagulation Therapy. *Genet Test Mol Biomarkers* **25**, 590-599 (2021). https://doi.org/10.1089/gtmb.2021.0025

Rangaraju, A. *et al.* Genetic variants in post myocardial infarction patients presenting with electrical storm of unstable ventricular tachycardia. *Indian Pacing Electrophysiol J* **18**, 91-94 (2018). https://doi.org/10.1016/j.ipej.2018.01.003

Eijgelsheim, M. *et al.* Genome-wide association analysis identifies multiple loci related to resting heart rate. *Hum Mol Genet* **19**, 3885-3894 (2010). https://doi.org/10.1093/hmg/ddq303

Faurholt-Jepsen, M., Kessing, L. V. & Munkholm, K. Heart rate variability in bipolar disorder: A systematic review and meta-analysis. *Neurosci Biobehav Rev* **73**, 68-80 (2017). https://doi.org/10.1016/j.neubiorev.2016.12.007

Carr, O., de Vos, M. & Saunders, K. E. A. Heart rate variability in bipolar disorder and borderline personality disorder: a clinical review. *Evid Based Ment Health* **21**, 23-30 (2018). https://doi.org/10.1136/eb-2017-102760

Stautland, A. *et al.* Heart rate variability as biomarker for bipolar disorder. *medRxiv*, 2022.2002.2014.22269413 (2022). https://doi.org/10.1101/2022.02.14.22269413

Dieckmann, L., Cole, S. & Kumsta, R. Stress genomics revisited: gene co-expression analysis identifies molecular signatures associated with childhood adversity. *Transl Psychiatry* **10**, 34 (2020). https://doi.org/10.1038/s41398-020-0730-0

Hesdorffer, D. C. *et al.* ADHD as a risk factor for incident unprovoked seizures and epilepsy in children. *Arch Gen Psychiatry* **61**, 731-736 (2004). https://doi.org/10.1001/archpsyc.61.7.731

Dolan, J. & Mitchell, K. J. Mutation of Elfn1 in mice causes seizures and hyperactivity. *PLoS One* **8**, e80491 (2013). https://doi.org/10.1371/journal.pone.0080491

Chawner, S. *et al.* Sleep disturbance as a transdiagnostic marker of psychiatric risk in children with neurodevelopmental risk genetic conditions. *Transl Psychiatry* **13**, 7 (2023). https://doi.org/10.1038/s41398-022-02296-z

Arns, M., Kooij, J. J. S. & Coogan, A. N. Review: Identification and Management of Circadian Rhythm Sleep Disorders as a Transdiagnostic Feature in Child and Adolescent Psychiatry. *J Am Acad Child Adolesc Psychiatry* **60**, 1085-1095 (2021). https://doi.org/10.1016/j.jaac.2020.12.035

Wang, Z., Zhou, X., Gui, Y., Liu, M. & Lu, H. Multiple measurement analysis of resting-state fMRI for ADHD classification in adolescent brain from the ABCD study. *Transl Psychiatry* **13**, 45 (2023). https://doi.org/10.1038/s41398-023-02309-5

Obermeyer, Z., Powers, B., Vogeli, C. & Mullainathan, S. Dissecting racial bias in an algorithm used to manage the health of populations. *Science* **366**, 447-453 (2019). https://doi.org/10.1126/science.aax2342

Dane, A. V., Schachar, R. J. & Tannock, R. Does actigraphy differentiate ADHD subtypes in a clinical research setting? *J Am Acad Child Adolesc Psychiatry* **39**, 752-760 (2000). https://doi.org/10.1097/00004583-200006000-00014

Alfano, C. A., Ginsburg, G. S. & Kingery, J. N. Sleep-related problems among children and adolescents with anxiety disorders. *J Am Acad Child Adolesc Psychiatry* **46**, 224-232 (2007). https://doi.org/10.1097/01.chi.0000242233.06011.8e

Beauchaine, T. P. & Thayer, J. F. Heart rate variability as a transdiagnostic biomarker of psychopathology. *Int J Psychophysiol* **98**, 338-350 (2015). https://doi.org/10.1016/j.ijpsycho.2015.08.004

Singh, I. & Rose, N. Biomarkers in psychiatry. *Nature* **460**, 202-207 (2009). https://doi.org/10.1038/460202a

Dunn, H. A., Patil, D. N., Cao, Y., Orlandi, C. & Martemyanov, K. A. Synaptic adhesion protein ELFN1 is a selective allosteric modulator of group III metabotropic glutamate receptors in trans. *Proc Natl Acad Sci U S A* **115**, 5022-5027 (2018). https://doi.org/10.1073/pnas.1722498115

Girgenti, M. J. *et al.* Transcriptomic organization of the human brain in post-traumatic stress disorder. *Nat Neurosci* **24**, 24-33 (2021). https://doi.org/10.1038/s41593-020-00748-7

Datta, D., Arion, D., Roman, K. M., Volk, D. W. & Lewis, D. A. Altered Expression of ARP2/3 Complex Signaling Pathway Genes in Prefrontal Layer 3 Pyramidal Cells in Schizophrenia. *Am J Psychiatry* **174**, 163-171 (2017). https://doi.org/10.1176/appi.ajp.2016.16020204

Yao, W. *et al.* ERVWE1 Reduces Hippocampal Neuron Density and Impairs Dendritic Spine Morphology through Inhibiting Wnt/JNK Non-Canonical Pathway via miR-141-3p in Schizophrenia. *Viruses* **15** (2023). https://doi.org/10.3390/v15010168

Deng, X. *et al.* Association study of polymorphisms in the neutral amino acid transporter genes SLC1A4, SLC1A5 and the glycine transporter genes SLC6A5, SLC6A9 with schizophrenia. *BMC Psychiatry* **8**, 58 (2008). https://doi.org/10.1186/1471-244X-8-58

Li, Y. X. *et al.* Inhibitors of the Neutral Amino Acid Transporters ASCT1 and ASCT2 Are Effective in In Vivo Models of Schizophrenia and Visual Dysfunction. *J Pharmacol Exp Ther* **367**, 292-301 (2018). https://doi.org/10.1124/jpet.118.251116

Lin, E., Lin, C. H. & Lane, H. Y. Logistic ridge regression to predict bipolar disorder using mRNA expression levels in the N-methyl-D-aspartate receptor genes. *J Affect Disord* **297**, 309-313 (2022). https://doi.org/10.1016/j.jad.2021.10.081

Weis, S. *et al.* Changes in region- and cell type-specific expression patterns of neutral amino acid transporter 1 (ASCT-1) in the anterior cingulate cortex and hippocampus in schizophrenia, bipolar disorder and major depression. *J Neural Transm (Vienna)* **114**, 261-271 (2007). https://doi.org/10.1007/s00702-006-0544-0

Lee, P. H. *et al.* Multi-locus genome-wide association analysis supports the role of glutamatergic synaptic transmission in the etiology of major depressive disorder. *Transl Psychiatry* **2**, e184 (2012). https://doi.org/10.1038/tp.2012.95

Lin, C. H., Huang, M. W., Lin, C. H., Huang, C. H. & Lane, H. Y. Altered mRNA expressions for N-methyl-D-aspartate receptor-related genes in WBC of patients with major depressive disorder. *J Affect Disord* **245**, 1119-1125 (2019). https://doi.org/10.1016/j.jad.2018.12.016

Malki, K. *et al.* Highly polygenic architecture of antidepressant treatment response: Comparative analysis of SSRI and NRI treatment in an animal model of depression. *Am J Med Genet B Neuropsychiatr Genet* **174**, 235-250 (2017). https://doi.org/10.1002/ajmg.b.32494

Hepbasli, D. *et al.* Genotype- and Age-Dependent Differences in Ultrasound Vocalizations of SPRED2 Mutant Mice Revealed by Machine Deep Learning. *Brain Sci* **11** (2021). https://doi.org/10.3390/brainsci11101365

Jia, Y. F. *et al.* Astrocytic Glutamate Transporter 1 (GLT1) Deficiency Reduces Anxiety- and Depression-Like Behaviors in Mice. *Front Behav Neurosci* **14**, 57 (2020). https://doi.org/10.3389/fnbeh.2020.00057

Löning, M. *et al.* sktime: A Unified Interface for Machine Learning with Time Series. *33rd Conference on Neural Information Processing Systems (NeurIPS 2019)* (2019).

Paszke, A. *et al.* PyTorch: An Imperative Style, High-Performance Deep Learning Library. *Adv Neur In* **32** (2019).

Oguiza, I. *tsai - A state-of-the-art deep learning library for time series and sequential data*, <https://github.com/timeseriesAI/tsai> (2022).

Marees, A. T. *et al.* A tutorial on conducting genome-wide association studies: Quality control and statistical analysis. *Int J Methods Psychiatr Res* **27**, e1608 (2018). https://doi.org/10.1002/mpr.1608

Purcell, S. *et al.* PLINK: a tool set for whole-genome association and population-based linkage analyses. *Am J Hum Genet* **81**, 559-575 (2007). https://doi.org/10.1086/519795

Chang, C. C. *et al.* Second-generation PLINK: rising to the challenge of larger and richer datasets. *Gigascience* **4**, 7 (2015). https://doi.org/10.1186/s13742-015-0047-8

Watanabe, K., Taskesen, E., van Bochoven, A. & Posthuma, D. Functional mapping and annotation of genetic associations with FUMA. *Nat Commun* **8**, 1826 (2017). https://doi.org/10.1038/s41467-017-01261-5

Demontis, D. *et al.* Genome-wide analyses of ADHD identify 27 risk loci, refine the genetic architecture and implicate several cognitive domains. *Nat Genet* **55**, 198-208 (2023). https://doi.org/10.1038/s41588-022-01285-8

Demontis, D. *et al.* Discovery of the first genome-wide significant risk loci for attention deficit/hyperactivity disorder. *Nat Genet* **51**, 63-75 (2019). https://doi.org/10.1038/s41588-018-0269-7

Di Tommaso, P. *et al.* Nextflow enables reproducible computational workflows. *Nat Biotechnol* **35**, 316-319 (2017). https://doi.org/10.1038/nbt.3820

Manrai, A. K., Ioannidis, J. P. A. & Patel, C. J. Signals Among Signals: Prioritizing Nongenetic Associations in Massive Data Sets. *Am J Epidemiol* **188**, 846-850 (2019). https://doi.org/10.1093/aje/kwz031

Kurilshikov, A. *et al.* Large-scale association analyses identify host factors influencing human gut microbiome composition. *Nat Genet* **53**, 156-165 (2021). https://doi.org/10.1038/s41588-020-00763-1

Frankish, A. *et al.* GENCODE: reference annotation for the human and mouse genomes in 2023. *Nucleic Acids Res* **51**, D942-D949 (2023). https://doi.org/10.1093/nar/gkac1071

Ochoa, D. *et al.* The next-generation Open Targets Platform: reimagined, redesigned, rebuilt. *Nucleic Acids Res* **51**, D1353-D1359 (2023). https://doi.org/10.1093/nar/gkac1046

Vosa, U. *et al.* Large-scale cis- and trans-eQTL analyses identify thousands of genetic loci and polygenic scores that regulate blood gene expression. *Nat Genet* **53**, 1300-1310 (2021). https://doi.org/10.1038/s41588-021-00913-z

Wang, D. *et al.* Comprehensive functional genomic resource and integrative model for the human brain. *Science* **362** (2018). https://doi.org/10.1126/science.aat8464

Bryois, J. *et al.* Cell-type-specific cis-eQTLs in eight human brain cell types identify novel risk genes for psychiatric and neurological disorders. *Nat Neurosci* **25**, 1104-1112 (2022). https://doi.org/10.1038/s41593-022-01128-z

Quinlan, A. R. & Hall, I. M. BEDTools: a flexible suite of utilities for comparing genomic features. *Bioinformatics* **26**, 841-842 (2010). https://doi.org/10.1093/bioinformatics/btq033

Rozowsky, J. *et al.* The EN-TEx resource of multi-tissue personal epigenomes & variant-impact models. *Cell* **186**, 1493-1511 e1440 (2023). https://doi.org/10.1016/j.cell.2023.02.018

Giambartolomei, C. *et al.* Bayesian test for colocalisation between pairs of genetic association studies using summary statistics. *PLoS Genet* **10**, e1004383 (2014). https://doi.org/10.1371/journal.pgen.1004383

Jansen, P. R. *et al.* Genome-wide analysis of insomnia in 1,331,010 individuals identifies new risk loci and functional pathways. *Nat Genet* **51**, 394-403 (2019). https://doi.org/10.1038/s41588-018-0333-3