Title

Cannabidiol blood metabolite levels after cannabidiol treatment are associated with broadband EEG changes and
improvements in visuomotor and non-verbal cognitive abilities in boys with autism requiring higher levels of support

Authors
Christian Cazares[1], Austin Hutton [1], Gisselle Paez[2], Doris Trauner[2], Bradley Voytek[1,3,4]

1 Department of Cognitive Science, University of California, San Diego, La Jolla, CA

2 Department of Neurosciences, University of California, San Diego School of Medicine, La Jolla, CA

3 Halıcıoğlu Data Science Institute, University of California, San Diego, La Jolla, CA, USA

4 Kavli Institute for Brain and Mind, University of California, San Diego, La Jolla, CA, USA

Corresponding Author: Christian Cazares, cazares@ucsd.edu

Abstract

Oral cannabidiol (CBD) treatment has been suggested to alleviate severe symptoms of autism spectrum disorder (ASD).
While many CBD preparations have been studied in clinical trials involving ASD, none has used purified CBD
preparations or preparations approved by the U.S. Food and Drug Administration, nor have they focused on children with
ASD with higher support needs. Previous studies have identified several candidate electrophysiological biomarkers of
cognitive and behavioral disabilities in ASD, with emerging biomarkers including periodic (oscillatory) and aperiodic
measures of neural activity. We analyzed electroencephalography (EEG) recordings from 24 boys with ASD and higher
support needs (aged 7-14 years) from a prior double-blind, placebo-controlled, crossover Phase II Clinical Trial
(NCT04517799) that investigated whether 8 weeks of daily CBD treatment (titrated to 20 mg/kg/day) improved severe
behavioral problems, measured at baseline, post-CBD, post-placebo, and post-washout. Using linear mixed effect
models, we found that aperiodic EEG measures varied with CBD metabolite levels in blood, as evidenced by a larger
aperiodic offset across the scalp and a decreased aperiodic exponent across occipital electrodes. Furthermore, CBD
metabolite levels in blood had a positive association with receptive vocabulary, nonverbal intelligence and visuomotor
coordination. Our data suggest that this daily CBD preparation and administration schedule produced mixed effects, with
some children showing improvements in cognitive and behavioral abilities while others demonstrated limited changes. Our
findings support the inclusion of aperiodic EEG measures alongside traditional oscillatory EEG measures as candidate
biomarkers for tracking the variable clinical impact of purified CBD treatment in children with ASD.

1. Introduction

Neurodevelopmental disorders (NDDs), including autism spectrum disorder (ASD), may be characterized by severe cognitive and behavioral dysfunction that impose significant burdens on caregivers, families, and finances [1]. The core symptoms of ASD include deficits in social communication and stereotyped repetitive behaviors, but other co-occurring behaviors such as self-injurious and aggressive behaviors [2], may require intervention. Current pharmacotherapies have limited efficacy, and progress in developing new treatments is hindered by a lack of reliable biomarkers that can identify patients who are most likely to benefit from investigational therapies in early life [3].

Local field potential signals, such as those recorded by scalp electroencephalography (EEG), comprise mixed periodic and aperiodic components. These components are thought to reflect synchronized and asynchronous neuronal firing in cortical networks, respectively, and can serve as putative indices of cortical excitation-inhibition (E:I) balance in health and disease [4–9]. These components correlate with cognitive processes and show alterations in a wide variety of conditions associated with E:I imbalance, including NDDs [7,10–15]. Task-free EEG has proven valuable for investigating neural correlates in intellectually impaired and minimally verbal pediatric populations [10,14,16,17]. EEG abnormalities in NDDs likely result from impaired neuronal maturation during early development [18–20]. Previous research has largely focused on periodic components, such as the oscillatory sensorimotor mu rhythm and visual cortical alpha rhythms, both in the ~8-12 Hz frequency range [21–23]. Peak alpha frequency (6-12 Hz) has been identified as a biomarker for non-verbal cognition in ASD [24], while delta rhythms (1-4 Hz) show disruptions in ASD [16,25]. Building on these findings, identifying complementary aperiodic EEG signal features that are thought to reflect cortical E:I balance could further inform the development of pharmacotherapies and biomarkers for severe ASD.

Medications like risperidone and aripiprazole show efficacy in improving ASD behavioral symptoms [26], but their adverse effects require close medical monitoring [27]. This has led to growing interest in cannabidiol (CBD), a non-psychoactive cannabis-derived compound that has shown promising anecdotal and clinical trial outcomes for children with intractable epilepsy and co-morbid ASD, with mild side effects [28–32]. This led to U.S. Food and Drug Administration (FDA) approval of Epidiolex®, a plant-based purified CBD medication for treatment of intractable childhood epilepsies. During the same time frame, there have been a number of investigations relating to the use of non-FDA-approved CBD-based medications in other conditions, including retrospective studies in children with ASD, where participants exhibited improvements in disruptive behavior when treated with CBD:THC (20:1) formulations [33], with parents reporting reduced aggression, hyperactivity and anxiety [34]. Social communication improvements have also been reported following 6 months of CBD:THC formulation in children and adolescents with ASD [35,36], with speculation that these improvements are mediated by oxytocin-dependent endocannabinoid signaling as shown in rodents [37,38]. If behavioral and social improvements are mediated through the endocannabinoid system, it raises the question whether purified CBD can lead to similar improvements without the need for a psychoactive THC component in its formulation. Recent evidence of clinical improvement with CBD comes from a clinical trial in which blinded clinicians observed reductions in aggressive behaviors and hyperactivity, as well as improvements in communication [39].

Single doses of CBD modulate GABA levels in prefrontal regions [40], potentially affecting cortical E:I balance through GABAergic inhibition, which has been proposed to be assessed from spectral parameterization of EEG signals [41–43]. Although CBD hasn't been directly studied in ASD animal models, studies in a Dravet syndrome mouse model showed improvements in social deficits [44], and CBD administration in adolescent mice yielded modest spatial memory improvements without negative impacts [45]. CBD-based medication is thus a viable candidate for investigating therapeutic effects on cognitive and behavioral symptoms in children with ASD, given its regulatory precedent, promising safety profile, and preliminary data from both human and mouse models. However, there has been no longitudinal, placebo-controlled monitoring of brain activity and cognitive-behavioral abilities in children with ASD requiring higher levels of support receiving CBD treatment.

Here we investigated periodic and aperiodic measures of EEG signals as potential biomarkers for CBD treatment effects in ASD. We hypothesized that EEG signal measures and cognitive-behavioral outcomes would be associated with CBD treatment responses, as indicated by active and inactive CBD metabolite levels in blood before and after an 8-week oral CBD regimen in boys with ASD and higher support needs [46–48].

2. Patients and Methods

85 Below is a condensed version of Patients and Methods. A detailed version can be found in the Supplementary Materials.

2.1. Experimental recordings and cognitive-behavioral testing

2.1.1. Participants

Our data was sourced from a clinical trial performed at the University of California, San Diego [39]. The trial was approved
by the institutional review board of the University of California, San Diego and registered with www.clinicaltrials.gov
(NCT04517799) to investigate Epidiolex® (CBD) effects on severe behavioral issues in boys with ASD. The dataset
included in this study consisted of 24 males (ages 7-14) who had prior clinical diagnoses of ASD using DSM-5 criteria.
The diagnoses were made by clinicians with expertise and experience in diagnosing and treating children with ASD. All
participants underwent Autism Diagnostic Observation Schedule, Second Edition (ADOS-2) assessments of symptoms
and exhibited severe behavioral issues including stereotypies, aggression, self-injury, and/or hyperactivity. Baseline
cognitive and behavioral assessments (Table 1) revealed the extent of impairments, with many participants unable to
attempt certain standardized assessments or performing below normative ranges, supporting their characterization as
having higher support needs. All participants were reported to require continuous supervision of activities of daily living
and personal safety. The participants were free of other neurological conditions, including epilepsy. Parental consent was
obtained, with child assent waived due to participants' cognitive limitations.

2.1.2 Experimental design

Our dataset consisted of assessments made via a randomized, two-arm design (CBD-Placebo or Placebo-CBD) with
treatment periods lasting 8 weeks, separated by a 4-week washout period. Cognitive-behavioral assessments and task-
free scalp EEG recordings were collected at baseline and after each timepoint (Figure 1A). CBD dosage was determined
based on participant weight starting with 5 mg/kg/day, with a gradual increase over the first two weeks to reach the
maximum dose (20 mg/kg/day, divided into two oral doses) for the remainder of the study period. Whole blood samples
analyzed in this study were collected near EEG and cognitive-behavioral assessments. CBD metabolite levels were
quantified alongside the major active metabolite 7-hydroxy-cannabidiol (7-OH-CBD) and inactive metabolite 7-carboxy-
cannabidiol (7-COOH-CBD) [46,48,49]. Analysis of CBD metabolite concentrations in blood samples following the 8-week
CBD treatment regimen revealed levels comparable to those found in a prior report using orally administered CBD
(Epidiolex®) for four days in healthy adults [46] (Figure Supplement 1A-C).

2.1.3 Cognitive-behavioral testing

Assessments included Autism Diagnostic Observation Schedule, Second Edition (ADOS-2) Modules 1-3, Test of
Nonverbal Intelligence, Fourth Edition (TONI-4) for nonverbal intelligence [50], Repetitive Behavior Scale-Revised (RBS-
R) for repetitive behaviors [51], Peabody Picture Vocabulary Test, Fourth Edition (PPVT-4) for receptive vocabulary [52],
Expressive One-Word Picture Vocabulary Test, Fourth Edition (EOWPVT-4) for expressive vocabulary [53], and Beery-
Buktenica Developmental Test of Visual-Motor Integration, Sixth Edition (Beery VMI-6) for visual-motor integration,
including supplemental subtests for visual perception (Beery VP) and motor coordination (Beery MC) [54]. Standard
scores were calculated where possible for sample characterization purposes at baseline, while raw scores were used for
model analyses to avoid floor effects and maintain sensitivity to change in this severely impaired population.

2.1.4 EEG recording setup

121 EEG data were recorded using a CGX Quick-20 Dry wireless headset with 20 channels in 10-20 montage, online
122 referenced to A1. Signals were sampled at 500 Hz (0-131 Hz bandwidth, 24-bit resolution), with electrode impedance
123 maintained below 500 kΩ prior to the start of each recording session based on the manufacturer's specifications. Task-
124 free EEG recordings were made during awake, passive viewing periods with the goal of capturing 5 to 10 minutes during
125 which the child remained calm by viewing a video of their choice.

2.2. Data analysis

2.2.1. Preprocessing

Analysis was performed in Python using numpy [55], scipy [56], autoreject [57], MNE [58], specparam (formerly known as
FOOOF) [59], and statsmodel [60]. EEG data was re-referenced offline to a linked ears reference (A1/A2) and
preprocessed using Autoreject [57] to establish global artifact rejection thresholds. Data was bandpass filtered (0.1-100
Hz) and processed with ICA ("extended infomax") to remove non-brain components per ICLabel criteria [61] (Figure
Supplement 1B). Final analysis included 89 sessions across timepoints (Baseline: 21, Post-CBD: 23, Post-Placebo: 23,
Post-Wash: 22), with 3-4 recordings per participant (Table 2).

2.2.2. Calculation of EEG signal features

Channels were grouped into frontal (F7, F3, Fz, F4, F8), central (Cz, C3, C4), and occipital (O1, O2) regions (Figure 1D).
The power spectral density (PSD) of EEG signals was calculated using Welch's method [62] (1.0s Hamming windows,
0.5s overlap, 0.1-50 Hz range). Spectral parameterization was limited to 0.5-13 Hz to minimize muscle artifacts [5], with
parameters: peak width limits (1, 12.0), maximum peaks: 6, minimum peak amplitude: 0.0, peak threshold: 2.0, aperiodic
mode: "fixed". Including frequencies above 13 Hz systematically increased model error and biased aperiodic parameters
by forcing the model to accommodate high frequency activity, which is particularly problematic in pediatric autism
populations where movement artifacts (>13 Hz) are prevalent (Figure Supplement 2). Analysis yielded aperiodic
exponent, offset, and aperiodic adjusted band powers (delta: 0.1-4 Hz, theta: 4-8 Hz, alpha: 8-13 Hz) (Figure 1E, Figure
Supplement 1E). Channels with model fits $R^2 <0.80$ were excluded. Aperiodic corrected oscillatory power was calculated
by subtracting the aperiodic fit from the PSD and finding the maximum corrected power within each band, then averaging
within channel group.

2.2.3. Statistical analysis

Blood metabolite levels were z-scored to address non-normality from detection thresholds. Our dataset showed variability
in sample sizes across assessments due to assessment administration difficulties, experimenter error, or low participant
cooperation. To account for missing data, we used Linear Mixed Effects (LME) models which are advantageous for
handling data where time points are nested within participants and can account for this type of data dependency, as used
in other studies where children were able to complete certain assessments but not others [63]. Our statistical approach
examined three distinct electrode groups (frontal, central, occipital) based on established functional neuroanatomy and
connectivity patterns. For each electrode group, we analyzed relationships with three biochemically distinct CBD
metabolites (CBD, 7-OH-CBD, 7-COOH-CBD), which have different pharmacological properties and half-lives. Each
model included random subject intercepts and fixed effects for randomization, timepoint, age, ADOS-2 module used to
assess participant as a proxy for participants' language ability, and days since last blood extraction for metabolite
measurements. Specifically, timepoint (Baseline, Post-placebo, Post-CBD, Post-wash) and Randomization order (CBD-
first or Placebo-first) were included to account for potential practice effects from repeated measurements. Three
metabolite models were applied to each electrode group (totaling 9 models) and cognitive-behavioral assessment (totaling
15 models). To further interpret our findings, we calculated the estimated marginal means (EMMs) for post-CBD
timepoints derived from significant model coefficients and analyzed the relationship between EMMs and changes in post-
CBD and baseline measures for participants that had data for both sessions included in the linear mixed model. The
number of sessions included in each model is listed in Table 2. Post-hoc regression analyses using estimated marginal
means were conducted to visualize the directionality and magnitude of significant mixed model relationships.

3. Results

3.1. Broadband and aperiodic adjusted task-free EEG features across the scalp had a positive association with CBD blood metabolite levels

We found a clear increase across all CBD blood metabolites levels following the 8-week CBD treatment period (Figure 1A,
B). Prior EEG studies on intellectually disabled and minimally verbal neurodevelopmental disorder populations have
shown heterogeneity in neural spectral power and aperiodic measures of scalp EEG activity grouped by frontal, central,
and occipital electrodes [10,14,16]. We first confirmed that post-processed EEG durations were similar across study time
points (Figure 1C) and used a similar approach by grouping channels into frontal, central, and occipital areas to
investigate the relationship between periodic and aperiodic EEG measures and CBD treatment (Figure 1D, E).
174

Our linear mixed effect models confirmed that CBD treatment increased metabolite levels in blood (p's < 0.001; Figure 2A,
C, F). CBD metabolite levels had varying relationships with periodic and aperiodic measures of EEG across brain regions.
In frontal electrodes, 7-COOH-CBD levels were associated with increased aperiodic offset (β = 0.090, 97.5% CI = [0.023,
0.158], p = 0.009; Figure 2A, bottom), as confirmed by post-hoc analysis of estimated marginal means from baseline to
post-CBD (n = 20, R^2 = 0.510, p < 0.001; Figure 2B). Central electrodes showed mixed effects, with higher CBD
metabolite levels associated with decreased aperiodic adjusted alpha power (β = -0.135, 97.5% CI = [-0.267, -0.002], p =
0.046; Figure 2C, top; post-hoc analysis of n = 20 participants, R^2 = 0.379, p = 0.004; Figure 2D), while 7-COOH-CBD
metabolite levels were associated with an increased aperiodic offset (β = 0.094, 97.5% CI = [0.026, 0.161], p = 0.006;
Figure 2C, bottom; post-hoc analysis of n = 20 participants, R^2 = 0.495, p < 0.001; Figure 2E). In occipital electrodes,
higher 7-COOH-CBD metabolite levels were associated with increased aperiodic offset (β = 0.099, 97.5% CI = [0.033,
0.165], p = 0.003; Figure 2F, bottom; post-hoc analysis of n = 20 participants, R^2 = 0.476, p < 0.001; Figure 2G) and
decreases in aperiodic exponent (β = -0.235, 97.5% CI = [-0.460, -0.010], p = 0.041; Figure 2F, bottom), though this latter
relationship was relatively modest in post-hoc analysis (n = 20 participants, R^2 = 0.058, p = 0.308; Figure 2H).

188 3.2. Measures of receptive vocabulary, nonverbal intelligence, and visuomotor coordination
189 had a positive association with CBD blood metabolite levels

Our linear mixed effect models showed a mixture of positive associations between CBD blood metabolite levels and
cognitive-behavioral performance measures after controlling for age, ADOS-2 module, and treatment order. Higher CBD
metabolite levels were associated with improved receptive vocabulary on the PPVT-4 (β = 0.004, 97.5% CI = [0, 0.007], p
= 0.039; Figure 3A, top), though this relationship was modest in post-hoc analysis (n = 12 participants, R^2 = 0.225, p =
0.119; Figure 3B). Similarly, increased 7-COOH-CBD levels were associated with higher nonverbal intelligence scores on
the TONI-4 (β = 0.012, 97.5% CI = [0.005, 0.019], p < 0.001; Figure 3C, bottom), with post-hoc analysis showing a
modest relationship (n = 10 participants, R^2 = 0.010, p = 0.780 ; Figure 3D). The strongest CBD treatment improvements
appeared in visuomotor integration abilities. Both 7-OH-CBD and 7-COOH-CBD blood metabolite levels were positively
associated with Beery-Buktenica VMI test performance, with 7-OH-CBD showing a modest effect (β = 0.034, 97.5% CI =
[0.002, 0.066], p = 0.038; Figure 3E, middle) with post-hoc analysis showing a modest relationship (n = 18 participants, R^2
= 0.136, p = 0.131; Figure 3F). Beery-Buktenica VMI scores particularly improved with higher 7-COOH-CBD levels (β =
0.036, 97.5% CI = [0.018, 0.054], p < 0.001; Figure 3E, bottom), confirmed by post-hoc analysis (n = 18 participants, R^2 =
0.257, p = 0.032; Figure 3G). ADOS-2 module used to assess participants showed a consistent negative relationship with
7-COOH-CBD blood metabolite levels across in the PPVT-4 (β = -0.194, 97.5% CI = [-0.384, -0.004], p = 0.046; Figure
3A, bottom) and TONI-4 (β = -0.192, 97.5% CI = [-0.372, -0.011], p = 0.037; Figure 3C, bottom) models. CBD treatment
showed no significant effects on repetitive behaviors (RBS assessment) or expressive vocabulary (EOWPVT-4
assessment) (Figure Supplement 3).

207 4. Discussion

Our secondary analyses from a double-blind, placebo-controlled trial provided longitudinal measures of oral CBD effects
on task-free scalp EEG signal features, CBD-related metabolism, and cognitive-behavioral assessments in children with
ASD requiring higher levels of support. Improvements in visuomotor coordination, alongside more modest gains in
nonverbal intelligence and receptive vocabulary abilities, but not in expressive language or repetitive behaviors, suggest
CBD treatment and related metabolites may differentially affect distinct aspects of brain function. The strongest
associations were observed in visuomotor tasks, potentially stemming from CBD's modulation of GABAergic signaling in
brain regions critical for motor control and visual processing [40,64,65]. The concurrent improvements in receptive
vocabulary and nonverbal intelligence, although more modest, align with previous CBD:THC preparation studies that
reported improvements in social and cognitive deficits in ASD patients [35,36]. While group-level analyses showed
positive trends for these measures, individual participants showed varying magnitudes of change, with some
demonstrating robust improvements while others showed minimal or negative changes in relation to CBD metabolite
levels, highlighting the importance of considering individual variability in treatment responses.
220
Our EEG findings provide insight into CBD treatment's potential impact on neural function. Specifically, the predominance
of aperiodic over periodic effects suggests CBD treatment influences brain activity through changes in basic cellular and
network properties rather than through specific excitatory or inhibitory circuits that mediate oscillatory activity. Aperiodic
measures of brain activity are thought to reflect changes in broadband power as a function of neuronal population spiking

[66,67]. This broadband change in neural activity aligns with magnetic resonance spectroscopy evidence showing CBD's
broad modulation of glutamate-GABA systems in ASD patient cortex [40]. The negative relationship between CBD
metabolite levels and the aperiodic exponent in occipital regions may reflect CBD's reported decrease of GABA+
macromolecules in ASD, as reduced inhibitory signaling typically decreases the aperiodic exponent. This relationship
could potentially explain the differential effects on behavioral measures, with tasks requiring extensive visuomotor
integration showing the strongest positive association with CBD blood metabolite levels, possibly due to their reliance on
GABAergic signaling in visual and motor circuits. The inverse relationship between CBD metabolite levels and aperiodic-
adjusted alpha power in central regions might similarly indicate a broad shift toward increased cortical excitability rather
than inhibition-mediated oscillatory signaling [40]. The robust spatial consistency of changes in aperiodic features across
the scalp, combined with their established test-retest reliability in ASD populations [8], suggests these measures may
provide complementary EEG biomarkers for tracking treatment outcomes alongside traditional oscillatory measures,
whose associations with CBD treatment were observed to be more spatially restricted.
237
Given that aperiodic measures are increasingly recognized as indicators of neurotypical brain development, their
alterations in relation to CBD blood metabolite levels here suggest their potential utility as biomarkers for treatment
responses in ASD, further extending previous work showing EEG features can serve as reliable biomarkers in human
neurodevelopmental disorders [10,14–16]. Specifically, we observed a negative relationship between levels of 7-COOH-
CBD metabolite in blood and the aperiodic exponent in occipital regions, indicating a decrease of the aperiodic exponent,
as reflected by a "flatter" slope of the power spectrum model fit after CBD treatment, which is observed in typically
developing children [59,68,69]. In contrast, we saw a positive relationship between levels of 7-COOH-CBD metabolite in
blood and the aperiodic offset, indicating an increase in broadband power that contrasts with the decreasing patterns seen
in typical neurodevelopment [68,69]. Our findings suggest that this CBD regimen may not induce all the expected
trajectories of aperiodic activity throughout neurodevelopment, mirroring how the treatment regimen did not improve all
cognitive-behavioral measures and had individual participant variability in responses.
249
Our restriction of spectral parameterization to 0.5-13 Hz aligns with established autism EEG biomarker research, where
peak alpha frequency (6-12 Hz) correlates with non-verbal cognition and delta rhythms (1-4 Hz) show characteristic
disruptions [16,24,25]. While this range excluded potentially relevant 25-35 Hz beta oscillations (also termed "iota") that
have been reported in healthy children and those with Fragile X syndrome [7,70], our focus on lower frequencies aimed to
capture the established biomarkers while maintaining interpretability of aperiodic parameters as indices of E:I balance.
This restriction also avoided distinguishing genuine high frequency oscillations from movement artifacts in our behaviorally
challenging population. Future studies could investigate CBD's effects on higher frequencies, but for our research
question about established lower-frequency markers of autism, the 0.5-13 Hz range provided robust model fits (mean R^2 =
0.987) while avoiding interpretational ambiguities.
259
Our treatment regimen was derived from prior CBD trials in epilepsy, but the optimal duration for treatment effects likely
differs [29,30]. The relationship between CBD metabolite levels in blood and cognitive-behavioral outcomes is notable,
particularly given that the most consistent effects were observed with 7-COOH-CBD, which doesn't cross the blood-brain
barrier. This suggests that the effects we observed may involve indirect peripheral mechanisms in addition to direct neural
modulation by CBD and 7-OH-CBD metabolites. Whether similar cognitive benefits would emerge with different treatment
durations (either shorter or longer than 8 weeks) remains an open question warranting future studies, as we are not aware
of any CBD-related clinical trials that have specifically examined these cognitive-behavioral measures. Longer-term
follow-up is necessary to determine if the observed cognitive improvements persist after treatment discontinuation. The
magnitude of effects we observed on visuomotor function compared to other domains may reflect neural circuit-specific
responses to CBD, but it could also reflect differential test sensitivity to change, or general attentional improvements. The
Beery VMI requires only drawing one additional line correctly for improvement, whereas language measures require
learning new vocabulary over 8 weeks, which is a much more demanding change for children with neurodevelopmental
conditions. Additionally, if CBD enhances attention and focus, this could more readily improve visuomotor tasks (where
attention to detail directly improves performance) than vocabulary tasks (where attention cannot compensate for unknown
words). Findings from the primary clinical trial support this interpretation, such that blinded clinicians observed reduced
aggression, hyperactivity, and anxiety in 68% of children during CBD treatment, with many showing calmer behavior and
improved communication [39]. These behavioral improvements suggest that CBD's cognitive effects may be mediated
through multiple mechanisms, including both enhanced attention and reduced disruptive behaviors during testing, as well
as potential direct domain-specific neural changes, which could all contribute to the observed improvements.

Our results revealed substantial individual variability in treatment responses in relation to CBD blood metabolite levels.
While group-level analyses showed positive associations, individual participants showed varying magnitudes of change,
with some reflecting no change or even decreases in outcome measures. This heterogeneity likely reflects the complex
pharmacology of CBD metabolism. The inactive 7-COOH-CBD metabolite has a longer half-life (>48 hours) compared to
CBD and 7-OH-CBD, which decrease rapidly after 2-8 hours post-administration [48]. Since assessments weren't
consistently conducted within 8-12 hours of blood draws, measuring 7-COOH-CBD provided more reliable treatment
indicators than CBD or 7-OH-CBD levels alone. We observed robust relationships between 7-COOH-CBD levels and both
EEG features and cognitive-behavioral measures, despite this metabolite's inability to cross the blood-brain barrier,
suggesting CBD's therapeutic effects involve both direct neural modulation via CBD or 7-OH-CBD and indirect peripheral
mechanisms. For future studies, we recommend conducting blood draws shortly before assessments to better capture
CBD and 7-OH-CBD effects. The absence of preserved weight data limited our reporting to weight-based dosing
protocols, so future work should document both weight-based and absolute dosing metrics to provide a more precise
measure of the actual amount of medication administered while incorporating longer intervention periods with validated
measures for detecting short-term changes. Additionally, recent evidence of sex-based differences in 7-COOH-CBD
metabolism [46] suggests that understanding metabolite relationships could help explain outcome variability and inform
personalized dosing strategies, particularly in clinical settings where precise timing is challenging with pediatric ASD
populations.
297

Conducting longitudinal studies in ASD children with higher support needs presented challenges that required flexibility in
EEG data collection approaches. In our study, children viewed self-selected videos during EEG recordings due to the
challenge of keeping them engaged, and differences in video content were not documented, leaving potential video
stimulus-related effects on brain activity unknown. However, prior work with similar populations suggests this passive
viewing approach does not confound interpretations of holistic neural activity measures [10]. This approach represented a
necessary compromise that enabled the study of neural activity in this historically understudied population. Furthermore,
the detection of scalp-wide aperiodic measure effects, which tend to be stable across behavioral conditions in pediatric
and ASD populations [5,8], suggests robust, global changes in neural activity rather than behavioral state artifacts that
might be induced by specific video content. In addition, our findings are limited to male children, which limits
generalizability to females with ASD. This was an intentional design choice in the original study, given the small sample
size recruited and known differences in autism symptomatology between males and females that may have confounded
findings [39].
310

A key limitation in interpreting our results stems from the scarcity of research validating cognitive and behavioral test
sensitivity within brief intervention periods, particularly for children with autism who have higher support needs. To our
knowledge, only one study has examined short-term changes in TONI-4 performance, finding small but detectable
changes in adult schizophrenia patients over just a two-week period [71]. While we observed improvements in visuomotor
coordination, receptive vocabulary and nonverbal intelligence measures in relation to CBD blood metabolite levels, the
clinical meaningfulness of these changes requires careful consideration. An additional limitation is that while our linear
mixed effects models detected significant associations when controlling for multiple covariates, several relationships
showed modest effect sizes in post-hoc linear regressions, including receptive vocabulary (PPVT-4), nonverbal
intelligence (TONI-4), and some EEG measures (aperiodic exponent in occipital electrodes). It is possible that these
represent subtle effects that emerge primarily through rigorous statistical control rather than strong standalone
associations, requiring cautious interpretation and replication in larger samples. The use of raw scores for EOWPVT-4,
PPVT-4, and Beery VMI-6 assessments rather than standardized scores helped avoid floor effects and maintain
sensitivity to detect changes, but this approach also makes it more challenging to contextualize the magnitude of
improvements against established clinical thresholds. Furthermore, individual variability in our results likely reflects both
genuine treatment response differences and measurement unreliability from variable attention and cooperation during
testing, with lower scores being more susceptible to such confounds than higher scores. Additionally, the 8-week
intervention period, while consistent with other pharmacological trials, may not have been optimal for capturing the full
extent of potential cognitive changes, as developmental improvements often unfold over longer timeframes. Our CBD
metabolite levels aligned with previous oral administration studies in healthy adults [41] and a child with Dravet Syndrome
[72], though age-related metabolic differences should be considered when interpreting these comparisons
331

Developing novel ASD treatments requires reliable methods to characterize associated brain activity changes. While task-free EEG measurements have shown promise as biomarkers of ASD severity, even from 3 months of age [24,73–75], these studies did not separate periodic and aperiodic components of neural power spectra, obscuring the specific neural processes affected. More recent work shows age-dependent variations: adult ASD populations show mixed differences in task-free EEG features [24,76], while in preterm children, aperiodic features are associated with increased ASD risk [77]. Notably, aperiodic measures of EEG activity in 6-8-year-old children with ASD demonstrates high test-retest reliability compared to healthy controls [8], supporting its potential as a developmental biomarker. Our study addresses these gaps by applying spectral parameterization to EEG recordings from near-adolescent ASD children with well-defined higher support needs.

In conclusion, we report that metabolic responses to an 8-week daily CBD regimen using a purified formulation with regulatory precedent were associated with improvements to select domains of cognitive and behavioral ability, such as visuomotor coordination, receptive vocabulary ability and nonverbal intelligence, and that these improvements coincided with neural activity changes primarily in the aperiodic component of task-free scalp EEG signals.

Data Availability Statement

The data presented in this study will be available on request once data analyses are completed from UC San Diego Center for Medicinal Cannabis Research (CMCR) through contacting Doris Trauner, M.D. The EEG and cognitive-behavioral assessment data are not publicly available to protect the children's and their families' privacy. The analysis code needed to reproduce the analysis and figures is provided here: https://github.com/voytekresearch/cbd-asd-eeg

References

1. GBD 2019 Mental Disorders Collaborators. Global, regional, and national burden of 12 mental disorders in 204 countries and territories, 1990-2019: a systematic analysis for the Global Burden of Disease Study 2019. Lancet Psychiatry. 2022;9:137–150.
2. American Psychiatric Association. Diagnostic and Statistical Manual of Mental Disorders. 5th ed. Arlington, VA: American Psychiatric Association; 2013.
3. Varcin KJ, Nelson CAI. A developmental neuroscience approach to the search for biomarkers in autism spectrum disorder. Current Opinion in Neurology. 2016;29:123.
4. Gao R. Interpreting the electrophysiological power spectrum. J Neurophysiol. 2016;115:628–630.
5. Schaworonkow N, Voytek B. Longitudinal changes in aperiodic and periodic activity in electrophysiological recordings in the first seven months of life. Dev Cogn Neurosci. 2020;47:100895.
6. Li W. Excitation and Inhibition Imbalance in Rett Syndrome. Front Neurosci. 2022;16:825063.
7. Wilkinson CL, Nelson CA. Increased aperiodic gamma power in young boys with Fragile X Syndrome is associated with better language ability. Mol Autism. 2021;12:17.
8. Levin AR, Naples AJ, Scheffler AW, Webb SJ, Shic F, Sugar CA, et al. Day-to-Day Test-Retest Reliability of EEG Profiles in Children With Autism Spectrum Disorder and Typical Development. Front Integr Neurosci. 2020;14:21.
9. Ostlund BD, Alperin BR, Drew T, Karalunas SL. Behavioral and cognitive correlates of the aperiodic (1/f-like) exponent of the EEG power spectrum in adolescents with and without ADHD. Developmental Cognitive Neuroscience. 2021;48:100931.
10. Roche KJ, LeBlanc JJ, Levin AR, O'Leary HM, Baczewski LM, Nelson CA. Electroencephalographic spectral power as a marker of cortical function and disease severity in girls with Rett syndrome. J Neurodev Disord. 2019;11:15.
11. Fries P. A mechanism for cognitive dynamics: neuronal communication through neuronal coherence. Trends in Cognitive Sciences. 2005;9:474–480.
12. Schnitzler A, Gross J. Normal and pathological oscillatory communication in the brain. Nat Rev Neurosci. 2005;6:285–296.
13. Kahana MJ. The Cognitive Correlates of Human Brain Oscillations. J Neurosci. 2006;26:1669–1672.
14. Sysoeva O, Maximenko V, Kuc A, Voinova V, Martynova O, Hramov A. Abnormal spectral and scale-free properties of resting-state EEG in girls with Rett syndrome. Sci Rep. 2023;13:12932.
15. Ostlund B, Donoghue T, Anaya B, Gunther KE, Karalunas SL, Voytek B, et al. Spectral parameterization for studying neurodevelopment: How and why. Developmental Cognitive Neuroscience. 2022;54:101073.
16. Neo WS, Foti D, Keehn B, Kelleher B. Resting-state EEG power differences in autism spectrum disorder: a systematic review and meta-analysis. Transl Psychiatry. 2023;13:389.
17. Van der Molen MJW, Van der Molen MW. Reduced alpha and exaggerated theta power during the resting-state EEG in fragile X syndrome. Biol Psychol. 2013;92:216–219.

18. Rubenstein JLR. Three hypotheses for developmental defects that may underlie some forms of autism spectrum
disorder. Curr Opin Neurol. 2010;23:118–123.
19. Sohal VS, Rubenstein JLR. Excitation-inhibition balance as a framework for investigating mechanisms in
neuropsychiatric disorders. Mol Psychiatry. 2019;24:1248–1257.
20. Voytek B, Knight RT. Dynamic network communication as a unifying neural basis for cognition, development, aging,
and disease. Biol Psychiatry. 2015;77:1089–1097.
21. Bender A, Voytek B, Schaworonkow N. Resting-state is not enough: alpha and mu rhythms change shape across
development, but lack diagnostic sensitivity. 2023:2023.10.13.562301.
22. Freschl J, Azizi LA, Balboa L, Kaldy Z, Blaser E. The development of peak alpha frequency from infancy to
adolescence and its role in visual temporal processing: A meta-analysis. Developmental Cognitive Neuroscience.
2022;57:101146.
23. Javitt DC, Siegel SJ, Spencer KM, Mathalon DH, Hong LE, Martinez A, et al. A roadmap for development of neuro-
oscillations as translational biomarkers for treatment development in neuropsychopharmacology.
Neuropsychopharmacol. 2020;45:1411–1422.
24. Dickinson A, DiStefano C, Senturk D, Jeste SS. Peak alpha frequency is a neural marker of cognitive function
across the autism spectrum. European Journal of Neuroscience. 2018;47:643–651.
25. Cornew L, Roberts TPL, Blaskey L, Edgar JC. Resting-State Oscillatory Activity in Autism Spectrum Disorders. J
Autism Dev Disord. 2012;42:1884–1894.
26. Ghanizadeh A, Sahraeizadeh A, Berk M. A Head-to-Head Comparison of Aripiprazole and Risperidone for Safety
and Treating Autistic Disorders, a Randomized Double Blind Clinical Trial. Child Psychiatry Hum Dev. 2014;45:185–
192.
27. Politte LC, McDougle CJ. Atypical antipsychotics in the treatment of children and adolescents with pervasive
developmental disorders. Psychopharmacology. 2014;231:1023–1036.
28. Porter BE, Jacobson C. Report of a parent survey of cannabidiol-enriched cannabis use in pediatric treatment-
resistant epilepsy. Epilepsy Behav. 2013;29:574–577.
29. Devinsky O, Marsh E, Friedman D, Thiele E, Laux L, Sullivan J, et al. Cannabidiol in patients with treatment-
resistant epilepsy: an open-label interventional trial. Lancet Neurol. 2016;15:270–278.
30. Devinsky O, Patel AD, Cross JH, Villanueva V, Wirrell EC, Privitera M, et al. Effect of Cannabidiol on Drop Seizures
in the Lennox-Gastaut Syndrome. N Engl J Med. 2018;378:1888–1897.
31. Kuester G, Vergara K, Ahumada A, Gazmuri AM. Oral cannabis extracts as a promising treatment for the core
symptoms of autism spectrum disorder: Preliminary experience in Chilean patients. Journal of the Neurological
Sciences. 2017;381:932–933.
32. Thompson MD, Martin RC, Grayson LP, Ampah SB, Cutter G, Szaflarski JP, et al. Cognitive function and adaptive
skills after a one-year trial of cannabidiol (CBD) in a pediatric sample with treatment-resistant epilepsy. Epilepsy
Behav. 2020;111:107299.
33. Aran A, Cassuto H, Lubotzky A, Wattad N, Hazan E. Brief Report: Cannabidiol-Rich Cannabis in Children with
Autism Spectrum Disorder and Severe Behavioral Problems-A Retrospective Feasibility Study. J Autism Dev Disord.
2019;49:1284–1288.
34. Barchel D, Stolar O, De-Haan T, Ziv-Baran T, Saban N, Fuchs DO, et al. Oral Cannabidiol Use in Children With
Autism Spectrum Disorder to Treat Related Symptoms and Co-morbidities. Front Pharmacol. 2019;9:1521.
35. Hacohen M, Stolar OE, Berkovitch M, Elkana O, Kohn E, Hazan A, et al. Children and adolescents with ASD treated
with CBD-rich cannabis exhibit significant improvements particularly in social symptoms: an open label study. Transl
Psychiatry. 2022;12:375.
36. Fleury-Teixeira P, Caixeta FV, Ramires da Silva LC, Brasil-Neto JP, Malcher-Lopes R. Effects of CBD-Enriched
Cannabis sativa Extract on Autism Spectrum Disorder Symptoms: An Observational Study of 18 Participants
Undergoing Compassionate Use. Front Neurol. 2019;10:1145.
37. Premoli M, Carone M, Mastinu A, Maccarinelli G, Aria F, Mac Sweeney E, et al. Cannabis Sativa Oil Promotes
Social Interaction and Ultrasonic Communication by Acting on Oxytocin Pathway. Cannabis Cannabinoid Res.
2024;9:1514–1523.
38. Wei D, Lee D, Cox CD, Karsten CA, Peñagarikano O, Geschwind DH, et al. Endocannabinoid signaling mediates
oxytocin-driven social reward. Proc Natl Acad Sci U S A. 2015;112:14084–14089.
39. Trauner D, Umlauf A, Grelotti DJ, Fitzgerald R, Hannawi A, Marcotte TD, et al. Cannabidiol (CBD) Treatment for
Severe Problem Behaviors in Autistic Boys: A Randomized Clinical Trial. J Autism Dev Disord. 2025. 24 May 2025.
https://doi.org/10.1007/s10803-025-06884-y.
40. Pretzsch CM, Freyberg J, Voinescu B, Lythgoe D, Horder J, Mendez MA, et al. Effects of cannabidiol on brain
excitation and inhibition systems; a randomised placebo-controlled single dose trial during magnetic resonance
spectroscopy in adults with and without autism spectrum disorder. Neuropsychopharmacol. 2019;44:1398–1405.
41. Isaacson JS, Scanziani M. How Inhibition Shapes Cortical Activity. Neuron. 2011;72:231–243.
42. Salvatore SV, Lambert PM, Benz A, Rensing NR, Wong M, Zorumski CF, et al. Periodic and aperiodic changes to
cortical EEG in response to pharmacological manipulation. J Neurophysiol. 2024. 7 February 2024.
https://doi.org/10.1152/jn.00445.2023.

43. Brake N, Duc F, Rokos A, Arseneau F, Shahiri S, Khadra A, et al. A neurophysiological basis for aperiodic EEG and
the background spectral trend. Nat Commun. 2024;15:1514.

44. Kaplan JS, Stella N, Catterall WA, Westenbroek RE. Cannabidiol attenuates seizures and social deficits in a mouse
model of Dravet syndrome. Proc Natl Acad Sci U S A. 2017;114:11229–11234.

45. Kaplan JS, Wagner JK, Reid K, McGuinness F, Arvila S, Brooks M, et al. Cannabidiol Exposure During the Mouse
Adolescent Period Is Without Harmful Behavioral Effects on Locomotor Activity, Anxiety, and Spatial Memory. Front
Behav Neurosci. 2021;15:711639.

46. Zhang Q, Melchert PW, Markowitz JS. Pharmacokinetic Variability of Oral Cannabidiol and Its Major Metabolites
after Short-Term High-Dose Exposure in Healthy Subjects. Med Cannabis Cannabinoids. 2024;7:1–9.

47. Tayo B, Taylor L, Sahebkar F, Morrison G. A Phase I, Open-Label, Parallel-Group, Single-Dose Trial of the
Pharmacokinetics, Safety, and Tolerability of Cannabidiol in Subjects with Mild to Severe Renal Impairment. Clin
Pharmacokinet. 2020;59:747–755.

48. Millar SA, Stone NL, Yates AS, O'Sullivan SE. A Systematic Review on the Pharmacokinetics of Cannabidiol in
Humans. Front Pharmacol. 2018;9:1365.

49. Ujváry I, Hanuš L. Human Metabolites of Cannabidiol: A Review on Their Formation, Biological Activity, and
Relevance in Therapy. Cannabis Cannabinoid Res. 2016;1:90–101.

50. Brown L, Sherbenou RJ, Johnsen SK. Test of nonverbal intelligence: TONI-4. 2010.

51. Bodfish JW, Symons FJ, Parker DE, Lewis MH. Varieties of repetitive behavior in autism: comparisons to mental
retardation. J Autism Dev Disord. 2000;30:237–243.

52. Dunn LM, Dunn DM. Peabody Picture Vocabulary Test--Fourth Edition. 2012.

53. Martin, Nancy A., and Rick Brownell. Expressive one-word picture vocabulary test-4 (EOWPVT-4). Academic
Therapy Publications; 2011.

54. Beery, K. E., Buktenica, N. A., & Beery, N. A. Beery-Buktenica Developmental Test of Visual-Motor Integration,
Sixth Edition (Beery VMI). Pearson; 2010.

55. Harris CR, Millman KJ, van der Walt SJ, Gommers R, Virtanen P, Cournapeau D, et al. Array programming with
NumPy. Nature. 2020;585:357–362.

56. Virtanen P, Gommers R, Oliphant TE, Haberland M, Reddy T, Cournapeau D, et al. SciPy 1.0: fundamental
algorithms for scientific computing in Python. Nat Methods. 2020;17:261–272.

57. Jas M, Engemann DA, Bekhti Y, Raimondo F, Gramfort A. Autoreject: Automated artifact rejection for MEG and
EEG data. Neuroimage. 2017;159:417–429.

58. Gramfort A, Luessi M, Larson E, Engemann DA, Strohmeier D, Brodbeck C, et al. MNE software for processing
MEG and EEG data. Neuroimage. 2014;86:446–460.

59. Donoghue T, Haller M, Peterson EJ, Varma P, Sebastian P, Gao R, et al. Parameterizing neural power spectra into
periodic and aperiodic components. Nat Neurosci. 2020;23:1655–1665.

60. Skipper S, Perktold J. Statsmodels: Econometric and Modeling with Python. 9th Python in Science Conference,
Austin, 28 June-3 July, 2010, 57-61. Proceedings of the 9th Python in Science Conference. 2010. 2010.
https://doi.org/10.25080/Majora-92bf1922-011.

61. Pion-Tonachini L, Kreutz-Delgado K, Makeig S. ICLabel: An automated electroencephalographic independent
component classifier, dataset, and website. NeuroImage. 2019;198:181–197.

62. Welch P. The use of fast Fourier transform for the estimation of power spectra: A method based on time averaging
over short, modified periodograms. IEEE Transactions on Audio and Electroacoustics. 1967;15:70–73.

63. Li B, Bos MG, Stockmann L, Rieffe C. Emotional functioning and the development of internalizing and externalizing
problems in young boys with and without autism spectrum disorder. Autism. 2020;24:200–210.

64. Mooney RA, Cirillo J, Byblow WD. GABA and primary motor cortex inhibition in young and older adults: a multimodal
reliability study. Journal of Neurophysiology. 2017;118:425–433.

65. Ip IB, Emir UE, Lunghi C, Parker AJ, Bridge H. GABAergic inhibition in the human visual cortex relates to eye
dominance. Sci Rep. 2021;11:17022.

66. Manning JR, Jacobs J, Fried I, Kahana MJ. Broadband shifts in local field potential power spectra are correlated
with single-neuron spiking in humans. J Neurosci. 2009;29:13613–13620.

67. Davis ZW, Muller L, Reynolds JH. Spontaneous Spiking Is Governed by Broadband Fluctuations. J Neurosci.
2022;42:5159–5172.

68. Hill AT, Clark GM, Bigelow FJ, Lum JAG, Enticott PG. Periodic and aperiodic neural activity displays age-dependent
changes across early-to-middle childhood. Dev Cogn Neurosci. 2022;54:101076.

69. Favaro J, Colombo MA, Mikulan E, Sartori S, Nosadini M, Pelizza MF, et al. The maturation of aperiodic EEG
activity across development reveals a progressive differentiation of wakefulness from sleep. NeuroImage.
2023;277:120264.

70. Snipes S. Iota oscillations (25–35 Hz) during wake and REM sleep in children and young adults. Journal of
Neurophysiology. 2025;134:1–9.

71. Chen K-W, Lee Y-C, Yu T-Y, Cheng L-J, Chao C-Y, Hsieh C-L. Test–retest reliability and convergent validity of the
test of nonverbal intelligence-fourth edition in patients with schizophrenia. BMC Psychiatry. 2021;21:39.

72. Pigliasco F, Malaca S, Lo Faro AF, Tini A, Cangemi G, Cafaro A, et al. Cannabidiol, Δ9-tetrahydrocannabinol, and
metabolites in human blood by volumetric absorptive microsampling and LC-MS/MS following controlled
administration in epilepsy patients. Front Pharmacol. 2022;13.
73. Bosl W, Tierney A, Tager-Flusberg H, Nelson C. EEG complexity as a biomarker for autism spectrum disorder risk.
BMC Med. 2011;9:18.
74. Bosl WJ, Tager-Flusberg H, Nelson CA. EEG Analytics for Early Detection of Autism Spectrum Disorder: A data-
driven approach. Sci Rep. 2018;8:6828.
75. Wilkinson CL, Levin AR, Gabard-Durnam LJ, Tager-Flusberg H, Nelson CA. Reduced frontal gamma power at 24
months is associated with better expressive language in toddlers at risk for autism. Autism Research.
2019;12:1211–1224.
76. Li Q, Weiland RF, Konvalinka I, Mansvelder HD, Andersen TS, Smit DJA, et al. Intellectually able adults with autism
spectrum disorder show typical resting-state EEG activity. Sci Rep. 2022;12:19016.
77. Shuffrey LC, Pini N, Potter M, Springer P, Lucchini M, Rayport Y, et al. Aperiodic Electrophysiological Activity in
Preterm Infants is Linked to Subsequent Autism Risk. Dev Psychobiol. 2022;64:e22271.

Acknowledgements

521 We'd like to thank Dillan Cellier, Dr. Sydney Smith, and Dr. Maribel Patiño for their thoughtful discussion.

Author Contributions

523 C.C.: Formal analysis, Investigation, Methodology, Visualization, Writing - original draft, Writing - review and editing.

524 G.P.: Methodology, Data acquisition, Writing - review and editing.

525 A.H.: Writing - review and editing.

526 D.T.: Conceptualization, Project administration, Funding acquisition, Investigation and Writing - review and editing.

527 B.V.: Supervision, Project administration, Funding acquisition, and Writing - review and editing.

Funding Statement

The research study was supported by a grant from the Ray and Ty Noorda Foundation and the Wholistic Foundation to
Dr. Trauner. Purified CBD (Epidiolex®) and identical placebo compounds were provided by GW Pharmaceuticals
(Cambridge, U.K.; now Jazz Pharmaceuticals). C.C. was supported by a NIH/NIMH Blueprint D-SPAN Award (K00
MH132569), a NIH/NIGMS IRACDA Award (K12 GM068524), and the Burroughs Wellcome Fund. A.H. was supported by
a NIH Blueprint BRAIN-ENDURE Award (R25NS119707). Research space for the EEG study was provided by the UC
San Diego Center for Medicinal Cannabis Research, I. Grant, Director. Blood for cannabinoid levels was drawn at the UC
San Diego Altman Clinical and Translational Research Institute (ACTRI). The project was partially supported by the
National Institutes of Health, Grant UL1TR001442 (to the ACTRI). The content is solely the responsibility of the authors
and does not necessarily represent the official views of the NIH.

Conflicts of Interest

539 The authors declare no conflict of interest.

Figure Legends

Figure 1. Overview of study design and spectral parameterization approach. (A) Schematic representation of data
included in our analyses, comprising four study time points for EEG, blood draws, and cognitive-behavioral assessments.
(B) Concentration of metabolite levels in blood post-CBD treatment. **(C)** Duration of post-processed EEG traces across
study time points. **(D)** Schematic of electrodes and electrode groups used in this study. **(G)** Representative power
spectrum of EEG channel 'O2' alongside extracted aperiodic exponent and offset values.

Table 1. Baseline participant characteristics. Raw score and standard score means with standard error of the mean
(SEM), ranges, and floor effects for cognitive and behavioral assessments at baseline. Raw Score N indicates participants
who attempted each assessment, while Standard Score N indicates participants who achieved valid standard scores, with
the gap between these values reflecting performance below normative ranges.

Table 2. Participant assessments included in linear mixed effects models. Mean and standard error of the mean
(SEM) for participant ages and assessment scores, alongside total number of participants (N) and total sessions included
in each linear mixed model (n), across study time points.

Figure Supplement 1. CBD blood metabolite levels and preprocessing of task-free EEG signals. (A) Concentration
of metabolite levels in blood, including CBD, **(B)** 7-OH-CBD and **(C)** 7-COOH-CBD, across all study timepoints. **(D)**
Representative traces of segmented EEG before and after artifact rejection. **(E)** Representative aperiodic exponent and
offset values for all EEG electrode channels in a single session for a single participant.

Figure Supplement 2. Spectral parameterization frequency range comparison. (A) Error distribution comparison
across all channels shows that 0.5-13 Hz fits (teal) produce systematically lower model errors compared to 0.5-50 Hz fits
(orange), with means of ~0.04 and ~0.06 respectively. (B) Box plot comparison showing the consistent reduction in model
error with 0.5-13 Hz frequency restriction. (C) Representative examples demonstrate how 0.5-50 Hz fits (right column)
attempt to model high-frequency artifacts, resulting in high frequency noise-biased aperiodic parameters compared to the
0.5-13 Hz fits (left column).

Figure 2. Relationships between task-free EEG power spectrum features and CBD blood metabolite levels.
Results from linear mixed models and linear regressions relating periodic and aperiodic measures of EEG signals to
levels of CBD, 7-OH-CBD, and 7-COOH-CBD metabolite in blood across groups of electrodes correspondingly located
above **(A, B)** frontal, **(C–E)** central, and **(F–H)** occipital regions. Visualization of each linear mixed model shows the z-
statistic for each model coefficient and a heatmap of the two-tailed p-value associated with the z-statistic. *p<0.05,
**p<0.01, ***p<0.001.

Figure 3. Relationships between cognitive-behavioral assessments and CBD blood metabolite levels. Results from
linear mixed models and linear regressions relating **(A, B)** receptive vocabulary ability, **(C, D)** non-verbal intelligence, and
(E–G) visuomotor coordination assessment scores to levels of CBD, 7-OH-CBD, and 7-COOH-CBD metabolite in blood.
Visualization of each linear mixed model shows the z-statistic for each coefficient and a heatmap of the two-tailed p-value
associated with the z-statistic. #p<0.06, *p<0.05, ***p<0.001.

Figure Supplement 3. Absent relationships between cognitive-behavioral assessments and CBD blood metabolite
levels. Results from linear mixed models and linear regressions relating (**A**) repetitive behavior, **(B)** expressive
vocabulary assessment scores to levels of CBD, 7-OH-CBD, and 7-COOH-CBD metabolite in blood. Visualization of each
linear mixed model shows the z-statistic for each coefficient and a heatmap of the two-tailed p-value associated with the
z-statistic. #p<0.06.

Figure 1

A 8 Weeks | 4 Weeks | 8 Weeks

Baseline | CBD Treatment | Post CBD | Wash | Post-Wash | Placebo Treatment | Post CBD
| Placebo Treatment | Post Placebo | | | CBD Treatment | Post Placebo

Assessment → Assessment → Assessment → Assessment

B Blood metabolite levels (Post-CBD)

Concentration (ng/mL)

CBD OHCBD COOHCBD

C Post-processed EEG duration (seconds)

Baseline Post-Placebo Post-CBD Post-Wash

D EEG channel groups

Frontal

Central

Occipital

E Spectral parameterization (channel 'O2')

log PSD

exponent = 1.42

offset = -10.12

log Frequency (Hz)
1   13

Figure 2

Figure 3

Figure Supplement 1

Figure Supplement 2

A

B

C

Figure Supplement 3

A

B

	Raw Score N	Raw Score Mean (SEM)	Raw Score Range	Standard Score N	Standard Score Mean (SEM)	Standard Score Range	Floor Effects n(%)
PPVT-4 (Receptive Vocabulary)	17	95.8 (14.7)	17 to 204	17	64.1 (6.2)	40 to 115	5 (29%)
TONI-4 (Nonverbal Intelligence)	14	19.3 (3.4)	2 to 37	14	86.8 (4.6)	61 to 121	0 (0%)
EOWPVT-4 (Expressive Vocabulary)	13	59.5 (11.8)	18 to 146	6	87.5 (9.0)	63 to 124	0 (0%)
Beery VMI (Visual-Motor Integration)	21	12.9 (1.6)	0 to 30	14	67.1 (5.8)	45 to 112	1 (7%)
Beery VP (Visual Perception)	21	9.6 (2.2)	0 to 29	7	88.1 (7.2)	62 to 118	0 (0%)
Beery MC (Motor Coordination)	21	11.4 (1.6)	1 to 30	10	64.1 (6.3)	45 to 109	2 (20%)
RBS-R Total (Repetitive Behaviors)	21	59.7 (4.6)	24 to 93	-	-	-	-
ADOS-2 Module	21 (1: 12, 2: 3, 3: 6)	-	-	-	-	-	-

	Baseline	Post-Placebo	Post-Wash	Post-CBD
Total participants (N)	**21**	**23**	**22**	**23**
Age mean	10.43	10.43	10.64	10.63
Age SEM	0.44	0.45	0.48	0.45
EEG participants (n)	21	23	22	23
RBS-R participants (n)	21	23	22	23
RBS-R mean	59.74	45.53	48.45	43.30
RBS-R SEM	4.57	3.67	3.70	3.81
PPVT-4 participants (n)	14	13	15	13
PPVT-4 mean	99.79	100.69	97.47	99.62
PPVT-4 SEM	15.50	16.87	15.81	17.81
TONI-4 participants (n)	12	13	14	13
TONI-4 mean	88.83	88.38	88.50	89.85
TONI-4 SEM	5.11	5.36	5.34	5.39
EOWPVT-4 participants (n)	13	13	12	12
EOWPVT-4 mean	59.54	57.92	57.17	61.50
EOWPVT-4 SEM	11.80	11.95	11.73	12.80
Beery-VMI participants (n)	21	23	22	21
Beery-VMI mean	12.90	13.87	14.45	13.57
Beery-VMI SEM	1.59	1.33	1.36	1.64
Beery-VP participants (n)	21	23	22	21
Beery-VP mean	9.62	11.04	12.32	9.90
Beery-VP SEM	2.19	2.27	2.27	2.29
Beery-MC participants (n)	21	23	22	21
Beery-MC mean	11.38	11.52	10.73	10.67
Beery-MC SEM	1.56	1.34	1.25	1.42