Title: Therapist perspectives on the clinical utility of hand performance information from at-home egocentric video in outpatient neurorehabilitation: A multi-methods evaluation study

Author names and affiliations:

Adesh Kadambi, BEng[1,2], Damian M. Manzone, PhD[1] and José Zariffa, PhD, PEng[1,2,3,4]

[1] KITE - Toronto Rehabilitation Institute, University Health Network, Toronto, Ontario, Canada;
[2] Institute of Biomedical Engineering, University of Toronto, Toronto, Ontario, Canada;
[3] Rehabilitation Sciences Institute, Temerty Faculty of Medicine, University of Toronto, Toronto, Ontario, Canada;
[4] Edward S. Rogers Sr. Department of Electrical and Computer Engineering, University of Toronto, Toronto, Ontario, Canada

Corresponding author(s):

Adesh Kadambi, BEng (adesh.kadambi@mail.utoronto.ca)
José Zariffa, PhD, PEng (jose.zariffa@utoronto.ca)

Abstract:

Background: Restoring hand function is a primary focus of rehabilitation after neurological injuries, such as stroke and spinal cord injury. However, monitoring hand use outside the clinic remains challenging. This study aims to evaluate how therapists perceive and would utilise information from a clinical decision support system (CDSS) that uses egocentric video to monitor patients' hand use at home.

Methods: Five patient-therapist dyads were recruited. Patients recorded daily activities using head-mounted cameras. Therapists reviewed dashboards of processed video data from their patients and completed semi-structured interviews and structured questionnaires. A multi-methods approach with thematic analysis was used to evaluate the CDSS's clinical usefulness.

Results: Qualitative analyses revealed four main themes: therapists strongly preferred observational video data over quantitative metrics, valued insights into home environments that cannot be captured in-clinic, identified ways the information could tailor therapy to real-world activities, but noted practical implementation challenges including time constraints and patient burden. The system achieved high usability scores (mean SUS: 80.0, 85-89th percentile).

Conclusions: This study demonstrates the potential of using egocentric video to inform clinical decision-making in neurorehabilitation, particularly for hand function. The strong preference for video over metrics suggests clinical decision support systems should prioritize interpretable, observation-based information aligned with clinical reasoning processes. Despite implementation challenges, therapists across technical familiarity levels expressed trust in the system and willingness to use it regularly. These findings indicate that egocentric video systems can bridge the clinic-home divide when designed to match interest-holder priorities.

Keywords

egocentric video; home monitoring; upper limb rehabilitation; stroke; spinal cord injury; usability; clinical decision support; wearable technology

Introduction

Neurological injuries and conditions commonly result in substantial loss of upper limb function, which greatly limits an individual's ability to perform activities of daily living (ADLs). Across diverse neurological populations, impaired hand function consistently ranks as a primary rehabilitation concern. For instance, individuals with spinal cord injury (SCI) consistently report regaining hand and arm function as the top recovery priority [1]. Similarly, for individuals who have experienced a stroke, data suggest upper-limb strength is a major predictor of quality of life [2]. Therefore, restoring hand function is a primary focus of rehabilitation efforts, as it directly impacts the ability to regain independence and quality of life.

Regular and accurate monitoring of hand function is essential for promoting recovery [3]. It enables clinical assessment and progress tracking, thereby allowing therapists to adjust treatment plans, and ensure therapy addresses real-world functional needs [3]. However, monitoring this function outside of the clinic remains a significant challenge [4, 5]. Conventional outpatient therapy is typically limited to face-to-face assessments and patient self-reports, which often fail to capture the complexity and variability of real-life hand use at home [6–9]. Patient self-reporting also relies on an individual's ability to recall activities performed since their last therapy session and is subject to reporting biases, such as cognitive deficits and social desirability [10–13]. These limitations prevent clinicians in outpatient settings from gaining a comprehensive understanding of how patients perform ADLs in their everyday environments, thus hindering the ability to tailor therapy plans on the basis of real-world functional performance.

Wearable technology offers an avenue for measuring hand function outside of the clinic. In particular, head-mounted egocentric cameras can provide rich, contextual insights into patients' hand function during ADLs [14–16]. However, clinicians have shown reluctance to adopting new technologies in their practices, citing concerns about time constraints and uncertainty regarding clinical value [5, 17–19]—factors identified by the Technology Acceptance Model [20] as primary barriers to adoption [21]. Digital tools, such as clinical decision support systems (CDSS), have demonstrated success in implementing evidence-based guidelines at the point of care and improving patient outcomes in various healthcare domains [22], including rehabilitation [23, 24], chronic disease management [25], and prescription and medication safety [26]. However, this success hinges on whether they present information that is useful, interpretable, and actionable for clinicians [27–30]. Therefore, understanding how clinicians perceive and value specific types of information from wearable devices is crucial for developing systems that address time and value concerns that have previously hindered the adoption of new technologies in clinical practice.

We recently co-designed a CDSS dashboard with clinicians to present information about patient hand use captured through egocentric video recordings in multiple formats, including quantitative metrics and video footage of functional activities [31]. Our current study aims to understand how this information from egocentric videos, presented through the dashboard interface [31], is perceived by therapists in the context of their clinical practice and decision making. Specifically, we sought to evaluate whether therapists find the information useful for better understanding patient hand performance at home and whether it could inform clinical decision-making. These aspects have not been explored in prior work with egocentric video in rehabilitation, which has primarily revolved around developing research tools to measure outcomes in clinical studies [15, 16, 32] and exploring patient perspectives [33, 34]. Due to the need for us to investigate

how this data might be useful to therapists in clinical practice, we evaluated this system with therapists treating patients with stroke and spinal cord injury as representative examples of acquired neurological conditions affecting upper limb function. Through this work, we seek to bridge the gap between digital health tool development and practical clinical application, ensuring that innovations can meaningfully enhance patient outcomes in neurorehabilitation settings.

Methods

This study was approved by the Research Ethics Board of the University Health Network (UHN; Study #21-5019) and conducted in accordance with the Tri-Council Policy Statement. We employed a multi-methods design to evaluate the clinical usefulness of the information presented to therapists [35]. Qualitative data from semi-structured interviews were analysed using a thematic analysis to explore therapist perspectives in depth. Quantitative data from structured questionnaires and the System Usability Scale (SUS) were analysed using descriptive statistics. These two data sources were analysed separately, with quantitative findings used to corroborate and contextualize qualitative themes. This approach allowed for a comprehensive understanding of the system's clinical utility and usability.

Participants

We recruited 5 patient-therapist dyads from the Toronto Rehabilitation Institute at UHN using a snowball sampling approach. This sample size was deemed appropriate for this study due to its specific research question and a single analytically relevant participant category (i.e., therapists assessing clinical utility) [36]. Therapist-participants were eligible if they had expertise in upper limb rehabilitation and were actively treating patients who were either adult stroke survivors or adults living with SCI. Patient-participants were eligible if they were over 18 years of age, actively receiving therapy at UHN from the therapist-participant, and had self-reported impairment of hand function from SCI or stroke while retaining the ability to use their hands in some capacity for at-home ADLs.

The therapist group included 4 occupational therapists and 1 physical therapist with 13.8 ± 5.5 years of clinical experience (range: 8-20 years). All therapists were female. The patient group consisted of 2 stroke survivors and 3 individuals living with SCI. This mix of stroke and SCI patients allowed us to explore the system's applicability across different types of upper limb impairments. All participants were assigned an alphanumeric code to anonymize transcript data and any excerpts in this article.

Data Collection

Our data collection process followed a three-stage pipeline consisting of patient video recordings, dashboard generation, and therapist interviews, as illustrated in Figure 1.

107
108 Figure 1: Overview of the data collection pipeline. Patient-participants recorded videos at home using a
109 head-mounted egocentric camera. A dashboard was generated from the recordings, which included both
110 quantitative metrics and manually curated video snippets. Therapist-participants reviewed the dashboard
111 generated for their patient for approximately 10 minutes before proceeding with the interview.

Video Recordings

Patient-participants were provided with head-mounted egocentric cameras (GoPro Hero 5, GoPro, Inc., San Mateo, CA, USA) to record their ADLs at home. The cameras recorded videos in .mp4 format at 1080p resolution and 30 frames per second. Patient-participants were instructed to record 1-hour sessions between 2-4 times per week for a minimum of 2 weeks. Following a previously reported protocol, recording times were agreed on by the investigator and participant in an initial interview [37]. The recording periods were selected to capture times when patient-participants performed tasks that were informative about their hand function, part of their regular routine, and did not raise privacy concerns. Examples included eating, food preparation, household tasks (e.g., laundry, putting away groceries, writing), or non-private grooming tasks (e.g., brushing teeth, combing hair). Participants had full control to start and stop recordings and could edit videos before returning them to the research team. The patient-participants recorded an average of 231.85 ± 75.49 minutes of video. Once the recordings were completed, the patient-participants returned the cameras to the researchers.

Dashboard Generation

The recorded videos were processed to generate a dashboard [31] that provided therapists with information about their patient's hand use at home. The complete dashboard generation and video processing pipeline is illustrated in Figure 2. Raw videos were segmented into 1-minute clips and sampled at 2 frames per second. We applied existing computer vision models to detect objects [38] and identify hand-object interactions (HOI) [39]. These detections were used to compute quantitative metrics and classify activities. *Minutes recorded per day* was computed using the metadata from the videos. HOI detections were used to compute *percentage interaction* per day (i.e., proportion of frames containing detected interactions), *number of interactions* per hour (i.e., distinct hand-object engagement events, defined as consecutive frames with detected interactions), and *average interaction duration* (i.e., temporal length of interaction events in seconds) [16]. The *activity breakdown* was generated using an ADL classification model [40] to categorize videos into seven ADL categories, and then manually verified for accuracy. *Video snippets* were manually curated from the 1-minute video segments by AK, with an intentional focus on tasks the patient struggled to complete or performed with compensatory strategies, as

139   these were deemed most clinically relevant for informing therapy planning through prior conversations with

140   therapists.

Figure 2: Dashboard generation pipeline. Raw egocentric videos were segmented into 1-minute clips and sampled at 2 frames per second. Object detection and HOI models identified active hand-object interactions. These features were used to automatically classify ADLs [40]. A sample dashboard for a patient-participant with synthetic data is displayed below the processing pipeline. Blue boxes show

146 *dashboard metrics and their derivation points. The red lines point to the specific user interface elements for*
147 *each metric.*

Therapist Interviews

The final stage of data collection involved therapist-participants reviewing and providing feedback on the
CDSS. This process began with a 10-minute review period where the therapist could familiarise themselves
with the dashboard and review the information provided for their patient. This was followed by a 45-minute
interview session consisting of structured questionnaires and a semi-structured interview. The structured
questionnaires included 5-point Likert scale questions to assess technological affinity and perceptions of
clinical utility and actionability, as well as a SUS questionnaire [41] to measure the usability of the CDSS.
The SUS is a widely validated, reliable 10-item scale for evaluating system usability across diverse
applications [42]. The semi-structured interview gathered in-depth qualitative feedback on the usefulness
of the information and its influence on clinical decision-making. Four of the five patient-participants had
completed their therapy block between one and three months prior to the therapist interview, while one
remained in active therapy. This retrospective evaluation allowed therapists to reflect on how the
information could have informed their clinical decision-making. All interviews were conducted by AK using
an interview guide developed for this study and provided in the supplementary material.

Data Analysis

We analysed qualitative and quantitative data separately and then presented findings together to provide a
comprehensive understanding of the system's clinical utility and usability. The data from the semi-
structured interviews were transcribed verbatim and analysed using reflexive thematic analysis [43, 44].
Reflexive thematic analysis is an iterative, interpretive approach where themes are actively constructed by
researchers through engagement with the data [44, 45]. This process involved two independent
researchers (AK and DMM) coding the data to identify key themes and patterns related to the clinical
usefulness and actionability of the information presented in the dashboard. Following initial coding, the
researchers engaged in consensus coding, where they met to compare their findings, discuss any
discrepancies, and reach agreement on the final set of themes and codes. JZ acted as an arbitrator for any
unresolved discrepancies. This approach was employed to reduce individual bias and enhance the
reliability and credibility of the analysis. The Likert scale questionnaire responses were analysed using
descriptive statistics. The SUS scores were used to assess the usability of the dashboard format for
presenting this information using the curved grading scale interpretation [46]. Quantitative findings were
used to corroborate and contextualize the qualitative themes identified through thematic analysis.

Results

Thematic Analysis

179 Four main themes were constructed through our reflexive thematic analysis: (1) Data Interpretation
180 Preferences, (2) Bridging the Clinic-Home Divide, (3) Tailoring Therapy Through Home Observations, and
181 (4) Practical Realities of Implementation. Table 1 presents an overview of these themes and their
182 subthemes.

183 *Table 1: Overview of themes and subthemes for thematic analysis from semi-structured interviews.*

Theme	Subthemes
Data Interpretation Preferences	1A. The Primacy of Functional Footage 1B. Navigating Novel Graph Presentations
Bridging the Clinic-Home Divide	2A. Beyond Clinic Walls 2B. Goal-Oriented Data Collection
Tailoring Therapy Through Home Observations	
Practical Realities of Implementation	4A. Clinical Integration 4B. Another Burden?

1. Data Interpretation Preferences

This theme explores how therapists interacted with and valued the different types of data presented to
them in the dashboard (i.e., graphs and videos). It highlights the distinction between more familiar,
observation-based representations of data and novel data visualisation techniques.

1A. The Primacy of Functional Footage

This subtheme captures therapists' strong preference for video data and its ability to provide rich,
functional insights into patients' real-world activities, emphasising its alignment with how they already
consume data they are presented with in the clinic (i.e., through patient observations). As one therapist
highlighted:

I would go right to the videos. The graphs were interesting in that I saw that he looked like he used his
left hand more, but that probably wouldn't give me that much information. I think the videos are the
most telling, at least for me. Visual, more direct application for me. (T05, OT)

This sentiment was also echoed by others: "To be honest, I really liked the videos, because that, as much
as the number tells you, it doesn't actually tell you the quality and the pattern that she's actually using her
hand" (T01, PT).

Overall, while the graphical data could be a good overview, therapists believed the real value came from
the videos: "The quantitative measures would give me an idea of how much he's progressing in terms of
quantity of affected hand use, [...] and then just yeah the videos give me a lot of rich information on how I
can adapt therapy" (T04, OT).

1B. Navigating Novel Graph Presentations

In contrast, therapists expressed challenges interpreting the information presented in the graphs and found
it difficult to integrate into their clinical reasoning or therapy planning in any way:

Normally I don't go through metrics or graphs or, this isn't a typical format for me to obtain or analyse
information being more like a frontline clinician. So just again, my eyes aren't as familiar with this as
from what my role has been in the past, like 20 years here. This isn't a regular part of how we're

interpreting our patient information. [...] I think there just needs to be more specification around [the
graphs] or just more defining parameters just for it to have any value. [...] like if we're using this at
[intake] and then discharge, I would want it to be as comparable as possible. (T02, OT)

2. Bridging the Clinic-Home Divide

This theme explores the unique insights that therapists gained from the CDSS about patient home
environments and daily activities beyond the clinical setting. It also highlights different data collection
approaches to make the information more useful to therapists.

2A. Beyond Clinic Walls

Therapists emphasised the value of seeing videos of their patients performing functional activities at home
"because the clinic is limited as a simulation of what occurs at home" (T04, OT). And since they "cannot go
to every patient's home, in a way [these videos] bring the patient's home to [them]" (T03, OT). As one
therapist noted:

Part of our OT assessment does include the physical environment, so we are asking about obviously
the basic layout, and how you're getting around, and how you're accessing different rooms, but as
far as, a lot of times the physical environment also does just limit the potential for how much they can
do. So, yeah, that's just an extra layer I would say. But as far as thinking about whatever
recommendations we might make for home modifications, or even how he most safely accesses his
space, it's just that opportunity to see a patient, again, outside of a very sort of generic clinic space or
setting. (T02, OT)

2B. Goal-Oriented Data Collection

While patients were instructed to record typical daily activities following the protocol outlined in [37],
therapists expressed a preference for focused data collection aligned with patients' specific goals and
priorities, while also being able to compare how a patient is able to perform different functional activities
across time points to better track progress over time:

It would be useful to me to be able to see the change from one time-point to the next when it's... like
if this was a pre-therapy video, then I would want to be able to compare it readily to mid-therapy and
post-therapy video. And I guess it would be nice to have some alignment between what his functional
goals are, because for instance one of his goals initially was to be able to like open a ring box so I
would have him videotape at home like how he's performing that goal, or his approximation of that
goal, like opening a box at home. I would want to see how he's progressing with things that are part
of his goals. (T04, OT)

3. Tailoring Therapy Through Home Observations

This theme discusses how the new insights gained from the CDSS could be used to adjust therapist
recommendations and therapy plans based on observed real-world patient behaviour. One therapist

mentioned the videos provide much greater detail than their initial assessments and could help with task
specific practice:

> 246 We have like tedious notes as far as our initial assessment, but we would still certainly not be able to
> 247 get that level of like, "So, okay, when you go in your kitchen...". This really gives, if we want to break
> 248 down the tasks even more, it definitely would lend itself really nicely to helping with more specific
> 249 practice therapy sessions, or breaking down the task and say, "Okay, now let's work on your left
> 250 hand reach and then reaching across your body with your right hand to open the microwave", or just
> 251 some of the details definitely could help with the problem solving or maybe modified devices. (T02,
> 252 OT)

253 Another therapist described how some of the activities she observed in the videos could be a safety hazard
254 and that they would modify exercises in clinic to prevent a fall in the future:

> 255 I love it [...] you open my horizon as a therapist. Wow, [patient name] is doing some... one big help
> 256 your project is... would be helpful in falls prevention because this patient is taking risk. I know the left
> 257 leg is not working, so why not move your walker closer and then you grasp... and if she had... so I'm
> 258 gonna modify my therapy by looking at those and improve my treatment so your project definitely will
> 259 help us to improve the results. (T03, OT)

260 **4. Practical Realities of Implementation**
261 This theme explores the practical considerations and potential barriers to implementing the dashboard in
262 clinical practice. It encompasses concerns about time constraints for clinicians and the additional burdens
263 placed on patients, highlighting the complexities of integrating new systems into existing clinical processes.
264

265 *4A. Clinical Integration*

266 Therapists expressed varied perspectives on how they could use this system in their current practice. One
267 therapist viewed the tool as being akin to reviewing patient charts in the electronic health record (i.e., Epic)
268 prior to a session:

> 269 I'd prefer to first look at it when I'm not in therapy. I think it's very quick to extract information from
> 270 this, so I don't think this takes me any longer than you know, logging into Epic and doing all these
> 271 things I have to do to document the session. [...] Yeah, I don't think it's too time-consuming. I think it's
> 272 very valuable information. (T04, OT)

While another believed they did not "have a ton of time for it" (T02, OT), but considered "[looking] at it with
the patient and [reviewing] it during the session" (T02, OT). As a result, one therapist questioned how often
they would even use it: "I don't know how often I would do it. If it was available, if there was something you
really wanted to know, how are you actually doing this? Then I could see that coming into play." (T05, OT).

4B. Another Burden?

Therapists also mentioned potential challenges for patients:

So for someone who doesn't really have any support, and for someone who doesn't have support
and a low level arm, like there's no way they're gonna get a GoPro on their head and strap it properly
and turn it on and make sure it's on and off, and you're not gonna get them to do that for four weeks.
So I think it's, it's great if, if you pick the right person to do it with, if not, I don't know how helpful it
would be. (T01, PT)

Quantitative Findings

In addition to the qualitative interviews, therapists completed a questionnaire about technology affinity and
the clinical usefulness of this system in their practice (Figure 3). The first six questions aimed to understand
a therapist's affinity towards technology, and were answered before the review session began, while the
remaining questions targeted their thoughts on our CDSS after the review period concluded. Regardless of
therapists having varying levels of technology use and affinity in their lives (Q1-6), they all found the CDSS
to be a useful tool for understanding patient performance at home (Q12), would use it regularly (Q7), and
indicated that it had the potential to influence therapy planning (Q15).

Figure 3: Therapist responses to the questionnaire regarding technology use and affinity (Q1-Q6) and clinical utility and actionability of the CDSS (Q7-Q16). Mean Likert scale scores are represented by black filled circles and the responses from each participant are represented by individually connected coloured

299 Our participants reported a mean SUS score of 80.0 (95% CI [65.95, 94.05]), placing our system in the 85-

300 89th percentile range [46]. All individual participants scored above 68 (Figure 4), which represents the 50th

301 percentile SUS score [46]. These results indicate above-average usability for our system overall.

Figure 4: System Usability Scale (SUS) results. The left panel displays the total SUS scores. The mean
score is represented by a black circle with error bars representing 95% CI. Individual participant scores are
represented by the smaller grey circles. The dashed line is displayed at a total SUS score of 68, which is
considered the average score at the 50th percentile [46]. The right panel displays the distribution of
responses for all participants and for all statements. Note that the titles over each graph have been
abbreviated for simplicity.

Discussion

This study aimed to evaluate whether information about hand use in the home environment derived from

egocentric video was perceived by therapists as being clinically useful in the context of outpatient upper

limb neurorehabilitation. The information presented was well-received and gave therapists the ability to

view patients performing functional activities and ADLs in their home environment, with the potential to

result in more tailored therapy in the clinic. However, there were clear preferences for specific types of
data and practical challenges for implementation.

Therapists consistently expressed a strong preference for video footage over graphical representations of
data. The videos enabled therapists to observe real-world hand use and understand patients' functional
capabilities in their home environments, allowing them to adapt therapy based on direct observation of
performance. For example, therapists reported the videos helped them break down tasks into specific
movements performed at home, or provide more specific recommendations based on safety concerns they
observed in the videos. This preference is supported by the high Likert scores related to the dashboard's
usefulness in assessing patient performance (Q7, Q12). This aligns with existing literature that underscores
the importance of direct observation in clinical decision-making, where functional video footage provides
critical context, such as movement quality, that is often lost in abstract numerical data [47, 48].

Conversely, several therapists found the graphical data more challenging to interpret and reported that the
metrics were unclear and less directly applicable to their clinical reasoning. Concerns were raised about
how quickly someone could learn how to use this information (Q14), suggesting a lack of clarity in how the
data were presented, and its potential to disrupt workflow (Q10). This preference for video over graphical
data likely reflects both the familiarity of video-based observation in clinical practice and the need for more
intuitive data visualization design.

Our prior work [16] had focused on developing novel outcome measures for clinical research and
demonstrated that quantitative metrics like *percentage interaction* and *number of interactions* were
correlated to higher Upper Extremity Measurement Scale (UEMS) [49], Spinal Cord Independence Measure
III (SCIM) [50], and Graded Redefined Assessment of Strength, Sensibility and Prehension (GRASSP) [51]
scores. However, therapists in the current study found these very metrics less clinically actionable than raw
video footage. This finding highlights the different considerations involved in using novel assessments as
tools for clinical research or for direct care provision. For the therapist-participants in this study, the most
useful aspect of this system was its ability to bridge the gap between the clinical and home environments.
Therapists unanimously agreed that the clinic could never fully simulate the complexity of patients' home
environments and while they "cannot go to every patient's home, in a way [these videos] bring the patient's
home to [them]" (T03, OT). Rather than presenting more sophisticated metrics, future systems may need to
facilitate efficient navigation and interpretation of video content itself to better align with clinicians' existing
workflows and mental models.

An important consideration in developing home monitoring systems is ensuring alignment between patient
and clinician perspectives. Our prior work has systematically explored these perspectives, first
investigating patient acceptability and preferences [33, 34], and now examining therapist perspectives on
clinical utility. The perspectives of patients from these prior studies and therapists from our current study
converged on recognizing the fundamental value of home observation, indicating that clinic settings cannot
fully capture home function, thereby supporting the core premise of home monitoring. Videos of patients
performing functional activities at home gave therapists crucial insights into the physical and environmental
barriers that patients encounter while performing real activities, which might not be evident in the controlled

setting of the clinic. This ability to remotely observe home environments offers a promising alternative to traditional home visits, which are often limited by time and resource constraints even though they result in better patient outcomes [52–55]. The high score for statements indicating trust in the information (Q9) and the belief that it would influence future therapy plans (Q15) demonstrates the overall utility of our system in clinical settings.

However, despite recognizing the clinical value of the information provided by this system, therapists also highlighted several practical challenges related to its implementation such as clinical time constraints and patient burden. While some therapists felt the dashboard was easy to integrate into their existing chart review time, others worried that reviewing the data would add too much to their already busy schedules (Q10). There were also concerns about the burden on patients with limited physical or technological support, which aligns with previous findings, where individuals living with SCI and stroke expressed preferences for cameras that were less noticeable and lighter to minimise self-consciousness and discomfort [33, 34]. Some therapists also noted that certain patients might struggle to use the necessary equipment to capture their daily activities. These concerns echo the existing broader implementation science literature on barriers to adoption of mobile health and remote patient monitoring devices in healthcare, with disruption to clinician workflows and patient training and/or willingness to use these technologies playing the largest role in adoption rates [56–59]. Therefore, future iterations will explore ways to automatically highlight interesting video snippets for therapists to view. Additionally, future work should aim to design egocentric cameras that are easier to use by individuals with hand impairments and increase their willingness to use and record (i.e., lighter, less bulky, etc.).

Limitations

While this study provides valuable insights into the clinical potential of the information presented by the CDSS, there are several limitations to consider. First, the small sample size of only five clinicians from a single hospital network limits the generalizability of our findings. This sample size was partly constrained by the number of eligible participants at our institution and the complexity of recruiting patient-therapist dyads, which is more challenging than recruiting therapists alone. Despite this limitation, the choice of 5 dyads was reasonable given that this study has a specific research question and a single analytically relevant participant 'category' (i.e., therapists) [36]. All participating therapists were female, and while we acknowledge that sex/gender may have influenced the findings of this study, our sample reflects the female-dominant nature of the field [60, 61]. Detailed patient demographics (e.g., age, time since injury, level of impairment) were also not collected, as this study focused on therapist perspectives regarding the clinical utility of the information system, evaluated in a manner agnostic to patient-specific characteristics.

Additionally, there is a potential for selection bias, as therapists more inclined to engage with and adopt new technologies may have been more likely to participate. While the quantitative questionnaire showed a wide range of responses in the technology affinity section (Q1-6), suggesting a variety of comfort levels with technology, it's noteworthy that even therapists with lower technology affinity provided positive feedback on the CDSS. This indicates that the CDSS's potential value was recognized across different levels of technological comfort. However, it's important to acknowledge that the overall positive reception

may not fully capture the perspectives of therapists who are generally less comfortable with technology
and chose not to participate in the study.

Finally, although the CDSS was well-received in interviews, therapists did not have extended opportunities
to integrate it into their daily practice. The video snippets were also intentionally curated to focus on tasks
patients struggled with. This approach may have biased the presentation toward deficits rather than
abilities, though therapists could also view the quantitative metrics which captured all hand use regardless
of difficulty. This limited exposure means that the feedback gathered may not fully capture the tool's long-
term usability or the practical challenges it might pose once embedded in everyday workflows.

Conclusion

This study demonstrates the potential of a system that uses egocentric video to provide therapists with
home-based hand use data about their patients to inform clinical decision-making in a neurorehabilitation
setting. While therapists found the information useful—particularly video data for assessing patient
performance in a naturalistic setting—challenges related to integrating the system into clinical workflows
were evident. The primary concerns centred around the clarity of graphical data and the perceived time
burden that regular use might impose on therapists. Despite these hurdles, therapists across varying levels
of technical familiarity expressed trust in the information provided by the system, indicated that they would
use it regularly to assess patient performance, and believed it could enhance their ability to tailor therapy
plans to real-world patient activities.

Reducing time demands through features like interpretations of graphical data or providing more emphasis
on video data may help mitigate the barriers identified. Ultimately, the adoption of wearable egocentric
cameras could facilitate a more seamless connection between face-to-face assessments and real-world
functionality, contributing to more effective and personalised rehabilitation. However, further research is
necessary to explore the long-term impact of this system in diverse clinical environments and its role in
broader healthcare settings.

List of abbreviations

SCI - Spinal Cord Injury
ADLs - Activities of Daily Living
CDSS - Clinical Decision Support System
UHN - University Health Network
SUS - System Usability Scale
OT - Occupational Therapist
PT - Physical Therapist
425
426

Declarations

Ethics approval and consent to participate

This study was approved by the Research Ethics Board of the University Health Network (Study #21-5019) and conducted in accordance with the Tri-Council Policy Statement: Ethical Conduct for Research Involving Humans (TCPS2). Informed consent was obtained from all individual participants included in the study.

Consent for publication

Not applicable

Availability of data and materials

The datasets generated and/or analysed during the current study are not publicly available due to privacy concerns regarding identifying information in home-recorded videos and interview transcripts but are available from the corresponding author on reasonable request.

Competing Interests

The authors declare that they have no competing interests.

Funding

This work was supported by the Praxis Spinal Cord Institute; the Ontario Early Researcher Award program under grant number ER16–12-013; and the Canadian Institutes of Health Research under grant number 13556838.

Authors' contributions

AK recruited all participants, conducted and transcribed all of the interviews, performed data analysis, and drafted the manuscript. DMM contributed to data analysis, created figures, and revised the manuscript. JZ contributed to study design and data interpretation, and revised the manuscript. All authors read and approved the final manuscript.

Acknowledgements

Not applicable

Clinical Trial Number

Not applicable

References

1. Anderson KD. Targeting recovery: priorities of the spinal cord-injured population. J Neurotrauma. 2004;21:1371–83.

2. van Lieshout ECC, van de Port IG, Dijkhuizen RM, Visser-Meily JMA. Does upper limb strength play a prominent role in health-related quality of life in stroke patients discharged from inpatient rehabilitation? Top Stroke Rehabil. 2020;27:525–33.

3. Burridge J, Alt Murphy M, Buurke J, Feys P, Keller T, Klamroth-Marganska V, et al. A systematic review of international clinical guidelines for rehabilitation of people with neurological conditions: What recommendations are made for upper limb assessment? Front Neurol. 2019;10:567.

4. Singh H, Musselman K, Colella TJF, McGilton KS, Iaboni A, Bayley M, et al. Exploring the perspectives of outpatient rehabilitation clinicians on the challenges with monitoring patient health, function and activity in the community. Disabil Rehabil. 2022;44:2858–67.

5. Jones M, Collier G, Reinkensmeyer DJ, DeRuyter F, Dzivak J, Zondervan D, et al. Big Data Analytics and Sensor-Enhanced Activity Management to Improve Effectiveness and Efficiency of Outpatient Medical Rehabilitation. Int J Environ Res Public Health. 2020;17.

6. Waddell KJ, Strube MJ, Bailey RR, Klaesner JW, Birkenmeier RL, Dromerick AW, et al. Does Task-Specific Training Improve Upper Limb Performance in Daily Life Poststroke? Neurorehabil Neural Repair. 2017;31:290–300.

7. Doman CA, Waddell KJ, Bailey RR, Moore JL, Lang CE. Changes in upper-extremity functional capacity and daily performance during outpatient occupational therapy for people with stroke. Am J Occup Ther. 2016;70:7003290040p1–7003290040p11.

8. Rand D, Eng JJ. Disparity between functional recovery and daily use of the upper and lower extremities during subacute stroke rehabilitation. Neurorehabil Neural Repair. 2012;26:76–84.

9. Fleming MK, Newham DJ, Roberts-Lewis SF, Sorinola IO. Self-perceived utilization of the paretic arm in chronic stroke requires high upper limb functional ability. Arch Phys Med Rehabil. 2014;95:918–24.

10. Adams SA, Matthews CE, Ebbeling CB, Moore CG, Cunningham JE, Fulton J, et al. The effect of social desirability and social approval on self-reports of physical activity. Am J Epidemiol. 2005;161:389–98.

11. Adams AS, Soumerai SB, Lomas J, Ross-Degnan D. Evidence of self-report bias in assessing adherence to guidelines. Int J Qual Health Care. 1999;11:187–92.

12. Tatemichi TK, Desmond DW, Stern Y, Paik M, Sano M, Bagiella E. Cognitive impairment after stroke: frequency, patterns, and relationship to functional abilities. J Neurol Neurosurg Psychiatry. 1994;57:202–7.

13. Sachdeva R, Gao F, Chan CCH, Krassioukov AV. Cognitive function after spinal cord injury: A systematic review. Neurology. 2018;91:611–21.

14. Cartas A, Radeva P, Dimiccoli M. Activities of Daily Living Monitoring via a Wearable Camera: Toward Real-World Applications. IEEE Access. 2020;8:77344–63.

15. Tsai M-F, Wang RH, Zariffa J. Validity of novel outcome measures for hand function performance after stroke using egocentric video. Neurorehabil Neural Repair. 2023;37:142–50.

16. Bandini A, Dousty M, Hitzig SL, Craven BC, Kalsi-Ryan S, Zariffa J. Measuring Hand Use in the Home after Cervical Spinal Cord Injury Using Egocentric Video. J Neurotrauma. 2022;39:1697–707.

17. Lang CE, Barth J, Holleran CL, Konrad JD, Bland MD. Implementation of wearable sensing technology for movement: Pushing forward into the routine physical rehabilitation care field. Sensors (Basel). 2020;20:5744.

18. Dunn J, Runge R, Snyder M. Wearables and the medical revolution. Per Med. 2018;15:429–48.

198 19. Morris J, Jones M, Thompson N, Wallace T, DeRuyter F. Clinician Perspectives on mRehab Interventions
199 and Technologies for People with Disabilities in the United States: A National Survey. Int J Environ Res
Public Health. 2019;16.

20. Holden RJ, Karsh B-T. The technology acceptance model: its past and its future in health care. J
Biomed Inform. 2010;43:159–72.

21. Wu J-H, Wang S-C, Lin L-M. Mobile computing acceptance factors in the healthcare industry: a
structural equation model. Int J Med Inform. 2007;76:66–77.

22. Sutton RT, Pincock D, Baumgart DC, Sadowski DC, Fedorak RN, Kroeker KI. An overview of clinical
decision support systems: benefits, risks, and strategies for success. NPJ Digit Med. 2020;3:17.

23. Qin Z, Armijo-Olivo S, Woodhouse LJ, Gross DP. An investigation of the validity of the Work
Assessment Triage Tool clinical decision support tool for selecting optimal rehabilitation interventions for
workers with musculoskeletal injuries. Clin Rehabil. 2016;30:277–87.

24. Chae SH, Kim Y, Lee K-S, Park H-S. Development and Clinical Evaluation of a Web-Based Upper Limb
Home Rehabilitation System Using a Smartwatch and Machine Learning Model for Chronic Stroke
Survivors: Prospective Comparative Study. JMIR Mhealth Uhealth. 2020;8:e17216.

25. Sperl-Hillen JM, Rossom RC, Kharbanda EO, Gold R, Geissal ED, Elliott TE, et al. Priorities Wizard:
Multisite web-based primary care clinical decision support improved chronic care outcomes with high use
rates and high clinician satisfaction rates. EGEMS (Wash, DC). 2019;7:9.

26. Shahmoradi L, Safdari R, Ahmadi H, Zahmatkeshan M. Clinical decision support systems-based
interventions to improve medication outcomes: A systematic literature review on features and effects. Med
J Islam Repub Iran. 2021;35:27.

27. Trinkley KE, Kahn MG, Bennett TD, Glasgow RE, Haugen H, Kao DP, et al. Integrating the Practical
Robust Implementation and Sustainability Model With Best Practices in Clinical Decision Support Design:
Implementation Science Approach. J Med Internet Res. 2020;22:e19676.

28. Newton N, Bamgboje-Ayodele A, Forsyth R, Tariq A, Baysari MT. A systematic review of clinicians'
acceptance and use of clinical decision support systems over time. NPJ Digit Med. 2025;8:309.

29. Khairat S, Marc D, Crosby W, Al Sanousi A. Reasons for physicians not adopting clinical decision
support systems: Critical analysis. JMIR Med Inform. 2018;6:e24.

30. Camacho J, Zanoletti-Mannello M, Landis-Lewis Z, Kane-Gill SL, Boyce RD. A Conceptual Framework to
Study the Implementation of Clinical Decision Support Systems (BEAR): Literature Review and Concept
Mapping. J Med Internet Res. 2020;22:e18388.

31. Kadambi A, Bandini A, Ramkalawan RD, Hitzig SL, Zariffa J. Designing an Egocentric Video-Based
Dashboard to Report Hand Performance Measures for Outpatient Rehabilitation of Cervical Spinal Cord
Injury. Top Spinal Cord Inj Rehabil. 2023;29 suppl:75–87.

32. Dousty M, Zariffa J. Tenodesis Grasp Detection in Egocentric Video. IEEE J Biomed Health Inform.
2020;PP.

33. Tsai M-F, Atputharaj S, Zariffa J, Wang RH. Perspectives and expectations of stroke survivors using
egocentric cameras for monitoring hand function at home: a mixed methods study. Disabil Rehabil Assist
Technol. 2024;19:878–88.

34. Bandini A, Kalsi-Ryan S, Craven BC, Zariffa J, Hitzig SL. Perspectives and recommendations of
individuals with tetraplegia regarding wearable cameras for monitoring hand function at home: Insights from
a community-based study. J Spinal Cord Med. 2021;:1–12.

35. Anguera MT, Blanco-Villaseñor A, Losada JL, Sánchez-Algarra P, Onwuegbuzie AJ. Revisiting the
difference between mixed methods and multimethods: Is it all in the name? Qual Quant. 2018;52:2757–70.

36. Nielsen J, Landauer TK. A mathematical model of the finding of usability problems. In: Proceedings of
the SIGCHI conference on Human factors in computing systems - CHI '93. New York, New York, USA: ACM

Press; 1993.

37. Tsai M-F, Bandini A, Wang RH, Zariffa J. Capturing Representative Hand Use at Home Using Egocentric
Video in Individuals with Upper Limb Impairment. J Vis Exp. 2020;:1–12.

38. Zhou X, Girdhar R, Joulin A, Krähenbühl P, Misra I. Detecting Twenty-Thousand Classes Using Image-
Level Supervision. In: Computer Vision – ECCV 2022. Springer Nature Switzerland; 2022. p. 350–68.

39. Shan D, Geng J, Shu M, Fouhey DF. Understanding human hands in contact at internet scale. In: 2020
IEEE/CVF Conference on Computer Vision and Pattern Recognition (CVPR). IEEE; 2020.

40. Kadambi A, Zariffa J. Detecting activities of daily living in egocentric video to contextualize hand use at
home in outpatient neurorehabilitation settings. IEEE Trans Neural Syst Rehabil Eng. 2025;33:1951–7.

41. Brooke J. SUS: A "quick and dirty" usability scale. In: Jordan PW, Thomas B, Weerdmeester BA,
McClelland AL, editors. Usability Evaluation in Industry. London: Taylor & Francis; 1996. p. 189–94.

42. Lewis JR. The system usability scale: Past, present, and future. Int J Hum Comput Interact.
2018;34:577–90.

43. Braun V, Clarke V, Hayfield N, Terry G. Thematic Analysis. In: Liamputtong P, editor. Handbook of
Research Methods in Health Social Sciences. Singapore: Springer Singapore; 2019. p. 843–60.

44. Braun V, Clarke V. Reflecting on reflexive thematic analysis. Qualitative Research in Sport, Exercise and
Health. 2019;11:589–97.

45. Braun V, Clarke V. One size fits all? What counts as quality practice in (reflexive) thematic analysis?
Qual Res Psychol. 2021;18:328–52.

46. Sauro J, Lewis JR. Quantifying the User Experience: Practical statistics for User Research. Oxford,
England: Morgan Kaufmann; 2016.

47. Haig J. Assessment tools used by occupational therapists with head injured patients in a rehabilitation
setting. Br J Occup Ther. 1997;60:541–5.

48. Robertson L, Blaga L. Occupational therapy assessments used in acute physical care settings. Scand J
Occup Ther. 2013;20:127–35.

49. Dunkleberger K, Slomine B, Salorio C, Salva E, Suskauer S, Christensen J. The upper extremity
measurement scale. J Head Trauma Rehabil. 2006;21:410.

50. Itzkovich M, Gelernter I, Biering-Sorensen F, Weeks C, Laramee MT, Craven BC, et al. The Spinal Cord
Independence Measure (SCIM) version III: reliability and validity in a multi-center international study. Disabil
Rehabil. 2007;29:1926–33.

51. Kalsi-Ryan S, Beaton D, Curt A, Duff S, Popovic MR, Rudhe C, et al. The Graded Redefined Assessment
of Strength Sensibility and Prehension: reliability and validity. J Neurotrauma. 2012;29:905–14.

52. Lockwood KJ, Harding KE, Boyd JN, Taylor NF. Home visits by occupational therapists improve
adherence to recommendations: Process evaluation of a randomised controlled trial. Aust Occup Ther J.
2020;67:287–96.

53. Read J, Jones N, Fegan C, Cudd P, Simpson E, Mazumdar S, et al. Remote Home Visit: Exploring the
feasibility, acceptability and potential benefits of using digital technology to undertake occupational therapy
home assessments. Br J Occup Ther. 2020;83:648–58.

54. Cumming RG, Thomas M, Szonyi G, Salkeld G, O'Neill E, Westbury C, et al. Home visits by an
occupational therapist for assessment and modification of environmental hazards: a randomized trial of
falls prevention. J Am Geriatr Soc. 1999;47:1397–402.

55. Gitlin LN, Hauck WW, Winter L, Dennis MP, Schulz R. Effect of an in-home occupational and physical
therapy intervention on reducing mortality in functionally vulnerable older people: preliminary findings:
Effect of occupational and physical therapy. J Am Geriatr Soc. 2006;54:950–5.

56. Gagnon M-P, Desmartis M, Labrecque M, Car J, Pagliari C, Pluye P, et al. Systematic review of factors
influencing the adoption of information and communication technologies by healthcare professionals. J
Med Syst. 2012;36:241–77.

57. Christodoulakis C, Asgarian A, Easterbrook S. Barriers to adoption of information technology in
healthcare. Conf Cent Adv Stud Collab Res. 2017;:66–75.

58. Zakerabasali S, Ayyoubzadeh SM, Baniasadi T, Yazdani A, Abhari S. Mobile health technology and
healthcare providers: Systemic barriers to adoption. Healthc Inform Res. 2021;27:267–78.

59. Khong PCB, Holroyd E, Wang W. A Critical Review of the Theoretical Frameworks and the Conceptual
Factors in the Adoption of Clinical Decision Support Systems. Comput Inform Nurs. 2015;33:555–70.

60. Ledgerd R, World Federation of Occupational Therapists. WFOT report: WFOT human resources project
2018 and 2020. World Fed Occup Ther Bull. 2020;76:69–74.

61. Rozier CK, Thompson ME, Shill J, Vollmar MG. Career paths of male physical therapist students entering
a female-dominant profession. J Phys Ther Educ. 2001;15:32–40.

Patient Video Recordings

1. Camera Handoff

2. Recording Protocol

3. Camera Return

Dashboard Generation

Quantitative Metrics
- **Minutes recorded per day**
- **Percentage interaction per day**
- **Number of interactions per hour**
- **Average interaction duration (seconds)**
- **Activity breakdown (minutes)**

Video snippets (manually curated)

Therapist Interviews

1. Structured Questionnaire (Q1-6)

2. Dashboard Review (10 min)

3. Interview (45 min)
  a. Structured Questionnaire (Q7-16)
  b. System Usability Scale
  c. Semi-Structured Interview

* (1) I predominantly deal with technical systems because I am forced to

(2) I enjoy spending time becoming acquainted with a new technical system

* (3) It is enough for me that a technical system works; I do not care how or why

(4) I try to understand how a technical system exactly works

* (5) It is enough for me to know the basic functions of a technical system

(6) I try to make full use of the capabilities of a technical system

(7) I would personally use this CDSS regularly to understand and assess patient performance

(8) I found the CDSS easy to use overall, and the various functions in this CDSS were well integrated

(9) I felt like I could trust the metrics and information provided to me by the CDSS

* (10) Using this CDSS regularly would substantially interrupt my workflow

* (11) I feel apprehensive about using this CDSS with future patients

(12) Using this CDSS is useful in understanding and assessing patient performance

* (13) I found the CDSS very cumbersome to use, and the presented metrics were not clear

(14) I would imagine that most people would learn to use this CDSS very quickly

(15) The information presented by the CDSS would potentially influence patient therapy plans

* (16) Presenting the information in this format is NOT useful to me or my assessment of patients

Statement

Technology use and affinity

CDSS clinical utility and acitonability

Mean
P5
P4
P3
P2
P1

Likert Scale Scores