Understanding Thromboembolus Transport Patterns In The Brain For Stroke In The Presence Of Carotid Artery Stenosis

Ricardo Roopnarinesingh[*1], Neel D. Jani [†2], Michelle Leppert [‡3] & Debanjan Mukherjee [§1]

[1]Paul M Rady Department of Mechanical Engineering, University of Colorado Boulder
[2]2Department of Neurology, University of Wisconsin, Madison
[3]Department of Neurology, University of Colorado Anschutz Medical Campus

Author preprint for medRxiv

Abstract

Deciphering the source of an embolism is a common challenge encountered in stroke treatment. Carotid stenosis is a key source of embolic strokes. Carotid interventions can be indicated when a patient has greater than 50% stenosis in the carotid ipsilateral to the cerebral infarction, which is designated as the symptomatic carotid. However, there are often significant number of cases where carotid emboli travel contralaterally leading to ambiguity regarding which carotid is symptomatic. We use a patient-specific computational embolus-hemodynamics interaction model developed in prior works to conduct an in silico experiment spanning 30 heart-to-brain arterial models with differing combinations of bilateral severe and mild stenosis degrees. We used these models to study source-to-destination transport of thromboemboli released from left/right carotid disease sites, and cardiogenic sources. Across all cases considered, thromboemboli from left and right carotid sources showed non-zero contralateral transport. We also found that cardiogenic thromboemboli do not have an altered hemisphere distribution or distinct transport preference dependent on stenosis degree, thus potentially making the underlying etiology more cryptic. In patients with carotid stenosis or chronic occlusion ipsilateral to the area affected by stroke, we have demonstrated that the presence of contralateral stenosis can cause emboli that travel across the Circle of Willis (CoW) which can potentially lead to ambiguity when deciding which carotid is truly symptomatic.

Keywords

Stroke; Embolus; Carotid Artery Disease; Carotid Revascularization; Stenosis; Circle of Willis

[*]ricardo.roopnarinesingh@colorado.edu

[†]NJani@uwhealth.org

[‡]Michelle.Leppert@CUAnschutz.Edu

[§]Corresponding Author: debanjan@colorado.edu

1 Introduction

Embolic stroke accounts for a majority of all ischemic stroke cases, with some studies categorizing up to 60% of these cases as Embolic Stroke of Undetermined Source ($ESUS$) [1–3]. Currently, the ability to identify and disambiguate an etiology of a stroke is limited for ESUS cases, which can restrict treatment efficacy and permit recurrent events [2]. The carotid arteries cause 15%-20% of all ischemic strokes due to build up of atherosclerotic plaques that can generate thromboemboli that move into the cerebral arteries [4]. These buildups vary in severity based upon the degree of stenosis and laterality of the stroke which informs potential interventions [5–8]. Carotid interventions aimed at revascularization, either through carotid endarterectomy (CEA) or carotid artery stenting (CAS) are performed when there is stenosis to restore perfusion and reduce future stroke risk [9]. Specifically, carotid intervention is indicated when a patient has a North American Symptomatic Carotid Endarterectomy Trial ($NASCET$) severity greater than 70% in the carotid ipsilateral to the stroke along with transient ischemic attacks [7]. Carotid interventions are considered if the patient has a severity between 50% and 69% along with symptoms, but are not required as the benefits may be limited [7, 9]. Atherosclerotic plaque can become vulnerable to embolization without easily decipherable markers, and these vulnerable plaques can subsequently lead directly to a cerebral occlusion [10, 11]. Although stenosis severity may be non-trivial to acquire, it may not always be possible to tell where an embolus originated from due to the complexities of physiological arterial hemodynamics, and the complex embolus-hemodynamics interaction mechanics. It is commonly believed that emboli originating from a carotid artery will travel into the Circle of Willis (CoW) and occlude a vessel within the ipsilateral hemisphere, but there have been known cases where the occlusion occurs in the contralateral hemisphere [12–14]. Although contralateral events are less frequent, these cases can contribute to the ambiguity in deciphering stroke etiology due to the non-intuitive pathways of occlusion. With both carotids being suspect to combinations of vulnerable severe and mild levels of plaque buildup, contralateral events could lead to patients who receive treatment for the ipsilateral non-symptomatic vessel thereby deriving less benefit. These contralateral events can cause the chance of missing secondary prevention opportunities or possible benefits from intervening on the contralateral carotid if that is the truly symptomatic vessel.

How contralateral carotid stenosis affects stroke risk is currently relatively unexplored, whereas this investigation can further illuminate mechanisms and risks of non-intuitive stroke in cases of bilateral stenosis [15, 16]. We additionally considered cases consisting of Contralateral Carotid Occlusions (CCO). Cases with CCO offer an interesting circumstance wherein each diseased carotid will be treated differently depending on which side is actively causing embolization. Understanding the probabilities of non-intuitive embolus source-to-destination movement for varying carotid stenosis cases can help to provide a more cohesive risk stratification across bilateral stenosis degrees. There is currently a lack of understanding in how patients with carotid disease may experience contralateral embolic events or non-intuitive embolic transport in various levels of carotid stenosis. Here, we present an *in silico* study analyzing how different combinations of varying carotid stenosis may present non-intuitive occlusion events and how the altered hemodynamics with the introduction of a stenosis can affect embolus transport.

2 Methods

2.1 Image-Based Modeling of Vascular Anatomy

A patient-specific vasculature model was created spanning arterial pathway from the aortic arch to the CoW using the open-source software tool Simvascular [17, 18]. The model was generated from a contrast enhanced computed tomography (CT) image, available from a set of CT images as part of the Institutional Review Board (IRB) approved Screening Technology and Outcome Project in Stroke ($STOP$-$Stroke$) database [19]. Since, we used the de-identified version of the CT image for secondary retrospective computational analysis, for this study no additional IRB approval was needed. The specific patient chosen for this study was based on the criteria that the patient exhibited a complete CoW anastomosis with healthy (*non-stenotic*) carotid

arteries (*see also our prior works [18, 20]*). Arteries from the aortic inlet to the cerebral arteries up until the M1, A1, and P1 segments of the MCA, ACA, and PCA, were segmented using planar 2D segmentation technique built into SimVascular. These segmentations were then lofted to generate the 3D solid model of the complete arterial pathway. Arterial stenosis was virtually modeled at specific carotid artery locations, using these healthy patient vascular segmentatios, and manually decreasing the segmentation sizes of the carotid bulb area to the corresponding NASCET severity. The stenosis degrees were categorized into three mild NASCET scores of [10%, 25%, 40%], and three moderate/severe NASCET scores of [50%, 70%, 85%]. These severity ranks were paired to where one carotid would have a mild stenosis and the other would have a moderate/severe stenosis for both the left and right carotid arteries permutations. Additionally, by virtually eliminating these segmentations locally at the carotid site, we modeled 100% occluded carotid across all 6 severities for both carotids to mimic cases of CCO. The length of stenosis affliction that spanned across the carotid artery varied proportionally with the severity of the stenosis. Length of stenosis is defined as the span of artery that has a disturbed cross-sectional area due to the stenosis. A more severe stenosis corresponds to a larger length. The equation of the stenosis length is defined in the Supplementary Material in greater detail. This led to 30 different models each with a unique pair of mild and moderate/severe stenosed carotids. The resulting models for the stenosis vascular anatomy for differing severities can be seen in Figure 1.

2.2 Hemodynamics Simulation

Hemodynamics across the arterial model was simulated using a stabilized finite element method for incompressible Navier-Stokes equations implemented in SimVascular [17]. Each model was discretized into a mesh comprising 9-10 million linear tetrahedral elements. The average element size across all the models was 0.65 mm. These parameters were chosen through mesh refinement analysis described in prior works [21]. Stenosed regions of the carotid arteries underwent additional element refinement due to the smaller cross-sectional area. This local refinement was guided by setting the element edge length to 1/10th of the smallest stenosis diameter, ensuring at least 10 or more elements within all cross-sectional slices of the stenosed region. A second local refinement operation was conducted for vessels in the CoW, with an average element size of 0.25 mm due to a smaller cross-sectional area relative to other vessels in the heart-brain arterial model. Blood was assumed to be an incompressible Newtonian fluid with average viscosity of 4.0 cP, and bulk density of 1.06 g/cc. Mathematical details of the underlying equations are provided in the Supplementary Material. A validated pulsatile flow profile with a cardiac cycle of 0.83 seconds was applied to the aorta inlet [22]. This inlet flow was set to a consistent cardiac output (CO) of 79 mL/s across all models. Resistance boundary conditions were assigned to each outlet of the model to account for the influence of the downstream vascular beds at the arterial outlets. For this purpose, Total Arterial Resistance (TAR) was estimated using a Mean Arterial Pressure (MAP) of 93.33 mmHg, which assumes that the patient exhibits an average systolic and diastolic blood pressure of 120 and 80 mmHg respectively. A proportion of the TAR was assigned to each vessel outlet based on target flow divisions as outlined in prior work [18, 20, 23]. For target flow division, 65% of the total CO was assumed to exit the descending aorta [24], flow rates exiting the six cerebral artery outlets were assigned based on measured MR data reported in [25], and the remainder volumetric flow was set to exit the external carotid and subclavian arteries proportional to their cross-sectional areas [26]. These resistance values were tuned to the targeted flow values through multiple steady-state flow simulations [23, 27]. Once obtained, the CO, TAR, MAP, and outlet resistances were kept the same across all models to enable controlled simulations of embolus transport. Hemodynamics was simulated for three cardiac cycles for each model, and computed velocity and pressure fields for the third cycle were used to conduct embolus transport simulations. Each of these hemodynamic simulations steps are outlined extensively in our previous works [18, 20, 23].

2.3 Embolus Transport Simulation

Embolus transport was modeled using a custom modified version of the Maxey-Riley equation [28] to represent embolus-hemodynamics interactions. This model estimates individual embolus velocity and position

as it traverses through blood flow based on embolus properties and hemodynamic features. To closely simulate arterial embolus transport, we modified the classical form of Maxey-Riley equations to account for: (a) lift forces produced by shear gradients near artery walls; (b) collisions between particles and the artery wall; and (c) elastohydrodynamic lubrication effects between embolus and artery wall. These forces were in addition to the forces of drag, added mass effects, and forces from the undisturbed background flow. Wall collisions were resolved as visco-elastic collisions with particle-to-particle collisions being neglected as each simulation comprises individual embolus transport one-by-one as a Monte Carlo sampling based approach. Mathematical details of the underlying equations are provided in the Supplementary Material. The emboli were assumed to be spherical particles with a one-way coupled fluid interaction, and individual emboli of $500\mu m$ were chosen, with mechanical properties of thromboembmoli [23]. This size was chosen such that we focus on small emboli whose transport can be more ambiguous to infer from imaging; while remaining within the bounds of the mathemaical approximations of one-way coupled embolus hemodynamics interactions, which break down as embolus size approaches vessel diameter [29]. The simulated flow field for the third cardiac cycle (see Section 2.2) was stitched together 5 times to generate a representative hemodynamic field spanning 5 cardiac cycles with no cycle-to-cycle variation. Individual thromboembolus trajectories were computed across these 5 cardiac cycles, using an Euler integration scheme with a numerical time-step of 0.05 ms.

2.4   Design of In Silico Experiments

We utilized the simulation methodology introduced in Sections 2.1- 2.3 to structure a parametric set of *in silico* experiments for investigating embolus transport in presence of varying carotid stenosis scenarios. The outline of these experiments is shown in Figure 2. To parameterize the stenosis severity, a total of 18 vascular models were created as a combination of one mild stenosis (*10%, 25%, and 40% NASCET severity*) and one moderate/severe stenosis (*50%, 70%, and 85% NASCET severity*) for each pair of left and right carotids respectively (*see also Figure 1*). An additional 12 models were designed with a fully occluded carotid artery, mimicking CCO cases, paired with one of each of the mild and severe stenosis degrees. For each of these 30 models, approximately 5,500 thromboemboli samples were released along the carotid artery walls, mimicking plaque thromboembolization at stenosis site. For the 12 CCO models, thromboemboli were released at the partially stenosed carotid site; while for the 18 non-CCO models, thromboemboli were releases from both carotid sites bilaterally. Additionally, for all 30 cases, an additional $\approx$ 5,500 samples of cardiogenic emboli were released at the aortic root inlet, to further illustrate how different combinations of stenosis severities may affect the pathway of cardiogenic emboli. In total, this design led to a set of 78 total embolus transport experiments, leading to combined distribution data spanning approximately 429,000 thromboemboli. For each emolus sample, their trajectory was tracked from release source to a destination vessel, and number fractions of distribution to the 6 cerebral arteries was computed and analysed for this study.

3   Results

3.1   Trends in Contralateral Distribution of Carotid Thromboemboli

Thromboembolus distribution from our simulations was computed as the percentage of emboli that exit each of the six outlets of the CoW. Figure 3 presents the proportion of emboli that were able to cross into the opposite hemisphere relative to the carotid release site for each pair of stenosis severities, illustrated for the 18 bilateral stenosis, non-CCO cases. Each model is labeled by their combination of left-right stenosis severity respectively (*i.e. 10L50R corresponds to the model with a 10% left carotid stenosis and a 50% right carotid stenosis*). These embolus distribution results therefore inform a probability of contralateral embolic events occurring with respect to stenosis severities across the right and left carotid arteries. We observed that, all simulations produced a non-zero contralateral distribution, meaning that all combinations of severe and mild stenosis showed the potential for contralateral occlusions. The largest extent of contralateral movement were found in the models with the most severe (*that is, 85%*) stenosed carotid, where contralateral movement

was highest from the mild carotid release sites. These models had a likelihood of contralateral distribution between 6% and 10%, which was notably higher than the other releases that ranged primarily between 0.5% and 4%. Statistically, the null hypothesis that contralateral movement does not occur from emboli released at both the left and right carotids (*that is the proportion of contralaterally traveling emboli = 0*) was rejected with p-values of 0.000025 and 0.00003 respectively (*based on a one-sample non-parametric Wilcoxon rank test*). We also observe that extent of contralateral movement of thromboemboli is generally greater for emboli released in right carotid and traveling towards the left hemisphere for severely stenosed left carotids, indicating an inherent right vs left asymmetry for the cases considered here. Generally, these simulation data revealed contralateral trans-hemispheric thromboembolus transport and distribution occurring at the CoW; with primary modality of embolis moving from the mild stenosis site towards the hemisphere ipsilateral to the severe stenosed site, and with the likelihood of contralateral movement generally increasing as the moderate/severe carotid stenosis increases in degree.

3.2 Trends in Cardiogenic Thromboembolus Distribution

In Figure 4 panel A, we present the cardiogenic thrombembolus distribution to each brain hemisphere relative to each severity combination, for the 18 bilateral stenosis, non-CCO cases. The individual models are labelled using the same scheme as in Figure 3. The cardiogenic distribution was acquired to illustrate how thromboembolus trajectory would be modified based upon stenosis severity. Thromboemboli traveling from the aortic root will be recruited into either the subclavian artery or the left/right common carotid artery and distribute into each hemisphere through the supplying cerebral arteries in the CoW. The resulting cardiogenic embolus distribution can help to illuminate preferential pathways in the presence of varying stenosis degrees for both the right and left carotid arteries that can, in turn, reveal non-intuitive pathways which potentially contribute to ESUS diagnosis. Our experiments reveal a rather consistent distribution of emboli across all models averaging 55% and 45% for the left and right hemispheres respectively (*standard deviation of 4.78%, coefficient of variation for left = 0.087, for right = 0.106*). These observations indicate that that stenosis severity does not potentially have a strong influence on distribution of cardiogenic thromboemboli across both hemispheres. This itself is counter-intuitive, as specifically for the high severity cases, it may seem expected that hemisphere ipsilateral to the high severity stenosis will not be a peferred cardiogenic embolus distribution. These observations therefore indicate the potential role of compensation from other vessels feeding the CoW, and the overall network anastomosis of CoW, by which cardiogenic thromboemboli are able to distribute into the hemisphere with a severely stenosed carotid artery. This is further explored next (Section 3.4) by comparing computed volumetric flow rates feeding into the cervical vessels that supply the CoW.

3.3 Thromboembolus Distribution in Contralateral Carotid Occlusion Cases

We further present thromboembolus transport and distribution in the 12 CCO models considered in our study. For this, we considered emboli releases from the carotid artery site that was not fully-occluded, and from aortic inlet (*that is, cardiogenic*). Figure 5 illustrates the proportion of carotid thromboemboli with contralateral trans-hemspheric distribution relative to the carotid release site for the 12 CCO models. The observations indicate a similar continuing trend compared to that observed in the results presented in Section 3.1, wherein increasing stenosis severity on one side permits increased contralateral movement for emboli released on the other side. The proportions indicated are for emboli reaching the hemisphere ipsilateral to the fully occluded carotid. The completely occluded cases allowed for a high proportion of emboli traveling contralaterally form the non-occluded carotid side with an average of 10.8% of contralateral emboli which is higher than all of the non-CCO simulation results presented earlier (*two-sample t-test between CCO cases and non-fully occluded cases: p-value < 0.001*). Furthermore, generally contralateral movement of carotid emboli in the simulated results are greater for 100% occluded left carotid compared to the right, indicating potential right vs left asymmetries as noted in Section 3.1. Furthermore, we also simulated cardiogenic thromboembolus and distribution for the 12 CCO cases, and the resulting distributions are presented in Figure 6, panel A. Cardiogenic thromboembolus distribution showed lesser variations in left

vs right hemisphere distribution when compared against carotid emboli, especially for the fully occluded right carotid cases. The fully occluded left carotid models saw an average of 57.4% of emboli going to the right hemisphere and 42.6% going to the left (*standard deviation of 7.95%, coefficient of variation for right and left = 0.14 and 0.19 respectively*) while the fully occluded right carotid models experienced an average 52.1% distributing to the left hemisphere and 47.9% distributing to the right (*standard deviation of 3.04%, coefficient of variation for right and left = 0.063 and 0.058 respectively*). Thus, while it was favorable for emboli to distribute to the hemisphere fed by a non-occluded CCA, the variability in the split of left and right hemisphere thromboembolus distributions remained low regardless of the stenosis severity of the non-occluded carotid. These observations again indicate potential role of flow pathways where the non-occluded cervical vessels (*e.g. left carotids and basilar arteries, for fully occluded right carotids*) feeding the CoW can recruit emboli towards the brain, leading to non-intuitive cardiogenic embolus transport patterns.

3.4 Flow Distribution in Vessels Feeding the Circle of Willis

We further explore the network effect of the vessel anastomosis connecting the cervical vessels to the Circle of Willis, by analysing estimated flow rates in the cervical vessels (*left and right internal carotid artery, and basilar artery*) for all the 30 models. These flow rates were estimated by integrating simulated blood flow velocity over a cross-sectional slice of that specific artery at each simulated time step. The total average flow rate was then completed by integrating the flow rates in time over one full cardiac cycle. We refer to Figure 4 panel B, for illustration of the computed average volumetric flow rate through the the basilar artery, the left internal carotid artery, and the right internal carotid artery - computed for the 18 non-CCO models with patent bilateral stenosis. Likewise Figure 6 panel B, presents the estimated flow rates at the basilar, and left and right carotid arteries for the 12 CCO cases. These flow rate variations with varying stenosis severity, indicate that owing to the network nature of the vascular anastomosis leading into the CoW, the three cervical vessels feeding the CoW can be engaged differently for varying stenosis severities to retain consistent volumetric flow into the brain. This provides one mechanism of non-intuitive distribution of emboli. Specifically, increasing right carotid flow for severely stenosed left carotids will lead more emboli from the right carotid site to reach the CoW, increasing the likelihood of their contralateral movement. This is despite there being a faster jet type flow at the severe stenosis site that does increase locally the speed of eny emboli released at the severe stenosis carotid site. Additionally, flow compensations within the basilar artery and the mildly stenosed carotid so as to preserve cerebral perfusion can pose as an alternate pathway of cardiogenic thromboemboli into the hemiisphere ipsilateral to the severely stenosed carotid. This indicates flow distribution and recruitment as a mechanism of consistent hemisphere distribution for cardiogenic emboli regardless of stenosis severity.

3.5 Circle of Willis Proximal Collateral Flow Routing for Varying Stenosis

In addition to varying extent of flow routing through the cervical vessels feeding the CoW, the communicating arteries in the CoW anastomosis can also recruit flow through them differently based on stenosis severity, so as to retain cerebral blood flow distribution. We investigated this proximal collateral flow routing patterns in the CoW communicating artery segments (*compared to distal collateral circulation through smaller leptomeningeal vessels*), by computing flow rate from simulation data at various locations along the CoW using cross-sectional integration of velocity, and time-averaging over a cardiac cycle (as described in Section 3.4). Figure 7, panel a. depicts these averaged volumetric flow rates, and the direction of the average flow, through the communicating artery segments in the CoW, for moderate left carotid stenosis paired with moderate right carotid stenosis. Figure 7, panel b. depicts the same for moderate right carotid stenosis paired with moderate left carotid stenosis. Additionally, for the 12 CCO cases, the corresponding CoW communicating artery segment flow maps are presented in Figure 8. Differential flow recruitment and changes in flow routing patterns based on stenosis severity is clearly observed across all of these illustrated flow data. The null hypothesis that communicating artery flows are same as those in the case of flow without any stenosis present was rejected with a p-value $\ll 0.005$ (*using a two-sample t-test*). Specifically, for instance, observing

the cases with moderate left carotid stenosis paired with severe right carotid stenosis, the anterior communicating artery (AcoA) switches direction, and AcoA flow increases towards the right with increasing right carotid stenosis severity. This trend is conversely true for cases with moderate right carotid stenosis paired with severe left carotid stenosis. Furthermore, for all CCO cases, the AcoA flow stays oriented towards the hemisphere ipsilateral to the occluded carotid. As another instance, we observe that with increasing flow recruitment through the Basilar Artery, the flow routed through the right P1 communicator (RP1) segment increases steadily with extent of right carotid stenosis severity (*and likewise for flow through the left P1 communicator for increasing left carotid stenosis severity*). This is also noted in the CCO cases, where with increasing severity of the left/right carotid stenosis, in presence of fully occluded right/left carotids, flow through the basilar artery into the LP1/RP1 segments increase. Indicating the inherent right-left asymmetry, we also note that flow through the right posterior communicator segment (R.Comm) switches direction for the two most severe right carotid stenosis models to compensate for RMCA and RACA flow while right carotid supply into the CoW is significantly reduced. Together, these results potentially illustrate pathways through the proximal collateral circulation at the CoW that can promote non-intuitive thromboembolus transport respective to the stenosis location and severity.

4   Discussion

Our investigation comprised a set of parametric simulations inquiring on how thromboembolus transport is affected by varying combinations of stenosis severities to specifically illuminate how these cases may contribute to elusive stroke etiologies through the mechanism of contralateral trans-hemispheric embolus movement. Fundamentally, vascular pathways feeding into the brain comprise a complex network. Flow can be routed through varying, often redundant, pathways to the major cerebral artery territories, in the event of a pathological disruption to flow in one or more vessels in the network. This manifests in differential flow recruitment in the cervical vessels, and proximal flow routing in the communicating vessels of the CoW. We note that while the role of CoW vessels in promoting proximal collateral flow routing for acute ischemic stroke has been discussed in several works [30–33]; but their role in routing emboli in the brain, and being a potential determinant for embolus source-destination relationship has been sparingly investigated. In a series of prior works, we have generated some of the first evidences on how embolus interaction with these varying hemodynamic pathways ultimately determines embolus fate [18, 20]. Here, we further expand upon this observation, and substantiate the role that proximal collateral flow routing may play in underlying non-intuitive embolus distribution in the brain from carotid artery sites.

For trans-hemispheric contralateral embolus distribution, a common pattern seen in our simulation data was that the majority of contralateral embolus movement occurred from the mildly stenosed carotid source to the hemisphere with the respective severe stenosis degree (*two-sample t-test between contralateral movement from the severe carotid release and the mild carotid release: p-value = 0.003*). This pattern can be informed by the altering proximal collateral flow routes in the CoW that is dependent on the pairing of stenosis severities. Based on observations presented in Sections 3.1, 3.3, 3.5, there are three major flow alterations seen for the increasing severity of the severely stenosed carotid (i.e. 10L50R, 10L70R, 10L85R), where the AcoA and posterior communicators play a key role (*as noted in Section 3.5*). First, the AcoA flow directs from the CoW hemisphere fed by a mild ipsilateral carotid stenosis to the contralateral severe carotid as the severe stenosis increases in degree. The AcoA flow rate also increases as the severity of the moderate/severe carotid stenosis degree increases. These two factors raise the potential of the AcoA recruiting emboli and enabling contralateral movement from the relative mild carotid hemisphere to the severe carotid hemisphere due the favorable AcoA flow rate and flow direction. Second, the flow rates in the internal carotid extension into the CoW increase in the hemisphere on the mild carotid stenosis side, and decrease in the hemisphere on the severe carotid stenosis side, allowing for more emboli to be recruited into the anterior region thus raising the probability of emboli being pulled into the AcoA. Third, as the stenosis severity increases, flow in the P1 communicating artery on the hemisphere ipsilateral to the mildly stenosed site increases and directs towards the anterior region. This is due to flow compensation of the basilar artery for the severely stenosed

region. Basilar artery carries flow into the posterior region of the CoW, thereby creating an overall increase of blood flow within the P1 and Comm. arteries in that respective hemisphere.

The CCO cases with fully occluded carotid arteries follow a similar pattern, with observably more pronounced contralateral flow and embolus routing. The AcoA notably plays a role here too in enabling thromboembolus movement from the hemisphere on the partially occluded carotid side to that on the fully occluded carotid side. Similar to non-CCO cases, flow routes through the cervical vessels to the side that has the severe or complete stenosis to compensate for the lack of flow, enabling the propensity of contralateral movement. Extrapolating purely based on non-CCO cases, we would expect then that complete carotid occlusions would increase the extent of contralateral severity, as more flow enters the CoW from the vertebrobasilar vessels, and the carotid that is not fully stenosed, which is also in accordance with observations presented in Section 3.3.

We also investigated cardiogenic emboli distributions to the CoW for varying stenosis severity to gain insights on source-destination propensities for cardiac vs carotid sources of thromboemboli. As the severity of carotid stenosis increases on one side (say, left), the less stenosed carotid on the other side (say, right) and the basilar arteries compensate for flow to maintain cerebral perfusion. As stenosis severity on the other side vary, resistance to flow through the stenosis overpowers, shifting the compensation balance further towards basilar artery flows alone. Consequently, again, a network based flow compensation effect, leads to observations on rather consistent cardiogenic embolus distributions across stenosis values in the non-CCO cases at an average of 55% and 45% to the left and right hemispheres respectively for the non-fully-occluded cases. This was also observed for the CCO models whereas there was a consistent distribution for both the left and right occlusion across the varying severities as seen in Figure 6 panel A.

These observations on contralateral thromboembolus distribution are mechanistically informed by the mildly stenosed carotid and the basilar artery recruiting more flow, and more emboli, towards the CoW, and subsequent trans-hemispheric transport of these emboli based on CoW proximal routing noted above pushing a fraction of these emboli into the hemisphere associated with the more severely stenosed carotid. Figure 9 presents a conceptual schematic summarizing these two effects, discussed extensively thus far, originating from two specific flow recruitment patterns in the cerebrovascular network: differential recruitment of flow at the cervical vessels, and proximal collateral flow routing at the CoW. This summary figure schematically illustrates a case of severe right carotid stenosis experiencing contralateral movement of carotid emboli, and non-intuitive transport of cardiogenic emboli, to the right hemisphere. From this perspective, for cardiogenic emboli with varying degrees of stenosis, we observe limited distinct preference in laterality of distribution for the simulations considered. Consequently, for a stroke patient with indicated carotid disease and suspected cardioembolic source, these results indicate that in the absence of quantitative source-destination transport patterns, disambiguating between cardiogenic and carotid etiologies would be challenging, which is commonly observed in ESUS cases.

The findings from this study provide criticial insights on embolus-hemodynamics interactions, and embolus source-destination relationship, in the context of deciphering etiology and intervention targets in potential embolic events with carotid stenosis. Carotid intervention is indicated when stenosis severity is higher than 70% in the ipsilateral carotid along with the correlating symptoms. Given our data suggesting an increased risk of embolus transport from a mildly stenosed contralateral carotid in the presence of severe ipsilateral carotid disease, carotid interventions could be indicated for the contralateral carotid to prevent stroke recurrence. Furthermore, in the presence of an ipsilateral complete carotid occlusion, we show non-negligible transport from the contralateral carotid. This finding could prompt consideration of intervention of the contralateral carotid for secondary stroke prevention in patients with chronic carotid occlusions. Our completed simulations show that up to 14% of embolization from the carotids can create contralateral events that may keep a carotid that is symptomatic or actively embolizing from being properly identified. This offers ambiguity in treatment as treatment depends directly on whether the carotid artery is symptomatic or

not. Through these results, contralateral, non-intuitive embolus distrbution is shown to have the potential to pose a prominent risk, and further risk stratification may be beneficial in deducing treatment for patients that are experiencing symptomatic carotid stenosis with a corresponding stroke.

This study was based on a few key assumptions. The underlying computational model relies on several assumptions to make high resolution simulations such as the ones presented here tractable, which have been discussed at length in our prior works [18, 20]. We assume that blood flow influences the emboli, but the emboli themselves do not significantly alter the flow. Each embolus is modelled independently, and does not interact with other emboli, and embolus sizes are small relative to the size of the arteries modeled; making it reasonable to assume one-way interaction of blood flow and embolus. Additionally, small embolus sizes avoid numerical issues with clogging the stenosis region, as we assumed a rigid particle and squeezing is not possible within our simulation scope. Second, we considered only a single embolus size in our study. This a meaningful simplification because we have extensively demonstrated how size-dependent embolus-hemodynamics interactions manifests in embolus source-to-destination distribution in a number of prior works [18, 20, 23, 27]. Hence, controlling the simulations to one embolus size enabled us to focus the computational efforts on stenosis severity and network flow routing effects which have been less investigated. Additionally, this enables us to understand the source-to-destination transport patterns for small emboli which plays a critical role in a range of complications such as microstrokes, lacunar and watershed infarcts, and cerebral small vessel disease, for which embolus etiology is often harder to disambiguate. Likewise, we used one base patient anatomy that was modified across multiple stenosis severities which allows for controlled anatomy within the simulations but does not give a very broad application across multiple patient anastamoses. A direct extension of this *in silico* analysis developed here would be to directly recreate stenosis geometries and inform potential release sites for a larger cohort of patients from images, and estimate thromboembolus distribution probabilities. Third, we have not accounted for specific cerebrovascular circulation features such as cerebral autoregulation and distal collateral flow for our study. This is primarily to capture general parametric trends with stenosis severity while maintaining the overall computational costs low. Integration of robust yet computationally efficient models for such processes within the *in silico* framework presented here remains an active area of investigation. Finally, our model focused on distribution trends for thromboemboli only. While stenosis sites can cause thrombogenesis and subsequent thromboembolization, ruptured plaques can often release cholesterol atheroemboli. Our analysis presented here can be directly extended to study cholesterol atheroemboli and atheroembolic showers in future investigations.

5   Concluding remarks

We have presented an *in silico* investigation on how varying degrees of bilateral carotid stenosis can induce non-intuitive embolus transport within the cerebral arteries that can ultimately pose challenges with disambiguating etiology and confusing treatment or intervention decisions. A total of 78 simulations were completed with 18 models of varying bilateral moderate/severe stenosis degrees and 12 models with contralateral carotid occlusion cases (CCO) with pairing of differing stenosed carotids on the non-occluded side. Contralateral embolus movement is the primary mechanism of non-intuitive transport within our simulations and all cases experienced a non-zero contralateral source-to-destination transport of thromboemboli. Models paired with the most severe partial occlusion degree (85%) were found to manifest the highest likelihood of contralateral thromboembolus movement, with fully-occluded CCO cases having on average a higher probability of contralateral movement than non-CCO cases. Cardiogenic thromboemboli did not exhibit a distinct hemisphere preference among all models, with there being consistent distributions to each side of the models regardless of stenosis degree due to flow compensation of the basilar artery and the less stenosed carotid artery. This was consistent for both non-CCO and CCO cases. Ultimately, these findings illustrate the capability of non-intuitive embolus travel from cardiogenic and contralateral carotid sources such that carotid embolus can seemingly source from the opposite side of severely stenosed carotid site, while cardiogenic emoblus can move to either carotid artery. We demonstrate the potential underlying hemodynamics based mechanism for this contralateral movement, comprising differential flow routing in the cervical vessels

feeding the Circle of Willis, and proximal collateral flow routing across the communicating artery segments of the Circle of Willis. With procedures such as CEA having a dependence on occlusion location, these findings can be beneficial for optimizing both treatment efficiency and efficacy in reducing recurrent events.

Conflicts of Interest

The Authors RR, ML, NJ, and DM have no conflicts of interest to report regarding this study.

Funding Acknowledgements

The Authors acknowledge funding support from National Institutes of Health Award: R21EB029736 for this study. This work utilized the Summit supercomputer, which is supported by the National Science Foundation (awards ACI-1532235 and ACI-1532236), the University of Colorado Boulder, and Colorado State University. The Summit supercomputer is a joint effort of the University of Colorado Boulder and Colorado State University.

Author Acknowledgements

Data Availability

All quantitative embolus distribution data for this study have been shared through the Open Science Framework as part of project titled, "Dataset: Transport and Distribution of Embolic Particles in Human Vasculature," which can be accessed at: `https://doi.org/10.17605/OSF.IO/CQKZT`. All image-based modeling and computational hemodynamics simulations were conducted using the open source software package SimVascular. The image-based models (segmentations, pathlines) and computational hemodynamics data will be made available through the Vascular Model Repository associated with the SimVascular project and can be accessed at `https://simvascular.github.io/`. The sharing process with Vascular Model Repository is ongoing, and when the models have been shared, they will be linked to the dataset released on Open Science Framework linked above. For any data access issues, or for additional access to scripts and tools, please contact the corresponding author directly via email, or via the contact page on the research team's official website: `https://www.flowphysicslab.com/`.

References

1. Ferro, J. M., Massaro, A. R. & Mas, J.-L. Aetiological diagnosis of ischaemic stroke in young adults. *The Lancet Neurology* **9,** 1085–1096 (2010).

2. Ntaios, G. Embolic stroke of undetermined source: JACC review topic of the week. *Journal of the American College of Cardiology* **75,** 333–340 (2020).

3. Ntaios, G. *et al.* Embolic strokes of undetermined source: a clinical consensus statement of the ESC Council on Stroke, the European Association of Cardiovascular Imaging and the European Heart Rhythm Association of the ESC. *European Heart Journal* **45,** 1701–1715 (2024).

4. Petty, G. W. *et al.* Ischemic stroke subtypes: a population-based study of incidence and risk factors. *Stroke* **30,** 2513–2516 (1999).

5. AbuRahma, A. F. *et al.* Society for Vascular Surgery clinical practice guidelines for management of extracranial cerebrovascular disease. *Journal of vascular surgery* **75,** 4S–22S (2022).

6. Meschia, J. F., Edwards, L., Roubin, G., Brott, T. G. & Lal, B. K. CREST-2 Commitment to Rigorous Assessment of Carotid Stenting for Primary Prevention of Stroke. *Stroke: Vascular and Interventional Neurology* **4,** e001227 (2024).

7. Roffi, M. & Yadav, J. S. Carotid stenting. *Circulation* **114,** e1–e4 (2006).

8. Jose, B. Guidelines for carotid endarterectomy: a statement for healthcare professionals from a special writing group of the Stroke Council, American Heart Association. *Stroke* **29,** 554–562 (1998).

9. Noiphithak, R. & Liengudom, A. Recent update on carotid endarterectomy versus carotid artery stenting. *Cerebrovascular Diseases* **43,** 68–75 (2017).

10. Hafiane, A. Vulnerable plaque, characteristics, detection, and potential therapies. *Journal of cardiovascular development and disease* **6,** 26 (2019).

11. Mughal, M. M. *et al.* Symptomatic and asymptomatic carotid artery plaque. *Expert review of cardiovascular therapy* **9,** 1315–1330 (2011).

12. Georgiadis, D., Grosset, D. G. & Lees, K. R. Transhemispheric passage of microemboli in patients with unilateral internal carotid artery occlusion. *Stroke* **24,** 1664–1666 (1993).

13. Mounier-Vehier, F., Leys, D. & Pruvo, J. Stroke patterns in unilateral atherothrombotic occlusion of the internal carotid artery. *Stroke* **26,** 422–425 (1995).

14. Dagirmanjian, A., Davis, D. A., Rothfus, W. E., Goldberg, A. L. & Deeb, Z. L. Detection of clinically silent intracranial emboli ipsilateral to internal carotid occlusions during cerebral angiography. *American Journal of Roentgenology* **174,** 367–369 (2000).

15. Maus, V. *et al.* Carotid artery stenosis contralateral to intracranial large vessel occlusion: an independent predictor of unfavorable clinical outcome after mechanical thrombectomy. *Frontiers in Neurology* **9,** 378608 (2018).

16. Viticchi, G. *et al.* Ipsilateral and contralateral carotid stenosis contribute to the outcome of reperfusion treatment for ischemic stroke. *Frontiers in Neurology* **14,** 1237721 (2023).

17. Updegrove, A. *et al.* SimVascular: an open source pipeline for cardiovascular simulation. *Annals of biomedical engineering* **45,** 525–541 (2017).

18. Mukherjee, D., Jani, N. D., Narvid, J. & Shadden, S. C. The role of circle of Willis anatomy variations in cardio-embolic stroke: a patient-specific simulation based study. *Annals of biomedical engineering* **46,** 1128–1145 (2018).

19. Smith, W. S. *et al.* Significance of large vessel intracranial occlusion causing acute ischemic stroke and TIA. *Stroke* **40,** 3834–3840 (2009).

20. Roopnarinesingh, R., Leppert, M. & Mukherjee, D. Evidence and mechanisms for embolic stroke in contralateral hemispheres from carotid artery sources. *Journal of the American Heart Association* **12,** e030792 (2023).

21. Mukherjee, D., Jani, N. D., Narvid, J. & Shadden, S. C. The role of circle of Willis anatomy variations in cardio-embolic stroke: A patient-specific simulation based study. *Annals of biomedical engineering* **46,** 1128–1145 (2018).

22. Olufsen, M. S. *et al.* Numerical simulation and experimental validation of blood flow in arteries with structured-tree outflow conditions. *Annals of biomedical engineering* **28,** 1281–1299 (2000).

23. Mukherjee, D., Jani, N. D., Selvaganesan, K., Weng, C. L. & Shadden, S. C. Computational Assessment Of The Relation Between Embolism Source And Embolus Distribution To The Circle Of Willis For Improved Understanding Of Stroke Etiology. *Journal of Biomechanical Engineering* **138,** 081008 (2016).

24. Carr, I. A., Nemoto, N., Schwartz, R. S. & Shadden, S. C. Size-dependent predilections of cardiogenic embolic transport. *American Journal of Physiology-Heart and Circulatory Physiology* **305,** H732–H739 (2013).

25. MacDonald, M. E. & Frayne, R. Phase contrast MR imaging measurements of blood flow in healthy human cerebral vessel segments. *Physiological Measurement* **36,** 1517 (2015).

26. Zamir, M., Sinclair, P. & Wonnacott, T. H. Relation between diameter and flow in major branches of the arch of the aorta. *Journal of biomechanics* **25,** 1303–1310 (1992).

27. Mukherjee, D. & Shadden, S. C. Inertial particle dynamics in large artery flows–Implications for modeling arterial embolisms. *Journal of Biomechanics* **52,** 155–164 (2017).

28. Maxey, M. R. & Riley, J. J. Equation of motion for a small rigid sphere in a nonuniform flow. *The Physics of Fluids* **26,** 883–889 (1983).

29. Mukherjee, D., Padilla, J. & Shadden, S. C. Numerical investigation of fluid–particle interactions for embolic stroke. *Theoretical and Computational Fluid Dynamics* **30,** 23–39 (2016).

30. Liebeskind, D. S. Collateral circulation. *Stroke* **34,** 2279–2284 (2003).

31. Romero, J. R. *et al.* Cerebral collateral circulation in carotid artery disease. *Current cardiology reviews* **5,** 279–288 (2009).

32. Faber, J. E., Chilian, W. M., Deindl, E., van Royen, N. & Simons, M. A brief etymology of the collateral circulation. *Arteriosclerosis, thrombosis, and vascular biology* **34,** 1854–1859 (2014).

33. Ginsberg, M. D. The cerebral collateral circulation: relevance to pathophysiology and treatment of stroke. *Neuropharmacology* **134,** 280–292 (2018).

34. Bonati, L. H. *et al.* Length of carotid stenosis predicts peri-procedural stroke or death and restenosis in patients randomized to endovascular treatment or endarterectomy. *International Journal of Stroke* **9,** 297–305 (2014).

Figures and Tables

Figure 1: *An illustration of the varying stenosis models for the right and left carotid. All stenosis degrees were imposed onto a single set of patient imaging data with pairings of non-severe and severe carotid stenosis degrees applied to each carotid. The 100% severity, or contralateral carotid occlusion cases, were paired with each of the non-severe and severe stenosis degrees.*

a) Embolus tranpsort simulation framework for partially-occluded models

b) Embolus tranpsort simulation framework for fully-occluded models

Figure 2: *Design of the in silico experiments for embolus transport. Two sets of models were created for these simulations. a) Partially-occluded models were built using pairing of non-severe and severe stenosis degrees across both the left and right carotid. Emboli were released from the both carotids and the face of the aortic root for cardiogenic cases. This totaled to 18 unique models with differing pairing of severe and non-severe stenosis degrees with embolus releases from these three sources. b) 12 models were created in which one of the carotid arteries were fully-occluded (100% NASCET degree) and the other carotid was partially occluded. These pairings were completed to where there was one fully-occluded carotid and the other carotid was partially-occluded using each degree of severe and non-severe NASCET rankings. Emboli were released from the partially occluded carotid and the face of the aortic root.*

Figure 3: *The proportion of emboli that completed contralateral movement for each pairing of stenosis degrees. The x-axis labels each model with their respective left and right stenosis degree (i.e. 10L50R indicates the model with a 10% left carotid stenosis and a 50% right carotid stenosis).*

Figure 4: *(A)Percentage distribution of cardiogenic emboli to each CoW hemisphere. The x-axis labels each model with their respective left and right carotid stenosis degree. (B) Percentage distribution of volumetric flow rate through each of the CoW-feeding vessels: the basilar artery, the left common carotid, and the right common carotid. These distributions are shown across all combinations of severe and non-severe stenosis degrees where the basilar artery and non-severely stenosed carotid artery compensate for flow as the severe stenosis increases in degree.*

Figure 5: *Proportion of emboli that distributed contralaterally within the models with a fully-occluded carotid artery (contralateral carotid occluson or CCO cases). The x-axis indicated the stenosis severity of non-fully-occluded carotid with each bar designating a fully occluded left or right carotid.*

Figure 6: *(A)The proportion of cardiogenic emboli that travel to each hemisphere of the CoW of the models with a fully-occluded carotid artery. The left plot shows the distribution for the models with a full-occluded left carotid and the right plot shows the models with a fully-occluded right carotid. The y-axis indicates the stenosis severity of the non-fully-occluded carotid. (B) Flow distribution through each of the three arteries that provide blood flow to the CoW for the models with fully-occluded carotid artery. Left plot shows the distribution for the models with a fully-occluded right carotid and the right plot shows the distribution for the models with a fully-occluded left carotid. The y-axis designates the stenosis severity of the carotid paired with the fully-occluded carotid.*

a) Communicating arteries' average volumetric flow rates (mL/min) in models with a high-moderate/severe right carotid stenosis and a moderate left carotid stenosis

b) Communicating arteries' average volumetric flow rates (mL/min) in models with a high-moderate/severe left carotid stenosis and a moderate right carotid stenosis

Figure 7: *This figure shows the average volumetric flow rate through each artery of the CoW in mL/min. Each diagram gives the flow rate for a stenosis model at each paired stenosis severity. a) The flow rates for each model with a severely stenosed right carotid and a non-severe left carotid. b) The flow rates for each model with a severly stenoesed left carotid and non-severe right carotid*

a) Average volumetric flow rates with Circle of Willis for models with a fully-occluded right carotid

a) Average volumetric flow rates with Circle of Willis for models with a fully-occluded left carotid

Figure 8: *Average volumetric flow rates for each of the communicating segments in the CoW within the models including a fully-occluded carotid artery. a) Diagrams for the models witha fully-occluded right carotid paired with a left carotid with the respective severe and non-severe stenosis degree. b) Diagrams for the models with a fully-occluded left carotid paired with a right carotid with the respective non-severe and severe stenosis degrees.*

Figure 9: *A schematic concept sketch that aims to summarize the observations on contralateral embolus source-to-destination transport based on potential hemodynamic interactions due to differential flow recruitment through the basilar artery and less stenosed carotid artery, and due to proximal collateral flow routing at the Circle of Willis*

Understanding Thromboembolus Transport Patterns In The Brain For Stroke In The Presence Of Carotid Artery Stenosis

Ricardo Roopnarinesingh, Neel Jani, Michelle Leppert, Debanjan Mukherjee

Supplementary Material Information

S1 Animation of stenosis flow and embolus dynamics

Here, we describe the simulation animations that have been provided as additional illustration of the simulation data to support discussions presented in the main manuscript.

Animation titled: ***V1-StenosisLocalFlow.mp4*** shows the flow disturbance caused by the severe stenosis degree of 85% across all three moderate stenosis degrees. This flow disturbance results in a jet impingement that varies in length with respoect to across the contralateral carotid stenosis degree. This stenosis results in differing carotid embolus recruitment from the cardiogenic source and altered downstream embolus dynamics.

Animation titled: ***V2-40R85L-AllEmbolusReleases.mp4*** shows how all three embolus releases travel into the cerebral arteries from respective surface, presented for the non-CCO 40% stenosed right carotid with 85% stenosed left carotid case. Dynamic source-to-destination transport patterns, trends in embolus recruitment in the vertebrobasilar pathway, and contralateral movement into the anterior CoW vessels can bee seen in the video.

Animation titled: ***V3-ContralateralMovement.mp4*** shows one instance of how contralateral trans-hemispheric embolus movement can be enabled through the AcoA, with emboli moving from the right carotid release to the left hemisphere. This for the case of 10% right carotid stenosis paired with 85% left carotid stenosis.

Animation titled: ***V4-ContralateralMovement-DualView.mp4*** shows thromboemboli released from two sources, both from the right carotid but of differing stenosis severity. This animation illustrates how the likelihood of contralateral movement is higher in cases with higher severity pairing where the amount of particles traveling into the left hemipshere is visibly higher in the model with a 40% right severity and a 85% left severity.

Animation titled: ***V5-CompleteOcclusionTransport-DualView.mp4*** compared two differing cases of CCO with one having a 10% severity and the other 85%. Both the cardiogenic and carotid releases are shown where we can note two distinct mechanics: (1) the recruitment of cardiogenic embolus through the vertebral arteries that supplements the occluded and highly stenosed carotid arteries, (2) the distinctly different hemodynamics within the 85% stenosed carotid where the embolus movement is much more erratic and less intuitive due to lack of flow and the jet impingement.

S2 Mathematical model details for hemodynamics simulation

Here we present the foundational equations for the stabilized finite element hemodynamic simulations. Blood was modeled as a Newtonian fluid with a bulk density of 1.06 g/cc and viscosity of 4.0 cP. whose momentum and mass balance obeys the standard Navier-Stokes and continuity equations stated in Eq. S1 and S2 as stated below:

$$\rho_f \left(\frac{\partial \mathbf{u}}{\partial t} + (\mathbf{u} \cdot \nabla \mathbf{u}) \right) = -\nabla p + \nabla \cdot \tau + \rho_f \mathbf{g} \tag{S1}$$

$$\nabla \cdot \mathbf{u} = 0 \tag{S2}$$

The term τ represents the viscous stresses in blood, which is modeled as per a linear Newtonian stress vs strain rate relation as follows:

$$\tau = 2\mu_f \mathbf{D}(\mathbf{u}) = \mu_f(\nabla\mathbf{u} + \nabla\mathbf{u}^T) \tag{S3}$$

These equations are solved using matrix systems of equations obtained by decomposing a stabilized Petrov Galerkin finite element variational formulation of these equations over a mesh composed of linear tetrahedral elements. This variational formulation is stated in Eq. S4 as follows:

$$\rho_f\left(\mathbf{w}, \frac{\partial\mathbf{u}}{\partial t}\right)_\Omega + \rho_f(\mathbf{w}, (\mathbf{u}\cdot\nabla)\mathbf{u})_\Omega + 2\mu_f(\mathbf{D}(\mathbf{w}), \mathbf{D}(\mathbf{u}))_\Omega$$

$$- \rho_f(\mathbf{w}, \mathbf{b})_\Omega - (\nabla\cdot\mathbf{w}, p)_\Omega - (\nabla q, \mathbf{u})_\Omega + (q, \mathbf{u}\cdot\mathbf{n})_\Gamma$$

$$- \underbrace{(\mathbf{w}, \mathbf{h}_{\Gamma_e\in\Gamma_B})}_{\text{traction boundaries}} + \underbrace{(\mathbf{w}, \mathbf{h}_{0D})_{\Gamma_B}}_{\text{outflow boundaries}} + \sum_{e=1}^{N_{el}}\left[\underbrace{\left(\tau_{\text{supg}}\left(\mathbf{u}^h\cdot\nabla\right)\mathbf{w}^h, \mathcal{R}^h\right)_{\Omega,e}}_{\text{SUPG stabilization}} + \underbrace{\left(\tau_{pspg}\nabla q^h, \mathcal{R}^h\right)_{\Omega,e}}_{\text{PSPG stabilization}}\right] = 0 \tag{S4}$$

In these equations, $\mathbf{u}$, p are blood flow velocity and pressure, $\mathbf{w}$, q are the two test functions employed in the finite element solver, μ_f is the averaged blood viscosity, ρ_f is the averaged density of blood, $\mathbf{g}$ designates gravity, Ω denotes the computational domain of the arterial network, Γ and the subscripts of Γ denote the boundary faces of the computational domain, $\mathcal{R}^h$ is the residual of the momentum equation, and $\tau_{supg/pspg}$ are the SUPG stabilization and PSPG stabilization factors used for convection and pressure stability respectively. $\mathbf{h}_{0D}$ represents the terms used to integrate the boundary conditions at the outlets and inlets of the arterial network; while $\mathbf{h}_{\Gamma_e\in\Gamma_B}$ represents the other boundary conditions that may be necessary to run physiologically realistic flow simulations.

S3   Mathematical model details for embolus dynamics simulation

Embolic particle simulations were conducted using a custom-modified version of the Maxey-Riley equation stated in Eq.S5 as below:

$$m_p\frac{d\mathbf{v}_p}{dt} = \underbrace{\frac{1}{2}\rho_f C_D\left(\pi R_p^2\right)\|\mathbf{u} - \mathbf{v}_p\|(\mathbf{u} - \mathbf{v}_p)}_{\text{drag force}} + \underbrace{C_{sl}\rho_f R_p^2\frac{\sqrt{\mu_f}}{\sqrt{\rho_f\|\omega\|}}\left[(\mathbf{u} - \mathbf{v}_p)\times\omega\right]}_{\text{lift force (shear-gradient)}}$$

$$+ \underbrace{\mathcal{V}_p(-\nabla p + \nabla\cdot\tau)}_{\text{fluid stresses, undisturbed flow}} + \underbrace{C_{\text{am}}\frac{\rho_f\mathcal{V}_p}{2}\left[\frac{D\mathbf{u}}{Dt} - \frac{d\mathbf{v}_p}{dt}\right]}_{\text{added mass}} + \underbrace{(m_p - \rho_f\mathcal{V}_p)\mathbf{g}}_{\text{buoyancy}} + \underbrace{\mathcal{I}(\mathcal{P},\mathcal{W})\left[\Delta\mathbf{v}_p\left(e_{\text{lub}}\right)\right]}_{\text{elastohydrodynamic wall-collisions}}$$

$$\tag{S5}$$

where m_p notes the particle mass, $\mathcal{V}_p$ is the particle volume, R_p is the particle radius, and $\mathbf{v}_p$ is the translational particle velocity. $C_a m$ is the added mass coefficient, C_D is the drag coefficient, and $C_s l$ denotes the shear-gradient lift force. $\omega = \nabla \times \mathbf{u}$ is the vortcity of the flow where the particle is located. $\mathcal{I}(\mathcal{P},\mathcal{W})$ is an indicator that has a value of 1 when a particle $\mathcal{P}$ comes into contact with the artery wall, the value of this function is 0 otherwise. Next, the contribution of the momentum changed created by the elastohydrodynamic lubrication and particle-wall collisions is denoted by the term $[\delta\mathbf{v}_p(e_l ub]$ where $e_l ub$ is the restitution coefficient.

S4   Length of stenosis region

The length of the stenosed region across the span of the carotid artery was calculated using the equation below:

$$Length = \alpha * (1 + NASCET) * D_{CCA} \tag{S6}$$

where α is a constant factor used across all stenosis applications as described in [34]. NASCET is the associated NASCET stenosis degree applied for that model. D_{CCA} is the diameter of the undisturbed common carotid artery at that ipsilateral carotid artery. This relation, for a chosen NASCET severity, was used to reduce the segmentation diameters and re-loft the models to create stenosed carotid artery models.

S5 Complete distribution dataset for emboli across all vessels

For each simulation, the total proportion of embolus samples released was calculated across each outlet of our heart-to-brain model. All embolus source-to-distribution statistics reported in the manuscript were derived from these distributions. While in the manuscript the focus was on contralateral trans-hemispheric distribution, the complete set of embolus distribution across all vessel outlets individually is not included in the main manuscript. We provide a visual depiction of this complete dataset for each embolus release source as additional reference in the supplementary figures below. The vessel outlets included the anterior cerebral arteries (ACAs), the posterior cerebral arteries (PCAs), the middle cerebral arteries (MCAs), the subclavians, the external carotids, and the descending aorta (DA).

Figure S1: *Complete simulated embolus distribution data across all left carotid embolus releases, for all possible vessel outlets in the heart-to-brain pathway modeled for this study.*

Figure S2: *Complete simulated embolus distribution data across all right carotid embolus releases, for all possible vessel outlets in the heart-to-brain pathway modeled for this study.*

Figure S3: *Complete simulated embolus distribution data across all cardiogenic embolus releases, for all possible vessel outlets in the heart-to-brain pathway modeled for this study.*

S6 Recruitment of cardiogenic emboli at the carotid bifurcation

Our results specifically illustrate how the proximal collateral flow of the CoW influences distribution of embolus to the cerebral arteries. These additional results presented here help to illuminate how emboli are transported into both of the common carotid arteries but do not directly reflect the distribution of embolus into either hemisphere of the CoW. It is commonly understood that the right common carotid artery has a higher susceptibility of recruiting cardiogenic embolus that is reflected in our results shown here in Figure S4. Although there is a preferential transport to the right common carotid, this is not reflected at the level of cerebral distribution as there is a 55% and 45% average of cardiogenic embolus to the left and right CoW hemipshere respectively for the partially-occluded models 4. This helps to highlight the importance of the proximal collateral flow and the basilar flow compensation in the presence of varying stenosis as the laterality of embolus uptake at the common carotid arteries does not fully dictate the hemisphere distribution at the cerebral arteries.

Figure S4: *Distribution of cardiogenic embolus into the left and right common carotid arteries (CCAs). The y-axis denotes the bilateral stenosis degrees with the x-axis providing the percentage of embolus distributing into both of the CCAs. Embolus have a preference of traveling into the right CCA versus the left.*