Human bioaccumulation of micro- and nanoplastics is primarily determined by the organs' vascular volume

Alfonso M. Gañán-Calvo[1,2*]

[1]Laboratory of Engineering for Energy and Environmental
Sustainability, Universidad de Sevilla, 41092 Sevilla, Spain.
[2]Departmento de Ingeniería Aeroespacial y Mecánica de Fluidos,
ETSI, Universidad de Sevilla, Camino de los Descubrimientos,
41092 Sevilla, Spain.

Corresponding author(s). E-mail(s): amgc@us.es;

Abstract

Recent studies have demonstrated the accumulation of micro- and
nanoplastics (MNPs) in human organs, particularly in the brain,
at an alarming rate (Campen et al., 2024, doi.org/10.21203/rs.3.rs-
4345687/v1). This study examines the phenomenon of bioaccumulation
in relation to up to six distinct biological and biomedical parameters
pertaining to vascular structure and composition in seven organ tis-
sues: the brain cortex, kidney, liver, testis, and three arteries (coronary,
carotid and aorta). As anticipated based on fundamental mechanis-
tic considerations of bioaccumulation (washing rate), the MNP content
appears to be proportional to the inverse of the capillary volume frac-
tion (V_c), exhibiting no correlation with any other geometric variable,
such as capillary diameter or capillary segment length. This finding
is consistent with the small size spectra of MNPs in comparison to
these dimensions. However, contrary to expectations, no correlation
was observed between MNP content and biological composition (lipid,
water, or carbohydrate content), while a likely non-causal correlation
was observed between MNP content and protein content, which may
be simply attributed to the non-linear correlation between protein
content and V_c. The conclusions that point to a possibly exceed-
ingly large number of MNP particles currently in our brains call for
urgent further investigation and the revision of the published data.

Keywords: MNP, microplastics, brain, vascular capillary volume, tissue washing rate

1 Introduction

In contrast to single-celled organisms, which rely entirely on the complex biochemistry and mechanics [1] of their surrounding membranes, organ tissues of multicellular organisms are intricate biological networks whose main common key to survival in their environment is their ability to maintain their biological structure and function over very long time scales (e.g. the lifetime of the global organism) compared to those of the residence time of vital fluids through the organ. These vital fluids, such as aqueous complex suspensions (blood, lymph, sap), seawater, or air, deliver nutrients in a timely manner and flush away metabolic products, unwanted compounds, and pathogens through specialized capillary networks whose geometric dimensions are determined by the function of each organ, as well as the diffusion and biochemistry of each nutrient and metabolite [2, 3]. These dimensions range from a few microns to millimeters. All living organisms have a multiplicity of protecting evolutionary mechanisms against the entry and accumulation of natural undesired particles below these dimensions. Among these mechanisms, continuous washout by vital fluids is prevalent in multicellular organisms.

In the natural environment, micro- and nanoplastics (MNP) are a new class of microscopic particles whose rapidly increasing presence and ubiquity in the troposphere may have the highest growth rate compared to any natural evolutionary time scale on Earth, with the exception of wide-range explosive aerosol releases due to giant volcanic activity or the impact of large extraterrestrial objects. Contrary to the higher mass density of the particles (minerals) released by these latter rare catastrophes, the density of the MNP particles (polymers such as polypropylene, polystyrene, PVC or SBD) is close to that of water and their mechanical diffusion is inextricably linked to this omnipresent vital fluid. As a consequence, on the one hand, the presence of MNP in the tissues of live organisms is currently unavoidable, and on the other hand, the biochemical abilities of the organs to get rid of these particles, which are even coronated by biological matter [4–8] that bio-interact with the organs, are hampered by the long evolutionary timescales. In addition, the toxicity of these particles[9], especially in mammals [10], is increasingly a matter of concern.

In practice, MNP enter the bloodstream through three primary routes: ingestion (gastrointestinal tract), inhalation (respiratory system), and dermal absorption (skin). Each route involves different absorption mechanisms that are influenced by the physicochemical properties of MNP and the biological barriers they encounter [5, 11]. Once in the bloodstream, and depending on several factors, including size, surface charge, lipid solubility, and the presence of specific transport mechanisms, probably the most concerning long-term issue

for our species is that the MNP can cross the blood-brain barrier (BBB) [12]. Regarding their evacuation and excluding microglial phagocytosis for obvious reasons, despite the sophisticated existing evacuation mechanisms[13] such as the glymphatic system, the meningeal lympatic vessels, or even transcytosis across the BBB, the brain lacks efficient excretion routes for MNP, especially for certain intermediate nanometric ranges (i.e. larger than about 100 nm). In general, the ease of transport and entry of MNP particles, their resistance to degradation, and their limited excretion pathways make them susceptible to bioaccumulation with no evolutionary time to compensate for their potential damage. Even so, for simple mechanistic reasons, organs with greater vascular clearance tend are expected to be less prone to accumulate.

Thus, while the specific biochemistry and biophysics associated with the interaction of MNP with different tissues may pose different aspects to consider, a natural hypothesis is whether the long-term human bioaccumulation of MNP could be mainly proportional to the inverse of the vascular capillary volume per unit volume, or simply the vascular volume fraction V_c of a given tissue.

In a recent development disclosure, Campen and colleagues[14] have prompted a tipping concern within the scientific community with the discovery of markedly elevated levels of MNP in the brain, kidneys, and liver, with an alarming increase observed from 2016 to 2024. In addition, other authors have almost simultaneously raised concerns about the levels of MNP found in canine and human testis [15] and in human arteries [16]. The data disclosed are presented at an opportune moment, allowing for a timely assessment of the preceding hypothesis.

2 Methods

In this study, we investigate the potential correlations between the concentration of MNP in seven distinct tissues and six primary variables: biochemical (lipid, protein and carbohydrate average content) and biomechanical (V_c, capillary diameter D and number density N_d of capillaries)[2, 3, 17, 18].

All MNP content values in tissue samples were determined by pyrolysis gas chromatography/mass spectrometry (Py-GC/MS), according to the detailed methods and supplementary material disclosed in each publication[14–16]. The values and statistics of the total mass concentration of MNP (ww) are taken directly from the data provided by the authors.

The data on the biochemical composition [19–36] and biological morphometry [35, 37–51] of the tissues analyzed and their statistical variability were comprehensively collected and rationalized from the established literature using currently available and highly reliable Large Languaje Models by Google and OpenAI, with double checking - two separate requests per variable evaluated. In particular, the values of total vascular volume V_c considered in this work include blood and lymphatic capillaries, as well as those capillaries related to the particular function of each organ, such as glomerular and

Fig. 1 The average MNP content and typical deviation found in 2016 [14], and in 2023-2024 [14?] versus the vascular capillary volume V_c for the seven organ tissues investigated. MNP is inversely proportional to V_c (dashed lines). For illustrative purposes, a typical deviation of ±20% is also provided for the V_c values. Solid and lighter tones of the same colors correspond to 2024 and 2016 from Campen et al. [14], respectively. The estimation for the concentration of MNP in the brain white matter (WM) based of the model proposed is also given.

peritubular capillaries, vasa recta (kidney); sinusoids (liver) and seminiferous tubules (testis).

3 Results

Figure 1 depicts the concentration of MNP in weight, as determined by Py-GC/MS, in tissues from distinct organs of decedent humans[14, 15] or patients with endarterectomy [16]. While the results generally support our hypothesis, the particularly precise correlation between MNP concentration and the inverse of V_c for the 2024 samples is noteworthy for brain, kidney, liver and testis tissues. The MNP concentrations from arteries show, as expected, slightly larger deviations from the nearly perfect correlation of the previous tissue data, since the arteria's values of V_c correspond to the highly heterogeneous but aggregated values of the arterial lumen and vasa vasorum, with very heterogeneous flushing rates and morphological structure. Nevertheless, the global trend, as postulated, is evident in all seven different tissues.

Figure 3, in contrast, demonstrates that, excluding protein concentration, there is no correlation between MNP content and the other variables, including the capillary diameter D. This suggests that the potential correlation between MNP aggregation and capillary clogging, which could be a concern for bioaccumulation, is unlikely to be a significant factor. The results indicate that the degree of aggregation is not sufficient to form large particles from the very small MNP size spectrum that eventually enters the circulatory system.

Figure 3 illustrates an intriguing anticipated correlation between V_c and protein content, which excludes the direct biochemical involvement of proteins in the correlation between their concentration and that of MNP. In reality, the critical and ubiquitous role of specific fibrin proteins in the lining (surface) of capillaries that create the vascular space provides an explanation for the salient proportionality between V_c and the protein mass concentration, with

Fig. 2 Plots of MNP content versus four additional variables. The double points (with the same color code as in fig 1) correspond to 2016 and 2024 data, where the lower MNP values represented by the same but lighter colors are always those of 2016. No correlations are observed.

Fig. 3 Protein content of the seven tissues investigated versus the V_c. Dashed line correspond to the linear expression in the legend.

the exception of brain tissue. In the latter case, the concentration of specific proteins (i.e., myelin) in neurons is associated with brain function but is not related to fluid circulation, which creates a clear deviation from the observed correlation for the other tissues.

4 Discussion

From the strong correlation found between MNP and the inverse of Vc, we estimate the MNP content of the brain white matter (BWM), whose V_c is roughly half that of the cortex, and found it to be about 0.8% by weight on average if the data for liver, kidney, testis and brain cortex measured in 2024 are extrapolated for the V_c of BWM. This means that about 10 grams of MNP could currently be present in the average elderly human brain. The extremely large number of MNPs resulting from this calculation, amounting to millions per neuron, and their very likely interference with cognitive functions[9, 52–60], even if this number were reduced by several orders of magnitude, require urgent further investigation.

Our findings also indicate that the brain and other organs with low vascular volume may undergo selective bioaccumulation of a purely mechanistic and therefore reliable nature when pharmaceuticals are delivered in controlled doses via slowly biodegradable polymers. This result has significant implications for the field of pharmacology.

Finally, it can be reasonably assumed that the observed bioaccumulation is linear in time, a conservative approach that allows for the possibility that the rate of accumulation may have been established within the observed time frame. This time frame encompasses the period of the global pandemic caused by the SARS-CoV-2 virus, as well as the extensive use of plastic fabric face masks worldwide[61]. The observed increase in bioaccumulation, which can be estimated to be approximately a doubling over the span of eight years (see figure 1), is consistent with the hypothesis that the highest rate of accumulation was established during this period.

5 Limitations

To obtain a more accurate and comprehensive evaluation of the proposed simple flushing mechanism, it is essential to extend the analysis to other organs, such as the pancreas, spleen, lungs, and muscles. Furthermore, a detailed mechanistic analysis is required to elucidate the underlying mechanisms responsible for the inverse proportionality between the observed vascular volume fraction and bioaccumulation. Additionally, this study has not addressed the biochemical aspects of MNP penetration through biological barriers, such as the BBB, cell membrane, or intestinal lining, which must be incorporated in subsequent studies. These additional insights would facilitate comprehension of the role of global statistics in the mechanisms of tissue bioaccumulation when new particles are introduced to biological organisms en masse. Furthermore, they would enable the understanding and exploitation of the potential significance of biochemical processes that may be pivotal in this context.

Declarations

- Funding: Partial funding from the the Spanish Ministry of Science and Innovation, grant no. PID2022-14095OB-C21.
- Conflict of interest/Competing interests (check journal-specific guidelines for which heading to use): The author declares no conflicts of interest.
- Ethics approval: Not applicable
- Consent to participate: Not applicable
- Consent for publication: Not applicable
- Availability of data and materials: Data and materials can be made available upon reasonable request.
- Code availability: Not applicable
- Authors' contributions: AMGC conception, calculations, writing, and funding.

References

[1] Flaum, E., Prakash, M.: Curved crease origami and topological singularities enable hyperextensibility of L. olor. Science **384**, 5511 (2024)

[2] Epp, R., Schmid, F., Weber, B., Jenny, P.: Predicting Vessel Diameter Changes to Up-Regulate Biphasic Blood Flow During Activation in Realistic Microvascular Networks. Front. Physiol. **11**, 566303 (2020)

[3] Celaya-alcala, J.T., Lee, G.V., Smith, A.F., Secomb, T.W.: Simulation of oxygen transport and estimation of tissue perfusion in extensive microvascular networks : Application to cerebral cortex. J. Cereb. Blood Flow Metab. **41**, 656–669 (2021)

[4] Gopinath, P.M., Saranya, V., Vijayakumar, S.: Assessment on interactive prospectives of nanoplastics with plasma proteins and the toxicological impacts of virgin , coronated and environmentally. Sci. Rep. **9**, 8860 (2019)

[5] Prata, J.C., João, P., Lopes, I., Duarte, A.C., Rocha-santos, T.: Science of the Total Environment Environmental exposure to microplastics : An overview on possible human health effects. Sci. Total Environ. **702**, 134455 (2020)

[6] Wang, L., Wu, W.-m., Bolan, N.S., Tsang, D.C.W., Li, Y., Qin, M., Hou, D.: Environmental fate , toxicity and risk management strategies of nanoplastics in the environment : Current status and future perspectives. J. Hazard. Mater. **401**, 123415 (2021)

[7] Ali, N., Katsouli, J., Marczylo, E.L., Gant, T.W., Wright, S., Bernardino, J., Serna, D.: Review The potential impacts of micro-and-nano plastics on various organ systems in humans. eBioMedicine **99**, 104901 (2024)

[8] Brouwer, H., Porbahaie, M., Boeren, S., Busch, M., Bouwmeester, H.: The in vitro gastrointestinal digestion - associated protein corona of polystyrene nano - and microplastics increases their uptake by human THP - 1 - derived macrophages. Part. Fibre Toxicol., 1–26 (2024)

[9] Casella, C., Ballaz, S.J.: Genotoxic and neurotoxic potential of intracellular nanoplastics : A review. J. Appl. Toxicol., 1–22 (2024)

[10] Banerjee, A., Shelver, W.L.: Science of the Total Environment Micro- and nanoplastic induced cellular toxicity in mammals : A review. Sci. Total Environ. **755**, 142518 (2021)

[11] Leslie, H.A., Velzen, M.J.M.V., Brandsma, S.H., Vethaak, A.D., Garcia-vallejo, J., Lamoree, M.H.: Discovery and quantification of plastic particle pollution in human blood. Environ. Int. **163**, 107199 (2022)

[12] Kopatz, V., Wen, K., Kov, T., Keimowitz, A.S., Pichler, V., Widder, J., Vethaak, A.D., Holl, O., Kenner, L.: Micro- and Nanoplastics Breach the Blood – Brain Barrier (BBB): Biomolecular Corona ' s Role Revealed. Nanomaterials **13**, 1404 (2023)

[13] Iliff, J.J., Wang, M., Liao, Y., Plogg, B.A., Peng, W., Gundersen, G.A., Benveniste, H., Vates, G.E., Deane, R., Goldman, S.A., Nagelhus, E.A., Nedergaard, M.: A Paravascular Pathway Facilitates CSF Flow Through the Brain Parenchyma and the Clearance of Interstitial Solutes, Including Amyloid β. Sci. Transl. Med. **4**, 147–111 (2012)

[14] Campen, M., Nihart, A., Garcia, M., Gonzalez-estrella, J.: Bioaccumulation of Microplastics in Decedent Human Brains Assessed by Pyrolysis Gas Chromatography-Mass Spectrometry, 1–11

[15] Hu, C.J., Garcia, M.A., Nihart, A., Liu, R., Yin, L., Adolphi, N., Gallego, D.F., Kang, H., Campen, M.J., Yu, X.: Microplastic presence in dog and human testis and its potential association with sperm count and weights of testis and epididymis. Toxicol. Sci. **200**, 235–240 (2024)

[16] Liu, S., Wang, C., Yang, Y., Du, Z., Li, L., Zhang, M., Ni, S., Yue, Z., Yang, K., Wang, Y., Li, X., Yang, Y., Qin, Y., Li, J., Yang, Y., Zhang, M.: Microplastics in three types of human arteries detected by pyrolysis-gas chromatography/mass spectrometry (Py-GC/MS). J. Hazard. Mater. **469**, 133855 (2024)

[17] Cassot, F., Lauwers, F., Fouard, C., Prohaska, S., Lauwers-Cances, V.: A Novel Three-Dimensional Computer-Assisted Method for a Quantitative Study of Microvascular Networks of the Human Cerebral Cortex. Microcirculation **13**, 1–18 (2006)

[18] Lauwers, F., Cassot, F., Lauwers-cances, V., Puwanarajah, P., Duvernoy, H.: Morphometry of the human cerebral cortex microcirculation : General characteristics and space-related profiles. Neuroimage **39**, 936–948 (2008)

[19] Wolinsky, H.: Response of the rat aortic media to hypertension: Morphological and chemical studies. Circulation Research **26**(4), 507–522 (1970)

[20] Bailey, A.J., Robins, S.P.: The chemical composition of atherosclerotic plaques. Atherosclerosis **22**(1), 77–94 (1975)

[21] Roach, M.R., Burton, A.C.: The reason for the shape of the distensibility curves of arteries. Canadian Journal of Biochemistry and Physiology **35**(8), 681–690 (1957)

[22] Dingemans, K.P., Teeling, P., Lagendijk, J.H., Becker, A.E.: Extracellular matrix of the human aorta in relation to age and atherosclerosis. Cardiovascular Research **46**(2), 345–355 (2000)

[23] Bailey, A.J.: The chemical changes occurring in the collagen of aging mammalian connective tissues. Mechanisms of Ageing and Development **14**(3–4), 255–268 (1980)

[24] Hayashi, K., Nagasawa, S.: Age-related changes in the mechanical properties of human major arteries. Journal of Applied Physiology **49**(1), 175–181 (1980)

[25] Setchell, B.P., Waites, G.M.H.: The blood-testis barrier. In: Astwood, E.B., Greep, R.O. (eds.) Handbook of Physiology, Section 7: Endocrinology, Volume V, Male Reproductive System, pp. 143–172 (1975)

[26] Fawcett, D.W., Phillips, D.M.: The fine structure and function of the sertoli cell. In: McKerns, K.W. (ed.) The Gonads, pp. 107–155 (1969)

[27] Russell, L.D., Griswold, M.D.: The Sertoli Cell. Cache River Press, Clearwater, FL (1993)

[28] Christensen, A.K.: Leydig cells. In: Astwood, E.B., Greep, R.O. (eds.) Handbook of Physiology, Section 7: Endocrinology, Volume V, Male Reproductive System, pp. 57–94 (1975)

[29] Mather, J.P., Phillips, D.M.: Primary culture of testicular somatic cells. Journal of Tissue Culture Methods **9**(1), 43–48 (1984)

[30] Wing, T.Y., Christensen, A.K.: Morphometric studies on rat leydig cells. Journal of Cell Biology **93**(1), 1178–1184 (1982)

[31] Alberts, B., Johnson, A., Lewis, J., Morgan, D., Raff, M., Roberts, K.,

Walter, P.: Molecular Biology of the Cell, 6th edn. Garland Science, New York (2015)

[32] Bloom, W., Fawcett, D.W.: A Textbook of Histology, 10th edn. W. B. Saunders Company, Philadelphia (1975)

[33] Syková, E., Nicholson, C.: Diffusion in brain extracellular space. Physiological Reviews **88**(4), 1277–1340 (2008)

[34] Nicholson, C., Syková, E.: Extracellular space structure revealed by diffusion analysis. Trends in Neurosciences **21**(5), 207–215 (1998)

[35] Kaissling, B., Le Hir, M.: The renal cortical interstitium: Morphological and functional aspects. Histochemistry and Cell Biology **130**(2), 247–262 (2008)

[36] Arias, I.M., Boyer, J.L., Chisari, F.V., Fausto, N., Schachter, D., Shafritz, D.A. (eds.): The Liver: Biology and Pathobiology, 5th edn. John Wiley & Sons, Hoboken, NJ (2009)

[37] Weibel, E.R.: Stereological Methods: Practical Methods for Biological Morphometry. Academic Press, London (1979)

[38] Duvernoy, H.M., Delon, S., Vannson, J.-L.: Cortical blood vessels of the human brain. Brain Research Bulletin **7**(5), 519–579 (1981)

[39] Braet, F., Wisse, E.: Structural and functional aspects of liver sinusoidal endothelial cell fenestrae: a review. Comparative Hepatology **1**(1), 1 (2002)

[40] Pallone, T.L., Edwards, A., Mattson, D.L., Cowley, A.W.: Countercurrent exchange in the renal medulla. American Journal of Physiology-Renal Physiology **284**(6), 1059–1072 (2003)

[41] Setchell, B.P.: The Mammalian Testis. Cornell University Press, Ithaca, NY (1978)

[42] Johnson, L., Neaves, W.B.: Age-related changes in the leydig cell population, seminiferous tubules, and sperm production in stallions. Biology of Reproduction **24**(4), 703–712 (1981)

[43] Zirkin, B.R., Papadopoulos, V.: Leydig cells: Formation, function, and regulation. Biology of Reproduction **99**(1), 101–111 (2018)

[44] Heistad, D.D., Marcus, M.L.: Role of vasa vasorum in nourishment of the aorta. Blood Vessels **16**(5), 225–238 (1979)

[45] Barger, A.C., Beeuwkes, R.: Reevaluation of the morphology of the

coronary arteries. Circulation Research **23**(1), 1–8 (1968)

[46] Gössl, M., Rosol, M., Malyar, N.M., Fitzpatrick, L.A., Beighley, P.E., Zamir, M., Ritman, E.L.: Functional anatomy and hemodynamic characteristics of vasa vasorum in the coronary arteries. Journal of the American College of Cardiology **42**(5), 1054–1060 (2003)

[47] Phelps, C.H., Kennedy, D.N.: Measurement of capillary density in cerebral cortex. Journal of Comparative Neurology **149**(3), 401–412 (1973)

[48] Vita, J.A., Treasure, C.B., Ganz, P., Fish, R.D., Gimbrone, M.A., Selwyn, A.P.: Vasa vasorum flow and aortic wall metabolism in experimental atherosclerosis. Circulation Research **66**(4), 932–944 (1990)

[49] Nichols, W.W., O'Rourke, M.F., Vlachopoulos, C.: McDonald's Blood Flow in Arteries: Theoretical, Experimental and Clinical Principles, 6th edn. CRC Press, Boca Raton, FL (2011)

[50] Standring, S. (ed.): Gray's Anatomy: The Anatomical Basis of Clinical Practice, 41st edn. Elsevier, Philadelphia (2016)

[51] Iliff, J.J., Wang, M., Liao, Y., Plogg, B.A., Peng, W., Gundersen, G.A., Benveniste, H., Vates, G.E., Deane, R., Goldman, S.A., Nagelhus, E.A., Nedergaard, M.: A paravascular pathway facilitates csf flow through the brain parenchyma and the clearance of interstitial solutes, including amyloid β. Science Translational Medicine **4**(147), 147–111 (2012)

[52] Liu, Z., Sokratian, A., Duda, A.M., Xu, E., Stanhope, C., Fu, A., Strader, S., Li, H., Yuan, Y., Bobay, B.G., Sipe, J., Bai, K., Lundgaard, I., Liu, N., Hernandez, B., Rickman, C.B., Miller, S.E., West, A.B.: Anionic nanoplastic contaminants promote Parkinson ' s disease – associated α-synuclein aggregation. Sci. Adv. (9), 8716 (2023)

[53] Sharma, A., Kaur, M., Sharma, K., John, P.: Neurotoxicology Nano polystyrene induced changes in anxiety and learning behaviour are mediated through oxidative stress and gene disturbance in mouse brain regions. Neurotoxicology **99**, 139–151 (2023)

[54] Jeong, A., Jung, S., Jeong, E., Won, K.: Nanoplastics exacerbate Parkinson ' s disease symptoms in C . elegans and human cells. J. Hazard. Mater. **465**, 133289 (2024)

[55] Li, G., Liu, X., Sun, X., Huang, L., Kuang, W., Ou, J., Zhang, J., Zhang, Z., Li, H., Tang, H., Feng, C., Gu, L.: Polystyrene microplastics induce anxiety via HRAS derived PERK-NF- κ B pathway. Environ. Int. **185** (2024)

[56] Paing, M.M.Y., Eom, Y., Bae, G., Kim, B., Gil, M., Hong, S., Hoon, S.: Science of the Total Environment Neurotoxic effects of polystyrene nanoplastics on memory and microglial activation : Insights from in vivo and in vitro studies. Sci. Total Environ. **924**, 171681 (2024)

[57] Sun, H., Yang, B., Zhu, X., Li, Q., Song, E., Song, Y.: Oral exposure of polystyrene microplastics and doxycycline affects mice neurological function via gut microbiota disruption : The orchestrating role of fecal microbiota transplantation. J. Hazard. Mater. **467**, 133714 (2024)

[58] Sun, M., Zhang, M., Di, F., Bai, W., Sun, J., Zhang, M., Sun, J., Li, M., Liang, X.: Ecotoxicology and Environmental Safety Polystyrene nanoplastics induced learning and memory impairments in mice by damaging the glymphatic system. Ecotoxicol. Environ. Saf. **284**, 116874 (2024)

[59] Wang, L., Li, S., Hao, Y., Liu, X., Liu, Y., Zuo, L., Tai, F., Yin, L., Young, L.J., Li, D.: Science of the Total Environment Exposure to polystyrene microplastics reduces sociality and brain oxytocin levels through the gut-brain axis in mice. Sci. Total Environ. **945**, 174026 (2024)

[60] Zheng, Y., Xu, S., Liu, J., Liu, Z.: The effects of micro- and nanoplastics on the central nervous system : A new threat to humanity ? Toxicology **504**, 153799 (2024)

[61] Zhao, B., Rehati, P., Yang, Z., Cai, Z., Guo, C., Li, Y.: Science of the Total Environment The potential toxicity of microplastics on human health. Sci. Total Environ. **912**, 168946 (2024)