Beyond Tumors: Reduced survival linked to pathogenic *PIK3CA* and *TP53* post-zygotic variants in the uninvolved breast tissue of recurrent cancer patients

Maria Andreou[1], Katarzyna Chojnowska[1], Natalia Filipowicz[1], Monika Horbacz[1], Piotr Madanecki[2], Katarzyna Duzowska[1], Urszula Ławrynowicz[1], Hanna Davies[3], Bożena Bruhn-Olszewska[3], Mikołaj Koszyński[1], Kinga Drężek-Chyła[1], Maciej Jaśkiewicz[1], Marcin Jąkalski[1], Anna Kostecka[1], Marta Drzewiecka-Kłysz[4], Magdalena Nowikiewicz[5], Manuela Las-Jankowska[6,7], Dariusz Bała[6,8], Jacek Hoffman[4], Ewa Śrutek[9,10], Michał Jankowski[6,8], Jerzy Jankau[11], Diana Hodorowicz-Zaniewska[12], Joanna Szpor[13], Łukasz Szylberg[10,14], Wojciech Zegarski[6,8], Tomasz Nowikiewicz[4,6], Patrick G. Buckley[15], Irene Tiemann-Boege[16], Jakub Mieczkowski[1], Magdalena Koczkowska[1], Jan P. Dumanski[1,3], Arkadiusz Piotrowski[1,*]

ABSTRACT

Histologically normal mammary tissue from breast cancer patients can harbor significant genetic alterations. We analyzed the spectrum of DNA variants in 408 matched histologically normal tissue and tumors from 77 poor-prognosis patients, 49 patients recruited without prognosis bias, and mammary gland samples from 15 non-cancerous individuals. Whole exome sequencing revealed a higher prevalence of pathogenic post-zygotic variants in cancer-associated genes: *AKT1*, *PIK3CA*, *PTEN*, *TBX3*, and *TP53*, affecting poor-prognosis patients (29%), compared to 12.5% in those without prognosis criteria (p=0.0008578). *PIK3CA* variants were recurrent across patients, while *TP53* variants were restricted to those with adverse prognoses. Duplex sequencing detected low-frequency pathogenic *PIK3CA* and *TP53* variants in distant normal tissues of poor-prognosis patients. Disease recurrence significantly reduced survival rates, with poor prognosis patients experiencing higher mortality within 24 months (p=0.0088), further worsened by the presence of pathogenic post-zygotic variants. These findings highlight the importance of genetic monitoring even in microscopically normal mammary tissue.

[1]3P-Medicine Laboratory, Medical University of Gdańsk, Gdańsk, Poland

[2]Department of Biology and Pharmaceutical Botany, Medical University of Gdańsk, Gdańsk, Poland

[3]Department of Immunology, Genetics and Pathology and Science for Life Laboratory, Uppsala

University, Uppsala, Sweden

[4]Department of Clinical Breast Cancer and Reconstructive Surgery, Oncology Center-Prof. Franciszek

Łukaszczyk Memorial Hospital, Bydgoszcz, Poland

[5]Department of Hepatobiliary and General Surgery, A. Jurasz University Hospital, Bydgoszcz, Poland

[6]Chair of Surgical Oncology, Ludwik Rydygier's Collegium Medicum, Bydgoszcz, Nicolaus Copernicus

University, Toruń, Poland

[7]Department of Clinical Oncology, Oncology Center-Prof. Franciszek Łukaszczyk Memorial Hospital,

Bydgoszcz, Poland

[8]Department of Surgical Oncology, Oncology Center-Prof. Franciszek Łukaszczyk Memorial Hospital,

Bydgoszcz, Poland

[9]Surgical Oncology, Ludwik Rydygier's Collegium Medicum, Bydgoszcz, Nicolaus Copernicus

University, Toruń, Poland

[10]Department of Tumor Pathology and Pathomorphology, Oncology Center-Prof Franciszek Łukaszczyk

Memorial Hospital, Bydgoszcz, Poland

[11]Department of Plastic Surgery, Medical University of Gdańsk, Gdańsk, Poland

[12]Ist Department of General Surgery, Jagiellonian University Medical College, Breast Unit University

Hospital, Krakow, Poland

[13]Department of Pathomorphology, Jagiellonian University Medical College, Kraków, Poland

[14]Department of Clinical Pathomorphology, Collegium Medicum in Bydgoszcz, Nicolaus Copernicus

University, Toruń, Poland

[15]Genseq, Dublin, Ireland

[16]Institute of Biophysics, Johannes Kepler University, Linz, Austria

A.P., J.P.D, and M. Koczkowska contributed equally.

*Correspondence: Arkadiusz Piotrowski, 3P-Medicine Laboratory, Medical University of Gdańsk,

Gdańsk, Poland, arkadiusz.piotrowski@gumed.edu.pl

Keywords: breast cancer, recurrence, post-zygotic variants, unfavorable outcome, uninvolved mammary

gland, mortality

Abbreviations

BCAP: Breast Cancer Adverse Prognoses

BCS: Breast-conserving surgery

BCUS: Breast Cancer Un-Selected

BL: Whole Peripheral Blood

PT: Primary Tumor

RM: Reduction Mammoplasty

SK: Skin

UM: Uninvolved mammary gland

UMD: Distal uninvolved mammary gland

WES: Whole Exome Sequencing

INTRODUCTION

Breast cancer accounts for 12.5% of global annual cancer diagnoses, with the incidence rate increasing by

0.5% annually in recent years[1,2] Despite an overall 42% reduction between 1989 and 2021, primarily

attributed to increased awareness and early detection, breast cancer still constitutes one of the leading

causes of death among women[3]. Notably, stage I, low-risk breast cancer cases still present a 15-20%

chance of recurrence even two decades after the initial diagnosis[4]. While 5-10% of breast cancer cases are

hereditary, with 25-30% of heritable breast cancer risk attributed to pathogenic variants in genes of high

and moderate penetrance[5], the majority of cases are considered sporadic[6,7].

Research has recently focused on the normal mammary gland within the affected breast for early

detection of tumor formation at the molecular level, preceding changes on imaging or palpative screens[8].

Concurrently, breast-conserving surgery (BCS), which aims to remove the tumor and preserve the

remaining healthy breast tissue, stands as the preferred approach[9,10]. However, mammary gland tissue

from breast cancer patients, although appearing normal, has been found to harbor significant genomic and

transcriptomic alterations[11–14]. In particular, non-tumorous tissue from patients undergoing BCS shows

clearly pathogenic low-level post-zygotic alterations in the *PIK3CA* and *TP53* genes, raising questions

regarding its oncogenic potential [15,16].

Nevertheless, the association between post-zygotic alterations in ostensibly normal mammary gland tissue

of breast cancer patients and their clinical utility remains unclear. Hence, we screened paired uninvolved

mammary gland (UM) and primary tumor (PT) samples of reportedly sporadic breast cancer patients with

adverse outcomes within 10 years post-original surgery for the presence of post-zygotic alterations. We

compared our findings with a second breast cancer cohort of reportedly sporadic patients recruited

without specific prognosis criteria. Additionally, normal mammary gland samples from individuals with

no personal or family history of cancer were included as controls (study overview in Supplementary

Figure 1).

Here, we demonstrate that pathogenic post-zygotic variants in cancer-associated genes are commonly

found in histologically normal mammary tissue of breast cancer patients with adverse prognoses. These

variants, often also found in corresponding primary tumors, are linked with patient survival, emphasizing

the need for molecular screening to improve the clinical management of patients.

MATERIALS & METHODS

Patient recruitment, sample collection, and DNA isolation

We carried out Whole Exome Sequencing (WES) on 408 samples from three distinct groups: two cohorts

of female breast cancer patients with differing prognostic outcomes and a control group composed of

female individuals who underwent mammoplasty surgery for non-cancer-related reasons. The assignment

of patients to the cohorts was determined by their clinical prognoses. The first cohort included 77

reportedly sporadic breast cancer patients with adverse outcomes. All individuals in this group

experienced either recurrent disease, such as local recurrence/metastasis to the breast or secondary organs

(n=40), developed a second independent tumor (n=18), or both (n=8), and/or succumbed to the disease

(n=45) within the proceeding 10 years (Breast Cancer Adverse Prognoses cohort, BCAP), (Table 1,

Supplementary Table 1). The second cohort included 49 individuals from the same ethnic population,

diagnosed with reported sporadic breast cancer but recruited without specific criteria related to prognosis

(Breast Cancer Un-Selected cohort, BCUS). Within this group, 5 out of 49 patients experienced

recurrence, and 3 of them died within 2 years post-surgery; however, the follow-up period for this cohort

was considerably shorter compared to the BCAP cohort (Table 1, Supplementary Table 1). The majority

of BCAP and BCUS patients were treated with BCS (n=63 and n=31 respectively) versus mastectomy

(n=12 and n=18 respectively) (Supplementary Table 1) (data missing for 2 BCAP patients). All recruited

individuals did not receive neoadjuvant therapy. The control group comprised 15 individuals who

underwent reduction mammoplasty surgeries and had no personal or familial history of cancer (Reduction

Mammoplasty cohort, RM). Written informed consent was obtained from all enrolled individuals. The

study was approved by the the Bioethical Committee at the Collegium Medicum, Nicolaus Copernicus

University in Toruń (approval number KB509/2010) and by the Independent Bioethics Committee for

Research at the Medical University of Gdańsk (approval number NKBBN/564/2018 with multiple

amendments), recruited and enrolled all donors under informed and written consent, collected, and stored

all tissue samples. A significant age difference was observed between BCAP, BCUS, and RM cohorts

(Kruskal-Wallis test, p=7.7e-05), with the BCAP and BCUS cohorts being significantly older than the

control group (Kruskal–Wallis H test, p=0.000034 and p=0.00036 for BCAP and BCUS respectively).

However, no significant difference was observed between the BCAP (median age: 62, range: 23-85) and

BCUS median age: 65, range: 37-84) cohorts (Kruskal–Wallis H test, p=0.082). A total of 415 samples,

including PT, UM, UMD, BL, or SK from all three cohorts were collected by the Oncology Centre in

Bydgoszcz, Jagiellonian University Hospital in Cracow, and the University Clinical Centre in Gdańsk,

with the necessary ethical approvals and written informed consent from participants and deposited in the

biobank of our unit at the Medical University of Gdańsk, along with clinical data, including follow-up

information (Table 1, Supplementary Table 1). Distal UM samples (UMD, 1.5-3 cm from PT, median

2.35 cm), available for 7 BCAP patients, were collected and included in the downstream confirmatory

analysis, however, they were not initially sequenced. For the RM cohort, sets of UM and BL samples

were included. UM samples were located at least 1 cm away from the corresponding PT. All collected

samples were frozen at -80°C. Detailed tissue-collecting protocols were previously described by

Filipowicz et al[17]. All fragments prepared for molecular analysis were histologically evaluated by expert

pathologists to identify tumor fragments (PT) and confirm the normal histology of UM and SK samples.

DNA isolation from tissue lysates and whole blood was performed as previously described[17].

Whole exome sequencing, data analysis, variant detection, and validation with independent

methods

WES analyses were performed using the Agilent SureSelectXT Human All Exon V7 capture kit for

sequencing library construction followed by 150 bp paired-end sequencing on the HiSeq Illumina

platform (Illumina, San Diego, CA), outsourced to Macrogen Europe (Amsterdam, The Netherlands).

Sequencing coverage was 200x on average, with at least 100x on target. The sequencing coverage and

quality statistics for each sample are summarized in Supplementary Table 2A.

FASTQ files were inspected and processed with *Trim Galore!* (v0.6.7)

(https://www.bioinformatics.babraham.ac.uk/projects/trim_galore/) to remove Illumina-specific adapter

sequences and poor-quality reads when necessary. After converting FASTQ files to BAM format and

extracting read groups from the raw data, reads were processed using GATK4 Best Practices (v4.0)

(https://github.com/gatk-workflows/seq-format-conversion,https://github.com/gatk-workflows/gatk4-

data-processing). The reads were mapped to the human genome (hg38) using the BWA-MEM tool

(http://bio-bwa.sourceforge.net). Octopus (v0.7.4), in cancer mode, was used for variant calling. The

cancer calling model can jointly genotype multiple samples from the same individual, using a reference

sample (whole peripheral blood or skin when blood was not available) to distinguish between post-

zygotic and germline variants (https://luntergroup.github.io/octopus/). Framesift insertions/deletions,

nonsense, and missense variants located in exons were included in the analysis. A random forest filtering

approach was implemented to minimize false calls. Variants in reads with poor mapping quality (<30),

and variants supported by high-quality bases ($\geq$30) in fewer than five reads were excluded from the

analysis. Variants were annotated using ANNOVAR (https://annovar.openbioinformatics.org/) (last

updated on 07.07.2020 and accessed between 06.2022-08.2022) and wANNOVAR

(https://wannovar.wglab.org/) (last accessed on 21.05.2024), using the MANE SELECT transcript for

investigated genes (https://www.ensembl.org/info/genome/genebuild/mane.html). A brief overview of

filtering strategies implemented for identifying post-zygotic and germline variants related to breast

cancer, within BCAP, BCUS, and RM cohorts and validation experiments for selected post-zygotic

variants of BCAP patients is available in Supplementary Figure 2.

Post-zygotic variants

Only variants with sequencing depth $\geq$ 50 and tissue allele frequency $\geq$ 0.03 were included in the analysis.

Variants were filtered based on their annotation in ClinVar and InterVar databases; variants reported as

"pathogenic", "likely pathogenic", "uncertain significance", or "conflicting interpretations of

pathogenicity" were included. In parallel, variants described in the COSMIC database

(Cosmic_95coding) were incorporated. Missense variants documented in the COSMIC database, but

annotated as "benign" or "likely benign" in ClinVar or InterVar databases, were excluded. Variants in

genomic regions with known read-through transcription between adjacent genes, or those spanning

multiple genes (e.g., *P2RY11; PPAN-P2RY11* and *KIR2DL1; KIR2DS5; LOC112267881*), were also

excluded.

The remaining variants were filtered by their frequency in the general population, retaining only those

with a minor allele frequency (MAF) ≤ 0.001 across all gnomAD populations ("popmax") or not listed in

gnomAD (v2.1.1) (Supplementary Table 3). Variants were further categorized as truncating

(Supplementary Table 4) or missense (Supplementary Table 5). Truncating variants were considered

pathogenic, regardless of their annotation in ClinVar or InterVar. For missense variants, *in-silico* analyses

using the REVEL tool[18] (with a threshold score of 0.75) were performed. In summary, truncating variants,

variants classified as "pathogenic" or "likely pathogenic" in ClinVar, and missense variants with

conflicting interpretations or annotated as "pathogenic" in ClinVar without assertion criteria, but having a

REVEL score ≥ 0.75 were deemed pathogenic.

Variants affecting breast cancer-related genes were prioritized for further investigation based primarily on

their ClinVar annotation. Variants annotated as "pathogenic" or "likely pathogenic" in ClinVar and

missense variants reported in ClinVar as variants of uncertain significance or with conflicting

interpretations (or without assertion criteria provided), with a REVEL score of ≥ 0.75 were included.

Truncating variants affecting known tumor suppressor genes (i.e. *KMT2C*[19], *TBX3*[20], and *TP53*[21]) were

also incorporated even if absent in ClinVar (Supplementary Table 6). UM, UMD, PT, and SK samples of

16 BCAP patients were subjected to further investigation (Sanger sequencing or High-Resolution

Melting, Duplex sequencing) to verify or explore the presence of selected variants (Table 2,

Supplementary Tables 7 and 8, Supplementary Figure 3).

Germline variants

For germline variant detection, only variants in high- and moderate-penetrance breast cancer

susceptibility genes were included in the study, as defined by the NCCN Clinical Practice Guidelines in

Oncology[22] (Version 1.2023, September 7, 2022). The list of genes of interest includes: *ATM* (MIM

*607585), *BRCA1* (MIM *113705), *BRCA2* (MIM *600185), *BARD1* (MIM *601593), *BRIP1* (MIM

*605882), *CHEK2* (MIM *604373), *CDH1* (MIM *192090), *PALB2* (MIM *610355), *PTEN* (MIM

*601728), *TP53* (MIM *191170), *NF1* (MIM *613113), *STK11* (MIM *602216), *RAD50* (MIM

*604040), *RAD51C* (MIM *602774), *RAD51D* (MIM *602954) and additionally *PIK3CA* (MIM

*171834). Variants were filtered based on their frequency in the general population: variants with minor

allele frequency (MAF) ≤ 0.01 across all gnomAD populations ("popmax") or not noted in the database

(gnomAD v2.1.1) were included. Evidence according to the American College of Medical Genetics and

Genomics and the Association for Molecular Pathology recommendations[23] was included to describe all

germline pathogenic variants. Specifically, the evaluation of identified *BRCA1* and *BRCA2* variants was

performed according to the Evidence-based Network for the Interpretation of Germline Mutation Alleles

(ENIGMA) *BRCA1* and *BRCA2* Variant Curation Expert Panel[24] (Version 1.1.0) (Clinical Genome

Resource, https://www.clinicalgenome.org/affiliation/50087/,

https://cspec.genome.network/cspec/ui/svi/doc/GN092,

https://cspec.genome.network/cspec/ui/svi/doc/GN097). Pathogenic germline variants meeting the study's

criteria, identified within breast cancer patients of the BCAP and BCUS cohorts are described in

Supplementary Table 9.

Duplex sequencing

UM samples from 11 BCAP patients were selected to investigate the presence of low-frequency *PIK3CA*

and *TP53* variants beyond the detection limits of Sanger sequencing and High-Resolution Melting and a

single, higher-frequency *TP53* variant, i.e. c.151G>T (p.Glu51*), located in a difficult GC-rich region.

(Table 2, Supplementary Table 8). Additionally, UMD samples, available for 6 of those patients, were

included to explore further the distribution of selected variants in a more distant seemingly normal

mammary tissue. Duplex sequencing was performed as previously described[15,25].

Duplex sequencing data analysis

217   Raw duplex sequencing data were analyzed using the Snakemake-based Duplex-seq-Pipeline (v1.1.4)

218   (https://github.com/Kennedy-Lab-UW/Duplex-Seq-Pipeline) as previously described[26]. The sequencing

219   coverage and quality statistics for each sample are summarized in Supplementary Table 2B.

Statistical analysis

All statistical analyses were carried out with in-house developed scripts using R studio version 4.1.2

(2021-11-01). Packages *pheatmap* (version 1.0.12) and *ggplot2* (version 3.4.1) were used for plotting.

Statistical significance of differences between two or multiple groups was tested using the Mann–

Whitney U test or the Kruskal–Wallis H test respectively. Statistical significance of features between

multiple groups was tested with the Hypergeometric test or Fisher's exact test. Hazard Ratios were

calculated using the *coxph* function from package *survival* (version 3.5-5). Kaplan-Meier analysis was

performed using the *survfit* and *ggsurvplot* functions from *survminer* package (version 0.4.9) and groups

were tested with log-rank test. Differences were considered significant at $p < 0.05$.

RESULTS

**Truncating post-zygotic variants in dosage-sensitive genes predominate in BCAP compared to
BCUS and RM cohorts**

We examined sets of UM and PT samples from all breast cancer patients to identify variants related to

breast cancer (Figure 1a). Whole peripheral blood (BL) or skin (SK), if blood was not available, served as

a reference sample, and was used to distinguish between post-zygotic and germline variants (Figure 1b).

The details of post-zygotic variants in BCAP, BCUS, and RM cohorts meeting the study's cut-off criteria

(Materials & Methods and Supplementary Figure 2) are summarized in Supplementary Table 3. Briefly,

all truncating variants were included for manual curation, while non-truncating variants were filtered

based on their annotation within ClinVar or InterVar databases as "pathogenic", "likely pathogenic",

"uncertain significance", or "conflicting interpretations of pathogenicity", or their presence in the

COSMIC database (cosmic_95coding). Additionally, variants were filtered by their frequency in the

general population including only those with a minor allele frequency (MAF) ≤ 0.001 across all gnomAD

populations ("popmax") or not noted in the database (gnomAD v2.1.1).

Within the BCAP cohort, 167 distinct variants were identified in UM samples of 41 patients. The

corresponding numbers for the BCUS were strikingly lower, i.e. 56 variants were identified in the UM

samples of 24 patients. The RM cohort presented 10 distinct variants in seven unrelated individuals, but

all were either variants of uncertain significance or not reported in the ClinVar/InterVar databases.

Variants were further categorized based on their impact on the encoded protein i.e. truncating and non-

truncating. Notably, truncating nonsense (n=25) and frameshift (n=12) variants, were exclusive to the

BCAP cohort (Supplementary Table 4). These alterations were responsible for transcript elimination via

nonsense-mediated mRNA decay[27] and were deemed pathogenic, irrespective of their annotation in the

ClinVar and InterVar databases. For missense variants in BCAP, BCUS, and RM UM samples, further

analyses were performed using multiple *in-silico* predictors (Supplementary Table 5).

In the BCAP cohort, 29% (49/167) of the identified variants were deemed pathogenic. This list included

truncating variants (n=37), missense variants annotated as pathogenic/likely pathogenic (n=8), and

missense variants reported as uncertain significance or pathogenic/likely pathogenic in ClinVar, lacking

assertion criteria but showing evidence of pathogenicity according to *in-silico* analyses (n=4)

(Supplementary Tables 4 and 5). *In-silico* analyses for missense variants were conducted using the

REVEL tool[18], with a threshold set to 0.75 (accessed between June 2022 and August 2022 via

ANNOVAR software). Notably, nearly one-quarter (24%, 12/49) of the pathogenic BCAP variants were

detected only in the UM samples and were absent in the corresponding PTs. In comparison, the BCUS

cohort had seven pathogenic variants, representing 13% of the total identified variants (n=56). These

included variants reported as pathogenic (n=4), and missense variants classified as uncertain significance

or pathogenic/likely pathogenic in ClinVar without assertion criteria, showing evidence of pathogenicity

according to REVEL (n=3) (Supplementary Table 5). Nearly half (43%, 3/7) of the pathogenic BCUS

variants were identified only in UM samples. The UM samples from the BCAP cohort were significantly

enriched for pathogenic post-zygotic variants (a combination of truncating variants and those non-

truncating variants, classified as pathogenic by ClinVar/REVEL) compared to those from the BCUS

cohort (Hypergeometric test, p=0.0008578).

To select post-zygotic variants potentially linked to breast cancer, we focused on pathogenic variants as

defined by the ClinVar database. Variants annotated as pathogenic or likely pathogenic in ClinVar,

reviewed by expert panels, or submitted by multiple parties without discordance are listed in

Supplementary Table 6 for both BCAP and BCUS cohorts. Missense variants classified as uncertain

significance, pathogenic/likely pathogenic without assertion criteria, or with conflicting interpretations in

ClinVar with a REVEL score ≥ 0.75 were also included. Truncating variants in known tumor-suppressor

genes implicated in breast cancer, such as *KMT2C*[28], *TBX3*[20], and *TP53*[21] were included in the same table

even if they were absent from the ClinVar database (Supplementary Table 6).

We identified several variants affecting dosage-sensitive genes. These included deleterious variants in the

tumor suppressors *KMT2C*[28], *PTEN*[29], *PTCH1*[30], *TBX3*[20], and *TP53*[21] as well as activating variants in

oncogenes *AKT1*[31] and *PIK3CA*[32] identified in BCAP UM samples (Supplementary Table 6). Oncogenes

such as *SF3B1*[33], *HRAS*[34], and *GNAS*[35], and genes with a dual role in cancer (*RUNX1*[36]) were affected

solely in the BCUS cohort, with only the latter two being dosage-sensitive (Supplementary Table 6).

PIK3CA was the only gene recurrently affected in both BCAP and BCUS cohorts.

Recurrence, coexistence, spatial distribution in the breast, and validation of variants

In the BCAP cohort, pathogenic variants in two driver genes, *PIK3CA* and *TP53*, were predominant

across all subtypes of invasive cancer. *PIK3CA,* which encodes the catalytically active p100alpha

isoform, is a key regulator of cell proliferation and growth receptor signaling cascades[37]. We detected

three distinct pathogenic post-zygotic *PIK3CA* variants in UM samples: c.1624G>A (p.Glu542Lys),

c.3140A>G (p.His1047Arg), and c.3140A>T (p.His1047Leu), found in two, four and one unrelated

individuals, respectively (Supplementary Table 5). Another *PIK3CA* variant, c.3012G>T (p.Met1004Ile),

was found in the UM of a single BCAP cohort individual. However, this variant reported only once as of

uncertain significance in the ClinVar database and with a REVEL score of 0.437 suggesting it might be

benign, was classified as a variant of uncertain significance due to limited evidence (Supplementary Table

5). *PIK3CA* c.3140A>G (p.His1047Arg) and c.3140A>T (p.His1047Leu) variants, located at the

commonly mutated *PIK3CA* site in breast cancer, were also observed in UMs of the BCUS cohort (Figure

2, Supplementary Table 5).

TP53 is the most commonly mutated gene in various human cancers and its normal protein function is

frequently compromised in many types of malignancies[38]. We detected seven *TP53* variants in the normal

mammary gland samples of nine BCAP patients, including two recurrent variants: six pathogenic or likely

pathogenic (c.151G>T [p.Glu51*], c.329G>C [p.Arg110His], c.637C>T [p.Arg213*], c.711G>A

[p.Met237Ile], c.1024C>T [p.Arg342*], and c.1025G>C [p.Arg342Pro]), and one frameshift variant

c.227del (p.Ala76Aspfs*47), not previously reported in the ClinVar database (Supplementary Table 6).

Importantly, no *TP53* variants were observed in the normal mammary gland samples of the BCUS cohort

(Figure 2). Pathogenic variants in *AKT1*, *PIK3CA*, *PTEN*, *TBX3*, and *TP53* in 16 BCAP patients were

selected for validation with independent methods (Table 2, Supplementary Figure 2). The presence of

PIK3CA variants c.3140A>G (p.His1047Arg) (in three patients), c.1624G>A (p.Glu542Lys) (in two

patients) and *TP53* c.711G>A (p.Met237Ile) (in a single patient) was confirmed by Sanger sequencing or

High-Resolution Melting in the UM samples of six BCAP patients (Table 2, Supplementary Table 7,

Supplementary Figure 3). Notably, the *PIK3CA* c.3140A>G (p.His1047Arg) and *TP53* c.711G>A

(p.Met237Ile) variants co-existed in the UM sample of patient BCAP15. Additionally, two pathogenic

variants, c.49G>A (p.Glu17Lys) in *AKT1* and c.388C>T (p.Arg130*) in *PTEN*, were identified in the UM

samples of three BCAP patients and subsequently confirmed via Sanger sequencing or High-Resolution

Melting (Table 2, Supplementary Table 7, Supplementary Figure 3). The presence of the *AKT1* variant

was confirmed in the UM samples of two patients and the UMD of one of them, indicating a broad spatial

distribution of this variant in that particular patient. Finally, the *TBX3* c.371_372insTGGT

(p.Ile125Profs*14) variant was confirmed in the UM sample of a single patient (Table 2, Supplementary

Table 7, Supplementary Figure 3). We further employed duplex sequencing to validate the presence of

extremely low-frequency *PIK3CA* and *TP53* variants and examine their tissue/spatial distribution. We

selected 11 individuals: BCAP01, BCAP31, BCAP36, BCAP38, BCAP45, BCAP47, BCAP48, BCAP53,

BCAP54, BCAP57, and BCAP58 based on the presence of *PIK3CA* and *TP53* variants in proximal UM

samples from WES data. UMD samples available for 6 BCAP patients, located at a greater distance from

the PTs and not included in the original WES run, were also analyzed (Table 2). Ultra-deep targeted

duplex sequencing with a mean coverage of 4789x confirmed the following low-frequency (low as

1.34%) pathogenic variants in *PIK3CA*: c.3140A>G (p.His1047Arg) and c.3140A>T (p.His1047Leu) in

the UM samples from two patients, and revealed the presence of c.3140A>G (p. His1047Arg) variant in

the UMD samples of another two patients (Table 2, Supplementary Table 8). Furthermore, duplex

sequencing confirmed the presence of six *TP53* variants (c.151G>T [p.Glu51*], c.227del

[p.Ala76Aspfs*47], c.329G>C [p.Arg110His], c.637C>T [p.Arg213*], c.1024C>T [p.Arg342*], and

c.1025G>C [p.Arg342Pro]) in UM tissues of eight patients, and revealed the presence of c.151G>T

(p.Glu51*), c.227del (p.Ala76Aspfs*47), and c.637C>T (p.Arg213*) variants in the paired UMD tissues

of four patients (Table 2, Supplementary Table 8). An overview of validated variants in *AKT1*, *PIK3CA*,

PTEN, *TBX3,* and *TP53* genes along with follow-up information for the corresponding patients is

provided in Figure 3.

Spectrum of germline pathogenic variants in the two breast cancer cohorts

All breast cancer cases included in our study were reported as sporadic based on the family history of the

patient, however, genetic testing results were not available prior to recruitment. We analyzed BL or SK

samples from each participant to screen for pathogenic or likely pathogenic germline variants across all

cohorts (Materials & Methods, Supplementary Figure 2). In the BCAP cohort, 14 of 77 individuals (18%)

carried germline pathogenic variants in known breast cancer-associated genes[22]. These included

c.4186C>T (p.Gln1396*), c.4689C>G (p.Tyr1563*), c.5179A>T (p.Lys1727*), and c.5266dup

(p.Gln1756Profs*74) in *BRCA1*, c.5645C>A (p.Ser1882*), c.6591_6592del (p.Glu2198Asnfs*4), and

c.9382C>T (p.Arg3128*) in *BRCA2*, c.172_175del (p.Gln60Argfs*7) and c.1671_1674del

(p.Ile558Lysfs*2) in *PALB2*, and c.3233_3236del (p.Lys1079Valfs*28) in *RAD50*. Only *BRCA1*

c.5266dup (p.Gln1756Profs*74) and *PALB2* c.172_175del (p.Gln60Argfs*7) were recurrent, observed in

four and two unrelated individuals, respectively. This incidence surpasses the rates from other studies

where up to approximately 10% of reportedly sporadic cases turn out hereditary after molecular testing

and likely reflects the aggressive outcomes in these cases[15,21,39]. Four individuals (4/14, 29%) with

germline pathogenic variants also carried pathogenic post-zygotic variants in known, curated cancer-

related genes[19] in their UM samples. The germline variants in these cases were found in *BRCA1* (four

cases) and *RAD50* (one case). The corresponding post-zygotic variants were identified in *PIK3CA* or

TP53. *BRCA1* c.5266dup (p.Gln1756Profs*74) was the only variant observed in a single patient from the

BCUS cohort. In the control group, no individuals were found to carry germline pathogenic or likely

pathogenic variants in genes associated with breast cancer. Pathogenic germline variants in high- and

moderate-penetrance breast cancer susceptibility genes identified in the BCAP and BCUS cohorts are

detailed in Supplementary Table 9.

Pathogenic post-zygotic variants in patients with recurrent disease affect the survival rate

We used Kaplan-Meier plots to evaluate survival probabilities and compare patients with recurrence

(n=53) to those without recurrence (n=72) in both the BCAP and BCUS cohorts. Overall, patients with

recurrence had significantly lower survival probabilities (log-rank test, p=0.00017) (Supplementary

Figure 4A), with a hazard ratio of 2.44 (95% CI: 1.07-5.54, p=0.0337), indicating more than twice the

risk of death compared to the non-recurrence group.

Due to the shorter follow-up period for the BCUS cohort (2 years) compared to the BCAP cohort (10

years), we focused on the first 24 months post-diagnosis. During this period, recurrence patients in both

cohorts (n=53) had significantly lower survival probabilities compared to non-recurrence patients (n=71)

(log-rank test, p<0.0001) (Supplementary Figure 4B), with a hazard ratio of 4.85 (95% CI: 1.4-16.25,

p=0.0105), indicating more than four times the risk of death for the recurrence group. Within the first 24

months, BCAP patients experienced significantly more recurrence events than BCUS patients (Fisher's

exact test, p=0.005488). In the BCAP cohort, patients with recurrence (n=48) had lower survival

probabilities throughout the follow-up period compared to those without recurrence (n=28) (log-rank test,

p=0.015). This pattern was also observed in the first 24 months (n=48 vs. n=27) (log-rank test, p=0.0088)

(Supplementary Figures 4C and 4D).

Additionally, we assessed the impact of pathogenic post-zygotic and germline variants on the survival of

patients with recurrent disease. Figure 4 displays Kaplan-Meier curves based on the presence and type of

pathogenic variants. Survival probabilities differ significantly across studied groups (log-rank test,

p=0.024), indicating that the combination of pathogenic post-zygotic variants, pathogenic germline

variants, and recurrence status significantly impact patient survival. Patients with pathogenic germline

variants (green) had the shortest recurrence-free survival, with most recurrences occurring within the first

60 months. Notably, patients with pathogenic post-zygotic variants in breast cancer-specific genes (blue)

also experienced recurrences, though less frequently and over a longer follow-up period, highlighting the

impact of these variants on recurrence risk, albeit to a lesser extent than germline variants. Patients

without pathogenic germline or postyzygotic variants (yellow) showed intermediate outcomes. In

conclusion, while pathogenic germline variants have the most pronounced effect on survival rates,

pathogenic post-zygotic variants seem to also play a notable role in influencing recurrence risk and

patient outcomes, highlighting their importance in understanding disease progression.

DISCUSSION

The early detection and treatment of breast cancer, including its precursors, have shifted research focus from tumors to the normal mammary gland to deepen our understanding of the disease's origins[40]. Genetic and transcriptomic studies have revealed a wide spectrum of alterations in critical breast cancer driver genes within the normal mammary gland of patients who have undergone BCS or mastectomy, compared to control tissues[11–15,41].

Recognizing that histologically normal tissue may harbor early genetic changes, we screened for post-zygotic alterations in two similarly aged breast cancer cohorts (BCAP and BCUS, Kruskal–Wallis H test, p=0.082) with differing survival outcomes. We also included a significantly younger control group of individuals treated surgically for non-cancerous reasons (RM cohort, Kruskal–Wallis H test, p=0.000034 and p=0.00036 for BCAP and BCUS cohorts respectively).

Truncating variants (nonsense and frameshift) were found exclusively in the UM samples of patients with adverse prognoses (BCAP cohort) (Supplementary Table 4). In contrast, BCUS patients and RM controls showed only missense variants (Supplementary Table 5). UM samples from BCAP patients were significantly enriched for pathogenic post-zygotic variants (Hypergeometric test, p=0.0008578), affecting several known cancer-related genes genes[19] i.e. *AKT1*, *KMT2C*, *PIK3CA*, *PTCH1*, *PTEN*, *TBX3*, and *TP53,* and almost a quarter were found exclusively in UM samples, absent from the corresponding PTs, suggesting early tumorigenic processes. These findings indicate that the presence of pathogenic post-zygotic alterations in UM samples could signal a higher risk for aggressive cancer, preceding clinical symptoms.

The high prevalence of *PIK3CA* and *TP53* variants in BCAP patients' UM samples highlights their pivotal role in oncogenesis. The *PIK3CA* gene encodes the p110α catalytic subunit of phosphoinositide 3-kinase (PI3K), a key regulator of the PI3K/AKT signaling pathway, essential for cellular growth, proliferation, and survival and has a known oncogene function[37,39]. Missense variants in *PIK3CA*, especially in the

accessory (p.Glu542Lys) and catalytic domains ([p.His1047Arg], [p.His1047Leu]), enhance kinase

activity and promote oncogenic signaling. The recurrence of the p.His1047Arg variant, critical for

PIK3CA function, underscores its significance in tumorigenesis. *PIK3CA* variants were common in both

BCAP and BCUS cohorts, suggesting a fundamental role in breast cancer development. In contrast, *TP53*

variants were found exclusively in the BCAP cohort, underscoring the gene's crucial role in maintaining

genomic integrity. *TP53* encodes the tumor protein p53, a key tumor suppressor involved in DNA repair,

apoptosis, and cell cycle regulation[21]. Loss-of-function variants in *TP53* can inactivate its tumor-

suppressing activity during oncogenesis[42–44]. The observed *TP53* variants included truncating variants

(nonsense and frameshift) as well as missense alterations affecting the DNA-binding domain

([p.Arg110His], [p.Met237Ile]) and the tetramerization motif (p.Arg342Pro), highlighting the various

ways p53 function can be disrupted, potentially leading to malignancy. The exclusive presence of *TP53*

variants in the BCAP cohort may suggest a more aggressive disease phenotype and a worse prognosis

associated with these variants.

PIK3CA and *TP53* variants co-occurred in three BCAP patients (BCAP15, BCAP45, BCAP54),

suggesting a synergistic role in cancer progression. In particular, patient BCAP15 exhibited concurrent

pathogenic variants in *PIK3CA* (p.His1047Arg), *TP53* (p.Met237Ile), and *PTEN* (p.Arg130*). *PTEN*,

another critical tumor suppressor, negatively regulates the PI3K/AKT pathway[29]. The presence of

alterations in all three genes within a single patient emphasizes the complex interplay of multiple

oncogenic and tumor-suppressive pathways in breast cancer pathogenesis. This confluence of variants

likely contributes to a more aggressive clinical course, underscoring the importance of comprehensive

genetic profiling in understanding individual tumor biology.

While post-zygotic variants in *TP53* or *PIK3CA* have been observed in breast tumors, their consequences

in normal mammary tissue are less clear. Some studies suggest a benign effect[45,46]. Healthy breast tissue

accumulates alterations with age at an accelerating rate[47] and is influenced by hormonal stimuli,

undergoing cycles of expansion during puberty, pregnancy, and lactation[48]. Estrogen and its metabolites

can cause DNA damage, increasing cellular stress and the risk of genetic alterations and cancer[48].

However, the accumulation of alterations alone does not cause cancer; observed tissue-specific patterns

and the "ground state" theory suggest that quiescent stem cells with oncogenic variants rarely transform

unless activated by developmental, aging, or injury factors, that normally resemble physiological

mammary gland conditions[49]. The presence of such variants in ostensibly normal mammary gland tissue

suggests they may represent early, pre-cancerous changes. Despite the mammary gland's inherent multi-

layer protection system against clonal expansions, surviving mutant clones can lead to large fields of

mutated cells, i.e. field cancerization, thereby increasing cancer risk[50].

A subset of the BCAP cohort (18%, n=14/77) and one patient from the BCUS cohort (2%, n=1/49)

carried germline pathogenic variants in breast cancer genes (Supplementary Table 9). Among the BCAP

patients with these variants, four had concurrent pathogenic post-zygotic variants in curated cancer genes,

whereas 14 had only post-zygotic alterations in curated cancer genes and genes implicated in breast

cancer (Supplementary Tables 6 and 9). Despite their differing genetic profiles, all BCAP patients

experienced adverse outcomes within ten years of surgery highlighting the impact of these genetic

variations on prognosis. The interaction between germline and post-zygotic variants remains unclear, as

recent research indicates that the influence of germline variants on tumor behavior can vary significantly

based on factors such as penetrance and lineage, with some variants exhibiting minimal or transient

effects on tumor development[51].

In the first 24 months post-diagnosis, the BCAP cohort had significantly more recurrence events than the

BCUS cohort (Fisher's exact test, $p = 0.005488$). Recurrence was associated with much lower survival

probabilities across both cohorts. While pathogenic post-zygotic variants alone did not drastically alter

survival rates, their impact became more severe when combined with disease recurrence, suggesting a

link to disease aggressiveness. This highlights the importance of comprehensive genetic screening and

vigilant monitoring of patients with pathogenic post-zygotic variants in disease-related genes to improve

outcomes.

Current diagnostics primarily focus on identifying germline pathogenic variants in known breast cancer-

associated genes to assess breast cancer risk and guide personalized therapy[22]. However, over 80% of

breast tumors are not caused by inherited alterations[9]. Our study reveals that pathogenic post-zygotic

variants, such as *PIK3CA* and *TP53* alterations, are often found in seemingly normal mammary gland

tissue left behind after BCS, with allele frequencies ranging from 0.03 to 0.28. In some instances, these

variants represent distinct clonal populations not present in the corresponding primary tumors

(Supplementary Table 6), indicating the independent evolution of cell lineages within the mammary tissue

and suggesting they are unlikely to be micrometastases. The timing of their emergence relative to the

primary tumor - whether during early tumor progression or later—remains uncertain, but in both

scenarios, they may contribute to recurrence in the breast or metastasis to other organs[52]. Understanding

these dynamics is essential for improving diagnostic approaches and tailoring effective treatments.

Our study comes with certain limitations, particularly regarding the notable age differences between the

breast cancer cohorts and individuals subjected to reduction mammoplasty surgeries. Recruiting age-

matched control individuals poses a challenge, as those opting for cosmetic surgical treatments are

typically younger. The incidence of breast cancer diagnosis among younger women is relatively

infrequent with only about one out of eight invasive breast cancers being diagnosed in women under the

age of 45[1]. Another limitation arises in recruiting healthy control individuals, given that approximately

13% of women are expected to develop invasive breast cancer during their lifetime and the precise onset

of carcinogenesis remains unclear[1,2]. Here, control normal mammary glands were sampled from

individuals without a personal or familial history of cancer undergoing plastic surgery. Hence, these

samples represent the most appropriate, available control samples from a biological standpoint.

Our findings demonstrate that pathogenic post-zygotic variants in breast cancer-associated genes are

significantly more prevalent in normal mammary tissues of patients with adverse outcomes, such as

recurrence or metastasis, compared to those without specific prognosis criteria or control individuals[15].

Monitoring these patients for nearly a decade after diagnosis allowed for a direct association between

these variants and clinical outcomes. These alterations were strongly linked to disease progression,

particularly recurrence, suggesting an increased risk of aggressive cancer before clinical symptoms

appear. This highlights the need for expanded genetic screening and enhanced surveillance to improve

personalized breast cancer management, particularly for patients with poor prognoses.

Competing interests

J.P.D. is a cofounder and shareholder in Cray Innovation AB. J.M. is a co-founder and shareholder of

Genegoggle sp. z o.o. The remaining authors have declared that no competing interests exist.

References

1. American Cancer Society. Breast Cancer Facts & Figures 2022-2024.Atlanta: American
Cancer Society, Inc. 2022. https://www.cancer.org/content/dam/cancer-org/research/cancer-
facts-and-statistics/breast-cancer-facts-and-figures/2022-2024-breast-cancer-fact-figures-
acs.pdf

2. American Cancer Society. Cancer Facts & Figures 2023.Atlanta: American Cancer Society;
2023. https://www.cancer.org/content/dam/cancer-org/research/cancer-facts-and-
statistics/annual-cancer-facts-and-figures/2023/2023-cancer-facts-and-figures.pdf

3. Howlader N, Noone AM, Krapcho M, Miller D, Brest A, Yu M, Ruhl J, Tatalovich Z,
Mariotto A, Lewis DR, Chen HS, Feuer EJ, Cronin KA (Eds). SEER Cancer Statistics
Review, 1975-2018, National Cancer Institute. Bethesda, MD,
Https://Seer.Cancer.Gov/Csr/1975_2018/, Based on November 2020 SEER Data
Submission, Posted to the SEER Web Site, April 2021.

4. Pan H, Gray R, Braybrooke J, Davies C, Taylor C, McGale P, Peto R, Pritchard KI, Bergh
J, Dowsett M, Hayes DF, EBCTCG. 20-Year Risks of Breast-Cancer Recurrence after
Stopping Endocrine Therapy at 5 Years. *N Engl J Med*. 2017;377(19):1836-1846.
doi:10.1056/NEJMoa1701830

5. Shiovitz S, Korde LA. Genetics of breast cancer: a topic in evolution. *Ann Oncol Off J Eur*
Soc Med Oncol. 2015;26(7):1291-1299. doi:10.1093/annonc/mdv022

6. Coughlin SS. Epidemiology of Breast Cancer in Women. *Adv Exp Med Biol*. 2019;1152:9-
29. doi:10.1007/978-3-030-20301-6_2

7. Kleibl Z, Kristensen VN. Women at high risk of breast cancer: Molecular characteristics,
clinical presentation and management. *Breast Edinb Scotl*. 2016;28:136-144.
doi:10.1016/j.breast.2016.05.006

8. Danforth DN. Genomic Changes in Normal Breast Tissue in Women at Normal Risk or at
High Risk for Breast Cancer. *Breast Cancer Basic Clin Res*. 2016;10:BCBCR.S39384.
doi:10.4137/BCBCR.S39384

9. Loibl S, Poortmans P, Morrow M, Denkert C, Curigliano G. Breast cancer. *Lancet Lond
Engl*. 2021;397(10286):1750-1769. doi:10.1016/S0140-6736(20)32381-3

10. Waks AG, Winer EP. Breast Cancer Treatment: A Review. *JAMA*. 2019;321(3):288-300.
doi:10.1001/jama.2018.19323

11. Gadaleta E, Fourgoux P, Pirró S, Thorn GJ, Nelan R, Ironside A, Rajeeve V, Cutillas PR,
Lobley AE, Wang J, Gea E, Ross-Adams H, Bessant C, Lemoine NR, Jones LJ, Chelala C.
Characterization of four subtypes in morphologically normal tissue excised proximal and
distal to breast cancer. *NPJ Breast Cancer*. 2020;6:38. doi:10.1038/s41523-020-00182-9

12. Aran D, Camarda R, Odegaard J, Paik H, Oskotsky B, Krings G, Goga A, Sirota M, Butte
AJ. Comprehensive analysis of normal adjacent to tumor transcriptomes. *Nat Commun*.
2017;8(1):1077. doi:10.1038/s41467-017-01027-z

13. Forsberg LA, Rasi C, Pekar G, Davies H, Piotrowski A, Absher D, Razzaghian HR,
Ambicka A, Halaszka K, Przewoźnik M, Kruczak A, Mandava G, Pasupulati S, Hacker J,
Prakash KR, Dasari RC, Lau J, Penagos-Tafurt N, Olofsson HM, Hallberg G, Skotnicki P,
Mituś J, Skokowski J, Jankowski M, Śrutek E, Zegarski W, Tiensuu Janson E, Ryś J, Tot T,
Dumanski JP. Signatures of post-zygotic structural genetic aberrations in the cells of
histologically normal breast tissue that can predispose to sporadic breast cancer. *Genome
Res*. 2015;25(10):1521-1535. doi:10.1101/gr.187823.114

14. Ronowicz A, Janaszak-Jasiecka A, Skokowski J, Madanecki P, Bartoszewski R, Bałut M,
Seroczyńska B, Kochan K, Bogdan A, Butkus M, Pęksa R, Ratajska M, Kuźniacka A,
Wasąg B, Gucwa M, Krzyżanowski M, Jaśkiewicz J, Jankowski Z, Forsberg L, Ochocka
JR, Limon J, Crowley MR, Buckley PG, Messiaen L, Dumanski JP, Piotrowski A.
Concurrent DNA Copy-Number Alterations and Mutations in Genes Related to
Maintenance of Genome Stability in Uninvolved Mammary Glandular Tissue from Breast
Cancer Patients. *Hum Mutat*. 2015;36(11):1088-1099. doi:10.1002/humu.22845

15. Kostecka A, Nowikiewicz T, Olszewski P, Koczkowska M, Horbacz M, Heinzl M,
Andreou M, Salazar R, Mair T, Madanecki P, Gucwa M, Davies H, Skokowski J, Buckley
PG, Pęksa R, Śrutek E, Szylberg Ł, Hartman J, Jankowski M, Zegarski W, Tiemann-Boege
I, Dumanski JP, Piotrowski A. High prevalence of somatic PIK3CA and TP53 pathogenic
variants in the normal mammary gland tissue of sporadic breast cancer patients revealed by
duplex sequencing. *Npj Breast Cancer*. 2022;8(1):1-10. doi:10.1038/s41523-022-00443-9

16. Pipek O, Alpár D, Rusz O, Bödör C, Udvarnoki Z, Medgyes-Horváth A, Csabai I, Szállási
Z, Madaras L, Kahán Z, Cserni G, Kővári B, Kulka J, Tőkés AM. Genomic Landscape of
Normal and Breast Cancer Tissues in a Hungarian Pilot Cohort. *Int J Mol Sci*.
2023;24(10):8553. doi:10.3390/ijms24108553

17. Filipowicz N, Drężek K, Horbacz M, Wojdak A, Szymanowski J, Rychlicka-Buniowska E, Juhas U, Duzowska K, Nowikiewicz T, Stańkowska W, Chojnowska K, Andreou M, Ławrynowicz U, Wójcik M, Davies H, Śrutek E, Bieńkowski M, Milian-Ciesielska K, Zdrenka M, Ambicka A, Przewoźnik M, Harazin-Lechowska A, Adamczyk A, Kowalski J, Bała D, Wiśniewski D, Tkaczyński K, Kamecki K, Drzewiecka M, Wroński P, Siekiera J, Ratnicka I, Jankau J, Wierzba K, Skokowski J, Połom K, Przydacz M, Bełch Ł, Chłosta P, Matuszewski M, Okoń K, Rostkowska O, Hellmann A, Sasim K, Remiszewski P, Sierżęga M, Hać S, Kobiela J, Kaska Ł, Jankowski M, Hodorowicz-Zaniewska D, Jaszczyński J, Zegarski W, Makarewicz W, Pęksa R, Szpor J, Ryś J, Szylberg Ł, Piotrowski A, Dumanski JP. Comprehensive cancer-oriented biobanking resource of human samples for studies of post-zygotic genetic variation involved in cancer predisposition. Kim IY, ed. *PLOS ONE*. 2022;17(4):e0266111. doi:10.1371/journal.pone.0266111

18. Ioannidis NM, Rothstein JH, Pejaver V, Middha S, McDonnell SK, Baheti S, Musolf A, Li Q, Holzinger E, Karyadi D, Cannon-Albright LA, Teerlink CC, Stanford JL, Isaacs WB, Xu J, Cooney KA, Lange EM, Schleutker J, Carpten JD, Powell IJ, Cussenot O, Cancel-Tassin G, Giles GG, MacInnis RJ, Maier C, Hsieh CL, Wiklund F, Catalona WJ, Foulkes WD, Mandal D, Eeles RA, Kote-Jarai Z, Bustamante CD, Schaid DJ, Hastie T, Ostrander EA, Bailey-Wilson JE, Radivojac P, Thibodeau SN, Whittemore AS, Sieh W. REVEL: An Ensemble Method for Predicting the Pathogenicity of Rare Missense Variants. *Am J Hum Genet*. 2016;99(4):877-885. doi:10.1016/j.ajhg.2016.08.016

19. Sondka Z, Bamford S, Cole CG, Ward SA, Dunham I, Forbes SA. The COSMIC Cancer Gene Census: describing genetic dysfunction across all human cancers. *Nat Rev Cancer*. 2018;18(11):696-705. doi:10.1038/s41568-018-0060-1

20. Liang B, Zhou Y, Qian M, Xu M, Wang J, Zhang Y, Song X, Wang H, Lin S, Ren C, Monga SP, Wang B, Evert M, Chen Y, Chen X, Huang Z, Calvisi DF, Chen X. TBX3 functions as a tumor suppressor downstream of activated CTNNB1 mutants during hepatocarcinogenesis. *J Hepatol*. 2021;75(1):120-131. doi:10.1016/j.jhep.2021.01.044

21. Cancer Genome Atlas Network. Comprehensive molecular portraits of human breast tumours. *Nature*. 2012;490(7418):61-70. doi:10.1038/nature11412

22. NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®) Genetic/Familial High-Risk Assessment: Breast, Ovarian, and Pancreatic Version 1.2023 — September 7, 2022 NCCN.org.

23. Richards S, Aziz N, Bale S, Bick D, Das S, Gastier-Foster J, Grody WW, Hegde M, Lyon E, Spector E, Voelkerding K, Rehm HL, ACMG Laboratory Quality Assurance Committee. Standards and guidelines for the interpretation of sequence variants: a joint consensus recommendation of the American College of Medical Genetics and Genomics and the Association for Molecular Pathology. *Genet Med Off J Am Coll Med Genet*. 2015;17(5):405-424. doi:10.1038/gim.2015.30

24. Walker LC, Hoya MDL, Wiggins GAR, Lindy A, Vincent LM, Parsons MT, Canson DM, Bis-Brewer D, Cass A, Tchourbanov A, Zimmermann H, Byrne AB, Pesaran T, Karam R,

Harrison SM, Spurdle AB, Biesecker LG, Harrison SM, Tayoun AA, Berg JS, Brenner SE, Cutting GR, Ellard S, Greenblatt MS, Kang P, Karbassi I, Karchin R, Mester J, O'Donnell-Luria A, Pesaran T, Plon SE, Rehm HL, Strande NT, Tavtigian SV, Topper S. Using the ACMG/AMP framework to capture evidence related to predicted and observed impact on splicing: Recommendations from the ClinGen SVI Splicing Subgroup. *Am J Hum Genet*. 2023;110(7):1046-1067. doi:10.1016/j.ajhg.2023.06.002

25. Salazar R, Arbeithuber B, Ivankovic M, Heinzl M, Moura S, Hartl I, Mair T, Lahnsteiner A, Ebner T, Shebl O, Pröll J, Tiemann-Boege I. Discovery of an unusually high number of de novo mutations in sperm of older men using duplex sequencing. *Genome Res*. 2022;32(3):499-511. doi:10.1101/gr.275695.121

26. Sanchez-Contreras M, Sweetwyne MT, Kohrn BF, Tsantilas KA, Hipp MJ, Schmidt EK, Fredrickson J, Whitson JA, Campbell MD, Rabinovitch PS, Marcinek DJ, Kennedy SR. A replication-linked mutational gradient drives somatic mutation accumulation and influences germline polymorphisms and genome composition in mitochondrial DNA. *Nucleic Acids Res*. 2021;49(19):11103-11118. doi:10.1093/nar/gkab901

27. Brogna S, Wen J. Nonsense-mediated mRNA decay (NMD) mechanisms. *Nat Struct Mol Biol*. 2009;16(2):107-113. doi:10.1038/nsmb.1550

28. Rampias T, Karagiannis D, Avgeris M, Polyzos A, Kokkalis A, Kanaki Z, Kousidou E, Tzetis M, Kanavakis E, Stravodimos K, Manola KN, Pantelias GE, Scorilas A, Klinakis A. The lysine□specific methyltransferase KMT 2C/ MLL 3 regulates DNA repair components in cancer. *EMBO Rep*. 2019;20(3):e46821. doi:10.15252/embr.201846821

29. Luongo F, Colonna F, Calapà F, Vitale S, Fiori ME, De Maria R. PTEN Tumor-Suppressor: The Dam of Stemness in Cancer. *Cancers*. 2019;11(8):1076. doi:10.3390/cancers11081076

30. Gailani MR, Bale SJ, Leffell DJ, DiGiovanna JJ, Peck GL, Poliak S, Drum MA, Pastakia B, McBride OW, Kase R. Developmental defects in Gorlin syndrome related to a putative tumor suppressor gene on chromosome 9. *Cell*. 1992;69(1):111-117. doi:10.1016/0092-8674(92)90122-s

31. Staal SP. Molecular cloning of the akt oncogene and its human homologues AKT1 and AKT2: amplification of AKT1 in a primary human gastric adenocarcinoma. *Proc Natl Acad Sci U S A*. 1987;84(14):5034-5037. doi:10.1073/pnas.84.14.5034

32. Bader AG, Kang S, Zhao L, Vogt PK. Oncogenic PI3K deregulates transcription and translation. *Nat Rev Cancer*. 2005;5(12):921-929. doi:10.1038/nrc1753

33. Kesarwani AK, Ramirez O, Gupta AK, Yang X, Murthy T, Minella AC, Pillai MM. Cancer-associated SF3B1 mutants recognize otherwise inaccessible cryptic 3' splice sites within RNA secondary structures. *Oncogene*. 2017;36(8):1123-1133. doi:10.1038/onc.2016.279

34. Hobbs GA, Der CJ, Rossman KL. RAS isoforms and mutations in cancer at a glance. *J Cell Sci*. 2016;129(7):1287-1292. doi:10.1242/jcs.182873

35. He X, Zhang L, Chen Y, Remke M, Shih D, Lu F, Wang H, Deng Y, Yu Y, Xia Y, Wu X,
Ramaswamy V, Hu T, Wang F, Zhou W, Burns DK, Kim SH, Kool M, Pfister SM,
Weinstein LS, Pomeroy SL, Gilbertson RJ, Rubin JB, Hou Y, Wechsler-Reya R, Taylor
MD, Lu QR. The G protein α subunit Gαs is a tumor suppressor in Sonic hedgehog-driven
medulloblastoma. *Nat Med*. 2014;20(9):1035-1042. doi:10.1038/nm.3666

36. Otalora□Otalora B, Henriquez B, Lopez□Kleine L, Rojas A. RUNX family: Oncogenes or
tumor suppressors (Review). *Oncol Rep*. Published online May 6, 2019.
doi:10.3892/or.2019.7149

37. Martínez-Sáez O, Chic N, Pascual T, Adamo B, Vidal M, González-Farré B, Sanfeliu E,
Schettini F, Conte B, Brasó-Maristany F, Rodríguez A, Martínez D, Galván P, Rodríguez
AB, Martinez A, Muñoz M, Prat A. Frequency and spectrum of PIK3CA somatic mutations
in breast cancer. *Breast Cancer Res*. 2020;22(1):45. doi:10.1186/s13058-020-01284-9

38. Hainaut P, Pfeifer GP. Somatic TP53 Mutations in the Era of Genome Sequencing. *Cold
Spring Harb Perspect Med*. 2016;6(11):a026179. doi:10.1101/cshperspect.a026179

39. Pereira B, Chin SF, Rueda OM, Vollan HKM, Provenzano E, Bardwell HA, Pugh M, Jones
L, Russell R, Sammut SJ, Tsui DWY, Liu B, Dawson SJ, Abraham J, Northen H, Peden JF,
Mukherjee A, Turashvili G, Green AR, McKinney S, Oloumi A, Shah S, Rosenfeld N,
Murphy L, Bentley DR, Ellis IO, Purushotham A, Pinder SE, Børresen-Dale AL, Earl HM,
Pharoah PD, Ross MT, Aparicio S, Caldas C. The somatic mutation profiles of 2,433 breast
cancers refine their genomic and transcriptomic landscapes. *Nat Commun*.
2016;7(1):11479. doi:10.1038/ncomms11479

40. Crosby D, Bhatia S, Brindle KM, Coussens LM, Dive C, Emberton M, Esener S, Fitzgerald
RC, Gambhir SS, Kuhn P, Rebbeck TR, Balasubramanian S. Early detection of cancer.
Science. 2022;375(6586):eaay9040. doi:10.1126/science.aay9040

41. Troester MA, Hoadley KA, D'Arcy M, Cherniack AD, Stewart C, Koboldt DC, Robertson
AG, Mahurkar S, Shen H, Wilkerson MD, Sandhu R, Johnson NB, Allison KH, Beck AH,
Yau C, Bowen J, Sheth M, Hwang ES, Perou CM, Laird PW, Ding L, Benz CC. DNA
defects, epigenetics, and gene expression in cancer-adjacent breast: a study from The
Cancer Genome Atlas. *NPJ Breast Cancer*. 2016;2:16007. doi:10.1038/npjbcancer.2016.7

42. Baugh EH, Ke H, Levine AJ, Bonneau RA, Chan CS. Why are there hotspot mutations in
the TP53 gene in human cancers? *Cell Death Differ*. 2018;25(1):154-160.
doi:10.1038/cdd.2017.180

43. Vogelstein B, Papadopoulos N, Velculescu VE, Zhou S, Diaz LA, Kinzler KW. Cancer
Genome Landscapes. *Science*. 2013;339(6127):1546-1558. doi:10.1126/science.1235122

44. The ICGC/TCGA Pan-Cancer Analysis of Whole Genomes Consortium, Aaltonen LA,
Abascal F, Abeshouse A, Aburatani H, Adams DJ, Agrawal N, Ahn KS, Ahn SM, Aikata
H, Akbani R, Akdemir KC, Al-Ahmadie H, Al-Sedairy ST, Al-Shahrour F, Alawi M,
Albert M, Aldape K, Alexandrov LB, Ally A, Alsop K, Alvarez EG, Amary F, Amin SB,
Aminou B, Ammerpohl O, Anderson MJ, Ang Y, Antonello D, Anur P, Aparicio S,

Appelbaum EL, Arai Y, Aretz A, Arihiro K, Ariizumi S ichi, Armenia J, Arnould L, Asa S,
Assenov Y, Atwal G, Aukema S, Auman JT, Aure MRR, Awadalla P, Aymerich M, Bader
GD, Baez-Ortega A, Bailey MH, Bailey PJ, Balasundaram M, Balu S, Bandopadhayay P,
Banks RE, Barbi S, Barbour AP, Barenboim J, Barnholtz-Sloan J, Barr H, Barrera E,
Bartlett J, Bartolome J, Bassi C, Bathe OF, Baumhoer D, Bavi P, Baylin SB, Bazant W,
Beardsmore D, Beck TA, Behjati S, Behren A, Niu B, Bell C, Beltran S, Benz C, Berchuck
A, Bergmann AK, Bergstrom EN, Berman BP, Berney DM, Bernhart SH, Beroukhim R,
Berrios M, Bersani S, Bertl J, Betancourt M, Bhandari V, Bhosle SG, Biankin AV, Bieg M,
Bigner D, Binder H, Birney E, Birrer M, Biswas NK, Bjerkehagen B, Bodenheimer T, et al.
Pan-cancer analysis of whole genomes. *Nature*. 2020;578(7793):82-93.
doi:10.1038/s41586-020-1969-6

45.   Kandel R, Li SQ, Ozcelik H, Rohan T. p53 protein accumulation and mutations in normal
and benign breast tissue. *Int J Cancer*. 2000;87(1):73-78. doi:10.1002/1097-
0215(20000701)87:1<73::AID-IJC11>3.0.CO;2-U

46.   Soysal SD, Ng CKY, Costa L, Weber WP, Paradiso V, Piscuoglio S, Muenst S. Genetic
Alterations in Benign Breast Biopsies of Subsequent Breast Cancer Patients. *Front Med*.
2019;6:166. doi:10.3389/fmed.2019.00166

47.   Cereser B, Yiu A, Tabassum N, Del Bel Belluz L, Zagorac S, Ancheta KRZ, Zhong R,
Miere C, Jeffries-Jones AR, Moderau N, Werner B, Stebbing J. The mutational landscape
of the adult healthy parous and nulliparous human breast. *Nat Commun*. 2023;14(1):5136.
doi:10.1038/s41467-023-40608-z

48.   Macias H, Hinck L. Mammary gland development. *Wiley Interdiscip Rev Dev Biol*.
2012;1(4):533-557. doi:10.1002/wdev.35

49.   Jassim A, Rahrmann EP, Simons BD, Gilbertson RJ. Cancers make their own luck: theories
of cancer origins. *Nat Rev Cancer*. 2023;23(10):710-724. doi:10.1038/s41568-023-00602-5

50.   Ciwinska M, Messal HA, Hristova HR, Lutz C, Bornes L, Chalkiadakis T, Harkes R,
Langedijk NSM, Hutten SJ, Menezes RX, Jonkers J, Prekovic S, Grand Challenge
PRECISION consortium, Wesseling J, Thompson AM, Nik-Zainal S, Sawyer EJ, Davies
HR, Futreal A, Navin NE, Hwang ES, Jonkers J, Van Rheenen J, Behbod F, Lips EH,
Schmidt M, Wessels LFA, Rea D, Bhattacharjee P, Stobart H, Collyar D, Pinto D,
Verschuur E, Van Oirsouw M, Simons BD, Scheele CLGJ, Van Rheenen J. Mechanisms
that clear mutations drive field cancerization in mammary tissue. *Nature*.
2024;633(8028):198-206. doi:10.1038/s41586-024-07882-3

51.   Srinivasan P, Bandlamudi C, Jonsson P, Kemel Y, Chavan SS, Richards AL, Penson AV,
Bielski CM, Fong C, Syed A, Jayakumaran G, Prasad M, Hwee J, Sumer SO, De Bruijn I,
Li X, Gao J, Schultz N, Cambria R, Galle J, Mukherjee S, Vijai J, Cadoo KA, Carlo MI,
Walsh MF, Mandelker D, Ceyhan-Birsoy O, Shia J, Zehir A, Ladanyi M, Hyman DM,
Zhang L, Offit K, Robson ME, Solit DB, Stadler ZK, Berger MF, Taylor BS. The context-
specific role of germline pathogenicity in tumorigenesis. *Nat Genet*. 2021;53(11):1577-
1585. doi:10.1038/s41588-021-00949-1

52.   Davis A, Gao R, Navin N. Tumor evolution: Linear, branching, neutral or punctuated?
Biochim Biophys Acta BBA - Rev Cancer. 2017;1867(2):151-161.
doi:10.1016/j.bbcan.2017.01.003

53.   Zhou X, Edmonson MN, Wilkinson MR, Patel A, Wu G, Liu Y, Li Y, Zhang Z, Rusch MC,
Parker M, Becksfort J, Downing JR, Zhang J. Exploring genomic alteration in pediatric
cancer using ProteinPaint. *Nat Genet*. 2016;48(1):4-6. doi:10.1038/ng.3466

Authors Contributions

Study design and conception: A.P., M.A.; Clinical data acquisition and interpretation: M.D., M.N., J.J.,

D.H.Z., J.S., E.S., M.L-J., J.H., D.B., M. Jankowski., W.Z., Ł.S., T.N., N.F., A.M., K.C., K.D-C., U. Ł.,

K.D., H.D., B. B-O.; Experiments: M.A., K.C., M. Koszyński, M. Jaśkiewicz., U. Ł., K.D., H.D., B. B-

O., A.K.; Methodology: I.T-B.; Data analysis and interpretation: A.P., M.A., M. Koczkowska, K.C.,

P.G.B., P.M., M.H., M. Jąkalski, J.M.; Funding acquisition: A.P., J.P.D., K.C., I.T-B.; Visualization:

M.A., M. Jąkalski; Manuscript writing – original: M.A., A.P.; Manuscript writing – review & editing:

M.A., A.P., J.P.D., N.F., M. Koczkowska, M.H, P.G.B., I.T-B., A.K; Supervision: A.P., J.P.D., M.

Koczkowska. All authors have read and agreed to the published version of the manuscript.

Ethics approval and consent to participate

733 **Consent for publication**

734 Not applicable.

Data availability

Raw duplex sequencing and WES data are available upon request in the EGA archive, under study IDs EGAS50000000538 and EGAS50000000539 respectively.

Funding

This research was supported by the Foundation for Polish Science under the International Research Agendas Program financed from the Smart Growth Operational Program 2014–2020 (Grant Agreement No. MAB/2018/6) to A.P. and J.P.D. Parts of the study were supported by The Swedish Cancer Society (No. 20 0889 PjF) and Swedish Medical Research Council (No. 2020-02010) to J.P.D., by The National Science Centre Poland Miniatura 4 (Project No. 2020/04/X/NZ2/02084) to K.C., and by the Austrian Science Fund FWF (P30867000) and the European Regional Development Fund (REGGEN ATCZ207) to I.T-B.

Corresponding author

Correspondence and requests for materials should be addressed to Arkadiusz Piotrowski.

Acknowledgments

The authors wish to thank Agata Wojdak for her help with administrative activities. We also would like to thank all the patients and volunteer control individuals for acceptance to participate in the study and sample contribution; hospital staff involved in the patient recruitment process in Oncology Center - Prof. Franciszek Łukaszczyk Memorial Hospital in Bydgoszcz, University Clinical Centre in Gdańsk and University Hospital in Cracow. We thank Dr. Leszek Kalinowski for the access to selected laboratory facilities. Figure 1 and Supplementary Figure 1 were partly generated using Servier Medical Art, licensed under a Creative Commons Attribution 3.0 Unported License (https://creativecommons.org/licenses/by/3.0/).

MAIN TABLES

758 **Table 1. Summarized clinicopathological features of breast cancer patients included in the Breast**

759 **Cancer Adverse Prognoses (BCAP) and the Breast Cancer Un-Selected (BCUS) cohorts.**

	BCAP cohort	BCUS cohort
Number of individuals	77	49
Age (median, range)	62, 23-85	65, 37-84
		p value = 0.082
Collected samples	238	147
Primary Tumor, PT	77	49
Uninvolved mammary gland, UM	77	49
Distal fragment of uninvolved mammary gland, UMD	7	-
Reference sample	77	49
(whole peripheral blood, BL or skin, SK)		
Histology		
Invasive ductal carcinoma, IDC	59	40
Invasive lobular carcinoma, ILC	3	4
IDC - ILC	6	1
other	9	4
Receptors		
Estrogen, ER (positive / negative / not available)	57 / 20	43 / 5 / 1
Progesterone, PR (positive / negative / not available)	43 / 34	44 / 4 / 1
HER2 (positive / negative / not available)	16 / 56 / 5	5 / 43 / 1
Subtype		

Luminal A	14	22
Luminal B	37	21
HER-2 enriched	9	2
Triple-negative breast cancer, TNBC	11	1
Not available	6	3
Follow-up information		
Recurrence (yes / no)	50 / 27	5 / 44
Second cancer (yes / no)	26 / 51	0 / 49
Death* (yes / no)	45 / 31	3 / 46

Table 1. Matched and primary tumor (PT) and uninvolved mammary gland (UM, ≥1 cm) samples were

collected from two breast cancer cohorts, i.e. 77 individuals characterized with adverse outcomes (BCAP

cohort) and 49 individuals recruited without any pre-selection criteria related to prognosis (BCUS cohort).

Whole peripheral blood (BL) or skin (SK) samples (if BL was not available) were collected as reference

samples to distinguish between post-zygotic and germline variants. Distal UM samples (UMD, 1.5-3 cm

from PT, median 2.35 cm), available for 7 BCAP patients, were included. The detailed sampling design is

described in Materials and Methods. An overview is also available in Figure 1. *Death status refers to

patients who succumbed to the disease (patient with ID BCAP61 died from non-oncological reasons).

Detailed clinicopathological information for BCAP and BCUS cohorts is provided in Supplementary

Table 1.

Table 2. Pathogenic post-zygotic variants within the uninvolved mammary gland (UM) samples of breast cancer patients with adverse
prognoses (BCAP cohort), selected for further investigation/validation.

Gene	Variant[a]	ClinVar[b]	COSMIC ID[c]	AVSNP150[d]	Individual ID and UM sample VAF	Confirmation
AKT1	c.49G>A (p.Glu17Lys)	Pathogenic	ID=COSV62571334	rs121434592	BCAP32 (0.6%), BCAP66* (0.7%)	SS / HRM
PIK3CA	c.1624G>A (p.Glu542Lys)	Pathogenic	ID=COSV55873227	rs121913273	BCAP56 (0.7%), BCAP45 (12%)	SS / HRM
PIK3CA	c.3140A>G (p.His1047Arg)	Pathogenic	ID=COSV55873195	rs121913279	BCAP15 (0.08%), BCAP31* (19%), BCAP36 (0.3%), BCAP53* (0.7%)	SS / HRM / DS
PIK3CA	c.3140A>T (p.His1047Leu)	Pathogenic	ID=COSV55873401	rs121913279	BCAP54* (0.5%)	DS
PTEN	c.388C>T (p.Arg130*)	Pathogenic	ID=COSV64288463	rs121909224	BCAP15 (0.7%)	SS / HRM
TBX3	c.371_372insTGGT (p.Ile125Profs*14)	n.a.	ID=COSV57471668	n.a.	BCAP44 (12%)	SS / HRM
TP53	c.151G>T (p.Glu51*)	Pathogenic	ID=COSV52694020	n.a.	BCAP58* (16%)	SS / HRM
TP53	c.227del (p.Ala76Aspfs*47)	n.a.	ID=COSV52728465	n.a.	BCAP54 (0.5%)	DS
TP53	c.329G>C (p.Arg110His)	Pathogenic	ID=COSV52668419	rs11540654	BCAP45 (0.8%)	DS
TP53	c.637C>T (p.Arg213*)	Pathogenic	ID=COSV52665560	rs397516436	BCAP01* (0.8%), BCAP48* (0.6%)	DS
TP53	c.711G>A (p.Met237Ile)	Pathogenic	ID=COSV52661887	rs587782664	BCAP15 (0.8%)	SS / HRM
TP53	c.1024C>T (p.Arg342*)	Pathogenic	ID=COSV52665487	rs730882029	BCAP38 (0.5%), BCAP47 (0.7%)	DS
TP53	c.1025G>C (p.Arg342Pro)	Pathogenic/ Likely pathogenic	ID=COSV52690857	rs375338359	BCAP57 (0.6%)	DS

Table 2. Presented variants, identified via Whole Exome sequencing, were corroborated with either Sanger sequencing/High-Resolution Melting,

or Duplex sequencing. [a]Variant annotation provided for the basic isoform of the transcript. [b]Pathogenicity classification according to the ClinVar

database. [c]ID of the variant in the COSMIC (Cosmic_95 coding) database. [d]rsIDs in dbSNP build 150. [e]Individual ID and Variant Allele

Frequency (VAF) for UM samples. Detailed description of selected post-zygotic variants is provided in Supplementary Table 6. Confirmation of

post-zygotic variants by Sanger sequencing/High-Resolution Melting, or Duplex sequencing is provided in Supplementary Tables 7 and 8,

respectively, and Supplementary Figure 3. SS – Sanger sequencing. HRM – High-Resolution Melting. DS – Duplex sequencing. n.a.- not

available. *variants were also detected in the distal uninvolved mammary gland sample (UMD) of selected patients.

 .MAIN FIGURES

Figure 1. Graphical representation of the sampling design used for variant investigation. (A)

Sampling design. Matched fresh-frozen uninvolved mammary gland (UM) and primary tumor (PT)

samples were collected from all breast cancer patients to identify variants according to study criteria

(Materials & Methods). UM samples were located at least 1 cm away from the corresponding PT. Control

mammary gland samples were obtained from individuals who underwent reduction mammoplasty

surgeries and had no history of cancer (Reduction Mammoplasty cohort, RM). (**B) Investigation of post-**

zygotic and germline variants. Matched peripheral blood (BL) or skin (SK) samples were collected for

each individual as reference samples to distinguish between post-zygotic and germline variants. Post-

zygotic variants were identified as those that were absent from the reference samples.

Figure 2. Post-zygotic (A) *PIK3CA* **and (B)** *TP53* **variants detected in the uninvolved mammary**

gland (UM) and primary tumor (PT) samples of breast cancer patients with adverse prognoses and

patients recruited without any prognoses bias (BCAP and BCUS cohorts, respectively). Lollipop

plots represent post-zygotic variants of *PIK3CA* and *TP53* genes detected by Whole Exome Sequencing

(WES). The upper panel represents variants detected in UM samples of BCAP patients. The lower panel

represents variants detected in UM samples of BCUS patients. *TP53* variants were found exclusively in

the BCAP cohort. All identified variants have been reported in the COSMIC database

(https://cancer.sanger.ac.uk/cosmic). Detailed descriptions of variants can be found in Supplementary

Table 6. p85 - p85 binding domain, RBD - Ras-binding domain, C2 - C2 domain, AD - accessory domain,

CD - catalytic domain. TAD1, TAD2 - transcription activation domain 1 and 2, DBD - DNA-binding

domain, DNA-binding sites are marked with green lines, TD - tetramerization domain. Lollipop plots

were prepared based on the images generated with the Protein paint application[53]. () indicates the number

of patients in whom the variant was identified.

Figure 3. Pathogenic post-zygotic variants in breast cancer patients with adverse prognoses.

Variants were confirmed via Sanger sequencing/High-Resolution Melting or Duplex sequencing

(Materials and Methods, Table 2, Supplementary Tables 7 and 8, Supplementary Figure 3). Variant

presence in the ClinVar database and follow-up information for the corresponding patients are included.

Detailed clinicopathological information for presented patients is provided in Supplementary Table 1. A

full description of detected variants is provided in Supplementary Table 6. PT – primary tumor. UMD –

uninvolved mammary gland at a further distance from corresponding PT.

Figure 4. Kaplan-Meier survival curves of breast cancer patients with pathogenic variants and

recurrent disease. The curves represent survival probabilities for different groups of patients from the

BCAP cohort (breast cancer patients with adverse prognoses) and the BCUS cohort (breast cancer

patients without specific prognosis criteria), stratified by the presence of recurrent disease and/or

pathogenic germline or post-zygotic variants in breast cancer-specific genes. Survival time was measured

from the date of diagnosis to death or the end of the follow-up period (10 years for BCAP and 2 years for

BCUS). Detailed information on recurrence status and patient outcomes can be found in Supplementary

Table 1. The x-axis represents time in months, and the y-axis represents the probability of survival.

Vertical ticks on the curves indicate censored death events.

(A)

UM

PT

Malignant transformation

- post-zygotic variant
- germline variant

(B)

SK OR BL

...ACTGCTC... ....TGATCGT...

UM ≥1 cm from PT

...ACTCCTC... ...TGATCGT...

PT

...ACTCCTC... ...TGATCGT...

post-zygotic variant *germline variant*