Associations Between Female Sex Hormones and Skeletal Muscle Ageing: The
Baltimore Longitudinal Study of Aging

Annabel J. Critchlow[a], Sarah E. Alexander[b], Danielle Hiam[a], Luigi Ferrucci[d], David Scott[a,c], Séverine
Lamon[a]

[a]School of Exercise and Nutrition Sciences, Institute for Physical Activity and Nutrition (IPAN),
Deakin University, Geelong, Australia.

[b] Cardiometabolic Health and Exercise Physiology, Baker Heart and Diabetes Institute, Melbourne,
Australia

[c]School of Clinical Sciences at Monash Health, Monash University, Clayton, Australia.

[d]National Institute on Aging, National Institutes of Health, Baltimore, Maryland, United States

Corresponding author:

A/Prof. Séverine Lamon

School of Exercise and Nutrition Sciences, Faculty of Health

Deakin University, 221 Burwood Hwy, Burwood 3125.

Australia

Ph (+61 3) 9244 5571

Email: severine.lamon@deakin.edu.au

Author contact details:

Annabel Critchlow: annabel.critchlow@deakin.edu.au

Dr Sarah Alexander: sarah.alexander@baker.edu.au

Dr Danielle Hiam: danielle.hiam@deakin.edu.au

Dr Luigi Ferruci: ferruccilu@grc.nia.nih.gov

A/Prof David Scott: d.scott@deakin.edu.au

Word count: 4452

Number of figures: 3

Number of tables: 2

Number of references: 39

Abstract

Background: To date, most research investigating the influence of circulating sex hormones on ageing female skeletal muscle has been cross-sectional and focused only on dichotomised young and old, or pre- versus post-menopausal groups. This excludes an important transitional period from high to low circulating oestrogen. Using secondary data from the Baltimore Longitudinal Study of Aging, this study aimed to investigate cross-sectional and longitudinal associations between circulating sex hormones and skeletal muscle mass and function across a continuum of ages.

Methods: Multiple and binomial linear regression was used to map cross-sectional (n=319) and longitudinal (n=83) associations between circulating sex hormones (oestradiol (E2), free oestradiol index (FEI), free (TT) and bioavailable testosterone (BioT), testosterone/oestradiol ratio (TT/E2)) and skeletal muscle mass and function in healthy females. Cross-sectional models analysed females across an ageing continuum (24-89 years) and longitudinal associations were tested across 4-6 years of ageing in females over 50 years old. Models were adjusted for age, height, physical activity, comorbidities, and ethnicity.

Results: Cross-sectionally, serum E2 and FEI were positively associated with relative appendicular lean mass (ALM; β=0.28 and 0.20, respectively, $p<0.05$) and thigh muscle percentage (β=0.19 and 0.15, respectively, $p<0.05$). E2 and FEI were negatively associated with total body fat percentage (β=-0.30 and -0.21, respectively, $p<0.05$). BioT was positively associated with absolute ALM (β=0.13, $p<0.05$) and total body fat percentage (β=0.18, $p<0.05$). TT was negatively associated with total body fat percentage (β=-0.14, $p<0.05$). The TT/E2 ratio was negatively associated with thigh muscle CSA (β=-0.08, $p<0.05$) and hamstring strength (β=-0.12, $p<0.05$). Across 4-6 years, a decrease in E2 and FEI were associated with a decrease in ALM (β=0.29 and 0.42, respectively, $p<0.05$), and decrease in FEI was associated with a decrease in handgrip strength (β=0.17, $p<0.05$). A decrease in TT and BioT were associated with an increase in relative ALM (β=-0.25 for both, $p<0.05$) and relative quadriceps strength (β=-0.12 and -0.27, respectively, $p<0.05$)

Conclusion: This study demonstrates novel associations between sex hormone levels and skeletal muscle in females across a wide continuum of ages. We also demonstrate that longitudinal fluctuations in circulating sex hormones must be considered to gain a comprehensive understanding of female muscle ageing.

Keywords: oestrogen, ovarian hormones, skeletal muscle, ageing

Introduction

The World Health Organization [1] predicts the global population above 60 years of age will double from 2019 to 2050. However, improvements to increase the lifespan have not been matched by advancements in the health span; that is, the length of time a person remains healthy, not just alive. While females live longer than males [2], those additional years are spent in significantly poorer health, highlighting a large disparity between the health span and lifespan in females compared to males [3].

Ageing is associated with a progressive loss of muscle mass and function which presents an increasingly substantial public health concern. Age-related muscle wasting is significantly associated with many adverse outcomes, including reduced independence and quality of life, and increased chronic disease, hospitalisation, and mortality [4]. While the prevalence of age-related muscle loss is seemingly similar between the sexes [5], evidence suggests that the patterns, and therefore the underlying molecular mechanisms of muscle ageing may be sex-specific. For example, females undergo a rapid decline in knee extensor strength ~20 years earlier than males [6]. In addition, while both sexes share the same direction of changes to the transcriptome with ageing, the magnitudes of change to differentially expressed genes differs between the sexes from young and old individuals [7].

Testosterone and oestrogen are the predominant sex hormones in males and females, respectively. While they are produced by both sexes, their profiles vary drastically across the lifespan. Females undergo a dramatic decline in endogenous oestrogen production starting ~40 years of age [8], meaning they can spend over a 3^{rd} of their life in an oestrogen-deficient state. A steep reduction in total and bioavailable (i.e. not bound to sex hormone-binding globulin (SHBG)) testosterone occurs in the early reproductive years, which stabilises around menopause [9]. In contrast, male testosterone production remains largely consistent across adult life with a gradual decline in later decades [9]. Oestrogen and testosterone elicit effects in multiple tissues, including skeletal muscle, by binding to the oestrogen (ER) and androgen receptor (AR), respectively [10]. In females, bioavailable testosterone is associated the maintenance of muscle mass [11], and oestrogen may regulate the maintenance of muscle mass, strength, and mitochondrial function [12].

A recent systematic review by our group found a consistent reduction in muscle mass and handgrip strength across the menopausal transition, however many studies show no association between oestradiol concentrations (E2, the major endogenous form of oestrogen) and muscle mass or function in pre- or postmenopausal females [13]. The cross-sectional nature and dichotomisation of age groups in this body of evidence is a limitation because it prevents conclusions to be drawn about the effects of changes in sex hormone concentrations on muscle health over time. Only one study has explored this association in a longitudinal manner, reporting no relationship between baseline circulating E2 or testosterone and the loss of appendicular lean mass or knee extensor strength over 3 years in

postmenopausal females [14]. However, it remains largely unknown how age-related fluctuations in
sex hormones impact the regulation of skeletal muscle across a broader age range, as associations
between the changes in sex hormones and skeletal muscle health across ageing and the menopausal
transition are still to be mapped.

This study used data from the Baltimore Longitudinal Study of Aging (BLSA, [15]) to address two
research questions. First, we aimed to determine the cross-sectional associations between circulating
female sex hormone concentrations and skeletal muscle mass and function across a continuum of ages
and hormonal status (24 to 89 years). Secondly, we aimed to investigate the longitudinal associations
between circulating sex hormone concentrations and the changes in muscle mass and function that
occurred over a 4–6-year period in females aged 50 to 87 years.

Methods

Study Population

This is a secondary analysis of data from the BLSA [15], an ongoing prospective study that aims to
extensively map the process of healthy ageing [15]. Testing is carried out at the National Institute of
Ageing Clinical Research Unit in Baltimore, Maryland. Healthy participants between 20-96 years
without any chronic disease or cognitive impairment were recruited. Data collected between 2003 and
2023 were utilised in this analysis. Participants under 60 years old were tested every 4 years, 60–79-
year-olds were tested every 2 years, and those above 80 years were tested every year.

Inclusion Criteria

Inclusion criteria for the present study included female BLSA participants with no missing variables
of interest (serum sex hormones, DXA body composition, handgrip, hamstring, and quadriceps
strength, and confounding variables). Data from females at their first visit was used for the cross-
sectional analysis if there were no missing variables (n=319). From these participants, females >50
years were included in the longitudinal analysis (n=83) if they had a follow up visit 4-6 years later
with all the required sex hormone and outcome variables. If they had multiple visits within this time
frame, data from the latest was used. As the BLSA recruit only healthy individuals at baseline, there
were no additional exclusion criteria specific to this study.

Measures

Serum Hormone Analysis

Fasted (12 h) blood samples were taken from participants in the morning of each visit and
immediately centrifuged at 4°C. Serum was aliquoted and stored at -80°C. Fasted serum hormone
concentrations of E2 (ng/dL) and total testosterone (TT, ng/dL) were determined by high-performance
liquid chromatography-tandem mass spectrometry (LC/MS; Esoterix part of LabCorp, Calabasas

Hills, CA). SHBG concentrations (nmol/L) were analysed via immunoradiometric assay (Esoterix part

of LabCorp). To determine bioavailable testosterone (BioT, ng/dL), ammonium sulphate was used to

separate SHBG-bound hormones from albumin-bound and free hormones (Esoterix part of LabCorp).

E2, TT, and BioT concentrations were converted to SI units (pmol/L, nmol/L and nmol/L,

respectively). The free oestrogen index (FEI) was calculated by dividing E2 (pmol/L) by SHBG

(nmol/L) and multiplying by 100. The testosterone to oestrogen (TT/E2) ratio was calculated by

dividing TT (nmol/L) by E2 (pmol/L).

Body Composition

Body composition measures including appendicular lean mass (ALM, kg), leg lean mass (LLM, kg),

and total body fat mass (TBFM, kg) were assessed via whole-body DXA (Prodigy Scanner, GE,

Madison, WI) scans with software version 10.51.006. The scanner was calibrated daily. Reliability has

been assessed in 12 older (>65 years) males, showing <1% difference in fat mass (kg) between 2

scans, 6 weeks apart [16]. Relative ALM ($kg.kg^{-1}$) was calculated by dividing appendicular lean mass

(kg) by total body fat mass (kg). Thigh muscle cross-sectional area (CSA, cm^2), subcutaneous fat area

(cm^2), intramuscular fat area (cm^2), and muscle density (HU) were assessed using computerised

tomography (CT) cross-sectional images of the mid-thigh (Somatom Sensation 10, Siemens, Malvern,

PA). Thigh muscle percentage was calculated by dividing total thigh CSA (cm^2) by muscle CSA

(cm^2). CT images were analysed and quantified with Geanie 2.1 software (BonAlyse Oy, Jyvaskla,

Finland).

Muscle Function

Handgrip strength (kg) was assessed using a Jamar Hydraulic hand dynamometer (Patterson Medical,

Warrenville, IL). Participants sat with their arm extended to 180° or shoulder height and were

instructed to squeeze the dynamometer as hard as possible. Maximum force was measured in three

trials on each hand, and the best of 6 were used in the analysis. The average of the left and right hand

was used in the analysis. Lower limb strength was measured using the Kin-Com isokinetic

dynamometer (Kin-Com model 125E version 3.2, Chattanooga Group, Chattanooga, TN) from 2003

to 2011, and the Biodex Multi-Joint System-Pro dynamometer (Biodex Medical System, Advantage

Software V4.X, Inc., Shirley, NY) from 2011 to 2023. Quadriceps strength (isokinetic knee

extension; Nm) and hamstring strength (isokinetic knee flexion; Nm) was measured at an angular

velocity of 30°/s and 180°/s. Participants were asked to extend/flex their knee as hard as possible

through a range of motion set between 100-160°. They completed two practice trials, followed by

three test trials. The highest peak torque value was included in the analysis. Peak torque normalised to

leg lean mass (Nm/kg) and thigh muscle CSA (Nm/cm^2), were calculated by dividing hamstring and

quadriceps peak torque (Nm) by leg lean mass (kg) and thigh muscle CSA (cm^2), respectively.

Physical Function

Gait speed was assessed by instructing participants to walk 400 m as fast as possible after a warm-up

of 2.5 minutes at a usual gait speed. Chair stand ability was measured by asking participants to fold

their arms and stand up from a chair 10 times. The pace of the first 5 chair stands (stands/s) was

recorded. Participants completed three tests (6 m walk test, chair stand test, and standing balance) to

comprise the Short Physical Performance Battery (SPPB), a validated assessment of mobility.

Performance in each of the 3 tests was scored from 0 (worst) to 4 (best) and summed to give a total

out of 12, as previously described [17]. Finally, participants were categorised as a faller if they had at

least one self-reported fall in the previous 12 months.

Potential confounding variables

Demographic information was collected from participants via an interview, including age, ethnicity,

comorbidities, and total physical activity in the previous two weeks. The age adjusted Charlson

comorbidity index (ACCI) was determined as previously described [18]. Briefly, a score is assigned to

each comorbid condition, with additional points for increasing age groups.

Statistical Analysis

Statistical analysis was completed using Stata software version 17.0 (StataCorp, College Station, TX).

Data are presented as mean $\pm$ SD and statistical significance was accepted as $p<0.05$. Participants with

values that were deemed physiologically implausible (>10 SDs) were excluded. One participant was

excluded from the cross-sectional analysis because handgrip strength was >300kg. Forty and 21

participants were excluded from the cross-sectional and longitudinal analysis, respectively, because

their hormone levels were below zero. Paired t-tests were performed to determine the differences

between variables at baseline and follow-up.

Multiple regression

Multiple linear regression was performed to determine the cross-sectional relationships between

circulating sex hormones, body composition (from DXA), and muscle function. Further outcome

variables were assessed in subsets of participants where measured, including thigh CT measures and

physical function. Additional models were performed to determine cross-sectional associations

between self-reported menopausal and hormonal supplementation status and all muscle outcomes. In

the longitudinal analysis, multiple regression models were performed to determine the relationship

between the change in sex hormone levels and the change in outcome measures across 4-6 years in

females over 50 years. Further analysis was completed to measure the longitudinal associations

between baseline menopausal status and HRT use and the change in muscle outcome measures.

Each model was adjusted for age, ethnicity, height, ACCI, and physical activity. Where significant and

opposing associations were detected for serum oestradiol (E2 or FEI) and testosterone (TT or BioT),

both hormones were added to the model to determine if they remained significantly and independently
associated with the outcome variable.

Multicollinearity was assessed by variation inflation factors with a threshold of 5. No variables
displayed collinearity. Normality of residuals was confirmed by plotting a histogram of studentised
residuals. Homoscedasticity was assessed by plotting fitted versus predicted residuals. Some
heteroscedasticity was observed, so robust standard errors were used. To determine the magnitude of
the relationships, the z-score was calculated for all independent (E2, FEI, TT, BioT, SHBG, and
TT/E2 ratio) and outcome measures (body composition, muscle function, and physical function),
where zero equals the mean, and 1 equals the standard deviation. Where the standardised coefficient
was <0.2, the effect size of the relationship was considered 'small', 0.2 to 0.5 denotes a 'medium'
effect size, and >0.5 represents a 'large' effect size.

Binomial linear regression

In females over 50 years of age, binomial linear regression models were used to determine the cross-
sectional relationship between sex hormone levels and dichotomous physical function outcome
measures (mobility disability and fallers).

Results

Participant characteristics

Characteristics of the participants included in the cross-sectional and longitudinal analysis are shown
in Tables 1 and 2, respectively. Self-reported menopausal status and hormonal supplementation is
shown in Supplementary Table S1. In the longitudinal analysis, there was a 50.8% reduction ($p<0.05$)
from baseline (mean age: 64.41 ± 10.99 years) to follow up (4.86 ± 0.88 years later) in circulating E2,
but not in other hormones. In body composition measures, there was a 4.6% reduction in thigh muscle
CSA ($p<0.05$), and a 3.9% reduction in thigh muscle density ($p<0.05$). Regarding muscle function,
there was a reduction in absolute and relative hamstring strength at 180 °/s (absolute: -12%, relative
to LLM: -11.2%, relative to muscle CSA: -8.4%, $p<0.05$), and absolute and relative quadriceps
strength at 180°/s (absolute: -8.0%, relative to LLM: -7.5%, $p<0.05$) and 30°/s (absolute: -8.2%,
relative to LLM: -8.1%, relative to CSA: -4.5%, $p<0.05$). There was a 9% increase in 400 m walk
time ($p<0.05$).

Cross-sectional analysis

Adjusted cross-sectional associations between circulating sex hormones and skeletal muscle outcome
measures in females between 24-89 years are displayed in Figure 1-3. Amongst the body composition
measures, circulating E2 and FEI were positively associated with relative ALM (kg, $p<0.05$). BioT
was positively associated with absolute ALM (kg, $p<0.05$) but negatively associated with relative

ALM (kg, $p<0.05$). E2, FEI, and TT were negatively associated with total body fat percentage
($p<0.05$), but there was a positive association for BioT ($p<0.05$). E2 and FEI were positively
associated with thigh muscle percentage ($p<0.05$), and TT/E2 ratio was negatively associated with
thigh muscle CSA (cm^2, $p<0.05$). BioT was positively associated with subcutaneous and
intramuscular fat (kg, $p<0.05$), while E2 was negatively associated with subcutaneous fat (kg,
$p<0.05$). In muscle function models, TT/E2 ratio was negatively associated with hamstring peak
torque (Nm) at 180°/s ($p<0.05$) and BioT was positively associated with 400 m walk time (s, $p<0.05$).
The binomial linear regression results are displayed in Table S2. There were no significant
associations between any hormones and indices of mobility disability or falls in females over 50 years
(n=238).

Cross-sectional associations between menopausal status, hormonal supplementation and muscle
outcomes are displayed in Table S3. Postmenopausal status was positively associated with total body
fat (%, $p<0.05$), subcutaneous fat (kg, $p<0.05$), and intramuscular fat (kg, $p<0.05$), and negatively
associated with relative ALM (kg, $p<0.05$) and thigh muscle CSA (cm^2, $p<0.05$). Amongst
postmenopausal females, use of HRT was positively associated with handgrip strength (kg, $p<0.05$).
In premenopausal females, use of the HCP was positively associated with relative quadriceps strength
normalised to leg lean mass at 30°/s (Nm/kg, $p<0.05$). It was also negatively associated with muscle
density (HU, $p<0.05$), absolute and relative hamstring strength (Nm and Nm/kg, $p<0.05$), and relative
quadriceps strength normalised to thigh muscle CSA at 30°/s (Nm/cm^2, $p<0.05$).

Where E2/FEI and BioT were found to have opposite associations with a muscle outcome, both
hormones were included in the model. BioT was added to the cross-sectional models investigating the
relationship between E2 and relative ALM, total body fat percentage, and subcutaneous fat, and the
relationship between FEI and relative ALM and total body fat. In all models, oestradiol (E2 or FEI)
remained significantly associated positively with relative ALM and negatively with body fat
percentage and subcutaneous fat, while BioT remained significantly associated in the opposite
direction ($p<0.05$; Table S4).

Longitudinal analysis

Adjusted longitudinal associations between the change in circulating sex hormones and the change in
skeletal muscle outcome measures in females over 50 years are displayed in Figures 1-3. Regarding
body composition, ΔE2 and ΔFEI were positively associated with the change in ALM (kg, $p<0.05$).
ΔTT and ΔBioT were negatively associated with the change in relative ALM (kg, $p<0.05$). ΔFEI was
negatively associated with the change in thigh muscle percentage ($p<0.05$), and positively associated
with the change in subcutaneous and intramuscular fat (kg, $p<0.05$). Regarding muscle function, ΔFEI
was positively associated with the change in handgrip strength (kg, $p<0.05$), but negatively associated
with the change in quadriceps strength (Nm) at 180°/s ($p<0.05$). Similarly, ΔTT and ΔBioT were

negatively associated with the change in relative quadriceps strength at 30°/s when normalised to leg

lean mass (Nm/kg, $p<0.05$). Finally, ΔBioT was positively associated with the change in 400 m walk

time (s, $p<0.05$).

Longitudinal associations between baseline menopausal status, HRT use and the change in muscle

outcomes across 4-6 years in females over 50 years old are displayed in Table S5. Postmenopausal

status at baseline was positively associated with the change in ALM (kg, $p<0.05$) and thigh muscle

CSA (cm^2, $p<0.05$). Being a current HRT user at baseline was negatively associated with the change

in handgrip strength (kg, $p<0.05$), while previous use of HRT was positively associated with the

change in intramuscular fat (kg, $p<0.05$) and relative hamstring strength normalised to muscle thigh

CSA (Nm/cm^2, $p<0.05$), and negatively associated with the change in thigh muscle CSA (cm^2,

$p<0.05$).

Discussion

281   To the best of the authors knowledge, this is the first study to 1) map cross-sectional associations

282   between sex hormones and muscle outcomes in a wide continuum of ages across the adult lifespan,

283   and 2) investigate how changes to circulating sex hormones beyond 50 years of age associate with

284   changes to muscle mass and function over 4-6 years.

Sex hormones and body composition:

Oestradiol

This study highlighted a positive cross-sectional association between serum E2 levels (total and FEI)

and measures of muscle mass (relative ALM and thigh muscle percentage) in females across the

lifespan, independent of age. Additionally, both total serum E2 and FEI were negatively associated

with total body fat percentage and thigh subcutaneous fat, suggesting a potential role for E2 in

maintaining the proportion of muscle mass throughout ageing. This is line with previous mechanistic

work in animals highlighting an anabolic role of E2 in skeletal muscle [12]. Interestingly, this

contradicts with many previous cross-sectional studies investigating samples of postmenopausal

females, which demonstrate no associations between serum oestradiol or FEI and muscle mass [13].

This is likely due to the inclusion of postmenopausal females that, by definition, have very low serum

E2 concentrations, so there is unlikely to be large variation between participants. In contrast, the

current cohort spans a wide continuum of adulthood (24 to 89 years) and E2 concentrations (5 to 1872

pmol/L), thereby allowing us to understand how large age-related fluctuations in oestrogens impact

the regulation of muscle mass over the lifespan.

In the longitudinal analysis of females over 50 years old, the change in serum E2 and FEI across the

5-year follow up was positively associated with the change in absolute ALM, suggesting females with

a larger decline in serum E2 and FEI have a greater decline in muscle mass. This further emphasises a

potential role for total and free E2 in anabolic regulation, in line with the cross-sectional findings. The

only longitudinal study to have previously investigated this found no association between serum E2

and the loss of ALM over 3 years in 49 postmenopausal females [14]. However, baseline hormone

concentrations were used in the model, as opposed to the change in hormone concentration from

baseline to follow up. This suggests that it may not be baseline E2 concentration, but the magnitude of

change in E2 concentration from baseline, that determines the extent of muscle loss post-menopause.

It is therefore important to understand the temporal nature of the relationship between E2 and skeletal

muscle, where the population is displaying rapid changes in both hormone concentrations and muscle

mass and function.

Surprisingly, the change in serum FEI was negatively associated with the change in thigh muscle

percentage, and positively associated with the change in thigh subcutaneous and intramuscular fat.

Together this suggests that a decline in serum FEI coincides with an increased relative proportion of

muscle in the thigh, potentially due to declines in both subcutaneous and intramuscular fat content,

rather than an inverse relationship between FEI and muscle mass. We suggest this is the case, as the

change in FEI was positively associated with absolute ALM. Although, DXA measures of ALM do

not appropriately distinguish muscle from intramuscular fat, which may have exaggerated this

positive association, highlighting the importance of including direct measures of muscle and fat

infiltration from CT scans.

Testosterone

In the cross-sectional analysis, serum TT concentrations were negatively associated with total body fat

percentage and no other compositional measures. This in line with previous studies showing a lack of

association between serum TT and lean mass or muscle CSA in both pre- [11, 19] and postmenopausal

[20-23] females, although this is the first study to investigate the association across the entire adult

lifespan. Instead, bioavailable (non-SHBG bound) or free (non-protein bound) forms of testosterone

appear to be more relevant in this context, as many studies have reported a positive association with

lean mass [11, 20, 21, 23-26] or muscle CSA [27-29].

In line with these studies, BioT concentration was positively associated with absolute ALM. However,

it was also positively associated with measures of fat mass (total body fat percentage, and thigh

subcutaneous and intramuscular fat), explaining the negative association between BioT and relative

ALM; an increase in fat tissue corresponds to a reduction in relative ALM, as relative ALM is indexed

to total body fat mass. While change in muscle and fat mass across 4-6 years of ageing may occur in

the same direction (i.e. both gained or lost), the resulting relative proportion of each tissue differs. A

similar trend was noted in the longitudinal analysis, as both the change in TT and BioT from baseline
to follow up were negatively associated with the change in relative ALM.

Sex hormones and muscle function:

Oestradiol

No significant cross-sectional associations were present between serum E2 or FEI and any measure of
muscle function. This contrasts with Pöllanen et al. [30] who found a positive association between
serum E2 and relative quadriceps femoris strength in an albeit much smaller sample (n=25) of pre-
and postmenopausal females. Importantly, age was not included as a confounding variable, therefore
these findings may just reflect the loss of both serum E2 and muscle strength with age. Indeed, when
age was included as a covariate, Bochud et al. [31] found no cross-sectional association between
urinary E2 concentrations and handgrip strength in a sample of 366 females between 18 and 90 years
of age.

The change in serum FEI was significantly and positively associated with the change in handgrip
strength across 4-6 years, suggesting that a greater decline in free oestradiol is related to a greater
decline in handgrip strength, a common proxy for overall muscle and physical function [32]. In
contrast, the change in serum FEI was negatively associated with the change in absolute quadriceps
strength. This finding is unexpected but may indicate the presence of other confounding factors that
are not yet identified. Rolland et al. [14] found no association between serum E2 and the loss of
muscle strength over 3 years in postmenopausal females, but higher serum oestrone (E1) was
similarly associated with a greater loss of muscle strength. E1 is an endogenous form of oestrogen
which becomes predominant after the menopause-related decline of serum E2 [30]. While less potent
than E2, its role within the muscle remains largely understudied, highlighting a target for future
research, particularly in postmenopausal females.

Testosterone

Similarly to E2, both serum TT and BioT were not cross-sectionally associated with any measures of
muscle strength. This is in line with other studies that investigated younger [11, 19, 27], older [33] or
both pre- and postmenopausal [30] females. To our knowledge, this is the first study to investigate
longitudinal associations between serum testosterone and muscle function in ageing females. The
change in both TT and BioT levels were negatively associated with the change in relative quadriceps
strength, suggesting that an increase in serum testosterone concentration coincides with a decrease in
lower limb strength. While this is unexpected, it is further supported by the positive cross-sectional
and longitudinal associations between BioT and gait speed. This suggests that greater concentrations
of, or changes to, circulating bioavailable testosterone associate with worse physical function in the
lower limbs of ageing females. This is surprising as testosterone supplementation in postmenopausal

females can elicit anabolic effects in skeletal muscle [34], although basal molecular regulation of

muscle by testosterone has not yet been investigated in females.

The TT/E2 ratio and skeletal muscle ageing:

When investigating their effects within skeletal muscle, circulating sex hormones have most often

been considered individually, but their production is closely linked due to common biosynthetic

pathways. For example, testosterone is converted to oestradiol via the enzyme aromatase; a process

that can be quantitatively assessed by the TT/E2 ratio, where a greater TT/E2 ratio indicates less

synthesis of E2 from testosterone, and therefore lower aromatase activity [35]. In premenopausal

females, the ovaries are responsible for ~95% of circulating oestradiol production, but the peripheral

conversion of testosterone to E2 becomes more physiologically relevant after menopause when the

ovaries cease E2 production [35, 36]. We observed a negative cross-sectional association between the

TT/E2 ratio and muscle mass (thigh muscle CSA) and strength (hamstring peak torque), suggesting

that a lower conversion rate of testosterone to E2 is concomitant to poorer muscle quality across

ageing. Further research should be conducted to understand the contribution of varying oestrogen

synthesis pathways in the regulation of skeletal muscle throughout ageing in females.

Limitations

While this study provides novel insights into the relationship between circulating sex hormones and

skeletal muscle ageing, it is not without limitations. Menstrual cycle phase has not been controlled

when scheduling premenopausal participant visits. Common measures of acute skeletal muscle

function are unlikely to be altered according to menstrual cycle phase [37], however, circulating E2

levels can vary from 70-1500 pmol/L between the early follicular and ovulatory phase [38].

Therefore, depending on the phase in which the blood sample was collected, the relationship between

circulating E2 and skeletal muscle outcomes may significantly differ. Future studies should test all

premenopausal participants within the early follicular phase as previously described [39], where

serum E2 is at its lowest or 'basal' level.

Secondly, premenopausal participants provided information on HCP use, but many other forms of

hormonal contraception are commonly used and provide external doses of oestrogen or progesterone,

including the intrauterine devices, implant, injections and skin patches. Therefore, a more

comprehensive understanding of exogenous sex hormone profiles is required to improve reliability of

the findings.

Finally, approximately one third of participants in this study were non-white, suggesting that these

findings are representative of a range of ethnicities. However, due to the inclusion criteria of the

BLSA, participants are likely healthier than the general population. Therefore, the generalisability of
these findings to females with multiple chronic health conditions is questionable.

Conclusion

This study has demonstrated novel and divergent cross-sectional and longitudinal associations
between circulating sex hormones and skeletal muscle outcomes in females across the ageing
continuum. This emphasises the need to consider the temporal aspect of sex hormone regulation in
ageing female muscle, particularly across the transitional period from a high to low state of circulating
oestrogen. Future research should aim to identify molecular mechanisms of oestrogen action in human
female skeletal muscle, to determine how ageing-induced oestrogen deficiency contributes to a loss of
muscle mass and function in the later decades of life.

Additional Information

Funding

SL is funded by an Australian Research Council Future Fellowship (FT210100278). DS is funded by a
National Health and Medical Research Council (NHMRC) Australia Investigator Grant
(GNT1174886). Supported in part by the Intramural Research Program of the National Institute on
Aging, NIH. Baltimore, MD, USA.

Ethical Standards

Written informed consent was provided by all participants at every visit and the BSLA protocol was
approved by the National Institute of Health (NCT00233272). This study (project number: 2023-029)
has been declared exempt from ethical review on 07/02/2023 by the Deakin University Human
Research Ethics Committee (DUHREC) as it comprises only of secondary data analysis.

Conflicts of Interest

Annabel Critchlow declares that she has no conflict of interest. Dr Sarah Alexander declares that she
has no conflict of interest. Dr Luigi Ferrucci declares that he has no conflict of interest. A/Prof David
Scott declares that he has no conflict of interest. A/Prof Severine Lamon declares that she has no
conflict of interest.

References

1. World Health Organization. Ageing and health. 2024. https://www.who.int/news-room/fact-
sheets/detail/ageing-and-health. Accessed 22/07/2024.

2. Australian Bureau of Statistics. Twenty years of population change. 2020.
https://www.abs.gov.au/articles/twenty-years-population-change. Accessed 22/07/2024.

3. Austad SN, Fischer KE. Sex differences in lifespan. Cell metabolism. 2016;23:1022-33.

4. Peterson SJ, Braunschweig CA. Prevalence of Sarcopenia and Associated Outcomes in the
Clinical Setting. Nutrition in Clinical Practice. 2016;31:40-8. doi:doi.org/10.1177/0884533615622537

5. Petermann-Rocha F, Balntzi V, Gray SR, Lara J, Ho FK, Pell JP, et al. Global prevalence of
sarcopenia and severe sarcopenia: a systematic review and meta-analysis. Journal of Cachexia,
Sarcopenia and Muscle. 2022;13:86-99. doi:doi.org/10.1002/jcsm.12783

6. Haynes EMK, Neubauer NA, Cornett KMD, O'Connor BP, Jones GR, Jakobi JM. Age and
sex-related decline of muscle strength across the adult lifespan: a scoping review of aggregated data.
Appl Physiol Nutr Metab. 2020;45:1185-96. doi:10.1139/apnm-2020-0081

7. de Jong J, Attema BJ, van der Hoek MD, Verschuren L, Caspers MPM, Kleemann R, et al.
Sex differences in skeletal muscle-aging trajectory: same processes, but with a different ranking.
Geroscience. 2023;45:2367-86. doi:10.1007/s11357-023-00750-4

8. Frederiksen H, Johannsen TH, Andersen SE, Albrethsen J, Landersoe SK, Petersen JH, et al.
Sex-specific Estrogen Levels and Reference Intervals from Infancy to Late Adulthood Determined by
LC-MS/MS. J Clin Endocrinol Metab. 2020;105:754-68. doi:10.1210/clinem/dgz196

9. Handelsman DJ, Sikaris K, Ly LP. Estimating age-specific trends in circulating testosterone
and sex hormone-binding globulin in males and females across the lifespan. Annals of Clinical
Biochemistry. 2016;53:377-84. doi:10.1177/0004563215610589

10.   Alexander SE, Pollock AC, Lamon S. The effect of sex hormones on skeletal muscle
adaptation in females. European Journal of Sport Science. 2022;22:1035-45.
doi:10.1080/17461391.2021.1921854

11.   Alexander SE, Abbott G, Aisbett B, Wadley GD, Hnatiuk JA, Lamon S. Total testosterone is
not associated with lean mass or handgrip strength in pre-menopausal females. Scientific Reports.
2021;11:10226. doi:10.1038/s41598-021-89232-1

12.   Pellegrino A, Tiidus PM, Vandenboom R. Mechanisms of Estrogen Influence on Skeletal
Muscle: Mass, Regeneration, and Mitochondrial Function. Sports Med. 2022;52:2853-69.
doi:10.1007/s40279-022-01733-9

13.   Critchlow AJ, Hiam D, Williams R, Scott D, Lamon S. The role of estrogen in female skeletal
muscle aging: A systematic review. Maturitas. 2023;178:107844.
doi:doi.org/10.1016/j.maturitas.2023.107844

14.   Rolland YM, Perry HM, 3rd, Patrick P, Banks WA, Morley JE. Loss of appendicular muscle
mass and loss of muscle strength in young postmenopausal women. J Gerontol A Biol Sci Med Sci.
2007;62:330-5. doi:10.1093/gerona/62.3.330

15.   Shock NW. Normal human aging: The Baltimore longitudinal study of aging. Washington
D.C.: National Institute of Health; 1984.

16.   Lindle RS, Metter EJ, Lynch NA, Fleg JL, Fozard JL, Tobin J, et al. Age and gender
comparisons of muscle strength in 654 women and men aged 20–93 yr. Journal of Applied
Physiology. 1997;83:1581-7. doi:10.1152/jappl.1997.83.5.1581

17.   Guralnik JM, Simonsick EM, Ferrucci L, Glynn RJ, Berkman LF, Blazer DG, et al. A short
physical performance battery assessing lower extremity function: association with self-reported
disability and prediction of mortality and nursing home admission. J Gerontol. 1994;49:M85-94.
doi:10.1093/geronj/49.2.m85

18.   Zhou S, Zhang X-H, Zhang Y, Gong G, Yang X, Wan W-H. The Age-Adjusted Charlson
Comorbidity Index Predicts Prognosis in Elderly Cancer Patients. Cancer Management and Research.
2022;14:1683-91. doi:10.2147/CMAR.S361495

19.   Alexander SE, Gatto B, Knowles OE, Williams RM, Fiebig KN, Jansons P, et al. Bioavailable
testosterone and androgen receptor activation, but not total testosterone, are associated with muscle
mass and strength in females. The Journal of Physiology. 2024;doi:https://doi.org/10.1113/JP286803

20.   Gower BA, Nyman L. Associations among oral estrogen use, free testosterone concentration,
and lean body mass among postmenopausal women. J Clin Endocrinol Metab. 2000;85:4476-80.
doi:10.1210/jcem.85.12.7009

21.   Bann D, Wu FC, Keevil B, Lashen H, Adams J, Hardy R, et al. Changes in testosterone
related to body composition in late midlife: Findings from the 1946 British birth cohort study. Obesity
(Silver Spring). 2015;23:1486-92. doi:10.1002/oby.21092

22.   Sipilä S, Heikkinen E, Cheng S, Suominen H, Saari P, Kovanen V, et al. Endogenous
Hormones, Muscle Strength, and Risk of Fall-Related Fractures in Older Women. The Journals of
Gerontology: Series A. 2006;61:92-6. doi:10.1093/gerona/61.1.92

23.   Rariy CM, Ratcliffe SJ, Weinstein R, Bhasin S, Blackman MR, Cauley JA, et al. Higher
serum free testosterone concentration in older women is associated with greater bone mineral density,
lean body mass, and total fat mass: the cardiovascular health study. J Clin Endocrinol Metab.
2011;96:989-96. doi:10.1210/jc.2010-0926

24.   Hwang AC, Liu LK, Lee WJ, Chen LY, Lin MH, Peng LN, et al. Association of androgen with
skeletal muscle mass and muscle function among men and women aged 50 years and older in Taiwan:
results from the I-Lan longitudinal aging study. Rejuvenation Res. 2013;16:453-9.
doi:10.1089/rej.2013.1442

25.   van Geel TA, Geusens PP, Winkens B, Sels JP, Dinant GJ. Measures of bioavailable serum
testosterone and estradiol and their relationships with muscle mass, muscle strength and bone mineral
density in postmenopausal women: a cross-sectional study. Eur J Endocrinol. 2009;160:681-7.
doi:10.1530/eje-08-0702

26.   Kong SH, Kim JH, Lee JH, Hong AR, Shin CS, Cho NH. Dehydroepiandrosterone Sulfate
and Free Testosterone but not Estradiol are Related to Muscle Strength and Bone Microarchitecture in
Older Adults. Calcif Tissue Int. 2019;105:285-93. doi:10.1007/s00223-019-00566-5

27.   Häkkinen K, Pakarinen A, Kallinen M. Neuromuscular adaptations and serum hormones in
women during short-term intensive strength training. Eur J Appl Physiol Occup Physiol. 1992;64:106-
11. doi:10.1007/bf00717946

28.   Häkkinen K, Pakarinen A, Kraemer WJ, Häkkinen A, Valkeinen H, Alen M. Selective muscle
hypertrophy, changes in EMG and force, and serum hormones during strength training in older
women. J Appl Physiol (1985). 2001;91:569-80. doi:10.1152/jappl.2001.91.2.569

29.   Yee ML, Hau R, Taylor A, Guerra M, Guerra P, Darzins P, et al. Sarcopenia in women with
hip fracture: A comparison of hormonal biomarkers and their relationship to skeletal muscle mass and
function. Osteoporos Sarcopenia. 2020;6:139-45. doi:10.1016/j.afos.2020.06.001

30.   Pöllänen E, Sipilä S, Alen M, Ronkainen PH, Ankarberg-Lindgren C, Puolakka J, et al.
Differential influence of peripheral and systemic sex steroids on skeletal muscle quality in pre- and
postmenopausal women. Aging Cell. 2011;10:650-60. doi:10.1111/j.1474-9726.2011.00701.x

31.   Bochud M, Ponte B, Pruijm M, Ackermann D, Guessous I, Ehret G, et al. Urinary Sex Steroid
and Glucocorticoid Hormones Are Associated With Muscle Mass and Strength in Healthy Adults. J
Clin Endocrinol Metab. 2019;104:2195-215. doi:10.1210/jc.2018-01942

32.   Vaishya R, Misra A, Vaish A, Ursino N, D'Ambrosi R. Hand grip strength as a proposed new
vital sign of health: a narrative review of evidences. J Health Popul Nutr. 2024;43:7.
doi:10.1186/s41043-024-00500-y

33.   Carcaillon L, Blanco C, Alonso-Bouzón C, Alfaro-Acha A, Garcia-García FJ, Rodriguez-
Mañas L. Sex differences in the association between serum levels of testosterone and frailty in an
elderly population: the Toledo Study for Healthy Aging. PLoS One. 2012;7:e32401.
doi:10.1371/journal.pone.0032401

34.   Huang G, Basaria S, Travison TG, Ho MH, Davda M, Mazer NA, et al. Testosterone dose-
response relationships in hysterectomized women with or without oophorectomy: effects on sexual
function, body composition, muscle performance and physical function in a randomized trial.
Menopause. 2014;21:612-23. doi:10.1097/gme.0000000000000093

35.   Cui J, Shen Y, Li R. Estrogen synthesis and signaling pathways during aging: from periphery
to brain. Trends Mol Med. 2013;19:197-209. doi:10.1016/j.molmed.2012.12.007

36.   Vrtačnik P, Ostanek B, Mencej-Bedrač S, Marc J. The many faces of estrogen signaling.
Biochemia medica. 2014;24:329-42.

37.   Colenso-Semple LM, D'Souza AC, Elliott-Sale KJ, Phillips SM. Current evidence shows no
influence of women's menstrual cycle phase on acute strength performance or adaptations to
resistance exercise training. Front Sports Act Living. 2023;5:1054542.
doi:10.3389/fspor.2023.1054542

38.   Stricker R, Eberhart R, Chevailler M-C, Quinn F, Bischof P, Stricker R. Establishment of
detailed reference values for luteinizing hormone, follicle stimulating hormone, estradiol, and
progesterone during different phases of the menstrual cycle on the Abbott ARCHITECT® analyzer.
Clinical chemistry and laboratory medicine : CCLM / FESCC. 2006;44:883-7.
doi:10.1515/CCLM.2006.160

39.   Elliott-Sale KJ, Minahan CL, de Jonge X, Ackerman KE, Sipilä S, Constantini NW, et al.
Methodological Considerations for Studies in Sport and Exercise Science with Women as
Participants: A Working Guide for Standards of Practice for Research on Women. Sports Med.
2021;51:843-61. doi:10.1007/s40279-021-01435-8

Table 1. Participant characteristics from cross-sectional analysis.

Participant Characteristics	n	Mean	SD	Range
General				
Age (years)	319	60.20	15.95	24.00 - 89.00
Sex hormone levels				
E2 (pmol/L)	319	122.78	234.64	5.14 - 1872.21
FEI	319	0.16	0.26	0.00 - 1.69
TT (nmol/L)	319	0.83	0.53	0.14 - 4.93
BioT (nmol/L)	319	0.13	0.10	0.02 - 0.80
SHBG (nmol/L)	319	81.65	45.49	13.87 - 379.00
TT/E2 ratio	319	33.43	30.55	0.45 - 158.33
Body composition				
BMI (kg/m^2)	319	26.51	5.02	17.83 - 45.74
ALM (kg)	319	17.49	2.78	10.93 - 25.76
Total body fat (%)	319	38.08	7.72	15.07 - 58.19
Relative appendicular lean mass (kg/kg total fat)	319	0.73	0.32	0.29 - 2.66
thigh muscle CSA (cm^2)	239	92.53	22.43	51.99 - 256.99
Thigh subcutaneous fat area (cm^2)	239	105.49	44.65	37.16 - 291.15
Thigh intramuscular fat area (cm^2)	239	10.31	4.18	1.17 - 29.45
Thigh muscle percentage (%)	239	41.60	8.10	22.95 - 63.96
Thigh muscle density (HU)	239	51.02	3.66	38.65 - 60.85
Muscle function				
Hand grip strength (kg)	319	25.56	6.83	10.00 - 53.00
Hamstring peak torque (Nm)				
180d/s	319	49.12	14.69	20.16 - 102.70
30d/s	319	46.89	15.11	11.30 - 106.80
Quadriceps peak torque (Nm)				
180d/s	319	63.07	21.73	23.70 - 120.40
30d/s	319	101.87	31.89	33.31 - 211.30
Hamstring relative peak torque to LLM (Nm/kg)				
180d/s	319	3.66	0.95	1.82 - 8.17
30d/s	319	3.49	0.98	0.81 - 8.02
Quadriceps relative peak torque to LLM (Nm/kg)				
180d/s	319	4.68	1.40	1.56 - 9.66
30d/s	319	7.56	1.97	2.19 - 13.21
Hamstring relative peak torque to muscle CSA (Nm/cm^2)				
180d/s	239	0.53	0.12	0.22 - 0.90
30d/s	239	0.51	0.14	0.13 - 0.93
Quadriceps relative peak torque to muscle CSA (Nm/cm^2)				
180d/s	239	0.69	0.18	0.21 - 1.29
30d/s	239	1.13	0.28	0.46 - 1.79
Physical function				
6-metre walk time (s)	203	5.17	0.91	3.31-7.92
400-metre walk time (s)	163	277.40	49.70	199.88 - 531.61
Time taken to complete 5 chair stands (s)	203	9.50	2.70	3.57 - 18.52
Mobility disability (%(n))	238			
Yes (SPPB ≤ 9)		1.3 (3)	-	-
No (SPPB > 9)		98.7 (235)	-	-
Falls in last 12 months (% (n))	238			
No falls		80.7 (192)	-	-
Falls		19.3 (46)	-	-

Physical function measures were only assessed in females ≥ 50 years old. ALM: appendicular lean mass; BioT: bioavailable testosterone; BMI: body mass index; CSA: cross-sectional area; E2: oestradiol; FEI: free oestradiol index; HU: Hounsfield unit; LLM: leg lean mass; SHBG: sex hormone-binding globulin; SPPB: short physical performance battery; TT: total testosterone; TT/E2: ratio of total testosterone to oestradiol.

Table 2. Participant characteristics from longitudinal analysis.

Participant Characteristics	n	Baseline	Follow up	Change	*p*-value
General					
Age (years)	83	64.41 ± 10.99	69.24 ± 10.95	4.83 ± 0.88	**<0.01***
Sex hormone levels					
E2 (pmol/L)	83	50.97 ± 93.99	25.1 ± 24.9	-25.87 ± 87.61	**<0.01***
FEI	83	0.08 ± 0.14	0.05 ± 0.08	-0.03 ± 0.14	0.08
TT (nmol/L)	83	0.76 ± 0.59	0.76 ± 0.43	0 ± 0.45	0.95
BioT (nmol/L)	83	0.13 ± 0.09	0.13 ± 0.09	0 ± 0.06	0.72
SHBG (nmol/L)	83	72.82 ± 38.55	70.09 ± 40.49	-2.73 ± 33.14	0.46
TT/E2 ratio	83	41.09 ± 29.77	45.86 ± 30.02	4.77 ± 21.83	0.05
Body composition					
BMI (kg/m^2)	83	27.23 ± 4.62	27.34 ± 5.13	0.1 ± 1.84	0.61
ALM (kg)	83	17.69 ± 2.55	17.65 ± 2.53	-0.04 ± 1.48	0.80
Total body fat (%)	83	39.87 ± 6.8	39.47 ± 7.98	-0.4 ± 3.55	0.31
Relative appendicular lean mass (kg/kg total fat)	83	0.65 ± 0.22	0.69 ± 0.32	0.04 ± 0.19	0.07
Thigh muscle CSA (cm^2)	61	91.87 ± 19.07	87.67 ± 19.89	-4.19 ± 6.13	**<0.01***
Thigh subcutaneous fat area (cm^2)	61	113.14 ± 40.96	111.27 ± 43.27	-1.87 ± 19.44	0.11
Thigh intramuscular fat area (cm^2)	61	11.57 ± 4.11	12.2 ± 4.87	0.54 ± 2.58	0.46
Thigh muscle percentage (%)	61	39.57 ± 6.89	38.87 ± 7.16	-0.7 ± 3.36	0.11
Thigh muscle density (HU)	61	50.56 ± 3.44	48.56 ± 4.28	-2 ± 2.81	**<0.01***
Muscle function					
Hand grip strength (kg)	83	24.39 ± 6.08	24.05 ± 6.76	-0.34 ± 4.8	0.52
Hamstring peak torque (Nm)					
180d/s	83	50.5 ± 15.71	44.44 ± 11.86	-6.06 ± 11.79	**<0.01***
30d/s	83	45.67 ± 14.58	44.57 ± 14.16	-1.11 ± 12.72	0.43
Quadriceps peak torque (Nm)					
180d/s	83	62.62 ± 20.49	57.58 ± 17.09	-5.04 ± 17.09	**<0.01***
30d/s	83	103.66 ± 31.29	95.12 ± 29.53	-8.54 ± 19.8	**<0.01***
Hamstring relative peak torque to LLM (Nm/kg)					
180d/s	83	3.69 ± 0.92	3.28 ± 0.72	-0.41 ± 0.9	**<0.01***
30d/s	83	3.35 ± 0.93	3.28 ± 0.92	-0.06 ± 0.97	0.56
Quadriceps relative peak torque to LLM (Nm/kg)					
180d/s	83	4.58 ± 1.29	4.24 ± 1.05	-0.35 ± 1.42	**0.03***
30d/s	83	7.58 ± 1.91	6.96 ± 1.76	-0.62 ± 1.67	**0.01***
Hamstring relative peak torque to muscle CSA (Nm/cm^2)					
180d/s	61	0.56 ± 0.12	0.51 ± 0.11	-0.05 ± 0.12	**<0.01***
30d/s	61	0.5 ± 0.11	0.52 ± 0.13	0.01 ± 0.13	0.51
Quadriceps relative peak torque to muscle CSA (Nm/cm^2)					
180d/s	61	0.69 ± 0.17	0.68 ± 0.16	-0.01 ± 0.18	0.78
30d/s	61	1.17 ± 0.25	1.11 ± 0.26	-0.05 ± 0.2	**0.05***
Physical function					
400m walk time (s)	50	261.19 ± 40.41	284.62 ± 63.5	23.44 ± 35.62	**<0.01***
Time taken to complete 5 chair stands (s)	74	9.83 ± 2.66	9.47 ± 3.03	-0.36 ± 2.29	0.18
Mobility disability (%(n))	83				
No (SPPB > 9)		97.59 (81)	93.98 (78)	-	-
Yes (SPPB ≤ 9)		2.41 (2)	6.02 (5)	-	-
Falls in last 12 months (% (n))	83				-
No falls		72.29 (60)	69.88 (58)	-	-
Falls		27.71 (23)	30.12 (25)	-	-

Physical function measures were only assessed in females ≥ 50 years old. *p*-value >0.05 indicates a statistically significant difference between baseline and follow up. ALM: appendicular lean mass; BioT: bioavailable testosterone; BMI: body mass index; CSA: cross-sectional area; E2: oestradiol; FEI: free oestradiol index; HU: Hounsfield unit; LLM: leg lean mass; SHBG: sex hormone-binding globulin; SPPB: short physical performance battery; TT: total testosterone; TT/E2: ratio of total testosterone to oestradiol. Data presented as mean ± SD. * Indicates a significant change from baseline to follow-up (*p*-value < 0.05).

Figures

Fig 1 A) Adjusted cross-sectional associations between serum oestradiol (E2) and free oestrogen index (FEI) and muscle outcomes. B) Adjusted longitudinal associations between the change in serum E2 and FEI and the change in muscle outcomes across 4-6 years in females ≥ 50 years old. Multiple regression models were adjusted for age, ethnicity, height, ACCI, and total physical activity levels in the past two weeks. Physical function measures were only assessed in females ≥ 50 years old. Relative hamstring and quadriceps peak torque (Nm/kg) have been normalised to leg lean mass (kg). ALM: appendicular lean mass; CSA: cross-sectional area; HU: Hounsfield unit. Data presented as beta coefficient (β) and 95% confidence interval. * Indicates a significant association (*p*-value < 0.05)

Fig 2 A) Adjusted cross-sectional associations between total serum testosterone (TT) and bioavailable testosterone (BioT) and muscle outcomes. B) Adjusted longitudinal associations between the change in TT and BioT and the change in muscle outcomes across 4-6 years in females ≥ 50 years old. Multiple regression models were adjusted for age, ethnicity, height, ACCI, and total physical activity levels in the past two weeks. Physical function measures were only assessed in females ≥ 50 years old. Relative hamstring and quadriceps peak torque (Nm/kg) have been normalised to leg lean mass (kg). ALM: appendicular lean mass; CSA: cross-sectional area; HU: Hounsfield unit. Data presented as beta coefficient (β) and 95% confidence interval. * Indicates a significant association (*p*-value < 0.05)

Fig 3 A) Adjusted cross-sectional associations between the total testosterone to total oestradiol (TT/E2) ratio and muscle outcomes. B) Adjusted longitudinal associations between the change in serum TT/E2 ratio and the change in muscle outcomes across 4-6 years in females ≥ 50 years old. Multiple regression models were adjusted for age, ethnicity, height, ACCI, and total physical activity levels in the past two weeks. Physical function measures were only assessed in females ≥ 50 years old. Relative hamstring and quadriceps peak torque (Nm/kg) have been normalised to leg lean mass (kg). ALM: appendicular lean mass; CSA: cross-sectional area; HU: Hounsfield unit. Data presented as beta coefficient (β) and 95% confidence interval. * Indicates a significant association (p-value < 0.05)