Title: Evaluating the performance and potential bias of predictive models for detection of transthyretin cardiac amyloidosis

Short Title: Comparing models for ATTR-CM detection

Authors:

Jonathan Hourmozdi, MD, MSAI, MA[1,2] Nicholas Easton, MSAI,[1,2] Simon Benigeri, MSAI,[1,2] James D. Thomas, MD,[1,2] Akhil Narang, MD,[1,2] David Ouyang, MD,[6] Grant Duffy, BS,[6] Ross Upton, PhD,[7] Will Hawkes, PhD,[7] Ashley Akerman, PhD,[7] Ike Okwuosa, MD,[1] Adrienne Kline, MD, PhD[2,3] Abel N. Kho, MD,[4,5] Yuan Luo, PhD,[4] Sanjiv J. Shah, MD,[1,2] Faraz S. Ahmad, MD, MS[1,2,4]

[1] Division of Cardiology, Department of Medicine, Northwestern University Feinberg School of Medicine, Chicago, Illinois

[2] Bluhm Cardiovascular Institute Center for AI, Northwestern Medicine, Chicago, Illinois

[3] Division of Cardiac Surgery, Department of Surgery, Northwestern University Feinberg School of Medicine, Chicago, Illinois

[4] Division of Statistics and Informatics, Department of Preventive Medicine, Northwestern University Feinberg School of Medicine, Chicago, Illinois

[5] Division of General Internal Medicine, Department of Medicine, Northwestern University Feinberg School of Medicine, Chicago, Illinois

[6] Department of Cardiology, Cedars-Sinai Medical Center, Los Angeles, California

[7] Ultromics Limited, Oxford, UK

JDT has received consulting fees from Abbott, EchoIQ, Eko Health, Caption Health, GE Healthcare. AN has received honoraria from Edwards, Bristol Myers Squibb, and Abbott. RU, WH, and AA are employed by Ultromics Ltd. IO has received honoraria from AstraZeneca. SJS was supported by research grants from the National Institutes of Health (U54 HL160273, R01 HL140731, R01 HL149423), American Heart Association (24SFRNPCN1291224), AstraZeneca, Boston Scientific, Corvia, Pfizer, and Tempus; and has received consulting fees from 35Pharma, Abbott, Alleviant, AstraZeneca, Amgen, Aria CV, Axon Therapies, BaroPace, Bayer, Boehringer-Ingelheim, Boston Scientific, BridgeBio, Bristol Myers Squibb, Corvia, Cyclerion, Cytokinetics, Edwards Lifesciences, Eidos, eMyosound, Imara, Impulse Dynamics, Intellia, Ionis, Lilly, Merck, MyoKardia, Novartis, Novo Nordisk, Pfizer, Prothena, ReCor, Regeneron, Rivus, SalubriusBio, Sardocor, Shifamed, Tectonic, Tenax, Tenaya, Ulink, and Ultromics. The remaining authors have nothing to disclose.

Funding: This work was support by the American Heart Association (AHA number 856917).

Corresponding Author:

Faraz S. Ahmad, MD MS

676 North Saint Clair Street, Suite 600

Chicago, IL 60611

Office: 312-695-4965; Fax: 312-926-8620

Email: faraz.ahmad@northwestern.edu; X handle: @FarazAhmadMD

Word Count: 4,378 /5,000 (Abstract: 245/250)

Tweet: In a comparison of 4 algorithms for detecting ATTR-CM, deep learning echo models outperformed others—including claims-based models and regression-based risk scoring —with low risk of harms due to racial bias. #AI #CardioTwitter #MedTwitter @Jhourmozdi @FarazAhmadMD @NMCardioVasc

Abstract

Background:

Delays in the diagnosis of transthyretin amyloid cardiomyopathy (ATTR-CM) contribute to the significant morbidity of the condition, especially in the era of disease-modifying therapies. Screening for ATTR-CM with AI and other algorithms may improve timely diagnosis, but these algorithms have not been directly compared.

Objectives:

The aim of this study was to compare the performance of four algorithms for ATTR-CM detection in a heart failure population and assess the risk for harms due to model bias.

Methods:

We identified patients in an integrated health system from 2010-2022 with ATTR-CM and age- and sex-matched them to controls with heart failure to target 5% prevalence. We compared the performance of a claims-based random forest model (Huda et al. model), a regression-based score (Mayo ATTR-CM), and two deep learning echo models (EchoNet-LVH and EchoGo® Amyloidosis). We evaluated for bias using standard fairness metrics.

Results:

The analytical cohort included 176 confirmed cases of ATTR-CM and 3192 control patients with 79.2% self-identified as White and 9.0% as Black. The Huda et al. model performed poorly (AUC 0.49). Both deep learning echo models had a higher AUC when compared to the Mayo ATTR-CM Score (EchoNet-LVH 0.88; EchoGo Amyloidosis 0.92; Mayo ATTR-CM Score

0.79; DeLong P<0.001 for both). Bias auditing met fairness criteria for *equal opportunity* among patients who identified as Black.

Conclusions:

Deep learning, echo-based models to detect ATTR-CM demonstrated best overall discrimination when compared to two other models in external validation with low risk of harms due to racial bias.

Keywords: amyloidosis, artificial intelligence, machine learning, heart failure, health care disparities

Abbreviation List:

AI = Artificial intelligence:

ATTR-CM = transthyretin amyloid cardiomyopathy

HF = heart failure

NMEDW = Northwestern Enterprise Data Warehouse

PYP = Pyrophosphate

TTE = transthoracic echocardiogram

AUC = area under the curve

AP = average precision

LVH = left ventricular hypertrophy

LVEF = left ventricular ejection fraction

Introduction

Using artificial intelligence (AI) and other machine-learning algorithms for the early detection or correcting misdiagnosis of rare diseases represents a major area of research and development.[1] This area of research is particularly important for patients with transthyretin amyloid cardiomyopathy (ATTR-CM), a highly morbid condition that is increasingly recognized as an important cause of heart failure (HF) and disproportionately impacts Black communities.[2] While the true prevalence of cardiac amyloidosis is unknown, recent data suggest that it is much more common than previously thought, comprising up to 17% of HF with preserved ejection fraction cases.[3] Despite early clues to diagnosis, delays in the diagnosis of ATTR-CM are extremely common, with 42% of cases diagnosed over 4 years after the development of cardiac symptoms and a median time from first hospitalization to diagnosis of 2 to 3 years.[4] Even prior to the development of clinical HF symptoms, subclinical changes in conduction, structure, and function often can be observed in cardiac testing, including routine electrocardiography and echocardiography.[5–8]

The emergence of transthyretin stabilizers, which can halt disease progression and dramatically improve morbidity for patients with ATTR-CM, as well as a pipeline of other disease-modifying therapies, have galvanized efforts to improve early and accurate diagnosis of ATTR-CM, including the use of AI and other algorithms for screening.[9] In order to better understand the potential implications of implementing algorithmic screening technologies for ATTR-CM in large health systems, we studied the performance of four models for the identification of cardiac amyloidosis in a retrospective cohort of patients treated across an integrated health system with 11 hospitals and over 200 sites. In addition to overall diagnostic performance, we assessed the

performance of each model with respect to selected fairness metrics to identify any potential risks to currently disadvantaged groups owing to possible algorithmic bias.

Methods

The data for this study will be made available by the corresponding author upon reasonable request and a data use agreement. The Northwestern University Institutional Review Board approved this study. This paper adheres to the TRIPOD-AI guidance for reporting of clinical prediction models (**See Supplemental Checklist**).[10]

Data Source and Participants

To identify patients with ATTR-CM, we first searched the Northwestern Enterprise Data Warehouse (NMEDW) for patients with at least 1 ICD code for amyloidosis, a pyrophosphate (PYP) scan, or medication order for Tafamidis from 1/1/2010 to 12/31/2022. For these patients, we performed a review of their records to confirm diagnosis of ATTR-CM, defined as either positive endomyocardial biopsy or PYP scan with a final reading of "strongly suggestive" and absence of serum evidence of amyloid light chain (AL) amyloidosis or negative bone marrow biopsy result. For a subset of patients with potential ATTR-CM based on our screening criteria but without available definitive testing in the NMEDW, we used data from the Northwestern Medicine Cardiac Amyloidosis Program registry (August 2018 to December 2022), which included the results of endomyocardial biopsies or PYP scans performed outside of the Northwestern Medicine system.

We then created a matched cohort of HF controls with at least one encounter ICD code for HF over the study period. We excluded potential cases of amyloidosis from this matched cohort, defined as at least one ICD code for amyloidosis and did not have definitive negative testing available, specifically a negative endomyocardial biopsy result or a PYP scan with a final reading of "Not Suggestive." All cases and controls were required to have a transthoracic echocardiogram (TTE) up to six months before or any time after definitive diagnosis by positive endomyocardial biopsy or PYP scan. The NMEDW query required that these TTEs also reported measurements for left ventricular ejection fraction (LVEF), posterior wall thickness, and relative wall thickness in a format available for automated data extraction and inclusion in the Mayo ATTR-CM Score model. If controls had multiple TTEs, the one nearest to the first HF diagnosis code was used for our model evaluation. We matched controls to cases by age, sex, and year that TTE was obtained using the PsmPy package in Python.[11] Logistic regression was performed to generate propensity logits using PsmPy's resampling method to account for class imbalance. Cases were then matched to controls in a 19:1 ratio using the nearest neighbor method without caliper, to target a prevalence of ATTR-CM of 5%, which was selected to mirror estimates in the HF population and reflect its potential clinical use as screening tool in a broad population.[11] We then excluded patients in which any of the model's predictions were not available due to missing data, image quality or other factors.

Cardiac Amyloidosis Risk Models

We compared the performance of four risk models with different inputs and model architectures (**Table 1**). The Huda et al. model[12] was developed to predict risk of wild-type ATTR-CM using databases of claims data and was validated on claims data and EHR data from patients with ICD

codes for amyloidosis (including for codes beyond the specific code for wild-type ATTR). The Huda et al. model uses tabular, ICD code data in a random forest model. The Mayo ATTR-CM Score was developed by Davies et al.[13] and uses tabular data that includes demographics (age, sex), history of hypertension (based on ICD codes for this analysis), and TTE measurements (left ventricular ejection fraction, wall thickness). The Mayo ATTR-CM Score is an integer-based score created from a linear regression model. Duffy et al.[14] developed the EchoNet-LVH model, which is a computer vision model that uses a convolutional neural network architecture. EchoNet-LVH generates a prediction for likelihood of cardiac amyloidosis (any type) from an echocardiogram study synthesizing information about wall thickness from the parasternal long axis view and additional information from the apical four chamber view. Lastly, EchoGo® Amyloidosis is another deep-learning classifier developed by Ultromics which was trained to predict a diagnosis of cardiac amyloidosis from a single apical 4 chamber view video clip and received FDA clearance for commercial use in November 2024.[15] EchoGo Amyloidosis produces a probability score (0-1) and an uncertainty score. If EchoGo Amyloidosis generated a probability score for multiple apical four chamber clips within an echo study, the least uncertain score was used.

Statistical Analysis

We used standard descriptive statistics to summarize baseline variables with categorical variables reported as counts and proportions and continuous variables reported as mean and standard deviation (SD). Comparisons of demographic and baseline clinical features between cases and controls were performed using Welch's T-test for continuous variables and Chi-squared testing for categorical variables. For each model, we report model discrimination with

receiver operating characteristics reported as area under the curve (AUC). We also report the area under the precision-recall curve as average precision (AP), a measure which may be more informative in datasets with class imbalance.[16] Where reported, 95% confidence intervals were calculated using the bootstrapping method. Pairwise comparisons of model discrimination with AUC between the Mayo ATTR-CM score were performed using the DeLong test.[17] In addition, we report F1 score, accuracy, sensitivity, selectivity, positive predictive value, negative predictive value, and false negative rate for all models based on published thresholds or recommendations from the model developers as shown in Table 1.

Sensitivity and Fairness and Bias Analyses

We conducted several sensitivity analyses to further examine the algorithmic performance under various clinical subpopulations and deployment decisions. For the clinical subgroup analyses, we first examined subgroups of patients with at least moderate left ventricular hypertrophy defined by posterior wall thickness measurements[18] and those with reduced LVEF. To evaluate the tradeoff between sensitivity and specificity in the two AI echo models, we examined differences in performance at different thresholds. Lastly, we evaluated the performance of the two AI echo models in a larger sample in which we imputed the value of 0 for patients in whom the AI echo models were unable to generate a score.

In conducting our fairness and bias analysis, we compared three metrics that address fairness across a protected attribute. A protected attribute refers to a feature or characteristic of a patient that should not be used as the basis of decision-making and should be generally protected from discrimination. These attributes can be defined according to anti-discrimination law or in accordance with institutional values.[19] We chose the protected attribute of self-identified race or

ethnicity given the previously described existing disparities in diagnosis of ATTR-CM for racial and ethnic minorities in the United States. From previously published fairness criteria,[19] we selected the following three to report in our primary analysis: *Demographic parity*, defined as the ratio of the proportion of patients predicted positive by the model between groups, *Predictive parity*, defined as the ratio of positive predictive value (or precision) of the model output between groups, and *Equal opportunity*, defined as the ratio of false negative error rate between groups. We chose to prioritize the equal opportunity fairness criteria in weighing the risk of harms due to the current imbalance in diagnostic error owing to under- or missed diagnosis in US Black individuals. All analyses were performed using the *Aequitas* package in Python 3.12.[20] The default fairness threshold of 0.2 was used to determine whether significant unfairness was present for each metric based on the "80% rule" of disparate impact.[20]

Results

Demographic Characteristics

240 cases of confirmed ATTR-CM were identified by NMEDW query and matched to 4560 patients for a total of 4800 patients. There was a total of 1,432 patients (42 cases) in which either EchoNet-LVH, EchoGo Amyloidosis, or both did not generate predictions due to factors including insufficient clip length, low image quality, insufficient videos by view classification, low predictive confidence, or uncertainty. The exclusion of these patients resulted in a final analytic cohort size of 3368 patients containing 176 confirmed cases of ATTR-CM matched to 3192 controls (**Figure 1**). In this cohort, mean age was 78.8 years (SD 9.5 years), 20.9% were female, 79.1% self-identified as White, and 9.0% self-identified as Black. Complete demographic and baseline clinical characteristics for case and control cohorts are shown in

Table 2. As compared to age and sex-matched controls with HF, the cohort of patients with

ATTR-CM had fewer patients who identified as White (58.5% vs. 80.3%) and more left

ventricular hypertrophy (LVH) as quantified by their baseline TTE (mean posterior wall

thickness of 15.6 mm vs. 11.1 mm). The demographics and clinical characteristics observed in

the subgroup of patients excluded due to missing or uncertain predictions are provided in

Supplemental Tables 1 and 2.

Overall Model Performance and Comparison

In terms of overall performance, the Huda et al. model failed to achieve meaningful

discriminatory performance with an overall AUC of 0.49 (95% CI, 0.44 to 0.54) and had

relatively poor performance on all other metrics. Compared to the ATTR-CM Score (AUC of

0.79, 95% CI, 0.76 to 0.83), EchoNet-LVH (AUC 0.88, 95% CI, 0.85 to 0.91) and EchoGo

Amyloidosis (AUC 0.92, 95% CI, 0.89 to 0.95) both outperformed the simpler model (both

DeLong P<0.001). A complete comparison of performance metrics for all four models can be

found in **Table 3**.

From the initial analysis of the primary model performance metrics, it was apparent that the

performance of the Huda et al. model was poor compared to the other two models. The AUC of

0.49 in the analytic cohort was much lower than the model's performance in their initial external

validation cohorts.[12] To determine whether insufficient number of diagnosis codes was

contributing to this decreased performance, we reassessed the Huda et al. model on the subset of

patients in the overall analytic cohort for whom at least 50 encounters were available (106 cases

and 774 controls). There was no significant improvement in the overall model performance

(AUC 0.56, 95% CI, 0.50 to 0.62, De long P=0.084, **Supplemental Table 3**). Because of this,

we removed the Huda et al. model from further analyses and focused on the other three models.

Clinical Subgroup and Additional Sensitivity Analyses

In the subgroup of patients with at least moderate left ventricular hypertrophy (n=504), the

prevalence of ATTR-CM was higher than in the overall cohort (23.8% vs 5.2%). Compared to

the overall cohort, the AI echo models had modest differences in AUC, sensitivity, and

specificity as demonstrated in **Table 4.** The lower AUC for the Mayo ATTR-CM score with non-

overlapping confidence interval suggests worse performance in this subgroup. In the subgroup of

patients with a reduced LVEF of lower than 40% (n=713, prevalence = 4.8%), overall similar

performance was again noted for the two AI echo models. The higher AUC for the Mayo ATTR-

CM score with non-overlapping confidence interval suggests an improvement in performance in

this subgroup (**Table 5**). An analysis of the two AI echo models at different thresholds highlights

the tradeoffs between sensitivity and specificity (**Supplemental Table 4**). The two AI echo

models have overall similar performance when set at the same threshold, with some tradeoff

between sensitivity and specificity. Lastly, we conducted a sensitivity analysis in those patients

in whom the computer vision models were unable to generate predictions in which we imputed

missing prediction as a negative (score=0) and found, as expected, both models had lower

sensitivity, false negative rate, and AUC (**Supplemental Table 5**).

Fairness and Bias Audit

Prevalence of ATTR-CM was highest in the group of patients who self-identified as Black (19%

compared to 3-5% in the other racial and ethnic groups). With respect to the protected attribute

of race, all models met the fairness criteria for equal opportunity in patients who self-identified their race as Black **(Figure 3)**. A summary of model performance with respect to all chosen fairness criteria is provided in **Table 6**. A complete comparison of all absolute metrics by protected attribute can be found in **Supplemental Figures 1 to 3**.

Discussion

In this study, we evaluated the potential screening performance of four available models to detect ATTR-CM in a large, integrated health system. We found that the Huda et al. model performed poorly compared to their previously published validation data and the other models.[12] The Mayo ATTR-CM score had lower sensitivity and higher specificity than previously reported, likely reflecting the differences in the cohort in this study and the derivation and validation cohorts for the score.[13,21] The AI echo models that we evaluated demonstrated similar performance in our validation cohort as compared to their initially reported results, including across a key subgroup as shown in the bias audit.[14,15] To our knowledge, this is the first comparison of multiple algorithmic approaches to screening for cardiac amyloidosis using different data inputs (diagnosis codes, structured clinical data, and echo images) and models (regression, random forest, and convolutional neural networks).

In addition to external validation, our study allowed us to directly compare a regression-based, simple score against more complex, computer vision models in our patient population and provides realistic estimates for predictive values beyond AUC alone. The deep learning vision-based models including EchoNet-LVH and EchoGo Amyloidosis, both performed better in terms of positive predictive value compared to Mayo ATTR-CM Score (0.66 for EchoNet-LVH vs.

0.34 for EchoGo Amyloidosis vs. 0.20 for Mayo ATTR-CM Score). Although the Mayo ATTR-CM Score include selected echo measurements, this analysis illustrates that AI echo models are likely integrating multiple features for cardiac amyloidosis that are likely more granular that our standard measurements. This has important implications for implementation as a reduction in unnecessary downstream testing due to false positive predictions could help to offset the higher upfront costs of adopting and integrating a more complex screening model. However, the optimal threshold for each model that balances between sensitivity and specificity remains uncertain, as evident by the different choice in threshold by the two AI echo model developers. Identifying the optimal threshold will require additional, prospective evaluation and expert consensus on the appropriate population for screening, approaches for implementation, and targets for sensitivity, positive predictive value, and other clinically informative metrics.

Our study also highlighted the technical challenge of AI model portability. Both computer vision- based models did not provide predictions in approximately 15-20% of patients, leading to the exclusion of ATTR-CM cases in our cohort. Thus, the sensitivity and other related performance metrics for the algorithm may be lower in real-world practice than estimated by our primary analysis as shown in the Supplement. Similarly, the data required to run the Mayo ATTR-CM Score likely does not exist in a structured form in many EHR data repositories, thus limiting its ability to be run at scale in a health system. The poor performance of the Huda et al. model underscores the potential limitations of training models using diagnosis codes alone, as demonstrated in another cardiac amyloidosis study.[22]

Risk of perpetuating bias

With any machine learning model, there is a risk that they can learn to exploit patterns in the training data that represent or perpetuate bias, including systemic and structural racism. These risks are exacerbated when there is underrepresentation of minority patient groups in training cohorts. In the United States, patients who identify as Black disproportionately bear the burden of missed and delayed diagnosis in ATTR-CM, leading to worse outcomes.[2] This is despite a higher prevalence of ATTR-CM in many Black communities in the US; in our cohort, the relative prevalence of ATTR-CM was about 5-fold higher for Black patients compared to all other racial groups. Such context is instrumental in interpreting fairness metrics related to any screening AI model, as there is a risk not only of differential predictive performance, but also of a differential distribution of error rate. Our findings support that there is a low risk of harm due to algorithmic bias from adopting the use of any of these machine learning models for screening in our integrated health system. This study also provides an example of how to apply the recently published TRIPOD-AI reporting guidance and bias and fairness metrics to external validation studies.

Study limitations

Importantly, a major limitation of our study is that we utilize retrospective data from a single large health system to perform our analyses. The prevalence in our cohort was artificially set to 5% based on realistic estimates of ATTR-CM prevalence in our HF patient population. Absolute measurements of predictive value vary depending on prevalence and may not generalize to all healthcare settings. However, comparisons of relative predictive value between models are more useful when considering realistic estimates of population prevalence.

Secondly, a limitation of our bias audit is that self-identified race and ethnicity represent a social construct. Concepts of race and ethnicity vary with respect to time and geography, and the results of our bias audit, while likely generalizable to our institution's population of patients with HF, should not be taken to represent a blanket determination of fairness with respect to race in all settings. Instead, we encourage institutions and constituents to partner with health equity researchers and consider similar methodology when evaluating models for potential deployment across a health system. Lastly, the study population differed from the validation population used by the different models by factors such as age cutoffs, clinical factors, and control selection. However, our study sample reflects the potential group in which a health system may elect to deploy a population screening strategy for cardiac amyloidosis and thus is clinically informative.

Conclusion

Machine learning techniques have shown promise when applied to imaging and EHR data to enhance the early detection or correct the misdiagnosis of diseases, such as cardiac amyloidosis. The use of such AI-assisted diagnostic tools in ATTR-CM could improve racial and ethnic disparities through the early identification and uptake of novel therapies in these patients. However, despite pioneering research and significant resources dedicated to this task, few algorithms have had a meaningful impact on the care of these patients in practice. The reasons for this are multi-factorial and include the lack of prospective, high quality evidence and consensus recommendations to support at scale implementation, barriers related to costs, infrastructure, and required personnel, and evolving regulatory guidance over the deployment and governance of AI and machine learning algorithms. Future work on algorithmic approaches for screening for cardiac amyloidosis should include evaluating the performance of using

multiple models in serial or fusion approaches. However, ultimately, to evaluate the value of these algorithms, we need randomized, effectiveness-implementation hybrid trials that leverage health equity conceptual models and frameworks to study the performance of the algorithms and the implementation strategies for successful deployment across diverse populations.[23–25]

Perspectives

Competency in medical knowledge: Algorithmic approaches to screening for ATTR-CM vary in their performance in single health system with AI echo models having the best overall performance. Both simple, regression-based score or AI echo models performed similarly in a fairness and bias analysis in patients who identify as Black.

Translational Outlook 1: Understanding the optimal thresholds for different algorithmic approaches and evaluating the impact of prospective deployment remain important questions.

Translational Outlook 2: Evaluation of multiple algorithmic screening approaches, either in serial or in a fusion approach, may lead to higher performance and requires further evaluation.

References:

1. Elias P, Jain SS, Poterucha T, et al. Artificial intelligence for cardiovascular care-part 1: Advances: JACC review topic of the week. *J Am Coll Cardiol*. 2024;83(24):2472-2486.

2. Spencer-Bonilla G, Njoroge JN, Pearson K, Witteles RM, Aras MA, Alexander KM. Racial and ethnic disparities in transthyretin cardiac amyloidosis. *Curr Cardiovasc Risk Rep*. 2021;15(6). doi:10.1007/s12170-021-00670-y

3. Kittleson MM, Maurer MS, Ambardekar AV, et al. Cardiac amyloidosis: Evolving diagnosis and management: A scientific statement from the American heart association. *Circulation*. 2020;142(1):e7-e22.

4. Lane T, Fontana M, Martinez-Naharro A, et al. Natural history, quality of life, and outcome in cardiac transthyretin amyloidosis. *Circulation*. 2019;140(1):16-26.

5. Bhogal S, Ladia V, Sitwala P, et al. Cardiac amyloidosis: An updated review with emphasis on diagnosis and future directions. *Curr Probl Cardiol*. 2018;43(1):10-34.

6. Maurer MS, Bokhari S, Damy T, et al. Expert consensus recommendations for the suspicion and diagnosis of transthyretin cardiac amyloidosis. *Circ Heart Fail*. 2019;12(9):e006075.

7. Phelan D, Collier P, Thavendiranathan P, et al. Relative apical sparing of longitudinal strain using two-dimensional speckle-tracking echocardiography is both sensitive and specific for the diagnosis of cardiac amyloidosis. *Heart*. 2012;98(19):1442-1448.

8. Sinha A, Zheng Y, Nannini D, et al. Association of the V122I transthyretin amyloidosis genetic variant with cardiac structure and function in middle-aged Black adults: Coronary

Artery Risk Development in Young Adults (CARDIA) study: Coronary artery risk development in young adults (CARDIA) study. *JAMA Cardiol*. 2021;6(6):718-722.

9. Donnelly JP, Hanna M. Cardiac amyloidosis: An update on diagnosis and treatment. *Cleve Clin J Med*. 2017;84(12 Suppl 3):12-26.

10. Collins GS, Moons KGM, Dhiman P, et al. TRIPOD+AI statement: updated guidance for reporting clinical prediction models that use regression or machine learning methods. *BMJ*. 2024;385:e078378.

11. Kline A, Luo Y. PsmPy: A package for retrospective cohort matching in python. *Annu Int Conf IEEE Eng Med Biol Soc*. 2022;2022:1354-1357.

12. Huda A, Castaño A, Niyogi A, et al. A machine learning model for identifying patients at risk for wild-type transthyretin amyloid cardiomyopathy. *Nat Commun*. 2021;12(1):2725.

13. Davies DR, Redfield MM, Scott CG, et al. A simple score to identify increased risk of transthyretin amyloid cardiomyopathy in heart failure with preserved ejection fraction. *JAMA Cardiol*. 2022;7(10):1036-1044.

14. Duffy G, Cheng PP, Yuan N, et al. High-throughput precision phenotyping of left ventricular hypertrophy with cardiovascular deep learning. *JAMA Cardiol*. 2022;7(4):386-395.

15. U.S. Food & Drug Administration. Re: K2340860. 2024. Available at: https://www.accessdata.fda.gov/cdrh_docs/pdf24/K240860.pdf. Accessed 16 Mar 2025.

16. Ozenne B, Subtil F, Maucort-Boulch D. The precision--recall curve overcame the optimism of the receiver operating characteristic curve in rare diseases. *J Clin Epidemiol*. 2015;68(8):855-859.

17. DeLong ER, DeLong DM, Clarke-Pearson DL. Comparing the areas under two or more correlated receiver operating characteristic curves: a nonparametric approach. *Biometrics*. 1988;44(3):837-845.

18. Marwick TH, Gillebert TC, Aurigemma G, et al. Recommendations on the use of echocardiography in adult hypertension: A report from the European association of cardiovascular imaging (EACVI) and the American society of echocardiography (ASE). *J Am Soc Echocardiogr*. 2015;28(7):727-754.

19. Chen RJ, Wang JJ, Williamson DFK, et al. Algorithmic fairness in artificial intelligence for medicine and healthcare. *Nat Biomed Eng*. 2023;7(6):719-742.

20. Saleiro P, Kuester B, Hinkson L, et al. Aequitas: A bias and fairness audit toolkit. *arXiv [csLG]*. Published online November 13, 2018. http://arxiv.org/abs/1811.05577

21. Bonfioli GB, Tomasoni D, Vergaro G, et al. The Mayo ATTR-CM score versus other diagnostic scores and cardiac biomarkers in patients with suspected cardiac amyloidosis. *Eur J Heart Fail*. Published online September 9, 2024. doi:10.1002/ejhf.3455

22. Vrudhula A, Stern L, Cheng PC, et al. Impact of case and control selection on training artificial intelligence screening of cardiac amyloidosis. *JACC Adv*. 2024;3(9):100998.

23. Curran GM, Bauer M, Mittman B, Pyne JM, Stetler C. Effectiveness-implementation hybrid designs: combining elements of clinical effectiveness and implementation research to enhance public health impact. *Med Care*. 2012;50(3):217-226.

24. Woodward EN, Matthieu MM, Uchendu US, Rogal S, Kirchner JE. The health equity implementation framework: proposal and preliminary study of hepatitis C virus treatment. *Implement Sci*. 2019;14(1):26.

25. Shelton RC, Chambers DA, Glasgow RE. An extension of RE-AIM to enhance sustainability: Addressing dynamic context and promoting health equity over time. *Front Public Health*. 2020;8:134.

Figure Legends

Figure 1: Flow diagram for analytic cohort creation. This figure shows the creation of the analytical dataset. NMEDW denotes Northwestern Medicine Enterprise Database Warehouse; ICD, international classification of diseases; TTE, transthoracic echocardiogram; PYP, pyrophosphate scan.

Figure 2: Comparison of model performance. Panel A shows receiver operator characteristic curves and Panel B shows the precision-recall curves (bottom) with bootstrap confidence intervals.

Figure 3: Parity metric treemaps for Mayo ATTR-CM Score model (A), EchoNet-LVH (B), and EchoGo Amyloidosis (C). Selected fairness metrics are expressed as a ratio of protected group: reference group. *Demographic parity* refers to positive prediction rate, *predictive parity* refers to positive predictive value (precision), and *equal opportunity* refers to false negative rate. Treemap squares are sized according to size of group in the cohort and colored red if results suggest disparity according to the "80% rule" of disparate impact.*denotes statistical significance of finding based on unadjusted alpha < 0.05.*

Central Illustration: Evaluating the Performance and Potential Bias of Predictive Models for the Detection of Transthyretin Cardiac Amyloidosis. In a diverse cohort of 3416 patients with heart failure including 177 patients with transthyretin amyloid cardiomyopathy (ATTR-CM), deep-learning echocardiography models were best able to detect ATTR-CM compared to

two other publicly available models and did not demonstrate any evidence of racial bias in a fairness audit. HF denotes heart failure; AUC, area under the receiver-operator characteristic curve; AP, average precision.

Table 1: Description of the evaluated machine-learning models for detection of cardiac amyloidosis.

Author (year)	Model Name	Model Architecture	Input Features	Outputs	Positive Case Threshold	Reported AUC[a]
Huda et al. (2021)	N/A	Random Forest	Medical Claims	Probability (0-1)	> 0.5	0.80
Davies et al. (2022)	Mayo ATTR-CM Score	Logistic Regression	Clinical Variables	Risk score (0-10)	≥ 6	0.84
Duffy et al. (2022)	EchoNet-LVH	ResNet3D	Echo Images	Probability (0-1)	≥ 0.8	0.83
Publication Pending; Validation data available on FDA website[15]	EchoGo Amyloidosis	Convolutional Neural Network	Echo Images	Probability (0-1); Label of Positive, Negative, or Uncertain because on probability and uncertainty scores	≥ 0.06	Not reported

[a]Values are from initial published performance in external validation cohorts.

AUC denotes area under the receiver-operating characteristic curve

Table 2: Demographic characteristics of the validation cohort.

	ATTR-CM (n=176)	HF Controls (n=3192)	P-value	Overall (n=3368)
Age, mean (SD)	78.6 (9.0)	78.8 (9.5)	0.805	78.8 (9.5)
Sex, n (%)			0.466	
Female	33 (18.8%)	670 (21.0%)		703 (20.9)
Male	143 (81.2%)	2522 (79.0%)		2665 (79.1)
Race/Ethnicity, n (%)			< 0.0001	
Black	59 (33.5)	243 (7.6)		302 (9.0)
Hispanic	< 10[a]	127 (4.0%)		-[a]
White	103 (58.5)	2563 (80.3)		2666 (79.2)
Other/Unknown	< 10[a]	259 (8.1%)		-[a]
LVEF, n (%)			0.142	
< 40%	34 (19.3%)	679 (21.2%)		713 (21.2%)
40% to < 50%	35 (19.9%)	451 (14.1%)		486 (14.4%)
≥ 50%	107 (60.8%)	2062 (64.6%)		2169 (64.4%)
LVH by Posterior Wall Thickness[b]			< 0.0001	
Mean, (SD)	15.6 (4.0)	11.1 (2.2)		11.3 (2.5)
Normal/Mild, n (%)	56 (31.8%)	2800 (87.8%)		2856 (84.8%)
Moderate, n (%)	57 (32.4%)	333 (10.5%)		390 (11.6%)
Severe, n (%)	63 (35.8%)	51 (1.6%)		114 (3.4%)
LVH by Septal Wall Thickness[b]			< 0.0001	
Mean, (SD)	15.9 (3.8)	11.7 (2.9)		12.0 (3.1)
Normal/Mild, n (%)	42 (23.9%)	2011 (63.0%)		2045 (60.7%)
Moderate, n (%)	48 (32.4%)	380 (15.3%)		428 (16.3%)
Severe, n (%)	57 (38.5%)	91 (3.7%)		148 (5.6%)
Relative Wall Thickness			< 0.0001	
Mean (SD)	0.78 (0.47)	0.47 (0.15)		0.49 (0.19)
> 0.57 (%)	137 (77.8%)	628 (19.7%)		765 (22.7%)
Hypertension, n (%)	144 (81.8%)	2560 (80.2%)	0.590	2704 (80.3%)

[a] Cell sizes not reported due to small cell size
[b] Sex-specific cutoffs were used from consensus recommendations[18]
ATTR-CM, transthyretin amyloid cardiomyopathy cases; HF, heart failure; LVEF, left ventricular ejection fraction; LVH, left ventricular hypertrophy; SD, standard deviation; IQR, interquartile range.

Table 3: Overall performance metrics with 95% CI for the four compared models for 176 cases and 3192 controls at specified threshold by model developer.

	F1	Acc	Sens	Spec	PPV	NPV	FNR	AP	AUC
Huda et al.	0.09 [0.07, 0.12]	0.66 [0.64, 0.67]	0.34 [0.26, 0.40]	0.68 [0.66, 0.69]	0.05 [0.04, 0.07]	0.95 [0.94, 0.96]	0.66 [0.60, 0.74]	0.06 [0.05, 0.08]	0.49 [0.44, 0.54]
Mayo ATTR-CM Score	0.31 [0.26, 0.35]	0.85 [0.83, 0.86]	0.65 [0.59, 0.73]	0.86 [0.84, 0.87]	0.20 [0.17, 0.23]	0.98 [0.97, 0.98]	0.35 [0.27, 0.42]	0.16 [0.13, 0.20]	0.79 [0.76, 0.83]
EchoNet-LVH	0.61 [0.55, 0.67]	0.96 [0.96, 0.97]	0.57 [0.50, 0.64]	0.98 [0.98, 0.99]	0.66 [0.58, 0.74]	0.98 [0.97, 0.98]	0.43 [0.35, 0.51]	0.62 [0.55, 0.69]	0.88 [0.84, 0.91]
EchoGo Amyloidosis	0.49 [0.43, 0.53]	0.91 [0.90, 0.92]	0.85 [0.79, 0.90]	0.91 [0.90, 0.92]	0.34 [0.30, 0.39]	0.99 [0.99, 0.99]	0.15 [0.11, 0.21]	0.67 [0.59, 0.74]	0.92 [0.89, 0.95]

CI, confidence interval; F1, F1-score; Acc, accuracy; Sens, sensitivity; Spec, specificity; PPV, positive predictive value; NPV, negative predictive value; FNR, false negative rate; AP, average precision; AUC, area under the receiver-operating characteristic curve.

Table 4: Clinical subgroup analyses for the 504 patients (120 cases, 384 controls) with at least moderate left ventricular hypertrophy by echo (posterior wall thickness ≥1.4 for men, ≥ 1.3 for women).

	Sensitivity [95% CI]	Specificity [95% CI]	AP [95%CI]	AUC [95%CI]
Mayo ATTR-CM Score	0.82 [0.74, 0.88]	0.40 [0.35, 0.44]	0.28 [0.24, 0.34]	0.60 [0.55, 0.65]
EchoNet-LVH	0.66 [0.57, 0.74]	0.96 [0.94, 0.98]	0.81 [0.74, 0.87]	0.91 [0.88, 0.94]
EchoGo Amyloidosis	0.92 [0.87, 0.96]	0.85 [0.82, 0.88]	0.83 [0.76, 0.89]	0.93 [0.90, 0.95]

CI, confidence interval; AP, average precision; AUC, area under the receiver-operating characteristic curve.

Table 5: Clinical subgroup analyses for Mayo ATTR-CM score, EchoNet-LVH, and EchoGo Amyloidosis in 713 patients (34 cases, 679 controls) with left ventricular ejection fraction < 40%.

	Sensitivity [95% CI]	Specificity [95% CI]	AP [95%CI]	AUC [95%CI]
Mayo ATTR-CM Score	0.71 [0.54, 0.86]	0.92 [0.9, 0.94]	0.24 [0.15, 0.36]	0.83 [0.74, 0.90]
EchoNet-LVH	0.59 [0.43, 0.77]	0.98 [0.97, 0.99]	0.65 [0.45, 0.79]	0.88 [0.79, 0.95]
EchoGo Amyloidosis	0.82 [0.69, 0.94]	0.88 [0.85, 0.9]	0.67 [0.49, 0.83]	0.89 [0.79, 0.95]

CI, confidence interval; AP, average precision; AUC, area under the receiver-operating characteristic curve.

Table 6: Summary of model performance across chosen fairness metrics stratified by protected attribute.

	Group	Prev	Sens	Spec	Demographic Parity	Predictive Parity	Equal Opportunity
Mayo ATTR-CM Score	White	0.04	0.64	0.86	ref	ref	ref
	Black	0.20	0.68	0.86	1.55	3.44	0.90[a]
	Hispanic	0.04	0.80	0.85	1.09	1.12	0.56
	Other	0.03	0.56	0.84	1.10	0.69[b]	1.24[b]
EchoNet-LVH	White	0.04	0.56	0.99	ref	ref	ref
	Black	0.20	0.54	0.96	3.85	1.26	1.05[a]
	Hispanic	0.04	0.60	1.00	0.64[b]	1.62	0.92
	Other	0.03	0.78	0.98	1.38	0.87	0.51
EchoGo Amyloidosis	White	0.04	0.83	0.92	ref	ref	ref
	Black	0.20	0.88	0.82	2.91	1.85	0.68[a]
	Hispanic	0.04	0.80	0.91	1.05	0.90	1.14
	Other	0.03	0.89	0.89	1.24	0.75[b]	0.64

Selected fairness metrics are expressed as a ratio of protected group: reference group. *Demographic parity* refers to positive prediction rate, *predictive parity* refers to positive predictive value (precision), and *equalized opportunity* refers to false negative rate.
[a] denotes statistical significance with an unadjusted alpha of <0.05 for parity or disparity.
[b] denotes disparity according to the "80% rule" of disparate impact.

CENTRAL ILLUSTRATION Evaluating the Performance and Potential Bias of Predictive Models for the Detection of Transthyretin Cardiac Amyloidosis

Methods

Retrospective Cohort

176 Confirmed Cases of ATTR-CM
3,192 Matched Controls with HF

Models Tested

Huda et al. Model

Mayo ATTR-CM Score

Echo AI Models

Results

Conclusions

 Deep learning echo-based models to detect ATTR-CM demonstrated best overall performance

 Bias auditing of the top three performing models met fairness criteria for equal opportunity among patients who identified as Black