Hospital *Enterococcus faecium* demonstrates distinct environmental and patient reservoirs: a

genomic point prevalence survey

Nenad Macesic*[1,2,3], Hugh Cottingham[1], Jessica A. Wisniewski[1], Luke V. Blakeway[1], Ravali

Theegala[1], Katherine Pragastis[1], Andrew Stewardson[1,2], Pauline Bass[2], Megan Gritt[2], Stephanie

Spilsbury[2], Denise Del Rosario-Kelly[2], Amanda Dennison[4], Denis W. Spelman[1,4], Adam W.J.

Jenney[1,4], Anton Y. Peleg*[1,3,5]

[1] Department of Infectious Diseases, The Alfred Hospital and School of Translational Medicine,

Monash University, Melbourne, Australia

[2] Infection Prevention & Healthcare Epidemiology, Alfred Health, Melbourne, Australia.

[3] Centre to Impact AMR, Monash University, Clayton, Australia

[4] Microbiology Unit, Alfred Health, Melbourne, Australia.

[5] Infection Program, Monash Biomedicine Discovery Institute, Department of Microbiology, Monash

University, Clayton, Australia

* Joint Corresponding Authors

Dr. Nenad Macesic, Department of Infectious Diseases, The Alfred Hospital and School of

Translational Medicine, Monash University, Level 1 Alfred Lane House, Alfred Hospital, 55

Commercial Rd, Melbourne, VIC 3004, Australia. Email: nenad.macesic1@monash.edu

Professor Anton Y. Peleg, Department of Infectious Diseases, The Alfred Hospital and School of

Translational Medicine, Monash University, Level 1 Alfred Lane House, Alfred Hospital, 55

Commercial Rd, Melbourne, VIC 3004, Australia. Email: anton.peleg@monash.edu

Word count: 1200 words

Abstract

Objectives

Enterococcus faecium is a common cause of hospital-acquired infections and is associated with

vancomycin resistance. The hospital environment is often a reservoir for *E. faecium* due to its ability

to survive on surfaces. In 2019 we noted increasing prevalence of *vanA E. faecium* causing

bloodstream infections. We aimed to assess the hospital environment as a reservoir of *vanA E.*

faecium.

Methods

We conducted a point prevalence survey of the environment at the Alfred Hospital, Melbourne,

Australia. We sampled all wards (n=12), 10 surface types and 357 surfaces. Resulting *E. faecium*

isolates underwent short-read whole genome sequencing and were compared to invasive infection

(bloodstream) isolates. The *van* operon, multi-locus sequence type (MLST) and core genome MLST

(cgMLST) were determined.

Results

Twenty percent (73/357) of surfaces isolated *E. faecium* with a median of 20% positivity per ward

(IQR 11.6-30.6%). *vanA* and *vanB* were detected in 12/73 (16%) and 58/73 (79%) environmental

genomes, respectively. In contrast, a higher proportion of bloodstream isolates carried *vanA* (32/72

[44%], *P*<0.001). Environmental genomes belonged to 10 MLSTs and 10 cgMLSTs, with ST796/ST78

accounting for 54/73 (74%) genomes. Clinical STs and cgMLSTs overlapped with environmental but

were more diverse. We noted 13 putative genomic transmission clusters. One cluster contained 25/44

(57%) *vanA* genomes, with the majority being clinical (22/26 genomes, 85%). Environmental-clinical

links were observed in 292/895 (33%) putative genomic transmission links. In completed assemblies,

the *vanA* and *vanB* operons were located on pRE25-like plasmids and the chromosome, respectively.

Conclusion

We noted a 20% prevalence of *E. faecium* environmental colonisation but the *vanA* operon was

detected in only 12/73 genomes. We identified distinct environmental and patient reservoirs, with

most environmental genomes carrying the *vanB* operon and having distinct STs/cgMLST clusters.

Environment-clinical reservoir spillover was detected but accounted for a minority of putative

transmission links.

Introduction

Enterococcus faecium is a common cause of hospital-acquired infections and is frequently associated with vancomycin resistance [1], resulting in high mortality, longer hospital stays and higher healthcare costs [2]. The hospital environment is a reservoir for E. faecium due to its ability to survive on surfaces for prolonged periods [1]. Prior studies have analysed patient and environmental reservoirs of E. faecium but have typically been limited to selected hospital wards in settings where either vanA or vanB vancomycin-resistant E. faecium (VREfm) predominate [3-5]. Australia has a specific VREfm epidemiology with more recent emergence of vanA VREfm on a background of vanB VREfm endemicity, with both continuing to circulate [6]. In 2019, we noted increasing prevalence of vanA VREfm bloodstream infections. We aimed to assess the hospital environment as a reservoir of vanA VREfm by conducting a point prevalence study and comparing environmental and contemporary clinical isolates using genomic analyses.

Methods

The study was approved by the Alfred Hospital Ethics Committee. The Alfred Hospital (Melbourne, Australia) is a 600-bed quaternary hospital with state referral services for burns and trauma, as well as stem cell and solid organ (heart/lung/kidney) transplant services. We conducted a point prevalence survey of the hospital environment. We sampled surfaces using FLOQswabs (Copan) over two months, which included all wards and the intensive care unit (ICU) (n=12), with 10 surface types sampled per ward/ICU. Samples were cultured on CHROMagar VRE media (Biomerieux) and species were identified using matrix-assisted laser desorption-ionisation time-of-flight (MALDI-TOF) (Bruker Daltonics). To assess for molecular epidemiological links between environmental and clinical isolates, we selected all E. faecium blood culture isolates from our established biobank from 6 months prior to 6 months post the environmental sampling (Supp. Figure 1 and Supp. Table 1). Clinical data were extracted from the electronic medical record.

All E. faecium isolates underwent short-read (Illumina) whole genome sequencing, as described previously [7]. We detected van operon presence and assigned multi-locus sequence type (MLST) and core genome MLST (cgMLST) [8]. For each cgMLST cluster, we assembled a long-read

reference genome and calculated within-cluster pairwise single nucleotide variant (SNV) distances.

Please see Supplementary Methods for full details.

Results

Environmental point prevalence survey

We swabbed 357 surfaces (median 31.5 per ward [range 10-60]). A total of 73/357 (20%) surfaces

isolated *E. faecium*, with median 20% positive per ward (IQR 11.6-30.6%) (Fig. 1A), indicating that *E.*

faecium was endemic in the hospital environment. Rates of *E. faecium* positivity varied widely, with

the highest rate in the Burns ward (10/10 swabs positive) and the lowest rate on Surgical Ward 3

(0/34 positive). Patient chairs had the highest rate of *E. faecium* positivity (13/25 [52%]) (Fig. 1B).

Environmental and clinical genome characteristics

All 72 *E. faecium* blood culture isolates that occurred in the 6 months pre- and post-environmental

sampling were sequenced and compared to the 73 positive environmental swabs (Fig. 1C, Supp Fig.

1). *vanA* and *vanB* were detected in 12/73 (16%) and 58/73 (79%) environmental genomes,

respectively (Fig. 1C). In contrast, 32/72 (44%) clinical genomes carried *vanA* ($P<0.001$) and 31/72

(43%) carried *vanB*. Environmental genomes belonged to 10 MLSTs and 10 cgMLSTs, with

ST796/ST78 accounting for 54/73 (74%) genomes (Fig. 1C, 1D). Clinical STs/cgMLSTs overlapped

with environmental genomes but were more diverse. Specifically, ST1421/1424 carried *vanA* and

were more frequent in clinical genomes (30/72 v 7/73 genomes, $P<0.001$). A single cgMLST cluster

(B) contained the majority of *vanA* genomes (25/44 genomes), while *vanB* genomes were noted

across two cgMLST clusters (A and C, 84/89 genomes) (Fig. 1D). STs/cgMLSTs carried either *vanA*

or *vanB* operons, except for 3 genomes in ST796/cgMLST cluster A that had concurrent *vanA* and

vanB carriage (Fig. 1C/1D).

Geography of E. faecium on hospital wards

We mapped *E. faecium* ST/cgMLST and presence of *van* operon across individual wards (Supp. Fig.

2). High-risk wards (ICU, Haematology/Oncology) had the majority of clinical genomes with a resultant

high diversity of STs/cgMLSTs (7 STs/10cgMLSTs for Haematology/Oncology and 7 STs/7 cgMLSTs

for ICU, respectively). The Haematology/Oncology ward had the highest proportion of *vanA* genomes

(12/25, 48%), with 7/12 of these being ST1424/cgMLST cluster B genomes. Other wards had uniform

ST/cgMLST composition with no differences between wards (Supp. Fig. 2).

E. faecium *transmission networks*

Using a 6 SNV cutoff [4], there were 895 putative genomic transmission links resulting in 13 clusters

(Fig. 2). While 292/895 (33%) links were environmental-clinical, of these only 88/292 (30%) were

between *vanA* genomes. Most clusters had a predominance of either environmental-environmental

links or clinical-clinical links (Supp. Table 2). A single cluster contained 25/44 (57%) *vanA* genomes,

with the majority being clinical (22/26, 85%). Environmental genomes in that cluster came from three

wards, while clinical genomes came from 11 wards. In contrast, *vanB* genomes belonged to three

major (>5 genomes) clusters, containing predominantly environmental genomes (42/56, 75%).

van *operon analysis*

In completed assemblies (n=26), the _vanA_ and *vanB* operons were located on pRE25-like plasmids

and the chromosome, respectively. The pRE25-like plasmids were heterogeneous and did not

indicate spread of a single plasmid across multiple cgMLST clusters (Supp. Figure 3).

Discussion

We noted a 20% prevalence of *E. faecium* environmental colonisation across a diverse range of

hospital surfaces, but only 12/73 (17%) environmental genomes carried *vanA*. This indicated that

while *E. faecium* was endemic in the hospital environment, it likely did not make a major contribution

to the increased *vanA* VREfm prevalence in bloodstream infection isolates. Indeed, we noted distinct

environmental and patient reservoirs: an environment dominated by *vanB* ST796/78 genomes, and a

clinical reservoir comprising *vanA* ST1421/1424 genomes. There was evidence of spill-over with

putative clinical-environmental links, but these were a minority (33%) and *vanA* putative genomic links

only contributed 88/895 (10%) of these in turn.

Our findings stand in contrast to prior work showing significant links between VREfm in the

environment and clinical colonisation and infection [3-5]. This difference may be due to these studies

describing non-outbreak settings and focusing on limited wards (ICU and Haematology/Oncology).

Potential contributors to our findings include our institution's differing infection prevention approaches

to *vanA* and *vanB* VREfm-colonised patients [9]: *vanA* patients are routinely placed in contact

precautions, while for *vanB* patients this occurs only on high-risk wards and if they have diarrhoea or

non-contained wounds. *E. faecium* ST796, a key *vanB* lineage in our study, has been shown to

develop biocide tolerance that may provide it with a fitness advantage in hospital environments [10].

Of note, the *vanA* clinical reservoir that prompted our investigation subsequently resolved during the

COVID-19 pandemic (Supp. Fig. 1), perhaps because of these factors.

Although the environment plays a major role in healthcare-associated VREfm spread, our study

shows that distinct clinical and environmental reservoirs of VREfm may exist, particularly in outbreak

settings where new clones are emerging and have not yet colonised the environment. This provides

support for use of molecular techniques to focus infection prevention strategies according to the

reservoir. While cleaning strategies aiming to reduce the environmental burden form an essential part

of any multifaceted approach to VREfm control [11], the clinical reservoir may require more active

attention through measures such as introduction of active surveillance, redoubled hand hygiene

efforts and more stringent application of contact precautions in order to prevent both between-patient

spread and future environmental colonisation.

Conflict of Interest Statements

Nil relevant.

Role of Funding Source

This work was supported by the National Health and Medical Research Council of Australia

(Emerging Leader 1 Fellowship APP1176324 to N.M., Practitioner Fellowship APP1117940 to

A.Y.P.).

The funders had no role in study design, data collection and interpretation, or the decision to submit

the work for publication.

Conference Presentation

Portions of this work were presented at European Society of Clinical Microbiology and Infectious

Diseases (ESCMID) Global 2024, held in Barcelona, Spain.

References

[1] X. Zhou, R.J.L. Willems, A.W. Friedrich, J.W.A. Rossen, E. Bathoorn, Enterococcus faecium: from
microbiological insights to practical recommendations for infection control and diagnostics, Antimicrob
Resist Infect Control 9(1) (2020) 130.
[2] M. Serra-Burriel, M. Keys, C. Campillo-Artero, A. Agodi, M. Barchitta, A. Gikas, C. Palos, G. López-
Casasnovas, Impact of multi-drug resistant bacteria on economic and clinical outcomes of healthcare-
associated infections in adults: Systematic review and meta-analysis, Plos One 15(1) (2020)
e0227139.
[3] L. El Haddad, B.M. Hanson, C.A. Arias, S.S. Ghantoji, C.P. Harb, M. Stibich, R.F. Chemaly,
Emergence and Transmission of Daptomycin and Vancomycin-Resistant Enterococci Between
Patients and Hospital Rooms, Clin Infect Dis 73(12) (2021) 2306-2313.
[4] T. Gouliouris, F. Coll, C. Ludden, B. Blane, K.E. Raven, P. Naydenova, C. Crawley, M.E. Torok,
D.A. Enoch, N.M. Brown, E.M. Harrison, J. Parkhill, S.J. Peacock, Quantifying acquisition and
transmission of Enterococcus faecium using genomic surveillance, Nat Microbiol 6(1) (2021) 103-111.
[5] A.S. Lee, E. White, L.G. Monahan, S.O. Jensen, R. Chan, S.J.v. Hal, Defining the Role of the
Environment in the Emergence and Persistence of vanA Vancomycin-Resistant Enterococcus (VRE)
in an Intensive Care Unit: A Molecular Epidemiological Study, Infect. Control Hosp. Epidemiology
39(6) (2018) 668-675.
[6] G.W. Coombs, D.A. Daley, P. Shoby, S. Mowlaboccus, Australian Group on Antimicrobial
Resistance (AGAR) Australian Enterococcal Surveillance Outcome Program (AESOP), Commun Dis
Intell (2018) 46 (2022).
[7] N. Macesic, J. Hawkey, B. Vezina, J.A. Wisniewski, H. Cottingham, L.V. Blakeway, T. Harshegyi, K.
Pragastis, G.Z. Badoordeen, A. Dennison, D.W. Spelman, A.W.J. Jenney, A.Y. Peleg, Genomic
dissection of endemic carbapenem resistance reveals metallo-beta-lactamase dissemination through
clonal, plasmid and integron transfer, Nat. Commun. 14(1) (2023) 4764.
[8] C. Higgs, N.L. Sherry, T. Seemann, K. Horan, H. Walpola, P. Kinsella, K. Bond, D.A. Williamson, C.
Marshall, J.C. Kwong, M.L. Grayson, T.P. Stinear, C.L. Gorrie, B.P. Howden, Optimising genomic
approaches for identifying vancomycin-resistant Enterococcus faecium transmission in healthcare
settings, Nat Commun 13(1) (2022) 509.
[9] D.J. Morgan, R. Murthy, L.S. Munoz-Price, M. Barnden, B.C. Camins, B.L. Johnston, Z. Rubin, K.V.
Sullivan, A.L. Shane, E.P. Dellinger, M.E. Rupp, G. Bearman, Reconsidering contact precautions for
endemic methicillin-resistant Staphylococcus aureus and vancomycin-resistant Enterococcus, Infect
Control Hosp Epidemiol 36(10) (2015) 1163-72.
[10] S.J. Pidot, W. Gao, A.H. Buultjens, I.R. Monk, R. Guerillot, G.P. Carter, J.Y.H. Lee, M.M.C. Lam,
M.L. Grayson, S.A. Ballard, A.A. Mahony, E.A. Grabsch, D. Kotsanas, T.M. Korman, G.W. Coombs,
J.O. Robinson, A. Gonçalves da Silva, T. Seemann, B.P. Howden, P.D.R. Johnson, T.P. Stinear,
Increasing tolerance of hospital Enterococcus faecium to handwash alcohols, Sci Transl Med 10(452)
(2018).
[11] G. De Angelis, M.A. Cataldo, C. De Waure, S. Venturiello, G. La Torre, R. Cauda, Y. Carmeli, E.
Tacconelli, Infection control and prevention measures to reduce the spread of vancomycin-resistant
enterococci in hospitalized patients: a systematic review and meta-analysis, J Antimicrob Chemother
69(5) (2014) 1185-92.
225

 Figure 1 – Summary of *Enterococcus faecium* environmental screening and contemporary
clinical isolates

A) Proportion of environmental screening swabs positive for *E. faecium* by ward and **B)** by surface
type. **C)** Summary of *E. faecium van* operon presence by multi-locus sequence types (MLST) and **D)**
core genome MLST (cgMLST) in clinical and environmental isolates.

235 **Figure 2 – Network analysis of *Enterococcus faecium* putative genomic transmission links**
236

237
238
239 Clinical and environmental *E. faecium* genomes are shown as nodes and putative genomic
240 transmission events (defined as pairwise single nucleotide variant distance ≤6) as edges. Edges
241 resulting from clinical-environmental links are shown in black, and clinical-clinical or environmental-
242 environmental in grey. A single major *vanA* cluster is noted with majority of clinical genomes. In
243 contrast, three major *vanB* clusters are noted with environmental genomes predominating.
244