Detection of Perivascular Spaces at the Gray-White Matter Interface Using Heavily T$_2$-weighted MRI at 7T

Gael Saib[*1], Zeynep H. Demir[*1], Paul A. Taylor[2], S. Lalith Talagala[3], Alan P. Koretsky[1]

[1]Laboratory of Functional and Molecular Imaging, National Institute of Neurological Disorders and Stroke, National Institutes of Health, Bethesda, Maryland, United States, [2]Scientific and Statistical Computing Core, National Institute of Mental Health, National Institutes of Health, Maryland, United States, [3]NIH MRI Research Facility, National Institute of Neurological Disorders and Stroke, National Institutes of Health, Bethesda, Maryland, United States. *These authors contributed equally to this work.

Corresponding Author

Gael Saib

Laboratory of Functional and Molecular Imaging, National Institute of Neurological Disorders and Stroke, National Institutes of Health

10 Center Drive, Building 10, Room B1D69

Bethesda, MD 20892, USA

email: saibgh@nih.gov

Abstract

Background: There is increasing interest in high-contrast cerebrospinal fluid (CSF) MRI for imaging perivascular spaces (PVSs). Dilated PVSs, associated with aging, dementia, and various other conditions, are readily detected within the white matter (WM), basal ganglia, and midbrain. While 7T MRI enables detection of smaller PVSs, cortical PVS burden has received limited attention despite its potential value for understanding neurological conditions.

Purpose: To investigate the detectability of cortical PVS segments in healthy participants using heavily T_2-weighted MRI at 7T.

Materials and Methods: A T_2-weighted 3D-TSE sequence was optimized at 7T to detect CSF with high resolution and contrast-to-noise ratio (CNR) while minimizing signal from surrounding tissues. A semi-automated pipeline was developed to extract PVSs and quantify their density in the whole brain, including the cortex.

Results: Seventeen healthy volunteers (40 ± 14 years) were scanned at 7T. Optimized TSE achieved a CSF-to-tissue CNR of ~180:1, enabling detection of small PVSs throughout the brain and leukocortical segments. About 20% of WM PVSs contain a leukocortical segment. WM PVSs with a leukocortical segment represented 70% of the total PVS volume. PVS density in the cortex was ~0.7% (~6-fold lower than WM), with highest in the insula and lowest in the auditory cortex.

Conclusion: High-resolution CSF imaging using optimized 3D-TSE MRI at 7T allows detection and quantification of leukocortical PVS segments at the gray-white matter interface in healthy individuals. This study lays the groundwork for exploring regional PVS changes related to the cortex and their potential use in diagnosis or prognosis of neurological diseases.

Keywords: perivascular spaces, cortex, leukocortical, gray-white matter interface, MRI, heavily T$_2$-weighted MRI, 7T, ultra-high field

Abbreviations

PVS: perivascular space, CSF: cerebrospinal fluid, TSE: turbo spin echo, ETL: echo train length, ES: echo spacing, FA: flip angle, BW: bandwidth, TI: inversion time, TA: acquisition time, TR: repetition time, TE: echo time, WM: white matter, GM: gray matter, CNR: contrast-to-noise ratio, SAR: specific absorption rate

1. Introduction

Perivascular spaces (PVSs), also known as Virchow-Robin spaces, are fluid-filled compartments surrounding blood vessels in the brain parenchyma (Durand-Fardel, 1843; Virchow, 1851). They serve as a pathway for cerebrospinal fluid (CSF) exchange with interstitial fluid (ISF) and play an important role in immune surveillance, brain homeostasis and the clearance of metabolic waste (Wardlaw et al., 2020). Emerging studies have shown that PVS dilatation is associated with aging, small vessel diseases, and neurodegenerative diseases (Francis et al., 2019). MRI has enabled the non-invasive characterization of dilated PVSs, notably along lenticulostriate arteries in the basal ganglia, medullary arteries in white matter (WM), and perforating arteries in midbrain (Kwee & Kwee, 2007). Large dilated PVSs associated with opercular and insular middle cerebral arteries (MCA) branches have also been identified in the anterior temporal lobe region and frontal operculum (McArdle et al., 2020) as well as along hippocampal perforating vessels (Yao et al., 2014). Heavily T_2-weighted MRI at high field ($\geq$3T) has provided superior CSF-to-tissue contrast to noise ratio (CNR) and finer spatial resolution (Daoust et al., 2017; Gao et al., 2014). 7T MRI has further improved the detection of PVSs at sub-millimeter spatial resolution (Barisano et al., 2019; Bouvy et al., 2014; Zong et al., 2016), enabling earlier identification of changes in smaller PVSs.

Most arterial PVSs in the basal ganglia and centrum semiovale are coated with two layers of leptomeninges (Pollock et al., 1997), in contrast to the cortex where the periarterial space is formed by a single layer of leptomeninges. In cortical penetrating arterioles and capillaries, this layer gets increasingly compressed by glia limitans and the pia mater against the vessel wall resulting in smaller PVSs (Morris et al., 2016; E. T. Zhang et al., 1990). For subcortical arteries

originating from the cortex, this transition occurs at the gray-white matter interface, defined as transition zone from cortical gray matter (GM) to WM (Song et al., 2025), and may be associated with regional differences in fluid drainage properties. As communication pathways between subarachnoid CSF and the cortex has been demonstrated (Eide et al., 2018; Ringstad et al., 2018), early changes in leukocortical PVS segments in the gray-white matter interface may offer valuable insight into neurological conditions. High-resolution MRI has enabled visualization of PVSs in this region in both healthy individuals (Liu et al., 2025) and in patients with cerebral amyloid angiopathy (Perosa et al., 2022). However, to the best of our knowledge, a detailed imaging study of cortical PVS segments has not previously been reported.

In a previous preliminary study, a high-resolution T_2-weighted 3D Turbo Spin Echo (TSE) sequence was optimized at 7T to maximize CSF-to-tissue CNR while minimizing signal from parenchyma, with a specific attention to CSF-to-GM contrast (Saib et al., 2023). In the present work, PVSs were characterized in the whole brain, with a focus on leukocortical segments in the gray-white matter interface, using a semi-automated processing technique that accounts for local variations in signal and background noise at 7T. Based on high-resolution MP2RAGE segmentation, PVS density was quantified in the whole brain and within 266 cortical areas for leukocortical PVS segments. This study lays the groundwork for assessing the diagnostic utility of detecting PVS burden in the gray-white matter interface as well as understanding the mechanisms for structural and contrast changes in the cortex in patients with neurological disorders.

2. Materials and Methods

2.1. Study design and participants

This study was conducted under a protocol (00-N-0082) approved by the Institutional

Review Board of the National Institutes of Health (Bethesda, Maryland, USA) and registered on

ClinicalTrials.gov (NCT 00004577). A total of 17 healthy volunteers (10 males and 7 females;

mean age 40 ± 14 years; age range 23-67 years) were recruited and signed an informed consent to

participate in the study. Participant identifiers were not known to anyone outside the research

group. The health status of the participants was assessed through screening, a standard

neurological examination performed by a Licensed Independent Practitioner from National

Institute of Neurological Disorders and Stroke and a clinical brain MRI. Individuals were

excluded if they were deemed to have a medical condition such as neurological disorders,

diabetes, uncontrolled hypertension, prior brain surgery or other relevant health issues. All

imaging was performed on a 7T Magnetom Terra (Siemens Healthineers, Erlangen, Germany)

using a 1Tx-32Rx head array (Nova Medical, Wilmington, USA). No dielectric pads were used.

2.2.MRI protocol optimization

The MRI sequence parameters that were developed and implemented are summarized in

Table 1. CSF-to-tissue CNR of a high-resolution (0.5 mm isotropic spatial resolution or 0.125

mm^3 voxel volume) 3D TSE sequence at 7T was optimized using different TEs and echo train

lengths (ETLs) while restricting the scan time to ~10 min. Intra-subject TSE scans were aligned

using MATLAB (R2022b, MathWorks, Natick, MA, USA) affine registration tool *imregister.*

Signal intensities in CSF ventricles, WM, and deep GM were estimated using four regions of

interest (ROIs) of varying sizes drawn in a central slice. The WM and deep GM ROIs were

placed in regions excluding PVSs but near the ventricles to minimize contrast variations due to

non-uniform B_1 distribution at 7T. The CNR was calculated as the mean signal difference

between CSF and tissue ROIs divided by the standard deviation of pixels within tissue ROIs.

Numerical simulations based on an Extended Phase Graph algorithm (Hennig et al., 2003) were

conducted to assess CSF and tissue signal evolution across varying TEs, refocusing flip angles

(FA) and TRs (Fig S1). Additionally, a high-resolution T_1-weighted MP2RAGE (Marques et al.,

2010) was acquired for tissue segmentation and a fluid attenuated inversion recovery (FLAIR)

(Hajnal et al., 1992) was performed to detect any WM hyperintensities (WMH) or lacunes.

	TSE	MPRAGE	FLAIR
Orientation	sagittal	sagittal	sagittal
Matrix size	400 x 400 x 320	454 x 480 x 320	272 x 272 x 224
Resolution (mm^3)	0.5 x 0.5 x 0.5	0.5 x 0.5 x 0.5	0.74 x 0.74 x 0.74
Refocusing FA (°)	100	-	variable
TR (ms)	3400/**2430**	5000	9000
TE (ms)	300/**500**/700	1.85	273
TI (ms)	-	740/2700	2300
BW (pixel/Hz)	500	360	681
ES (ms)	4.64	7.3	3.9
ETL	151/**108**	264	165
IPAT (GRAPPA)	4	3	6
Slice partial Fourier	-	6/8	-
Elliptical scanning	yes	no	no
TA (min:sec)	10:45	14:00	10:00

Table 1: Protocol sequence parameters for 7T MRI scans conducted in this study. The CSF-to-

tissue CNR of 3D TSE was assessed at three varying TEs and image sharpness was evaluated at

echo train length (ETL) 108 and 151 using 4.64 ms echo spacing (ES), respectively

corresponding to a ETL duration of ~500 ms and ~700 ms. Optimal TSE parameters are

highlighted in bold. A high-resolution MP2RAGE was acquired to segment brain tissues and

cortical areas with a high accuracy. A FLAIR was also performed to detect potential WMH and

lacunes that may be confounded with PVSs.

2.3.Processing pipeline and PVS quantification

An outline of the semi-automated PVS processing framework is illustrated in Figure 1. The

MP2RAGE and FLAIR images were aligned to the TSE images using a linear affine

transformation with *3dAllineate* in AFNI (R. W. Cox, 1996). Alignment was guided using a

combination of the local Pearson absolute (lpa) and lpa+ZZ cost functions, which apply a

combination of non-linear local weightings to derive matches based on the global set of details

(Saad et al., 2009). For computational efficiency of aligning the high-resolution datasets,

downsampling in place by a factor of 3 in all dimensions was used in the initial alignment, where

a larger phase space of parameters is searched, with subsequent refinement at full resolution. The

first inversion MP2RAGE images were aligned to the TSE images, and the same transformation

was applied to the second inversion and UNI images, which combine the first and second

inversion dataset. Next, the FLAIR image was aligned to the already aligned second inversion

image.

Figure 1: Flow chart of semi-automated PVS processing. PVS extraction consisted of applying a

vesselness Frangi filter to the denoised TSE images and calculation of a Z-score map of each

vessel-like structures based on their average surrounding noise. In addition, the TSE images were

used to extract a bulk CSF mask while skull stripping and WM segmentation was conducted on

MP2RAGE high-resolution anatomical images with FSL. GM mask was generated by extracting

the region between WM and bulk CSF masks. A cortical atlas was then generated using GM

mask. PVS density was calculated by the ratio of PVS voxels detected with a Z-score > 2.58 in

tissue mask divided by the total number of voxels in the tissue mask or the cortical area. Note

that intracortical PVS structures were excluded from PVS masks, retaining only detected cortical

PVS voxels intersecting any PVS clusters connected to WM.

For tissue segmentation, MP2RAGE second inversion datasets were skull stripped using

FSL-BET (Brain Extraction Tool) (Jenkinson et al., 2005) to produce a brain mask. The brain-

masked T_1-weighted MP2RAGE UNI images were used to generate a WM mask using FSL-

FAST (FMRIB's Automated Segmentation Tool) (Y. Zhang et al., 2001). A flood-fill operation

was performed on the WM mask to remove any imperfections arising from large WM PVSs

mistaken as GM tissue during segmentation. The heavily T_2-weighted TSE images were

denoised by applying a block-matching 4D filtering (BM4D) to further improve the CNR and

ease CSF/PVS segmentation (Maggioni et al., 2013). A bulk CSF mask was extracted from the

denoised TSE images by adaptive thresholding using the MATLAB function *adaptthresh* with a

cubic neighborhood of 35 voxels and selection of largest connected cluster with MATLAB

function *bwconncomp*. Dilatation of 1 voxel was applied to the CSF mask to exclude potential

residual partial volume contamination of bulk CSF signal with tissue. The GM mask was

generated by extracting the region between the WM and CSF masks. Additionally, a multi-modal

cortical parcellation atlas with 180 areas per hemisphere (Glasser et al., 2016) was aligned to

MP2RAGE UNI images and refined using AFNI's *3dROIMaker* and *3dcalc* to ensure ROI

boundaries after mapping were constrained within the subject's GM mask and that resampling

left no gaps among ROIs. Nonlinear alignment from the MNI 2009c asymmetric template was

performed using AFNI's *SSwarper* program, which wraps around *3dQwarp* (R. Cox & Glen,

2013). Again, results of alignment were checked visually for quality, and the estimated

transformation was applied to map the reference atlas to each subject's TSE image space. The

separation of the cortex and deep GM was performed by removing the atlas region from the GM

mask.

PVS mapping consisted of calculating the Z-scores in each vessel-like structure detected in

the denoised TSE images based on their local noise in a surrounding region. The vesselness of

each voxel was estimated by 3D Frangi filtering (Frangi et al., 1998) using default parameters

(α=0.5, β=0.5, c=500). The Frangi filter scale was set to a large range of 1 to 11 voxels with a

step size of 1 voxel to maximize the inclusion of potentially very small PVSs. To estimate the

local noise, a 3D spherical region with 2 mm radius was defined around each potential PVS by

applying a morphological dilation in 3D, excluding any overlapping vessel-like structures within

this region. The local Z-score was calculated as the signal difference between the potential PVS

signal and the mean signal of the surrounding region, divided by the standard deviation of the

signal within that surrounding region. The Z-score map was thresholded at 2.58, corresponding

to the 99.5th percentile of one-sided Gaussian distribution, to generate the PVS mask. This

threshold was empirically determined based on visual inspection and provided a conservative

local CNR criterion ensuring that the PVS voxel signal intensity level was clearly above the

mean local noise. Subsequently, non-PVS voxels included in the PVS mask, such as residual

bulk CSF and potential WMH or lacunes, were manually removed. Isolated intracortical PVS

structures were also excluded because they were small which made reliable identification as

vessel-like structures difficult. Whole-brain PVS clusters were identified using *bwconncomp*, and

WM PVS clusters that intersected any cortical PVS voxel were retained in full and classified as

PVS with a leukocortical segment.

The PVS densities in the WM, deep GM, and cortical GM masks were calculated as the

percentage of the number of voxels in the PVS masks divided by the total number of voxels in

the tissue masks. We then calculated the percentage of WM PVS clusters and WM PVS volume

fraction with a leukocortical segment. The region below the top of cerebellum commonly

affected by signal reduction due to the use of head arrays at 7T was excluded from the

calculation. Scan-rescan data were acquired in 7 subjects within the same session without

repositioning in the magnet. The reproducibility of PVS density measurements was assessed by

calculating the intraclass coefficients (ICC) using one-way random effects model (1,1) and by

Bland-Altman analysis. The PVS density was also quantified in 266 cortical areas (in both

hemispheres) of the brain atlas within the brain region selected for analysis.

3. Results

3.1.TSE parametrization for optimal PVS detection

Figure 2A shows heavily T_2-weighted TSE images acquired at varying TE values for a

representative subject; corresponding simulations are provided in Fig. S1A-C. A reduced and

constant refocusing flip angle was found superior to variable flip angle approaches to reduce

blurring due to shorter ETL needed to maximize the CNR within our scan time limit of 10 min

(Fig. S1D-F). This resulted in a sharper point spread function, effectively enhancing the

visualization of small PVSs. CSF-filled PVSs can be identified as hyperintense punctate or

tubular structures, depending on their orientation in presented axial images. As predicted by

simulations, TSE images acquired at TEs of 500 and 700 ms using a ETL of 151 (echo train

duration 700 ms) exhibited strong CSF signal against a markedly suppressed background signal.

Measured CSF-to-tissue CNR (~100:1) was optimal for both WM and GM at TE = 500 ms (Fig

2B). Maximum intensity projection (MIP) of these images shows elongated features that are

PVSs (Fig 2C). Use of a protocol with a shorter ETL of 108 (echo train duration 500 ms)

maintaining 500 ms TE improved the overall image sharpness, allowing better depiction of small

PVSs. CSF-to-tissue CNR decreased to ~54:1 with shorter ETL of 108 due to the use of a shorter

TR required to maintain scan time under 10 min, as expected from simulations. The CSF-to-

tissue CNR of denoised TSE images at shorter ETL was ~180:1, corresponding to a 230%

enhancement compared to the original image.

Figure 2: TSE optimization with (A) TSE images acquired at varying effective TEs in a

representative subject and (B) corresponding CNR (mean std) determined across four subjects.

PVSs can be visualized as bright punctate and linear structures as pointed by the cyan arrows.

Note that the contrast is optimal at TE = 500 ms (TR = 3.4 s and ETL = 151) for both the WM

and GM. (C) Comparison of 5.5 mm MIP of TSE images (TE = 500 ms) acquired with a ETL of

151 (TR = 3.4 s) and 108 (TR = 2.4 s). PVSs can be identified as linear structures with high

signal intensity. Use of a reduced ETL improved the visualization of small PVSs as pointed by

the cyan arrows.

3.2. Whole-brain semi-automated PVS quantification

PVSs were segmented with the data processing pipeline described in Fig. 1 using TSE

images acquired with the optimal parameters listed in Table 1. TSE image (Fig 3A) and its

associated thresholded PVS Z-score map (Fig. 3B) identify signal intensities exceeding 2.58

standard deviation above local background noise. This enabled detection of PVS signal with a

strong local CNR corresponding to the 99.5 quantile of a Gaussian distribution. The Z-score

threshold reflects a conservative local contrast criterion and not a statistical confidence interval.

Figure 3: Optimal TSE acquisition with resulting PVS Z-score segmentation. (A) displays the

TSE image acquired with optimal parameters listed in Table 1, along with (B) overlay of the Z-

score maps (in color Z>2.58) generated with the pipeline presented in Figure 1. As expected,

many PVSs appearing as discrete spots in single slice are revealed to be elongated vessel-like

structures in (C,D) corresponding maximum intensity projection (MIP) across 11 slices. Note

that bulk CSF was removed from MIPs to improve visualization of the PVSs. Moreover, the PVS

segmentation was performed excluding the region below the top of cerebellum commonly

affected by signal reduction due to the use of head arrays at 7T.

Denoising effectively reduced background tissue noise while preserving CSF and PVS sharp

appearance. The noise-only background limited the risk of structure smoothing. An example of

the relationship between the TSE signal, before and after denoising, with corresponding PVS Z-

score is shown in Figure S2.

PVSs exhibited Z-score values as high as 114 ± 63 across subjects with a mean Z-score of

6 ± 1. Z-scores close to CSF-to-tissue CNR measured in ventricles were likely associated with

large PVSs while lower Z-scores were attributed to partial volume effects of small PVSs with

surrounding tissues. As expected, most of extracted PVSs appearing as discrete structures were

revealed to be elongated vessel-like structures, when viewed in the corresponding MIPs (Fig 3C,

D). The overlay of the magnitude gradient of the CSF image is shown on aligned MP2RAGE

UNI image for a representative subject in Figure 4A. The edge of the bulk CSF perfectly aligns

with the border between CSF and gray matter in the MP2RAGE image, indicating excellent

registration of the anatomical image on the TSE image (additional examples in Fig S3). This

allowed precise localization of each PVS within anatomical regions, as illustrated by the PVS Z-

score map overlaid on the MP2RAGE anatomical image in Figure 4B.

Figure 4: Example of TSE and MP2RAGE alignment with PVS localization in anatomical

regions. (A) Overlay of the edges of the TSE image (in cyan) precisely aligns with the interface

between the dark CSF border and gray matter in the registered MP2RAGE UNI image,

indicating excellent anatomical alignment. (B) PVS Z-score map (in color Z > 2.58) overlaid on

the MP2RAGE UNI image, highlighting extracted PVSs position in anatomical regions.

3.3. PVS characterization at the gray-white matter interface

Using the acquisition and data processing protocol described above, approximately 20 3%

of WM PVS clusters across subjects were found to have a leukocortical segment. The portion of

WM PVSs with a leukocortical segment accounted for 74 5% of the total PVS volume. This

indicates that WM PVSs connected to the gray-white matter interface accounted for majority of

the PVS volume, while a substantial fraction of WM PVSs corresponded to small isolated PVS

structures. Figure 5 presents representative examples of PVSs at the gray-white matter interface,

likely associated with medullary arteries, across different individuals. Figure 5A and 5B show

examples of individual PVSs with a leukocortical segment, while Figure 5C illustrates a case of

PVS appearing as a cluster with multiple leukocortical segments within a common cortical area.

Leukocortical PVS segments get increasingly smaller in the cortex likely due to compression

around cortical penetrating vessels, resulting in a Z-score below threshold. The potential

connection of PVSs with the subarachnoid space was difficult to assess, notably due to partial

volume effects induced by bulk CSF within GM regions. Although the PVS sections in WM can

be identified as hypointense on MP2RAGE images (Fig 5, column 2), the connecting

leukocortical PVS segments could only be reliably detected with processing of the TSE images

(Fig 5 column 5). Most PVSs were not visible on FLAIR images but FLAIR confirmed that

PVSs were not confounded by WMH or lacunes. GM mask outer surface did not coincide

perfectly with the FLAIR outer edge due to its lower spatial resolution.

Figure 5: Representative examples of PVSs with a leukocortical segment in the gray-white

matter interface. The first column shows anatomical locations on M2RAGE images across

different subjects, with corresponding zoomed regions in each row for the UNI MP2RAGE,

FLAIR and TSE images. In each scan, the segmented cortical boundary is overlaid as a solid

cyan line while the dotted line delineates outer cortical GM surface at the GM-CSF interface.

The 5^{th} column shows the PVS masks overlaid on TSE images with WM PVS segments

highlighted in cyan and their leukocortical segments in magenta. PVSs with a leukocortical

segment were frequently observed in the gray-white matter interface either as individual

structure (A,B) or clusters (C). These PVSs were likely associated to medullary arteries

originating from small penetrating arterioles in the adjacent cortical areas. Large PVSs at the

boundary may have affected the definition of the cortical boundary on MP2RAGES images.

Unfortunately, FLAIR was not sufficient to identify the true location of the cortical boundary.

Note that the outer surface of the mask did not coincide perfectly with the outer edge of the

FLAIR images because of its lower spatial resolution (0.74 mm isotropic resolution) compared to

MP2RAGE and TSE (0.5 mm isotropic resolution).

Figure 6 presents additional examples of PVSs likely associated with large insular arteries.

As expected, the leukocortical segments of these PVSs exhibited high Z-score values (69 ± 16),

suggesting the presence of large PVS compartments. The morphology of these PVSs often

appeared as dense groups or clusters in both hemispheres, consistent with the anatomy of

perforating arteries. A slight denoising-induced smoothing resulted in a patch-like appearance in

PVS masks. FLAIR contrast was further affected by B_1 inhomogeneities in this inferior region

but confirm the absence of confounding effects.

Figure 6: Additional examples of PVSs exhibiting a leukocortical segment in the gray-white matter interface. The first column shows anatomical locations on axial MP2RAGE images across subjects, with corresponding zoomed regions of the insula displayed in each row. Segmented cortical boundary is overlaid as a solid cyan line while the dotted line delineates outer cortical GM surface at the GM-CSF interface. In this region, PVSs were frequently observed as clusters, likely associated with groups of perforating arteries branching from the MCA. In the fifth column, PVS masks overlaid on TSE images highlight large leukocortical PVS segments in the insula (magenta) associated to WM PVSs (cyan). Note that PVS clusters were systematically found in both hemispheres. Large PVS clusters may further alter T_1-contrast in MP2RAGE images, affecting cortical boundary delineation. FLAIR contrast was further degraded due to B_1 inhomogeneities in this inferior region.

In some cases, PVS-induced signal reduction on MP2RAGE images may have locally impacted the definition of the cortical boundary in tissue segmentation, especially for large PVSs

and dense PVS clusters (Fig S4). Line-scan analysis across contrasts supported that leukocortical

PVS segments presented in Figure 5C have a portion within cortex as signal was observed to

spread deeper within the cortical region, towards the CSF-GM interface (Fig S4A). Subsets of

leukocortical PVS segments in Figure 6 may have been overestimated due to potential errors in

tissue masks. As shown in line-scan analysis of Figure S4C, the precise location of the boundary

in this region was hard to assess in both MP2RAGE and FLAIR images, yet PVSs with a lower

signal intensity in the TSE likely did not affect WM MP2RAGE signal and appear to be located

within the cortex.

3.4. PVS density in WM, deep GM, and cortex

The average PVS density was found to be 4.3±1.5% in WM, 2.6±1.7% in the deep GM

and 0.74±0.4% in the cortex (Fig S5). The PVS density in the cortex represents only the

segments in the gray-white matter interface. The scan-rescan reliability showed good intrasubject

reproducibility of repeated measurements with an ICC of 0.93, 0.98, and 0.63 respectively for

WM, deep GM, and cortex. The 95% limit of agreement in Bland-Altman analysis was slightly

exceeded in two subjects in the WM, likely due to motion artifacts in rescan TSE acquisitions.

PVS density in WM and deep GM varied with age (Fig S5A), whereas it appeared to be more

stable in the cortex.

Figure 7 shows the 20 cortical areas with the highest PVS density measured across the 17

subjects. Among 266 specific cortical areas analyzed above the cerebellum, top PVS densities

(~2.7-6.2%) were found related to the anterior agranular insula complex (AAIC) and the

posterior insular areas (PoI1, PoI2) supplied by the branches of the MCA. The primary auditory

cortex (A1) and the lateral belt complex (LBelt) were found to have lowest PVS densities (<0.04

%). Table S1 summarizes all regional PVS densities of the 266 cortical areas averaged across all

the participants.

Figure 7: Twenty highest cortical PVS density in atlas regions for both hemispheres. Highest

PVS density (~3-6%) was found in anterior agranular insula complex (AAIC) and posterior

insular areas (PoI1, PoI2) while lowest (< 0.04%) were found in primary auditory cortex (A1)

and the lateral belt complex (LBelt).

4. Discussion

This study introduces an optimized heavily T_2-weighted 3D-TSE sequence at 7T that

achieves high CSF-to-tissue CNR while minimizing background tissue signal, enabling high-

resolution whole-brain imaging of PVSs in the gray-white matter interface. Leukocortical

segments associated with a majority of WM PVS volumes were readily detected, as the TSE

contrast was tailored to maximize CSF-to-GM CNR. Using the semi-automated PVS processing

framework developed here, PVS density was quantified across individual cortical areas

throughout the brain, allowing assessment of leukocortical PVS burden in the gray-white matter

interface.

Brain tissue suppression with TSE was excellent at TE=500 ms when using a constant

refocusing flip angle of 100° and a TR of 2.4 s. Employing a short ETL and reduced flip angle

resulted in lowering the echo signal variation across the ETL, leading to minimal blurring as well

as lower SAR. Reduced and constant refocusing flip angle was found to increase image

sharpness compared to variable flip angle strategies, such as sampling perfection with

application-optimized contrasts using different flip-angle evolutions (SPACE) (Mugler et al.,

2000), which required extended ETL to achieve similar CNR. A very high CSF-to-tissue CNR of

180:1 at 0.5 mm isotropic spatial resolution was achieved by combining optimal TSE parameters

with denoising, facilitating the PVS quantification in the GM cortical regions. The semi-

automated processing method developed in this study relies on the vesselness filtering and

calculation of PVS local Z-score. Use of a local Z-score accounts for spatial variation of noise

well known to be prominent at 7T (Maximov et al., 2012). A Z-score threshold of 2.58 enabled

extraction of PVSs with conservative local CNR criterion providing confidence that each PVS

signal intensity was clearly above the mean local noise.

The majority of the leukocortical PVS segments detected were likely associated to

medullary arteries as described previously (Kwee & Kwee, 2007). These PVSs were sometimes

organized in dense clusters at the gray-white matter interface across various anatomical locations

(Fig S4). Groups of large PVSs with leukocortical PVS segments were consistently found within

the insula, associated to M2 branches of the MCA, which involve larger arteries that abuts and

indents the cortex (McArdle et al., 2020; Varnavas & Grand, 1999). Alignment of TSE images

with high-resolution angiography will provide a better understanding of their association with the

underlying cerebral vasculature. Intracortical PVSs were occasionally detected by Frangi

filtering, however, they lost their vessel-like appearance after Z-score thresholding due to their

small size. Therefore, it was difficult to identify these PVSs reliably. Future improvements in

MRI sensitivity will be necessary to assess intracortical structures. PVS detection may be

enhanced by restoring the longitudinal magnetization between TRs, implementing zero-filling

approaches and minimizing background tissue noise further along using advanced denoising

methods. These strategies and application of prospective motion correction techniques to reduce

image artifacts will be investigated in future studies.

The average WM PVS density (~4.3%) was found to be within the range reported in

previous studies (Barisano et al., 2021; Cai et al., 2015). The cortical PVS density of segments

connected to WM PVSs was about ~0.7% which was, as expected, much lower compared to WM

PVS density. However, the actual cortical PVS density may be higher since possible intracortical

PVSs were not included in the quantification. The highest PVS density was found in the insula

(AAIC, PoI1 and PoI2) while lowest density was found in the auditory cortex (LBelt and A1). In

agreement with previous studies (Chen et al., 2011; Zhu et al., 2011), PVS density within WM

and deep GM appeared to increase with age. However, the PVS density in the cortex was found

to be stable across individuals. Since the PVS density has been found to depend on multiple

factors even in the healthy population (Barisano et al., 2021), a larger cohort will be required to

robustly assess age-related PVS volume changes.

The brain MRI alignment and segmentation pipeline allowed for the extraction of bulk

CSF, WM, and GM as well as the precise definition of 266 cortical areas for the quantification of

the PVS densities. Large PVSs or dense PVS clusters near the cortical boundary could affect

local boundary delineation due to signal reduction in WM on MP2RAGE images (Fig S4).

Therefore, some extracted leukocortical PVS segments may include a greater contribution from

WM. This effect may lead to an overestimation of the PVS density in some cortical regions.

Although better definition of the boundary would help precisely isolate the cortical segments of

these leukocortical PVSs, many PVSs in the gray-white matter interface were observed deeper

within cortex, supporting the presence of a distinct cortical segment. A FLAIR protocol with

improved gray-white matter contrast may provide additional insight into the definition of the

boundary between the tissues. Moreover, due to reduced MP2RAGE contrast at 7T, part of the

thalamus was not properly segmented and was included into WM mask, which may have

affected deep GM quantification. MRI contrast may be improved with parallel transmission RF

pulses at 7T (Gras et al., 2017). This will improve tissue and PVS segmentation in inferior brain

regions while minimizing TSE contrast variability across the brain, which should enable

quantitative volumetric PVS mapping.

The amount of CSF in any PVS voxel can be estimated by using the relationship that the

TSE signal intensity of a 100 % CSF voxel corresponds to a voxel volume of 125 nL (0.125

mm^3). The mean CSF signal measured within fully CSF-filled voxels in the ventricles was

330$\pm$51 across subjects, while the mean background tissue signal was 8.3$\pm$1.8. For chosen

detection threshold (Z=2.58), the corresponding signal intensity was ~13 (8.3+2.58*1.8). This

threshold resulted in CSF detection limit of approximatively 5 nL (125 nL*13/330). Potential

effects such as T_2 may also influence PVS signal and will have to be accounted for accurate PVS

volumetry.

In conclusion, this study optimized a high-resolution (0.125 mm^3 voxels) T_2-weighted

3D-TSE sequence at 7T providing a CSF-to-tissue CNR of ~180:1 with minimal tissue

contribution. In addition, this work developed a processing pipeline enabling robust PVS

segmentation and quantification in the whole brain. PVSs with a leukocortical segment were

consistently identified at the gray-white matter interface in healthy individuals and their density

was analyzed across multiple cortical regions. Changes in morphology or density of

leukocortical PVS segments may reflect properties of the associated cortical areas and could help

characterize various neurological conditions.

Data and Code availability

The dataset, including high-resolution 7T MRI scans (T_2-weighted 3D TSE, MP2RAGE

and FLAIR) from 17 healthy participants, as well as retest data from 7 participants, and

associated PVS masks, is available at https://figshare.com/s/6d97007e79a12b7bc47d. Computer

codes for image alignment and segmentation are available at https://afni.nimh.nih.gov. Frangi

filter implementation used for PVS extraction is available on MATLAB File Exchange

(https://www.mathworks.com/matlabcentral/fileexchange/24409-hessian-based-frangi-

vesselness-filter). Other scripts used in the manuscript are available upon request.

Author Contributions

GS and ZHD have equal first authorship. ZHD conceived of the presented idea and

developed the theory, as well as leading subject recruitment. GS performed technical design,

optimizing MRI sequences and performed majority of the data processing. GS conducted the

experiments and data acquisitions with ZHD. GS and ZHD analyzed and interpreted the data

together. GS wrote the manuscript, with contributions from ZHD. PAT handled MRI image

alignments and provided input on technical aspects. SLT provided technical insight on MRI

methods, contributed to data analysis and writing of manuscript. APK provided insights on study

design and data processing, supervised the project and contributed to the manuscript.

Funding

This research was supported by the Intramural Research Program of the National

Institutes of Health (NIH). The contributions of the NIH author(s) were made as part of their

official duties as NIH federal employees, are in compliance with agency policy requirements,

and are considered Works of the United States Government. However, the findings and

conclusions presented in this paper are those of the author(s) and do not necessarily reflect the

views of the NIH or the U.S. Department of Health and Human Services. GS, ZHD, SLT and

APK were supported by the Intramural Research Program of the National Institute of

Neurological Disorders and Stroke at the NIH. PAT was supported by the Intramural Research

Program of the National Institute of Mental Health. This work utilized the computational

resources of the NIH HPC Biowulf cluster (https://hpc.nih.gov).

Declaration of Competing Interests

All authors declare that no competing interest exist.

References

Barisano, G., Sepehrband, F., Ma, S., Jann, K., Cabeen, R., Wang, D. J., Toga, A. W., & Law, M. (2019). Clinical 7 T MRI: Are we there yet? A review about magnetic resonance imaging at ultra-high field. *The British Journal of Radiology*, *92*(1094), 20180492. https://doi.org/10.1259/bjr.20180492

Barisano, G., Sheikh-Bahaei, N., Law, M., Toga, A. W., & Sepehrband, F. (2021). Body mass index, time of day and genetics affect perivascular spaces in the white matter. *Journal of Cerebral Blood Flow & Metabolism*, *41*(7), 1563–1578. https://doi.org/10.1177/0271678X20972856

Bouvy, W. H., Biessels, G. J., Kuijf, H. J., Kappelle, L. J., Luijten, P. R., & Zwanenburg, J. J. M. (2014). Visualization of Perivascular Spaces and Perforating Arteries With 7 T Magnetic Resonance Imaging: *Investigative Radiology*, *49*(5), 307–313. https://doi.org/10.1097/RLI.0000000000000027

Cai, K., Tain, R., Das, S., Damen, F. C., Sui, Y., Valyi-Nagy, T., Elliott, M. A., & Zhou, X. J. (2015). The feasibility of quantitative MRI of perivascular spaces at 7 T. *Journal of Neuroscience Methods*, *256*, 151–156. https://doi.org/10.1016/j.jneumeth.2015.09.001

Chen, W., Song, X., Zhang, Y., & for the Alzheimer's Disease Neuroimaging Initiative. (2011). Assessment of the Virchow-Robin Spaces in Alzheimer Disease, Mild Cognitive Impairment, and Normal Aging, Using High-Field MR Imaging. *American Journal of Neuroradiology*, *32*(8), 1490–1495. https://doi.org/10.3174/ajnr.A2541

Cox, R., & Glen, D. (2013). Nonlinear warping in AFNI. *Proceedings of the 19th Annual Meeting of the Organization for Human Brain Mapping*.

Cox, R. W. (1996). AFNI: Software for Analysis and Visualization of Functional Magnetic

Resonance Neuroimages. *Computers and Biomedical Research*, *29*(3), 162–173.

https://doi.org/10.1006/cbmr.1996.0014

Daoust, A., Dodd, S., Nair, G., Bouraoud, N., Jacobson, S., Walbridge, S., Reich, D. S., &

Koretsky, A. (2017). Transverse relaxation of cerebrospinal fluid depends on glucose

concentration. *Magnetic Resonance Imaging*, *44*, 72–81.

https://doi.org/10.1016/j.mri.2017.08.001

Durand-Fardel, M. (1843). *Traité du ramollissement du cerveau*.

Eide, P. K., Vatnehol, S. A. S., Emblem, K. E., & Ringstad, G. (2018). Magnetic resonance

imaging provides evidence of glymphatic drainage from human brain to cervical lymph

nodes. *Scientific Reports*, *8*(1), 7194. https://doi.org/10.1038/s41598-018-25666-4

Francis, F., Ballerini, L., & Wardlaw, J. M. (2019). Perivascular spaces and their associations

with risk factors, clinical disorders and neuroimaging features: A systematic review and

meta-analysis. *International Journal of Stroke*, *14*(4), 359–371.

https://doi.org/10.1177/1747493019830321

Frangi, A. F., Niessen, W. J., Vincken, K. L., & Viergever, M. A. (1998). Multiscale vessel

enhancement filtering. In W. M. Wells, A. Colchester, & S. Delp (Eds.), *Medical Image*

Computing and Computer-Assisted Intervention—MICCAI'98 (Vol. 1496, pp. 130–137).

Springer Berlin Heidelberg. https://doi.org/10.1007/BFb0056195

Gao, K. C., Nair, G., Cortese, I. C. M., Koretsky, A., & Reich, D. S. (2014). Sub-millimeter

imaging of brain-free water for rapid volume assessment in atrophic brains. *NeuroImage*,

100, 370–378. https://doi.org/10.1016/j.neuroimage.2014.06.014

Glasser, M. F., Coalson, T. S., Robinson, E. C., Hacker, C. D., Harwell, J., Yacoub, E., Ugurbil, K., Andersson, J., Beckmann, C. F., Jenkinson, M., Smith, S. M., & Van Essen, D. C. (2016). A multi-modal parcellation of human cerebral cortex. *Nature*, *536*(7615), 171–178. https://doi.org/10.1038/nature18933

Gras, V., Vignaud, A., Amadon, A., Bihan, D., & Boulant, N. (2017). Universal pulses: A new concept for calibration free parallel transmission. *Magnetic Resonance in Medicine*, *77*(2), 635–643. https://doi.org/10.1002/mrm.26148

Hajnal, J. V., Coene, B. D., Lewis, P. D., Baudouin, C. J., Cowan, F. M., Pennock, J. M., Young, I. R., & Bydder, G. M. (1992). High Signal Regions in Normal White Matter Shown by Heavily T2-Weighted CSF Nulled IR Sequences: *Journal of Computer Assisted Tomography*, *16*(4), 506–513. https://doi.org/10.1097/00004728-199207000-00002

Hennig, J., Weigel, M., & Scheffler, K. (2003). Multiecho sequences with variable refocusing flip angles: Optimization of signal behavior using smooth transitions between pseudo steady states (TRAPS). *Magnetic Resonance in Medicine*, *49*(3), 527–535. https://doi.org/10.1002/mrm.10391

Jenkinson, M., Pechaud, M., & Smith, S. (2005). *BET2: MR-based estimation of brain, skull and scalp surfaces*. Eleventh Annual Meeting of the Organization for Human Brain Mapping.

Kwee, R. M., & Kwee, T. C. (2007). Virchow-Robin Spaces at MR Imaging. *RadioGraphics*, *27*(4), 1071–1086. https://doi.org/10.1148/rg.274065722

Liu, S., Li, J., Hua, R., Xing, Y., Wu, J., Lin, J., Wang, J., Shan, Y., Xu, L., Shi, F., & Zeng, M. (2025). Visualization of perivascular spaces in the human brain with 5-T magnetic resonance imaging. *BMC Neuroscience*, *26*(1), 18. https://doi.org/10.1186/s12868-025-00925-z

Maggioni, M., Katkovnik, V., Egiazarian, K., & Foi, A. (2013). Nonlocal Transform-Domain

Filter for Volumetric Data Denoising and Reconstruction. *IEEE Transactions on Image*

Processing, *22*(1), 119–133. https://doi.org/10.1109/TIP.2012.2210725

Marques, J. P., Kober, T., Krueger, G., Van Der Zwaag, W., Van De Moortele, P.-F., & Gruetter,

R. (2010). MP2RAGE, a self bias-field corrected sequence for improved segmentation

and T1-mapping at high field. *NeuroImage*, *49*(2), 1271–1281.

https://doi.org/10.1016/j.neuroimage.2009.10.002

Maximov, I. I., Farrher, E., Grinberg, F., & Jon Shah, N. (2012). Spatially variable Rician noise

in magnetic resonance imaging. *Medical Image Analysis*, *16*(2), 536–548.

https://doi.org/10.1016/j.media.2011.12.002

McArdle, D. J. T., Lovell, T. J. H., Lekgabe, E., & Gaillard, F. (2020). Opercular perivascular

cysts: A proposed new subtype of dilated perivascular spaces. *European Journal of*

Radiology, *124*, 108838. https://doi.org/10.1016/j.ejrad.2020.108838

Morris, A. W. J., Sharp, M. M., Albargothy, N. J., Fernandes, R., Hawkes, C. A., Verma, A.,

Weller, R. O., & Carare, R. O. (2016). Vascular basement membranes as pathways for

the passage of fluid into and out of the brain. *Acta Neuropathologica*, *131*(5), 725–736.

https://doi.org/10.1007/s00401-016-1555-z

Mugler, J., Kiefer, B., & Brookeman, J. (2000). Three-Dimensional T2-Weighted Imaging of the

Brain Using Very Long Spin-Echo Trains. *Proceedings of the 8th Annual Meeting of*

ISMRM.

Perosa, V., Oltmer, J., Munting, L. P., Freeze, W. M., Auger, C. A., Scherlek, A. A., Van Der

Kouwe, A. J., Iglesias, J. E., Atzeni, A., Bacskai, B. J., Viswanathan, A., Frosch, M. P.,

Greenberg, S. M., & Van Veluw, S. J. (2022). Perivascular space dilation is associated

with vascular amyloid-β accumulation in the overlying cortex. *Acta Neuropathologica*, *143*(3), 331–348. https://doi.org/10.1007/s00401-021-02393-1

Pollock, H., Hutchings, M., Weller, R. O., & Zhang, E.-T. (1997). Perivascular spaces in the basal ganglia of the human brain: Their relationship to lacunes. *Journal of Anatomy*, *191*(3), 337–346. https://doi.org/10.1046/j.1469-7580.1997.19130337.x

Ringstad, G., Valnes, L. M., Dale, A. M., Pripp, A. H., Vatnehol, S.-A. S., Emblem, K. E., Mardal, K.-A., & Eide, P. K. (2018). Brain-wide glymphatic enhancement and clearance in humans assessed with MRI. *JCI Insight*, *3*(13), e121537. https://doi.org/10.1172/jci.insight.121537

Saad, Z. S., Glen, D. R., Chen, G., Beauchamp, M. S., Desai, R., & Cox, R. W. (2009). A new method for improving functional-to-structural MRI alignment using local Pearson correlation. *NeuroImage*, *44*(3), 839–848. https://doi.org/10.1016/j.neuroimage.2008.09.037

Saib, G., Demir, Z., Taylor, P. A., Talagala, L. S., & Koretsky, A. P. (2023). Visualization of Perivascular Spaces using an Optimized 3D-TSE sequence with Reduced Flip Angle at 7T. *Proceedings of the Annual Meeting of ISMRM*.

Song, J. Y., Fleysher, R., Ye, K., Kim, M., Zimmerman, M. E., Lipton, R. B., & Lipton, M. L. (2025). Characterizing the microstructural transition at the gray matter-white matter interface: Implementation and demonstration of age-associated differences. *NeuroImage*, *306*, 121019. https://doi.org/10.1016/j.neuroimage.2025.121019

Varnavas, G. G., & Grand, W. (1999). The Insular Cortex: Morphological and Vascular Anatomic Characteristics. *Neurosurgery*, *44*(1), 127–136. https://doi.org/10.1097/00006123-199901000-00079

Virchow, R. (1851). Ueber die Erweiterung kleinerer Gefäfse: Hierzu Tab. IV. *Archiv für*

Pathologische Anatomie und Physiologie und für Klinische Medicin, *3*(3), 427–462.

https://doi.org/10.1007/BF01960918

Wardlaw, J. M., Benveniste, H., Nedergaard, M., Zlokovic, B. V., Mestre, H., Lee, H., Doubal,

F. N., Brown, R., Ramirez, J., MacIntosh, B. J., Tannenbaum, A., Ballerini, L., Rungta,

R. L., Boido, D., Sweeney, M., Montagne, A., Charpak, S., Joutel, A., Smith, K. J., …

colleagues from the Fondation Leducq Transatlantic Network of Excellence on the Role

of the Perivascular Space in Cerebral Small Vessel Disease. (2020). Perivascular spaces

in the brain: Anatomy, physiology and pathology. *Nature Reviews Neurology*, *16*(3),

137–153. https://doi.org/10.1038/s41582-020-0312-z

Yao, M., Zhu, Y.-C., Soumaré, A., Dufouil, C., Mazoyer, B., Tzourio, C., & Chabriat, H. (2014).

Hippocampal perivascular spaces are related to aging and blood pressure but not to

cognition. *Neurobiology of Aging*, *35*(9), 2118–2125.

https://doi.org/10.1016/j.neurobiolaging.2014.03.021

Zhang, E. T., Inman, C. B., & Weller, R. O. (1990). Interrelationships of the pia mater and the

perivascular (Virchow-Robin) spaces in the human cerebrum. *Journal of Anatomy*, *170*,

111–123.

Zhang, Y., Brady, M., & Smith, S. (2001). Segmentation of brain MR images through a hidden

Markov random field model and the expectation-maximization algorithm. *IEEE*

Transactions on Medical Imaging, *20*(1), 45–57. https://doi.org/10.1109/42.906424

Zhu, Y.-C., Dufouil, C., Mazoyer, B., Soumaré, A., Ricolfi, F., Tzourio, C., & Chabriat, H.

(2011). Frequency and Location of Dilated Virchow-Robin Spaces in Elderly People: A

Population-Based 3D MR Imaging Study. *American Journal of Neuroradiology*, *32*(4),

709–713. https://doi.org/10.3174/ajnr.A2366

Zong, X., Park, S. H., Shen, D., & Lin, W. (2016). Visualization of perivascular spaces in the

human brain at 7 T: Sequence optimization and morphology characterization.

NeuroImage, *125*, 895–902. https://doi.org/10.1016/j.neuroimage.2015.10.078

Supplementary Material

Figure S1: T_2-weighted MRI sequence simulations using extended phase graphs algorithm. (A) Signal evolution of CSF and tissue for a constant refocusing flip angle (FA) with an echo train length (ETL) of 151 (~700 ms), for a TR of 3400 ms (solid lines) and TR infinite (dotted line). Achievable FA and echo spacing (ESP) were respectively restricted to 100° and 4.24 ms due to SAR limits. (B) TR as a function to ETL for a total acquisition time (TA) of 10 min, defined as $TR(s) = TA.ETL/(N_y.N_z)/IPAT$. Elliptical scanning reduced scan duration by ~ 120 sec. (C) Contour plot of normalized CSF-to-GM contrast as a function of refocusing FA and TE. (D) Variable refocusing FA (VFA) along ETL was extracted from Siemens SPACE sequence using CSF $T_1 = 4300$ ms and $T_2 = 2000$ ms, with desired sequence requirements. In this case, the ETL and ESP were respectively adapted to 198 and 4.34 ms to achieve a TE of 500 ms, while TR was set to 4.5 s to limit total TA to ~10 min. (E) Signal evolution of CSF and tissue for constant refocusing FA (solid line) and VFA (dotted line). The use of a constant refocusing FA enables shorter ETL to suppress background tissue signal more effectively compared to VFA. At 500 ms TE, the TSE led to an increase in CSF signal, contrast, and sharpness, as illustrated in (C) by resulting point spread function, where TSE is represented by the solid line and SPACE by the dotted line.

Figure S2: PVS signal intensity profile and corresponding Z-score. (A) A line profile (white line), drawn on TSE image, intersects two leukocortical PVS segments (cyan arrows) and one WM PVS (light gray arrow). (B) TSE signal intensity along this line (black line) shows peaks corresponding to bulk CSF and PVSs. Denoising effectively reduced tissue-related noise while largely preserving the CSF signal (light gray line). Despite a mild smoothing effect of PVS signal, denoising enabled CSF-to-tissue CNR to improve by ~230%. (C) Corresponding Z-score profile highlights detected PVSs with a Z>2.58. Note that signal peaks not visible in the Z-score plot were either not identified by the Frangi filter or were below the Z-score detection threshold (dotted line).

Figure S3: Representative examples showing the alignment of MP2RAGE images to TSE images in 4 additional subjects. Rigid-body registration was performed using AFNI's *3dAllineate*. The cyan contour overlaid on the MP2RAGE images highlights CSF magnitude gradient from the TSE images and aligns precisely with the interface between the dark CSF border and gray matter in the MP2RAGE scans. This demonstrates the high accuracy of the image registration confirming a strong alignment between PVSs and anatomical regions.

Figure S4: Effect of PVS clusters on cortical boundary delineation in different anatomical locations. MP2RAGE UNI images (first column) show localized blurring of the cortical boundary in areas containing PVSs, visible on TSE images (third column), likely associated to medullary arteries (A-B) and insular arteries (C). PVS structures could not be identified clearly on FLAIR images (second column) and assessment of tissues in these regions was difficult due lower resolution and reduced contrast at 7T. According to MP2RAGE segmentation, these PVS clusters had a subset within cortex (blue arrows). In graphs, normalized line scans along yellow dotted lines in each image show corresponding signal intensity evolution in these areas (from left to right). The black dotted vertical lines represent the segmented cortical boundary while the solid vertical lines correspond to the outer surface of the GM mask at the CSF-GM interface. MP2RAGE signal decreases from the cortical boundary towards the CSF-GM interface, while FLAIR signal tends to increase almost linearly across this region. MP2RAGE signal effect is consistent with presence of large PVSs while FLAIR signal appears insensitive to these PVSs. Although large PVSs may have artificially displaced the boundary into WM, small subsets of these PVSs segments deeper into cortex have limited impact on the MP2RAGE signal. Boundary PVSs exhibited higher signal, with gradual decrease toward the cortex, suggesting a likely cortical portion.

Figure S5: PVS quantification with (A) average density extracted in the 17 subjects within WM, deep GM and cortex with (B) Bland-Altman analysis showing the reproducibility of PVS density measurements in these regions for 7 subjects. The black dotted lines represent 95% interval of confidence while the red dotted line shows the mean of the difference between scan and rescan measurements for each subject. Note that the differences in density found between the scans and rescans slightly exceeded 95% confidence interval in the WM for two subjects. This difference was due to motion artefacts within the second scan.

Area Name	Area Description	Right PVS density (%)	Left PVS density (%)
AAIC	Anterior AgranularInsula Complex	6.16 ± 5.3	5.76 ± 4.7
PoI2	Posterior Insular Area 2	5.21 ± 3.4	3.91 ± 2.8
PoI1	Area PosteriorInsular 1	4.24 ± 3.7	2.65 ± 1.7
PIr	PIrform Cortex	3.29 ± 2.7	2.08 ± 1.7
6d	Dorsal area 6	2.07 ± 2.9	1.23 ± 2.0
RSC	RetroSplenial Complex	2.05 ± 0.8	2.16 ± 1.0
55b	Area 55b	1.87 ± 2.5	0.83 ± 1.1
33pr	Area 33 prime	1.83 ± 1.1	1.65 ± 1.4
6ma	Area 6m anterior	1.81 ± 2.3	1.72 ± 2.3
MI	Middle Insular Area	1.71 ± 2.1	1.18 ± 0.9
6mp	Area 6mp	1.68 ± 2.1	0.99 ± 1.6
FEF	Frontal Eye Fields	1.54 ± 2.0	0.61 ± 0.8
SFL	Superior FrontalLanguage Area	1.30 ± 1.7	1.16 ± 1.6
8BL	Area 8B Lateral	1.26 ± 2.0	1.58 ± 2.9
i6-8	Inferior 6-8Transitional Area	1.24 ± 1.8	0.85 ± 0.9
PEF	Premotor Eye Field	1.21 ± 1.6	0.40 ± 0.5
p24	Area posterior 24	1.14 ± 0.5	1.37 ± 1.0
9a	Area 9 anterior	1.10 ± 1.3	0.92 ± 1.1
9m	Area 9 Middle	1.09 ± 1.2	1.05 ± 1.1
10d	Area 10d	1.08 ± 1.4	0.73 ± 0.8
8Av	Area 8Av	1.07 ± 1.6	1.06 ± 1.0
9p	Area 9 Posterior	1.07 ± 1.4	1.17 ± 1.3
6v	Ventral Area 6	1.07 ± 1.2	0.83 ± 1.1
8BM	Area 8BM	1.01 ± 1.3	0.83 ± 2.0
s6-8	Superior 6-8Transitional Area	0.98 ± 1.2	1.10 ± 1.5
FOP3	Frontal OPercularArea 3	0.98 ± 1.0	0.71 ± 0.7
PF	Area PF Complex	0.96 ± 0.9	0.74 ± 0.8
9-46d	Area 9-46d	0.87 ± 1.0	0.76 ± 0.9
OP4	Area OP4/PV	0.87 ± 0.8	0.40 ± 0.4
p9-46v	Area posterior 9-46v	0.84 ± 1.1	0.48 ± 0.6
pOFC	posterior OFC Complex	0.84 ± 0.9	0.49 ± 0.4
1	Area 1	0.84 ± 0.8	0.60 ± 0.8
a24pr	Anterior 24 prime	0.81 ± 1.1	0.42 ± 0.4
4	Primary MotorCortex	0.81 ± 1.0	0.51 ± 0.5
25	Area 25	0.80 ± 0.7	0.41 ± 0.5
a32pr	Area anterior 32prime	0.79 ± 1.0	0.68 ± 0.8
a24	Area a24	0.77 ± 0.7	0.57 ± 0.5
IFSp	Area IFSp	0.74 ± 1.5	0.22 ± 0.3
SCEF	Supplementary andCingulate Eye Field	0.74 ± 1.4	0.59 ± 0.8
8Ad	Area 8Ad	0.74 ± 1.1	0.64 ± 0.7
46	Area 46	0.74 ± 0.8	0.62 ± 0.7
PSL	PeriSylvian LanguageArea	0.74 ± 0.7	0.53 ± 0.3
8C	Area 8C	0.73 ± 0.9	0.37 ± 0.2
PFop	Area PF opercular	0.73 ± 0.8	0.71 ± 0.8
STV	Superior TemporalVisual Area	0.70 ± 0.9	0.47 ± 0.5
2	Area 2	0.67 ± 0.8	0.51 ± 0.7
52	Area 52	0.67 ± 0.8	0.29 ± 0.3
FOP4	Frontal OPercularArea 4	0.67 ± 0.8	0.54 ± 0.4
PFt	Area PFt	0.67 ± 0.7	0.59 ± 0.6
6a	Area 6 anterior	0.65 ± 0.7	0.25 ± 0.3
7PC	Area 7PC	0.64 ± 1.0	0.48 ± 0.8

d32	Area dorsal 32	0.64 ± 0.8	0.62 ± 0.6
3b	Primary SensoryCortex	0.63 ± 0.8	0.35 ± 0.5
6r	Rostral Area 6	0.62 ± 0.5	0.56 ± 0.4
PFm	Area PFm Complex	0.61 ± 0.6	0.57 ± 0.5
RI	RetroInsular Cortex	0.59 ± 0.6	0.27 ± 0.4
FOP2	Frontal OPercularArea 2	0.58 ± 0.5	0.61 ± 0.6
A4	Auditory 4 Complex	0.57 ± 0.6	0.16 ± 0.3
p10p	Area posterior 10p	0.56 ± 0.8	0.78 ± 0.8
p24pr	Area Posterior 24prime	0.56 ± 0.7	0.40 ± 0.4
PGi	Area PGi	0.53 ± 0.9	0.36 ± 0.2
LIPv	Area LateralIntraParietal ventral	0.53 ± 0.8	0.42 ± 0.4
TPOJ1	AreaTemporoParietoOccipital Junction 1	0.51 ± 1.1	0.19 ± 0.2
7Am	Medial Area 7A	0.51 ± 0.4	0.53 ± 0.7
IP2	Area IntraParietal 2	0.50 ± 0.8	0.56 ± 0.8
AVI	Anterior Ventral Insular Area	0.50 ± 0.7	0.25 ± 0.3
43	Area 43	0.49 ± 0.5	0.45 ± 0.4
STGa	Area STGa	0.47 ± 0.8	0.90 ± 1.4
PI	Para-Insular Area	0.47 ± 0.8	0.20 ± 0.4
AIP	Anterior IntraParietalArea	0.46 ± 0.7	0.44 ± 0.7
5L	Area 5L	0.45 ± 0.4	0.71 ± 1.4
OP2-3	Area OP2-3/VS	0.45 ± 0.4	0.33 ± 0.2
44	Area 44	0.44 ± 0.3	0.35 ± 0.2
PFcm	Area PFcm	0.43 ± 0.5	0.59 ± 1.1
FOP5	Area FrontalOpercular 5	0.42 ± 1.0	0.46 ± 0.4
IFSa	Area IFSa	0.39 ± 0.5	0.23 ± 0.2
Ig	Insular GranularComplex	0.39 ± 0.3	0.25 ± 0.4
POS2	Parieto-Occipital Sulcus Area 2	0.38 ± 0.5	0.36 ± 0.5
OP1	Area OP1/SII	0.38 ± 0.5	0.34 ± 0.7
FOP1	Frontal OPercularArea 1	0.38 ± 0.4	0.34 ± 0.6
7Pm	Medial Area 7P	0.38 ± 0.3	0.64 ± 0.7
p32pr	Area p32 prime	0.37 ± 0.5	0.63 ± 1.4
v23ab	Area ventral 23 a+b	0.36 ± 1.0	0.13 ± 0.2
5m	Area 5m	0.36 ± 0.6	0.17 ± 0.1
7AL	Lateral Area 7A	0.36 ± 0.5	0.39 ± 0.6
45	Area 45	0.36 ± 0.5	0.29 ± 0.3
TA2	Area TA2	0.34 ± 0.8	0.21 ± 0.6
a9-46v	Area anterior 9-46v	0.33 ± 0.3	0.52 ± 0.7
11l	Area 11l	0.31 ± 1.0	0.19 ± 0.3
10pp	Polar 10p	0.31 ± 0.4	0.16 ± 0.3
V6A	Area V6A	0.30 ± 0.9	0.05 ± 0.1
A5	Auditory 5 Complex	0.30 ± 0.5	0.28 ± 0.3
PCV	PreCuneus VisualArea	0.30 ± 0.2	0.38 ± 0.6
7Pl	Lateral Area 7P	0.29 ± 0.3	0.16 ± 0.2
PGs	Area PGs	0.29 ± 0.3	0.34 ± 0.2
VIP	Ventral IntraParietal Complex	0.28 ± 0.4	0.35 ± 0.6
IP1	Area IntraParietal 1	0.28 ± 0.4	0.24 ± 0.2
5mv	Area 5m ventral	0.28 ± 0.3	0.24 ± 0.3
IFJp	Area IFJp	0.28 ± 0.3	0.30 ± 0.4
IFJa	Area IFJa	0.27 ± 0.5	0.22 ± 0.3
MIP	Medial IntraParietal Area	0.27 ± 0.4	0.36 ± 0.5
23d	Area 23d	0.27 ± 0.3	0.19 ± 0.2
24dd	Dorsal Area 24d	0.27 ± 0.3	0.18 ± 0.3

a47r	Area anterior 47r	0.27 ± 0.3	0.14 ± 0.1
3a	Area 3a	0.26 ± 0.4	0.18 ± 0.1
24dv	Ventral Area 24d	0.26 ± 0.2	0.31 ± 0.4
47l	Area 47l (47 lateral)	0.24 ± 0.3	0.18 ± 0.3
TPOJ3	AreaTemporoParietoOccipital Junction 3	0.23 ± 0.4	0.34 ± 0.3
STSdp	Area STSd posterior	0.23 ± 0.3	0.16 ± 0.2
LIPd	Area LateralIntraParietal dorsal	0.23 ± 0.2	0.35 ± 0.5
STSva	Area STSv anterior	0.20 ± 0.6	0.02 ± 0.1
IPS1	IntraParietal Sulcus Area 1	0.20 ± 0.5	0.18 ± 0.2
7m	Area 7m	0.19 ± 0.2	0.19 ± 0.3
31a	Area 31a	0.19 ± 0.2	0.27 ± 0.4
s32	Area s32	0.17 ± 0.6	0.10 ± 0.2
PBelt	ParaBelt Complex	0.17 ± 0.3	0.11 ± 0.1
MBelt	Medial Belt Complex	0.17 ± 0.3	0.03 ± 0.0
p32	Area p32	0.17 ± 0.2	0.32 ± 0.5
p47r	Area posterior 47r	0.17 ± 0.2	0.27 ± 0.2
23c	Area 23c	0.16 ± 0.2	0.20 ± 0.2
10v	Area 10v	0.16 ± 0.2	0.17 ± 0.4
a10p	Area anterior 10p	0.16 ± 0.2	0.19 ± 0.2
47s	Area 47s	0.16 ± 0.2	0.21 ± 0.4
d23ab	Area dorsal 23 a+b	0.14 ± 0.2	0.12 ± 0.1
31pv	Area 31p ventral	0.14 ± 0.2	0.11 ± 0.1
31pd	Area 31pd	0.13 ± 0.3	0.21 ± 0.2
10r	Area 10r	0.13 ± 0.2	0.19 ± 0.5
13l	Area 13l	0.11 ± 0.2	0.07 ± 0.1
47m	Area 47m	0.06 ± 0.1	0.12 ± 0.3
STSda	Area STSd anterior	0.06 ± 0.1	0.02 ± 0.0
OFC	Orbital Frontal Complex	0.04 ± 0.0	0.03 ± 0.1
A1	Primary Auditory Cortex	0.03 ± 0.1	0.00 ± 0.0
LBelt	Lateral Belt Complex	0.03 ± 0.0	0.04 ± 0.1

Table S1: PVS densities in the 266 cortical areas for both right and left hemisphere across the 17 subjects. The cortical areas are arranged in descending order based on PVS density in the right hemisphere. The AAIC exhibits the highest density, whereas the L3belt and A1 areas show the lowest densities in the right and left hemispheres, respectively.