Prayer, Politics, and Policy Related to Age-Adjusted Cancer, Heart Disease, Infant Mortality, and
COVID-19 Death Rates, U.S. States 2018-2021

Leon S. Robertson

Professor Adjunct

Yale University School of Public Health

Email: leon.robertson@yale.edu

Abstract

The role of religion and politics in the responses to the coronavirus pandemic raises the

question of their influence on the risk of other diseases. This study focuses on age-adjusted death

rates of cancer, heart disease, and infant mortality per 1000 live births before the pandemic

(2018-2019) and COVID-19 in 2020-2021. Eight hypothesized predictors of health effects were

considered by examining their correlation to age-adjusted death rates and indicators of health

behavior among U.S. states, percentage who pray once or more daily, Republican attitudes and

influence on state health policies as indicated by the percentage vote for Trump in 2016, percent

of household incomes below poverty, median family income divided by a cost-of-living index,

the Gini income inequality index, urban concentration of the population, physicians per capita,

and public health expenditures per capita. Since prayer for divine intervention is common to

otherwise diverse religious beliefs and practices, the percentage of people claiming to pray daily

in each state was used to indicate potential religious influence. Based on collinearity, inequality

was chosen for inclusion over poverty, and the prayer and political variables were analyzed

separately. All of the death rates were higher in states where more people claimed to pray daily.

Only cancer and COVID-19 were correlated significantly with Trump's percentage of the vote.

Lower death rates from cancer and heart disease are associated with more public health

expenditures, but not for infant mortality or COVID-19 deaths. COVID-19 death rates were

lower in states with more physicians per capita, but that variable was not significantly associated

with the other death rates. Heart disease, infant mortality, and COVID-19 death rates were higher

in states with more income inequality. All rates except infant mortality were lower in states

where a greater percentage of the population resides in urban areas. The correlation between

daily prayer and smoking cigarettes, as well as the neglect of public health recommendations for

fruit and vegetable consumption and COVID-19 vaccination, suggests that reliance on prayer

may be a factor in neglect of preventive practices.

Prayer, Politics, and Policy Related to Age-Adjusted Cancer, Heart Disease, Infant Mortality, and

COVID-19 Death Rates, U.S. States 2018-2021

Introduction

The coronavirus pandemic tested medical, political, and economic resources worldwide.

Despite its economic and scientific resources, the U.S. had among the highest death rates in

democratic countries. The U.S. Centers for Disease Control and Prevention's response mainly

consisted of recommendations regarding distancing, mask use, testing, and consultations with

local health officials [1]. Unreliable testing equipment sent to the States had to be recalled,

delaying the testing of the sick [2]. Masking and distancing recommendations were widely

resisted, especially where they were mandated [3]. On-demand testing was associated with

increased virus spread as negative tests were promoted to make travel decisions, with insufficient

emphasis on isolation if one tested positive (4). The public health system, underfinanced in many

states, was unprepared to identify and contact persons exposed to those infected. On 70 percent

of the days in U.S. states in 2020-2021, there were not enough tracing personnel to interview the

infected, much less trace their contacts [5]. People with negative tests traveled and came in

contact with infected people who would not isolate or had no symptoms [4]. The major

accomplishment of the federal government was the effort to develop vaccines that turned out to

be effective beyond expectations. Getting people to use them was more difficult [5,6].

After a CDC spokesperson issued severe warnings about the virus in February 2020, then-

President Trump took charge of televised public briefings, claiming that the virus would

disappear and endorsing treatments lacking scientific support [7]. Following the initial

nationwide shutdown in spring 2020, authority over restrictions on travel, gathering sizes, mask

mandates, and vaccination requirements shifted to the state and local governments, as did the

procurement of protective gear for healthcare workers [8]. These issues became politicized.

Testing and political impacts were separate but additive [9]. When accounting for crowding in

various venues and other risk factors, COVID-19 death rates were higher in counties with more

negative tests per capita and, independently, in those with a greater percentage of Trump votes

[10]. Most Republican governors, with some exceptions, were less likely to impose restrictions,

and a few even questioned vaccine effectiveness [11]. Fox News and right-wing radio broadcasts

promoted skepticism of science [12]. A poll showed that confidence in the CDC's accuracy

among Fox News viewers dropped from about 85 percent in early 2020 to 19 percent two years

later, while remaining above 80 percent among CNN and MSNBC viewers. By 2022, only 16

percent of Fox News viewers reported wearing masks, compared to 60 percent among viewers of

other networks [13]. In the Summer of 2021, around forty percent of surveyed healthcare

professionals received threats [14]. Restricting attendance at religious services was especially

controversial [15], with some religious leaders opposing mask use and vaccines [16].

In recent decades in the U.S., religious fundamentalists have increasingly influenced

Republican politics. Republican operatives and a fundamentalist evangelist formed an

organization called The Moral Majority that persuaded many evangelistic clergy and their

followers to support Republican candidates. However, most African American fundamentalists

and some religious liberals remained loyal to the Democratic Party [17]. By the second decade of

the 21st Century, Republicans had about a 20-percentage-point advantage among Protestants

[18,19].

The scientific study of religious beliefs and practices regarding diseases and injuries mainly

focuses on whether individual risk is associated with prayer or attending religious gatherings.

The results are mixed. A large-sample longitudinal study of cardiovascular disease among

women found that, adjusted for other risk factors, prayer and reading religious literature were

associated with subsequent increased cardiovascular risk [20]. A study of African-American

church attendance and subsequent mortality found that deaths were lower among regular

attendees [21]. Church attendees were found to live longer in another study, but shorter lives

were associated with private religious practices [22]. Deaths per population less than 75 years of

age were substantially correlated with the percentage of people who say they pray daily among

U.S. states before and during the pandemic [6]. The focus on individual behavior and outcomes

ignores the role of government in improving public health historically and the role of religion

and science in the process [6].

There is a large variation in recent leading causes of death (cancer, heart disease, and COVID-

19) among U.S. states. Data obtained for this study showed, in 2018-2019, the range of pooled

age-adjusted rates among the states was: cancer (120-180), heart disease (120-248). In 2020-

2021, the range for COVID-19 deaths was 10-fold (14-140). This paper reports a cross-sectional

analysis of U.S. state differences in the mentioned age-adjusted death rates by cause before the

pandemic (2018-2019) and COVID-19 during the first two years of the pandemic (2020-2021),

correlated to prayer, politics, economic factors, physician accessibility, public health

expenditures per capita, and urban residency. Infant mortality per 1000 live births was added as it

has been correlated to inequality [23]. Although there is a huge variation in ideology,

organization, and ceremonial attendance among religions, respondents to a US survey found 30-

42 percent of Catholics, Protestants, and "other" religions pray for health [24]. The variations in

the rates among 50 U.S. states are examined separately for each cause of death to test the

hypothesis that prayer is associated with death risk, controlling statistically for the other factors.

Correlations of daily prayer with health-related behaviors (smoking, diet, and COVID-19

vaccinations) were also examined using the same methodology.

Materials and Methods

Eight hypothesized predictors of health effects were examined for their correlation to age-

adjusted state death rates and behavioral risk factors – percent who pray once or more daily,

Republican influence on state health policies as indicated by the percentage vote for Trump in

2016, percent households in poverty, median family income divided by a cost-of-living index,

the Gini income inequality index, urban concentration of the population, physicians per capita,

and public health expenditures per capita. The references to data sources of the mentioned

variables are noted in Table 1.

Table 1. Data Sources

| Daily Prayer | Lipka and Wormald, 2016, Pew Research Center [25] |
| Trump Vote 2016 | CNN, 2016 [26] |

Median family Income/cost of living	Moneyrates.com, 2019 [27]
Income Inequality (Gini)	U.S. Census Bureau, 2019 [28]
Physicians per population	National Center for Health Statistics, 2019 [29]
Percent households with poverty income	U.S. Census Bureau, 2020 [30]
Percent Urban	U.S. Census Bureau, 2024 [31]
Public Health Expenditures Per Population	Shadac.org, 2024 [32]
Age-adjusted Mortality Rates	National Institute of Minority Health and Health Disparities, 2018-2021 [33]
Infant Mortality Rate	National Center for Health Statistics, 2018-2019 [34]
Smoking, vegetable, and fruit consumption	Centers for Disease Control Behavioral Risk Factor Surveillance System, 2024 [35]

After examination of the correlations of potential predictor variables for collinearity of the

mentioned predictor variables, least-squares regression estimates of their potential effects on

death rates and behavioral risk factors were calculated. The form of the equation is:

Rate = a + b1(percentage who pray daily) + b2(public health expenditures per population) +

b3(physicians per population) + b4(income per cost of living) + b5(income inequality) +

b6(percent urban) + e (residual variation)

where "Rate" is a death rate or percent health behavior, as appropriate. In a separate analysis, the

percentage of the vote captured by Trump in 2016 was substituted for the percentage who pray

daily in the equations.

The Pew survey of religious behavior is based on a sample of about 35000 adults, about 700

per state, which produces a percentage for each state. The Behavioral Risk Factor survey is based

on interviews with 2500 or more adults in each state. All of the data in this study were statistics

downloaded from publicly available sources. There are no ethical risks to individuals.

Results

The means and standard deviations of the variables included in the analysis are shown in

Table 2. The standard deviations indicate substantial variation in each variable among the 50

U.S. states.

Table 2. Means and Standard Deviations of Analyzed Variables from 50 U.S. States

Variable	Mean	Standard Deviation
Percent Pray Daily	54.2	8.8
Percent Trump Vote 2016	49.7	10.2
Median Income/cost of living	906.6	101.5
Inequality (Gini Index) x 100	46.4	1.8

Percent Poverty	12.1	2.7
Physicians/100000 Population	272.6	55.4
Public Health Expenditures/pop ($)	42.1	29.2
Percent Urban	72.4	14.8
Cancer deaths/100000 population	149.0	14.2
Heart disease deaths/ 100000 population	169.6	30.6
COVID-19 deaths / 100000 population	68.3	28.3
Infant Mortality/ 1000 live births	5.8	1.1
Percent Who Smoke Cigarettes	18.5	3.5
Percent Consuming Recommended Fruit	11.7	1.8
Percent Consuming Recommended Vegetables	9.0	1.8
Percent Fully Vaccinated Against COVID-19	67.7	9.5

Correlations among the potential predictor variables were examined to avoid regression

coefficients distorted by collinearity (Table 3). Most are not statistically significant. The most

important are the correlations among physicians per population, the percentage who pray daily,

poverty, and the Trump vote. On average, there are fewer physicians for potential access by the

ill in states with more Trump voters and people who say they pray daily. The urban percentage is

less correlated with the percentage of daily prayer than the Trump vote percentage. The percent

who pray daily is strongly correlated to the percent in poverty, precluding their use as predictors

in the same equation. While death rates are well known to be higher among the impoverished

aged 40 and older [36], the percentage of the population who pray daily is three times that of the

percentage in poverty (Table 2), encompassing a far larger population. The Gini inequality index

is correlated to percent poverty, but not significantly to the other variables. It controls partly for

poverty in the regression analysis.

Table 3. Least-squares correlation coefficients among predictor variables, N = 50 US states

	Trump Vote 2016	Income/ Cost of Living	Gini Index	Percent in Poverty	Physicians /pop	Public Health Expend/Pop	Urban
Percent Pray Daily	.59	-.17 (ns)	.33 (ns)	.71	-.57	-.13 (ns)	-.13 (ns)
Percent Trump Vote 2016		.05 (ns)	-.12 (ns)	.45	-.68	-.25 (ns)	-.54
Median Income/cost of living			-.23 (ns)	-.49	.09 (ns)	-.36	.17 (ns)
Inequality				.54	.21 (ns)	-.08 (ns)	.26 (ns)
Percent Poverty					.31	-.02 (ns)	.19 (ns)
Physicians/ Population						.10 (ns)	.25 (ns)

Public Health Expenditures/ population							.04 (ns)

146   ns = not statistically significant at p=.05, two-tailed.

147   Table 4. Least-squares regression coefficients and robust standard errors on predictor variables

148   and selected age-adjusted mortality rates per 100,000 population, U.S. States 2018-2019,

149   COVID-19 in 2020.

	Cancer		Heart Disease		Infants		COVID-19	
	Prayer	Trump	Prayer	Trump	Prayer	Trump	Prayer	Trump
Percent Pray Daily	.538* .205		1.493* .440		.087* .015		1.141* . .327	
Percent Trump Vote		. 462* .222		.967 .502		.036 .021		.791* .037
Public Health Expenditures/pop	-.147* .050	- .136* .051	-.229* .107	-.207 .116	-.007 .037	-.006 .005	-.045 .079	-.027 .086
Physicians per Population	.015 .034	.006 .035	-.047 .072	-.107 .079	.001 .002	-.005 .003	-.206 * .054	- .246* .059
Income/Cost of Living	-.006 .015	-.014 .016	-.023 .032	-.039 .036	-.001 .001	-.002 .001	-.047 .024	- .060* .027
Inequality (Gini)	.998 .948	1.937 .838	4.298* 2.034	7.193* 1.893	.064* .070	.124 .078	5.502 * 1.512	7.667* 1.403
Percent Urban	-.563*	.448*	-.718*	-.501	-.013	-.007	-.244 *	-.061

	.100	.122	.208	.275	.007	.011	- .155	.204
Intercept	121.8	85.4	-15.7	.275	6.1	1.9	-130.9	-199.7
Adjusted R-squared	.60	.63	.60	.53	.62	.37	.74	.70

•   P<.05, two-tailed.

Table 4 presents the regression coefficients of age-adjusted death rates per 100,000 population

of the four selected causes of death with robust standard errors beneath the least-squares

estimates. The left column for each cause of death contains the coefficients when prayer is

included as a predictor, and the right column is when the percent Trump vote is substituted for

the percent daily prayers. Contrary to the hypothesis that prayer reduces risk, consistently for

each cause of death, more deaths per capita occur in states where people more often pray daily.

Prayer is the only variable significantly predictive of all four fatality rates. The Trump vote is not

significantly associated with heart disease or infant mortality. Lower death rates from cancer and

heart disease are associated with more public health expenditures, but not for infant mortality or

COVID-19 deaths. COVID-19 death rates were lower in states with more physicians per capita,

but that variable was not significantly associated with the other death rates. Heart disease, infant,

and COVID-19 death rates were higher in states with more income inequality. All of the rates

except infant mortality were lower in states where a greater percentage of the population lives in

urban areas, but were only significantly predictive of cancer in the model that replaced the prayer

variable with the percentage Trump vote.

The association of the predictor variables with important behavioral risk factors is displayed

in Table 5. The greater the percentage of people who say they pray daily in a state, the more who

report smoking cigarettes, consuming less than recommended fruits and vegetables, and having a

 lower percentage of the population vaccinated against COVID-19. Those behaviors were also

 reported by a greater percentage of people in states where Trump received a larger percentage of

 the vote and by a smaller percentage in states with more urban populations. Public health

 expenditures and physicians per capita were associated with a greater percentage of the

 population vaccinated against COVID-19, but not the other risk factors.

 Table 5. Least-squares regression coefficients and robust standard errors on predictor variables

 and the percent of selected health behaviors, N = 50 US states

	Smoke Cigarettes		Consume Fruit		Consume Vegetables		COVID-19 Vaccinated	
	Prayer	Trump	Prayer	Trump	Prayer	Trump	Prayer	Trump
Percent Pray Daily	.132* .052		-.119* .023		-.110* .027		-.333* . .108	
Percent Trump Vote		.198* .050		-.082* .028		-.076* .031		-.550* .094
Public Health Expenditures/pop	-.011 .013	-.006 .012	-.001 .006	-.003 .007	.004 .006	.002 .007	052* .026	.037 .022
Physicians per Population	-.002 .009	.006 .008	.002 .004	.006 .004	-.002 .004	.002 .005	.095* .018	.073* .015
Income/Cost of Living	-.001 .004	-.004 .004	-.004 .002	.000 .002	-.007* .002.	-.005* 002	-.001 .008	.009 .007
Inequality (Gini)	.338 .242	.490* .192	.024 .107	-.202 .107	-.078 .123	-.288* .117	.031 .497	.304 .353

Percent Urban	-.142*	-.087*	.056*	.037*	.054*	.037*	.115*	-.040
	.025	.028	.011	.015	.013	.017	.051	.051
Intercept	7.3	-5.3	14.0	5.5	21.2		48.32	
Adjusted R-square	.56	.63	.68	.56	.56		.75	

Discussion

The large variation in death rates among the states and their strong association with prayer

frequency suggest that higher death rates could be at least partly because prayerful people are

more likely to neglect preventive and ameliorative behavior. They may also be praying more due

to poorer health. The correlation between daily prayer and cigarette smoking, as well as the

inverse correlations between daily prayer and public health-recommended fruit and vegetable

consumption, and COVID-19 vaccination, support the neglect hypothesis. The extent to which

the neglect is an active rejection of public health recommendations or a lack of consideration of

the evidence cannot be determined from aggregated data. As noted in the introduction, studies of

health outcomes related to religious practices are mixed. The result on resistance to COVID-19

vaccinations is similar to that found in a review of twelve studies, finding that the religious are

less likely to be vaccinated [37]. Thousands of lives were shortened because of vaccine

resistance, and thousands more annually would be longer and healthier if there were less

smoking and less consumption of foods that increase the risk of cancer and heart disease [38,39].

It is possible that the correlation between frequent prayer and death rates in this study partly

results from prayerful people's tendency to support state and federal governments that neglect

less fortunate people and oppose environmental protection standards. For example, in a 2016

survey, the more prayerful were more likely to agree that aid to the poor "does more harm than

good" and that "stricter environmental laws and regulations cost too many jobs and hurt the

economy" [40]. Extensive research indicates that job loss related to environmental regulation is

largely offset by jobs created to comply with and enforce the regulation [41].

Public health expenditures per capita are correlated with lower cancer and heart disease death

rates, but not with reported health behaviors other than vaccinations. Since cancer and heart

disease develop over extensive periods, current surveys of behavior may not reflect past changes,

such as the large reduction in smoking [42]. In 2019, a state and local public health agency

survey noted the percentage of particular programs. The most prominent were tobacco, alcohol,

and drugs (74%), emergency preparedness (62%), infectious disease and vaccination (60%), food

safety (48%), obesity/physical activity (45%), and wastewater/sanitation (39%). The agencies

that served larger populations had higher percentages of each effort. State and local health

departments can call on the Centers for Disease Control and Prevention when illnesses of

unknown origin occur. Revenues to support public health departments are diversified – federal

direct and passthrough (27%), local sources (25%), state sources (21%), Medicare/Medicaid

(10%), and others combined (17%). These also vary by the size of the population served; state

sources per capita are used more in less populated areas [43]. Therefore, it is not surprising that

partisan politics is less correlated to public health expenditures than one might think. The anti-

public health rhetoric and threats during the pandemic seem to have accelerated the trend of state

and local public health professionals to retire or seek alternative careers. A study of changes in

state and local public health staff trends from 2017 to 2021 does not portend well for the health

of their states. Nearly half left the agencies, including three of every four younger than thirty-six

[44].

During the first months of the newly elected Trump administration in 2025, Trump issued numerous executive orders that indiscriminately fired thousands of federal employees working in public health and other humanitarian efforts. Thousands of federally funded health-related research grants and contracts were canceled, continuing the anti-science trend in Republican politics [45]. The KFF Health Tracking Poll found that 72 percent of Republicans approved of the cuts [46].

Inferences from the lack of correlation between physicians per capita and death rates other than COVID-19 should be tempered by the possibility that the coefficients may be distorted by collinearity. Many medical prescriptions and procedures have been repeatedly verified for efficacy and safety in controlled clinical trials, but that does not guarantee that they are applied appropriately or that the recipients of treatment comply with instructions. Each must be judged based on evidence of correct application and compliance. The efficacy of medical care is partly offset by medical errors, deaths from which are estimated at more than 200,000 per year in the U.S. [47], competing with unintentional injuries as the third leading cause of death after heart disease and cancer.

The correlation of economic factors with death rates varies by mode of death. Corrected statistically for the other factors, income corrected for cost of living explains no variance. Income inequality is associated with each except for heart disease. Numerous studies have found higher death rates associated with poverty. COVID-19 deaths per capita were substantially higher among the poor and people engaged in essential low-wage work that could not be done remotely [48], but also people employed in health care [49].

A major limitation of this study is the inference of individual behavior from aggregate data. Occasionally, such inferences are found invalid. Also, there is the possibility that inclusion of

one or more unmeasured variables would alter the results. While conclusive inferences of

causation cannot be based on cross-sectional correlations, if prayer were efficacious in

preventing deaths from the conditions examined, the correlations should be negative, not positive

as they are. Of course, prayer does not cause the higher death rates in the more prayerful states,

and increased prayer can be the consequence of chronic disease. An inference that a reduction in

prayer, per se, would reduce disease would be false. The issue is the degree to which people

neglect science-based prevention and treatment and support anti-science public policy because of

religious beliefs. The correlations are of sufficient magnitude to warrant more detailed research

on religious beliefs and practices regarding health maintenance, preventive strategies, treatment,

and politics. Research on the effectiveness of eliciting clergy to promote improved health is also

needed [50].

References

1.   Wozniczka Dl, Demeke HB, Thompson-Paul AM, et al. Real-Time CDC consultation

during the COVID-19 Pandemic—United States, March–July, 2020. International Journal of

Environmental Research and Public Health 18: 7251, 2021.

https://doi.org/10.3390/ijerph18147251

2.   Kaplan S. C.D.C. Labs Were Contaminated, Delaying Coronavirus Testing, Officials Say.

New York Times, May 7, 2020. https://www.nytimes.com/2020/04/18/health/cdc-coronavirus-

lab-contamination-testing.html

3.   Rains SA, Colombo PM, Quick BL, Kriss LA, State mask mandates and psychological

reactance theory: The role of political partisanship and COVID-19 risk in mask adoption and

resistance. Social Science & Medicine 314:115479, 2022.

4.   Robertson LS. Unanticipated behavioral consequences of on-demand COVID-19 testing

policy in US states. Disaster Medicine and Public Health Preparedness. 19: e7, 1–5.

https://doi.org/10.1017/dmp.2024.344

5.   Robertson LS. Roads to COVID-19 Containment and Spread. Second Edition. Amazon,

2025.

6.   Robertson LS. Dysfunctional: Scientists, Fundamentalists, and Politicians. Amazon,

2024.

7.   Cathey L. Timeline: Tracking Trump alongside scientific developments on

hydroxychloroquine. ABC News. August 8, 2020. https://abcnews.go.com/Health/timeline-

tracking-trump-alongside-scientific-developments-hydroxychloroquine/story?id=72170553

8.   Lewis T. How the U.S. pandemic response went wrong—and what went right—during a

year of COVID. Scientific American. 2021. https://www.scientificamerican.com/article/how-the-

u-s-pandemic-response-went-wrong-and-what-went-right-during-a-year-of-covid/

9.   Shear MD. Inside the failure: 5 takeaways on Trump's effort to shift responsibility. New

York Times. 2020. https://www.nytimes.com/2020/07/18/us/politics/trump-coronavirus-failure-

takeaways.html

10.   Robertson, LS. Association of COVID-19 mortality with politics and on-demand testing

in 217 U.S. counties. BMC Public Health. 2021. 21:2203.

https://link.springer.com/content/pdf/10.1186/s12889-021-12063-2.pdf

11. Wang G, Devine RA, Molina-Sieiro G. Democratic governors quicker to issue stay-at-home

orders in response to COVID-19. Leadership Quarterly. 2021.

283 gohttps://pmc.ncbi.nlm.nih.gov/articles/PMC9759985/#:~:text=We%20found%20that%20govern

284 ors'%20political,different%20party%20than%20the%20governor's.

285 12. Mann ME and Hotez P. Science Under Siege. New York: Public Affairs. 2025.

286 13. Jackson C, Newall M, Diamond J, et al. Snapshot of America at the two-year pandemic

287 mark. 2022. https://www.ipsos.com/en-us/news-polls/two-years-of-COVID

288 14. Horney JA, Harjivan A., Stone KW, et al. Threats to public health workers. Public Health

289 in Practice 6, 2023. https://www.sciencedirect.com/science/article/pii/S2666535223000812

290 15. Ballotpedia. Arguments against religious service restrictions during the coronavirus

291 (COVID-19) pandemic, 2020.

292 https://ballotpedia.org/Arguments_against_religious_service_restrictions_during_the_coronaviru

293 s_(COVID-19)_pandemic,_2020

294 16. Reeve E, Guff S, Waldrop E, et al. Many Evangelicals say they won't be vaccinated

295 against Covid-19. Some experts say distrust and misinformation have played a role. CNN. 2021.

296 https://www.cnn.com/2021/04/14/us/covid-vaccine-evangelicals

297 17. Steensland B and Wright EL. American Evangelicals and conservative politics: past,

298 present, and future. Sociology Compass 8:705–717, 2014.

299 https://compass.onlinelibrary.wiley.com/doi/pdf/10.1111/soc4.12175?casa_token=RH8bQoURV2

300 QAAAAA%3Am5qsHjEc-4-

301 LfOC0NTBjI5w3jsIO8L07TP2mDMNmcHpnKQX19AAGYtmgLhLYtQVQMUEDmJguO5HP

302 isRI

18.   Allen LG and Olson SF. Racial attitudes and political preferences among black and white

evangelicals. Politics and Religion 15:631–648, 2022.

19.   Pew Research Center. Party identification among religious groups and religiously

unaffiliated voters. 2024. https://www.pewresearch.org/politics/2024/04/09/party-identification-

among-religious-groups-and-religiously-unaffiliated-voters/

20.   Salmoirago-Blotcher E, Fitchett G, Hovey KM, et al. Frequency of private spiritual

activity and cardiovascular risk in postmenopausal women: the Women's Health Initiative.

Annals of Epidemiology. 23:239-245, 2013. https://doi.org/10.1016/j.annepidem.2013.03.002.

21.   McCaffrey AM, Eisenberg DM, Legedza ATR, et al. Prayer for health concerns: results of

a national survey on prevalence and patterns of use. Archives of Internal Medicine. 164:858–862,

2004. https://jamanetwork.com/journals/jamainternalmedicine/fullarticle/216980

22.   Musik MA, House JS, Williams DR. Attendance at religious services and mortality in a

national sample. Journal of Health and Social Behavior. 45:198-213. 2004.

https://journals.sagepub.com/doi/pdf/10.1177/002214650404500206?casa_token=TZHa4NY--

KEAAAAA:VxmDQ9UT55GraJbjrxNDnpdEnanL_BiuBPFwlqUa7nOPokcs9906GJTOb-

51cJH3yZgaVcd5ItrOyA

23.   MacDorman MF. Race and ethnic disparities in fetal mortality, preterm birth, and infant

mortality in the United States: An overview. Seminars in Perinatology. 35:200-208. 2011.

https://doi.org/10.1053/j.semperi.2011.02.017.

24.   McCaffrey AM, Eisenberg DM, Legedza ATR, Davis RB, Phillips RS. Prayer for health

concerns: results of a national Survey on prevalence and patterns of use. Archives of Internal

Medicine. 164:858–862, 2004.

https://jamanetwork.com/journals/jamainternalmedicine/fullarticle/216980

25.   Lipka M, Wormald B. How religious is your state? Pew Research Center, 2016.

https://www.pewresearch.org/short-reads/2016/02/29/how-religious-is-your-state/?state=alabama

26.   CNN. Results of the 2016 presidential elections in the United States. 2016.

https://www.statista.com/statistics/630799/preliminary-results-of-the-2016-presidential-election/

27.   Moneyrates.com. Best states to make a living in the United States in 2019. https://www-

statista-com.yale.idm.oclc.org/statistics/226377/most-affordable-states-in-the-us/

28.   U.S. Census Bureau. Gini index of income inequality, 2019.

https://data.census.gov/table/ACSDT1Y2019.B19083?q=gini%20&g=010XX00US$0400000&t

p=true&tid=ACSDT1Y2019.B19083

29.   National Center for Health Statistics. Number of physicians in patient care per 100,000

resident population, by state: United States, 2018. Health, United States, 2019.

https://www.ncbi.nlm.nih.gov/books/NBK569310/table/ch2.tab16/

30.   U.S. Census Bureau. Percentage of people in poverty by state, 2019-2020, 2021.

https://www.census.gov/library/publications/2021/demo/p60-

273.html#:~:text=The%20official%20poverty%20rate%20in,and%20Table%20B%2D1).

31.   U.S Census Bureau. The Urban Population as a Percentage of the Total Population by

U.S. Region and State (1790-2020). 2024.

https://en.wikipedia.org/wiki/Urbanization_in_the_United_States

32. Shadac.org. Per person state public health funding. 2019.

https://statehealthcompare.shadac.org/rank/117/per-person-state-public-health-

funding#2,3,4,5,6,7,8,9,10,11,12,13,14,15,16,17,18,19,20,21,22,23,24,25,26,27,28,29,30,31,32,3

3,34,35,36,37,38,39,40,41,42,43,44,45,46,47,48,49,50,51,52/a/27/154/false/location

33. National Institute of Minority Health and Health Disparities, 2018-2021. Mortality.

https://hdpulse.nimhd.nih.gov/data-

portal/mortality/table?cod=247&cod_options=cod_15&ratetype=*&ratetype_options=ratetype_2

&race=00&race_options=race_6&sex=0&sex_options=sex_3&age=001&age_options=age_11&

ruralurban=0&ruralurban_options=ruralurban_3&yeargroup=5&yeargroup_options=yearmort_2

&statefips=00&statefips_options=area_states&county=01000&county_options=counties_alabam

a&comparison=counties_to_us&comparison_options=comparison_state&radio_comparison=are

as&radio_comparison_options=cods_or_areas

34. National Center for Health Statistics. Infant Mortality Rates by State. 2018-2019.

https://www.cdc.gov/nchs/pressroom/sosmap/infant_mortality_rates/infant_mortality.htm

35. Centers for Disease Control and Prevention (CDC). Behavioral Risk Factor Surveillance

System Survey Data. Atlanta, Georgia: U.S. Department of Health and Human Services, 2024.

https://hdpulse.nimhd.nih.gov/data-

36. Brady D, Kohler U, Zheng H. Novel estimates of mortality associated with poverty in the

US. JAMA Internal Medicine. 183(6):618–619. 2023.

https://jamanetwork.com/journals/jamainternalmedicine/fullarticle/2804032

37.   Tiwana MH and Smith J. Faith and vaccination: a scoping review of the relationships

between religious beliefs and vaccine hesitancy. BMC Public Health. 2024. 24: 1806.

https://www.ncbi.nlm.nih.gov/pmc/articles/PMC11227154/

38.   Diab A, Nedda Dastmalchi L, Gulati M and Michos ED. 2023. A heart-healthy diet for

cardiovascular disease prevention: where are we now? Vascular Health and Risk Management,

237-253. https://doi.org/10.2147/VHRM.S379874

39.   Grosso G, Bella F, Godos J, Sciacca S, Del Rio D, Ray S, Galvano F , Giovannucci EL.

Possible role of diet in cancer: systematic review and multiple meta-analyses of dietary patterns,

lifestyle factors, and cancer risk. 2017. Nutrition Reviews, 75(6):405–419.

https://doi.org/10.1093/nutrit/nux012

40.   Pew Research Center. Frequency of prayer. 2024b.

https://www.pewresearch.org/religious-landscape-study/database/compare/frequency-of-

prayer/by/state\

41.   Hafstead MAC and Williams III RC. Jobs and environmental regulation. In Kotchen MJ,

Deryugina T, and Wolfram CD (eds.) Environmental and Energy Policy and the Economy.

Chicago: University of Chicago Press, 2020.

https://www.journals.uchicago.edu/doi/full/10.1086/706799

42.   Burns D. How far we have come in the last 50 years in smoking attitudes and actions.

Annals of the American Thoracic Society. 2014. 11:224-226.

https://doi.org/10.1513/AnnalsATS.201308-258PS

portal/knowledge/map?race=00&race_options=race_6&sex=0&sex_options=sex_3&statefips=0

0&statefips_options=area_us&knowledgetopic=070&knowledgetopic_options=knowledge_9&s

creening=92&screening_options=vax_2&datatype=010&datatype_options=datatypedirect_1

43   National Association of County and City Health Officials. National Profile of Local

Health Departments. 2019. https://www.naccho.org/uploads/downloadable-

resources/Programs/Public-Health-Infrastructure/NACCHO_2019_Profile_final.pdf

44.   Leider JP, Castrucci BC, Robins M, et al. The exodus of state and local public health

employees: separations started before and continued throughout COVID-19. Health Affairs.

2023. https://www.healthaffairs.org/doi/full/10.1377/hlthaff.2022.01251

45.   Hotez PJ. The Deadly Rise of Anti-science. Baltimore: Johns Hopkins University Press.

46.   Sparks G, Montero A, Presiado M, Montalvo III J, Kirzinger A, and Hamel L. KFF

Health Tracking Poll April 2025: Public's View on Major Cuts to Federal Health Agencies. 2025.

https://www.kff.org/medicaid/poll-finding/kff-health-tracking-poll-april-2025-publics-view-on-

major-cuts-to-federal-health-

agencies/#:~:text=Most%20of%20the%20public%20(61,are%20in%20support%20(32%25).

47.   Rodziewicz TL, Houseman B, Vaqar S, et al. Medical Error Reduction and Prevention.

[Updated 2024 Feb 12]. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing;

2024. https://www.ncbi.nlm.nih.gov/books/NBK499956/

48.   Matthay EC, Duchowny KA, Riley AR, et al. Occupation and educational attainment

characteristics associated with COVID-19 mortality by race and ethnicity in California. JAMA

Network Open. 5(4):e228406. 2022. doi:10.1001/jamanetworkopen.2022.8406

49.   Hawkins D, Davis L, Kriebel D. COVID 19 deaths by occupation, Massachusetts,

March 1–July 31, 2020. American Journal of Industrial Medicine. 64:238–244. 2020.

https://doi.org/10.1002/ajim.23227

50.   Bopp, M and Meghan M. Leading their flocks to health? Clergy health and the role of

clergy in faith-based health promotion interventions. Farm Community Health. 2013. 36:182-

192.