Roles of community and sexual contacts as drivers of clade I mpox outbreaks

Hiroaki Murayama[1†], Toshiaki R. Asakura[2,3,4†], Borame L. Dickens[5], Dorothy Boyle[5], Jen Han Foo[5], Shihui Jin[5], Patrick K. Mukadi[6,7,8], Keisuke Ejima[9,10], Sung-mok Jung[11], Akihiro Nishi[12,13], Kiesha Prem[3,5], Audry M. Wakamba[14], Diafuka Saila-Ngita[15], David Niyukuri[16,17,18], Akira Endo[2,3,4,5*]

1. School of Medicine, International University of Health and Welfare, Narita, Japan
2. Department of Infectious Disease Epidemiology, London School of Hygiene & Tropical Medicine, London, UK
3. Centre of Mathematical Modelling of Infectious Diseases, London School of Hygiene & Tropical Medicine, London, UK
4. School of Tropical Medicine and Global Health, Nagasaki University, Nagasaki, Japan
5. Saw Swee Hock School of Public Health, National University of Singapore, Singapore
6. Graduate School of Biomedical Sciences, Nagasaki University, Nagasaki, Japan
7. Department of Clinical Infectious Diseases, Institute of Tropical Medicine, Nagasaki University, Nagasaki, Japan
8. Department of Epidemiology and Global Health, Institut de Recherche Biomédicale, Kinshasa, Democratic Republic of Congo
9. Lee Kong Chian School of Medicine, Nanyang Technological University, Singapore, Singapore
10. National Centre for Infectious Diseases, Singapore, Singapore
11. Carolina Population Center, University of North Carolina at Chapel Hill, Chapel Hill, NC, USA
12. Department of Epidemiology, Fielding School of Public Health, University of California, Los Angeles, CA, USA
13. California Center for Population Research, University of California, Los Angeles, CA, USA
14. Ministry of Health, National Expanded Program for Immunization, Kinshasa, DRC
15. Department of Basic Sciences, Faculty of Veterinary Medicine, Université de Kinshasa (UNIKIN), Kinshasa, DRC
16. Doctoral School, University of Burundi, Bujumbura, Burundi
17. Department of Mathematics, University of Burundi, Bujumbura, Burundi
18. The South African Department of Science and Technology-National Research Foundation (DST-NRF) Centre of Excellence in Epidemiological Modelling and Analysis (SACEMA), Stellenbosch University, Cape Town, South Africa

* Correspondence to: aendo@nus.edu.sg
† These authors contributed equally.

Abstract

Initial investigation into the emerging mpox outbreak of novel clade Ib in eastern Democratic Republic of the Congo has identified signs of sustained human-to-human transmission and epidemiological links to sexual contacts involving female sex workers (FSWs)[1], which have not been observed in previous clade Ia outbreaks. Using mathematical models incorporating age-dependent contact patterns, we quantified the role of frequent sexual interactions as opposed to community contacts in clade Ib's dynamics and found that this additional mode of transmission could explain its increased outbreak potential compared with clade Ia. As with the globally-circulating clade IIb, transmitted predominantly among men who have sex with men[2], our findings reinforce the importance of protecting key population groups, specifically FSWs for clade Ib, in controlling ongoing mpox outbreaks.

Main text

Since September 2023, a sustained local transmission of a novel subclade of monkeypox virus (MPXV), classified as clade Ib, has been observed in the Democratic Republic of the Congo (DRC) along with its spread to multiple other countries[3]. In response to the escalating situation, the World Health Organization (WHO) declared a public health emergency of international concern (PHEIC) on 14 August 2024[4]. Clade Ib was first identified from a cluster of cases in the Kamituga health zone of the South Kivu province, eastern DRC, where an initial investigation found epidemiological links to sexual contacts and a significant proportion of female sex workers (FSWs) among reported cases[5]. Genetic analyses have indicated a distinct profile from clade Ia circulating in other parts of DRC, characterised by the accumulation of apolipoprotein B mRNA editing enzyme catalytic subunit 3 (APOBEC3)-associated mutations[5]. Those mutations suggest sustained human-to-human transmission and were also observed in the globally-circulating MPXV clade IIb, which primarily affected men who have sex with men (MSM)[2]. Over the summer of 2024, clade Ib showed steady growth within many health zones in the North and South Kivu provinces and introductions to other provinces of DRC[6]. Additionally, international spread to neighbouring countries (Burundi, Uganda, Rwanda, Kenya, Zambia and Zimbabwe) and beyond the African continent (Sweden, Thailand, India, Germany and UK) had been confirmed by October 2024[7], of which Burundi had reported the largest number of cases (1,509 as of 27 October 2024[8]). Thereafter, the mpox outbreak has continued to expand predominantly within Africa, with nearly 20,000 cases in 2025 alone as of May[9]. The number of countries ever reporting clade Ib cases have increased to 12 within Africa and 29 globally, including sporadic suspected local transmission events outside of Africa[9].

MPXV clade Ia has been circulating in DRC and other Central African countries for over 50 years[10] via zoonotic transmission followed by self-limiting chains of human-to-human transmission through close contact, mostly within households[11]. Since the cessation of the smallpox vaccination programme in 1980, which had provided cross-protection against mpox[12–17], DRC has seen a consistent rise in suspected mpox cases[18,19], from 3.0 per 100,000 people in 2010 to 11.5 per 100,000 people in 2023[20]. Despite this increase, zoonotic spillover remained as the primary driver of outbreaks[21,22], and there has been no evidence of large clade I outbreaks solely sustained by human-to-human transmission before the emergence of clade Ib in 2023. In contrast, the clade Ib outbreak in the South Kivu province has shown a clear exponential growth trend, suggesting human-to-human contact as the primary mode of transmission[23,24]. The rapid rise of clade Ib cases with a potential increase in the contribution of human-to-human transmission relative to clade Ia is of growing global concern, especially given that the outbreak is not restricted to specific at-risk populations but is affecting broader groups, including children[25].

The extensive clade Ib case profiles including both children and adults along with
documented occupational risks among FSWs may reflect its mixed transmission routes
through regular contacts in the community and through sexual contacts. Sexual contact as a
novel route of transmission has also been documented in the ongoing global mpox outbreak
caused by clade IIb since 2022 and was suggested to have contributed to the rapid spread
among MSM[26,27]; meanwhile, sustained transmission of clade IIb through community or
heterosexual contact routes have not been observed and cases were mostly restricted to the
MSM group[28]. If clade Ib, transmissible through heterosexual contact, continues to grow into
another international mpox outbreak following clade IIb, the potentially more emphasised
role of community contacts in its transmission than clade IIb may enable it to affect a larger
susceptible population and pose challenges for control. Although the involvement of sexual
contacts among FSWs and their clients has been well documented in the initial spread of
clade Ib in the Kamituga health zone in South Kivu[25], little is currently understood on the
relative contribution of community and sexual contacts to subsequent clade Ib dynamics in
broader settings not limited to Kamituga, and how they may explain the distinct
epidemiological features of clade Ib compared to clade Ia. With the gradual expansion of
clade Ib within DRC and beyond borders, analytical efforts to identify its plausible
transmission mechanisms and priority groups for prevention should leverage the best
available evidence, if limited, to inform and support effective control policies.

In this study, we hypothesised that the rapid growth of the clade Ib outbreak could be
explained by its entry into a key population group with a high rate of sexual contact. We
developed a mathematical model for MPXV transmission dynamics incorporating both
community contact route and sexual contact route involving FSWs. We first used a model
incorporating only a community contact route to replicate the age profiles of the current and
historical clade Ia outbreak datasets. We then expanded the model to include a sexual contact
route to quantify the contribution of each route to the overall clade Ib transmission dynamics.

Characterising community contact transmission patterns in historical and current clade
Ia outbreaks
The age-dependent contact matrix, whose entries represent the average number of contacts
(typically defined as either conversational[a] or physical contacts) that an individual of a
specific age group has with other age groups per unit time, is a common tool for modelling
transmission dynamics of directly-transmitted infections across multiple age groups.
Assuming that the risk of transmission is proportional to the frequency of relevant types of
contacts one experiences with infectious individuals, the contact matrix is associated with the
so-called next generation matrix, a matrix that is multiplied by a vector containing the
number of cases by age groups in one generation to produce a vector for the next
generation[29]. During the exponential phase of an outbreak, the age distribution of cases would
quickly converge to the dominant eigenvector of the next generation matrix and the dominant
eigenvalue would define the effective reproduction number[29]. We modelled the next
generation matrix by combining a contact matrix informed by empirical contact survey data
from Zimbabwe[30] and age-dependent susceptibility that we assumed to reflect potentially
higher susceptibility among children aged 0–4 years[31] and lower susceptibility among
smallpox-immunised birth cohorts born in or before 1980[12–17].

This simple model, validated by hold-out data (empirical Kullback-Leibler divergence <

---

[a] Contact involving a minimum threshold of words exchanged during a two-way conversation without any physical contact

0.01; Extended Data Table 2 and Figure S1), well described the age distribution for both the
historical clade Ia cases in the Tshuapa province (from a well-documented clade Ia outbreak
between 2011-2015[32]), DRC and the recent cases in six provinces of DRC where clade Ia has
been endemic, with a decade's shift in the age of smallpox-immunised cohorts from 34 and
above to 45 and above (Figures 1A and B). Our model employed four contact matrices with
different contact definitions; since all four models were overall similar in goodness-of-fit, we
used model averaging[33] to pool the model outputs (Table 1). The estimated susceptibility
among children aged 0-4 (model averaging estimate: 1.5 [95% credible interval: 1.05–2.1])
and smallpox-immunised cohort (0.26 [0.17–0.38]) relative to other age groups was
consistent with literature: ratios of 1.3–1.6 between the historical secondary attack risk
estimates for clade Ia among age 0–4 and 5–14[31] and the estimated effectiveness of past
smallpox vaccines against clade IIb of around 70–75%[17,34] combined with high historical
coverage in DRC (89–97%)[35].
155

Figure 1. Observed and modelled age distributions of mpox cases. (A) Clade Ia cases with suspected human-to-human exposure from the Tshuapa province, DRC, 2011–2015. Those aged 34 and older were assumed to be smallpox-immunised. (B) Cases from clade Ia-endemic provinces in DRC, January–mid

August 2024. Those aged 45 and older were assumed to be immunised. (C)–(F) Clade Ib cases from multiple settings in DRC, January–mid August 2024 and Burundi, mid September–October 2024. Those aged 45 and older were assumed to be immunised. Half-transparent lines represent the model fit without assuming transmission over sexual contact. Dots and whiskers represent the observed age distributions among cases and their 95% confidence intervals.

Table 1. Relative contribution of sexual contacts to the transmissibility of clade Ib in DRC and Burundi

Model	Contact type	Model weights[*]	Estimated relative susceptibility[**]		Estimated fraction of R_{eff} attributable to sexual transmission[***]			
			Children aged 0–4	Immunised cohort	North and South Kivu	Kamituga health zone	Other health zones, South Kivu	Burundi
1	All contacts[†]	0.21	1.9 (1.6–2.2)	0.27 (0.18–0.38)	41% (36–45%)	60% (54–66%)	14% (9–21%)	16% (12–20%)
2	Physical contacts only	0.33	1.6 (1.4–1.9)	0.30 (0.21–0.42)	42% (37–46%)	61% (55–66%)	15% (9–21%)	16% (13–20%)
3	All contacts at home	0.29	1.3 (1.1–1.6)	0.22 (0.15–0.32)	43% (38–47%)	63% (57–69%)	12% (5–19%)	15% (11–20%)
4	Physical contacts at home	0.17	1.2 (0.96–1.4)	0.25 (0.17–0.36)	42% (38–47%)	63% (57–68%)	12% (4–19%)	15% (11–19%)
Model average	—	—	1.5 (1.05–2.1)	0.26 (0.17–0.38)	42% (37–47%)	62% (55–68%)	14% (6–21%)	16% (11–20%)

Posterior median estimates and 95% credible intervals are shown.
* Based on the Watanabe-Akaike weights[36]
** Referenced to pre-immunisation cohort aged 5 or older
*** Defined as the relative reduction in R_{eff} when sexual contacts were excluded from the next generation matrix
† Physical + conversational contacts. Physical contacts are defined as any skin-to-skin contact and conversational contacts are defined as in-person contacts with a two-way exchange of at least three words.

Modelling sexual contact transmission for clade Ib outbreaks

Assuming that the community contact transmission patterns relevant to clade Ia would also apply to clade Ib transmission, we modelled a sexual contact transmission route among heterosexual individuals with high sexual activity (who we assume primarily represent FSWs and clients) alongside the community contact matrix to represent the transmission dynamics of clade Ib outbreaks. We considered simple proportionate mixing between age groups, where sexual partners are chosen randomly among those in the high-activity group of the opposite sex. We parameterised the age-dependent proportion of male and female high-activity individuals (aged 15–19, 20–29, 30–39 and 40–49) and their mean neighbour degrees[37] over the infectious period of mpox to construct the age-, sex- and transmission-route (community vs sexual contact)-dependent next generation matrix. We fitted our model to the age-sex distribution of clade Ib cases in multiple geographical settings: the North and South Kivu provinces (referred to as the Kivus hereafter), the Kamituga health zone, other health zones of South Kivu, and Burundi. Across these geographical settings, we assumed that age-dependent relative susceptibility was identical to what we estimated from clade Ia datasets.

Our model reproduced the observed age-sex distributions of clade Ib cases generally well, including age groups not assumed to be involved in high-activity sexual contact (Figure 1C–F). This supports the hypothesis that the transmission dynamics of clade Ib could be explained by a combination of community contact transmission patterns similar to clade Ia and additional sexual contact transmission among the high-activity group. The estimated role of transmission over sexual contact varied between geographical settings, potentially

reflecting different sexual behaviours among high-activity individuals. The estimated relative
contribution of sexual contact to the overall effective reproduction number was higher in the
Kivus (model averaging estimate: 42% [95% credible interval: 37-47%]), which is most
likely attributable to the dynamics in Kamituga health zone (62% [55-68%]) (Table 1). In the
other health zones of South Kivu (14% [6-21%]) and Burundi (16% [11-20%]), sexual
contact was suggested to play a smaller (although non-negligible) role. A recent study on the
Kamituga outbreak associated clade Ib spread in the region with a high number of bars
serving as places for commercial sex services in the context of the local gold mining
industry[38]. Frequent engagement in commercial sex associated with mining work[39] in
Kamituga and potentially other nearby mining areas[38] may account for the distinct
epidemiological characteristics of clade Ib in the Kivus. These differences in the relative roles
between community and sexual contacts could shape contrasting age-dependent transmission
patterns between regions. We reconstructed the frequency of transmission across age groups
from our modelled next generation matrix assuming proportionate mixing (Figure 2A).
Compared with clade Ia in the endemic provinces, clade Ib in the Kivus was estimated to be
more frequently transmitted among sexually-active age groups (Figure 2A, left vs middle).
Such excess in estimated transmission among the sexually-active age groups was also present
but less pronounced in Burundi (Figure 2A, right).

Linking transmission patterns and outbreak potential of clade I

Our models allowed us to project the time-evolving effective reproduction numbers (R_y) of
MPXV clade Ia and Ib (Figure 2B). We constructed R_y to reflect the transmission potential in
a hypothetical 'pre-outbreak' state, i.e. accounting for immunity from the historical smallpox
immunisation but not from infections or mpox vaccination, to focus on the effect of changing
smallpox immunity landscape associated with demographic shift. The projection suggested a
9.8% increase in R_y of clade Ia over the last decade in DRC, reflecting the ageing of the
smallpox-immunised cohorts. When assuming an R_y of 0.82 for clade Ia in DRC, 2015[21], this
increase translates into an R_y of 0.90 in 2024, which aligns with our estimates from the pre-Ib
mpox incidence data in DRC (Figure S2). As R_y gradually approaches 1, the expected final
size of an outbreak initiated by a single zoonotic infection, given as $1/(1-R_y)$[40], would have
increased from 5.6 in 2015 to 10.0 in 2024, which may also explain the recent rise in clade Ia
cases in the endemic areas in DRC[41].

If the same community contact transmission patterns estimated for clade Ia also apply to
clade Ib, the overall R_y in the Kivus combining both community and sexual contact routes
would be around 1.5 in our projection. This is similar to our estimate of the initial
reproduction number from suspected clade Ib cases in South Kivu (Figure S2). The increase
in R_y attributable to sexual contact, resulting in an overshoot of the epidemic threshold of 1,
could explain the stable exponential growth of clade Ib cases observed in the Kivus over the
summer of 2024. In Burundi, our model projected that R_y for clade Ia would be 4% higher
than DRC in 2024 reflecting a slightly different population age distribution. With an
additional sexual contact transmission route, R_y for clade Ib in Burundi may have also
exceeded the epidemic threshold of 1 (with a point estimate of 1.08), which explains the
major spread of clade Ib in Burundi, observed the earliest among the countries neighbouring
DRC.

Focused/mass vaccination strategies

Using the Kivus as a case study, we compared the possible impact of FSW-focused and mass
vaccination strategies (Figure 2C). Given the age eligibility for vaccines (18 years and above
for JYNNEOS[42]) and the previously smallpox-immunised cohort, we considered adults aged

20–39 as the primary group for mass vaccination—our results also suggested that this group
is more likely to contribute to onward transmission in the Kivus (Figure 2A). Under current
global vaccine shortages, prioritising FSWs, typically accounting for < 1% of the total
population[43], may allow for the efficient use of available supply. In the Kivus, where sexual
contact was estimated to play a substantial role in the overall dynamics, achieving high
vaccine uptake among FSWs would reduce the doses required among general adults for
control (Figure 2C). Realistically, both strategies may be pursued in tandem, considering
uncertainty and a possible shift in the relative role of sexual and community contact
transmission, as well as challenges in approaching FSWs facing social barriers and stigma[44].
Regardless of the vaccine target population, clear and supportive communication and de-
stigmatisation are essential[45]. Existing community support networks for FSWs and other
populations at risk, including HIV/AIDS programmes, could play a key role in public health
outreach[45,46].
261

Conclusions
This study quantified the role of community and sexual contacts in MPXV clade I outbreaks.
The results suggest that past and current clade Ia outbreaks are primarily driven by
community contacts, whereas clade Ib is additionally transmitted through contacts among
individuals with high sexual activity such as FSWs and clients. This additional transmission
route may be responsible for the increased effective reproduction number of clade Ib in the
Kivus compared with clade Ia in other endemic regions. The estimated relative contribution
of sexual contact was particularly pronounced in the Kamituga health zone and less so
elsewhere, suggesting that the current epidemiology of clade Ib in the Kivus may be largely
attributable to sexual contact patterns specific to Kamituga and that generalisability to other
settings should be explored in conjunction with the local sociobehavioural contexts. Our
results also suggest that, with the ageing of smallpox-immunised cohorts over decades, the
reproduction number of MPXV clade I has already been approaching the epidemic threshold
of 1 even without additional involvement of sexual contact. In this regard, a small increment
in transmission pathways, including but not limited to sexual contact, may sustain a clade I
outbreak, particularly in countries with lower historical vaccine coverage.
278

The current MPXV clade Ib outbreak has exhibited a distinct epidemiological pattern from
past clade Ia outbreaks, characterised as more rapid spread involving a novel mode of human-
to-human transmission through sexual contact. A similar sudden emergence of sustained
human-to-human transmission was the nature of the global clade IIb outbreak in 2022
onwards, which was primarily (if not exclusively) transmitted through sexual contacts among
MSM[47]. Our previous modelling work offered a plausible explanation for this transition that
clade IIb had established itself in a densely-connected portion of the MSM sexual network,
allowing for the rapid spread that was not sustained elsewhere[26]. Likewise, the emergence of
clade Ib may be another example of MPXV's entry into a key population group where it can
sustain itself. Whilst extensive transmission among FSWs and their clients was not
specifically reported in the clade IIb global outbreak, sexual transmission of clade II
involving FSWs had already been reported from 2017[48] and the potential risk has been
discussed since the initial phase of the global outbreak[49,50]. It remains unclear whether only
clade Ib is able to be readily transmitted within FSW-related networks or it was just the first
to have an opportunity among the clades with similar potential. Recent studies on co-
circulating clades Ia and Ib in Kinshasa, DRC[51] and on the clade IIb outbreak in Sierra
Leone[52] both report overrepresentation of sexually active age groups of both sexes among
cases, as was typically observed in clade Ib outbreaks, although without direct evidence of
transmission through sexual contacts. Further investigation is warranted to assess and monitor

the potential shift in the role of sexual contacts in broader mpox outbreak settings.
Regardless, investigation and control efforts for mpox continue to require a tailored and
inclusive approach, with a particular focus on the most vulnerable groups including children,
MSM, FSWs and other key populations at sexually transmitted infection (STI) risks[53,54].
302
Limitations & sensitivity analyses
This study holds several limitations. First, we assumed that contact survey data sufficiently
represented community contacts relevant to MPXV transmission. Some forms of
transmission, including fomite or nosocomial routes, may not be fully captured by existing
contact data from population samples. Contact matrices also represent the mean contact rates
between age groups and neglect possible heterogeneity beyond this. We however believe our
eigenvector approach with hold-out validation (Figures S1C and S1E) helped to ensure
robustness. We used the empirical contact data from a different country (Zimbabwe) as a
proxy since neither empirical contact data from DRC nor Burundi was available.
Nevertheless, the model using Zimbabwe contact data outperformed the model using a
synthetic contact matrix for DRC (Extended Data Table 2 and Figure S1) and our findings
remain overall consistent between these models (Figure S6).
315
Second, our model of the sexual contact route for clade Ib was simplistic due to limited
sexual behaviour data, e.g. proportionate mixing with age-invariant contact rates. However,
the estimated relative contributions of the community and sexual contact routes, the primary
interest of this study, would be relatively robust to these assumptions as long as our model
can capture the excess in mixing patterns attributable to sexual contacts (Figure S7). We
assumed sexual contacts outside of the high-activity groups were captured by contact survey
data and had limited contributions to the difference in dynamics between clades Ia and Ib.
Despite our sensitivity analyses, additional data on local sexual behaviours including the
detailed distribution of sexual engagement rates in FSWs and clients could further inform the
analysis. We also limited our analysis to heterosexual contacts and did not consider MSM
networks. The reported clade Ib case profile with near-even sex balance as of the time of
analysis (Table S1) does not suggest significant transmission within the MSM group[55], and
our additional analysis also suggests that infection- or vaccine-derived immunity in this group
may lower the risk of sustained clade Ib transmission in populations previously affected by
the global clade IIb outbreak (Figure S9). However, this does not exclude the future
possibility of clade Ib outbreaks among MSM, e.g. in countries without previous clade IIb
circulation (including DRC and Burundi[56]) or after herd immunity is lost[57].
333
Third, our model primarily focused on a few epidemiological variables (age, sex and sexual
activity) as key population attributes and other potential risk factors including malnutrition[58],
co-infection with other STIs (particularly HIV[59]) and living conditions were not explicitly
considered. Although some of these factors may be indirectly accounted for by the modelled
components, such as age-dependent susceptibility and sexual or community contact matrices,
future studies are required to better characterise the role of those risk factors in mpox
outbreaks. Biological factors have not been considered either, including possible viral
adaptation that might breach our assumption of the same community transmission patterns
between clades Ia and Ib. Although our model suggested that the increase in the estimated
reproduction number for clade Ib can be explained without requiring evolutionary changes,
this does not necessarily exclude their potential role in the clade Ib dynamics.
345
Finally, case data used for the analysis may have been subject to uncertainty and biases due
to challenges in healthcare access, testing or reporting in an outbreak, especially when

coinciding with civil conflict and natural disasters[60,61]. Most importantly, the potential
underreporting of cases among FSWs, who are facing criminalisation and stigma[62] may have
resulted in an underestimation of the relative contribution of sexual contacts to the clade Ib
dynamics.
352
353

354
Figure 2. Transmission dynamics of MPXV clade I outbreaks. (A) Age-specific transmission patterns
inferred for clade Ia in endemic provinces of DRC and clade Ib in North and South Kivu, and Burundi.
Colours represent daily transmission rates between infector-infectee pairs in given age groups according to
the estimated next generation matrices. (B) Projected effective reproduction number (R_y) for clades Ia and
Ib in DRC (blue lines) and Burundi (red lines), 2010–2030. Lines and shaded areas represent median
estimates and 95% credible intervals of R_y. The estimated reproduction number of 0.82 from clade Ia case
data between 2013-2017 in Tshuapa province, DRC[21] was used as a reference value for 2015 (the midpoint
of the study period). The empirical estimates of the effective reproduction number from the DRC incidence
data in the pre-clade Ib period in 2023 (pre-Ib) and from the South Kivu incidence data (SK) in 2024, as
proxy data for clade Ia and Ib, respectively, are displayed as dots for comparison. Paired dots (jittered
horizontally for visual aid) denote different serial interval distributions used, which were estimated

elsewhere from two separate sets of infector-infectee pairs (distribution 1: mean 17.5 days; distribution 2:
mean 11.4 days)[63]. (C) Contour plots for estimated impact of different combinations of FSW-focused and
mass vaccination strategies on the effective reproduction number in the Kivus.
369
Methods
Data sources
We used the numbers of confirmed mpox cases by age and sex in both historical and current
outbreaks from public sources[8,32,64,65]. We used reported cases from 1 January to 18 August
2024 for the current outbreak in DRC for its stable growth trend and the data availability at
the subprovincial level (Kamituga and other health zones of South Kivu) during that period.
For Burundi which only reported cases from late July 2024 onwards, we used case data from
16 September to 27 October, roughly corresponding to the second half of the stable growth
phase until its peak, to exclude potential influences of reporting bias or initial transients.
Further details of the datasets are available in Extended Data Table 1. Weekly incidence data
for DRC and South Kivu were collected from public reports[23,24,66]. Empirical contact survey
data from Manicaland, Zimbabwe[30] was retrieved via {socialmixr} R package[67]. Population
age distributions in DRC, Burundi and Manicaland from relevant years were collected from
censuses (where available)[68,69] and the World Population Prospects 2024[70]. We also used
synthetic contact matrix data published by Prem et al.[71].
385
386 **Extended Data Table 1. MPXV clade I outbreak datasets.**

ID	Location	Year	Clade	Exposure type	N	Reference
1	Tshuapa province, DRC	2011-2015	Ia	Zoonotic*	308	[32]
2	Tshuapa province, DRC	2011-2015	Ia	Human-to-human**	279	[32]
3	Six endemic provinces***, DRC	2024†	Ia	Not classified	482	[64]
4	North and South Kivu provinces, DRC††	2024†	Ib	Not classified	1,377	[64]
5	Kamituga health zone, South Kivu††	2024†	Ib	Not classified	725	[64]
6	Other health zones, South Kivu††	2024†	Ib	Not classified	510	[64]
7	Burundi	2024†††	Ib	Not classified	945	[8,65]

* Defined by reporting only exposure to an animal within 3 weeks before onset
** Defined by reporting only exposure to a human with a rash illness only within 3 weeks before onset
*** Equateur, South and North Ubangi, Sankuru, Thsopo and Tshuapa
† As of 18 August 2024
†† North and South Kivu provinces data also contains cases in Kamituga and other health zones of South
Kivu.
††† 16 September–27 October 2024
394
395 *Model structure*

396　We assumed that the observed number of mpox cases by age or by age and sex follows a
397　multinomial distribution with probabilities proportional to the elements of the dominant
398　eigenvector of the next generation matrix.

$$i \sim \text{Multinom}\left(\sum i, v(K)\right), \tag{1}$$

where $v(K)$ is the normalised dominant eigenvector of the next generation matrix K. We
constructed the next generation matrix as a product of transmissibility β, age-dependent
relative susceptibility σ_a and a contact matrix C. We used age-disaggregated case data and
contact matrices for the analysis of clade Ia; i.e. $C = \{c_{ab}\}$, where c_{ab} represents the mean
number of community contacts (conversational and/or physical, depending on the type of
contact matrix used) an individual from age group b has per unit time with those from age
group a. We used multiple contact matrices potentially relevant to mpox transmission
(Extended Data Table 2). For each contact matrix, β was adjusted so that the dominant
eigenvector of the next generation matrix estimated for the historical Tshuapa dataset is equal
to the previously estimated reproduction number of 0.82[21].

For the analysis of clade Ib, we used age-and-sex-disaggregated case data and expanded the
contact matrix to account for heterosexual transmission dynamics in addition to community
contacts. We stratified male and female populations into high and low sexual activity groups.
Following the approach in Endo et al.[26], we further stratified cases in the high-activity groups
by their route of exposure (community- vs sexually-associated exposures) and allowed
sexually-acquired cases to have higher risks of onward sexual transmission because of a
substantial variance in sexual behaviour[72,73]. Namely, in a highly heterogeneous sexual
network, transmissions through sexual contact tend to concentrate among individuals with the
highest rates of sexual activity; this renders sexual exposure an indicator of even higher
contact rates than the average among the high-activity groups. We assumed that this variance
effect is negligible among low-activity groups (such that stratification by exposure route can
be disregarded) and that their sexual contacts are reflected in the existing contact matrices,
whose definition of contacts includes sexual contacts. Despite its simplicity, our approach of
stratifying sexual contacts by two activity classes and two exposure routes could well capture
the stationary growth pattern of mpox transmission (see Supplementary Information for a
proof-of-concept network model).

Let subscripts H and L denote high- and low-activity groups and M/m and F/f represent male
and female cases with sexual (upper-case) and community exposures (lower-case),
respectively. Vectors i_X represent the age-stratified number of mpox infections in category X.
The generation-wise reproduction process of i_X's is then described in a block matrix format
(each block is an 8-by-8 matrix accounting for age group stratification) as:

$$\begin{bmatrix} i_{M_H}^{t+1} \\ i_{m_H}^{t+1} \\ i_{m_L}^{t+1} \\ i_{F_H}^{t+1} \\ i_{f_H}^{t+1} \\ i_{f_L}^{t+1} \end{bmatrix} = B_\sigma \begin{bmatrix} O & O & O & S_{MF} & \Sigma_{MF} & O \\ PC_{MM} & PC_{MM} & PC_{MM} & PC_{MF} & PC_{MF} & PC_{MF} \\ (I-P)C_{MM} & (I-P)C_{MM} & (I-P)C_{MM} & (I-P)C_{MF} & (I-P)C_{MF} & (I-P)C_{MF} \\ S_{FM} & \Sigma_{FM} & O & O & O & O \\ QC_{FM} & QC_{FM} & QC_{FM} & QC_{FF} & QC_{FF} & QC_{FF} \\ (I-Q)C_{FM} & (I-Q)C_{FM} & (I-Q)C_{FM} & (I-Q)C_{FF} & (I-Q)C_{FF} & (I-Q)C_{FF} \end{bmatrix} \begin{bmatrix} i_{M_H}^{t} \\ i_{m_H}^{t} \\ i_{m_L}^{t} \\ i_{F_H}^{t} \\ i_{f_H}^{t} \\ i_{f_L}^{t} \end{bmatrix}, \tag{2}$$

where O and I are zero and unit matrices, respectively. S_{MF} and S_{FM} are block contact
matrices representing heterosexual transmission from sexually-acquired high activity
individuals. Similarly, Σ_{MF} and Σ_{FM} represent heterosexual transmission from community-

acquired high activity individuals. Again, note that we expect S_X to be larger than Σ_X because
sexually-acquired cases are more likely to have more sexual partners than the population
average. $P = (p_a)$ and $Q = (q_a)$ are diagonal matrices representing the age-dependent
proportions of high-activity individuals for males and females, respectively. C_X is a sex-
specific contact matrix and is assumed to be half of the sex-aggregated contact matrix
hereafter for simplicity. We define B_σ as a diagonal block matrix whose diagonal blocks are
an 8-by-8 diagonal matrix of the age-specific susceptibility σ_a multiplied by transmissibility
β.

Noting that for $t \geq 1$, $i_{m_H} = P(i_{m_H} + i_{m_L})$ and $i_{f_H} = Q(i_{f_H} + i_{f_L})$, the above relationship can
be simplified into

450
$$
\begin{bmatrix} i_{M_H}^{t+1} \\ i_m^{t+1} \\ i_{F_H}^{t+1} \\ i_f^{t+1} \end{bmatrix} = B_\sigma \begin{bmatrix} O & O & S_{MF} & \Sigma_{MF}Q \\ C_{MM} & C_{MM} & C_{MF} & C_{MF} \\ S_{FM} & \Sigma_{FM}P & O & O \\ C_{FM} & C_{FM} & C_{FF} & C_{FF} \end{bmatrix} \begin{bmatrix} i_{M_H}^t \\ i_m^t \\ i_{F_H}^t \\ i_f^t \end{bmatrix},
\tag{3}
$$

452 where $i_m = i_{m_H} + i_{m_L}$ and $i_f = i_{f_H} + i_{f_L}$.

454 We assumed proportionate mixing to model S_{MF}, S_{FM}, Σ_{MF} and Σ_{FM}; further details on
455 parameterisation are described in the next section.

457 We constructed a multinomial likelihood for the observed number of male and female cases
458 by age group:

459
$$
i \sim \text{Multinom}\left(\sum i, [i_{M_H} + i_m; i_{F_H} + i_f]\right),
\tag{4}
$$

460 where $[i_{M_H} + i_m; i_{F_H} + i_f]$ is a vector obtained by sex-wise aggregation of the normalised
461 dominant eigenvector of the next generation matrix in Equation 3.

463 **Extended Data Table 2. Community contact matrices used for analysis.**

ID	Contact type	Contact setting	Data type	Country	Reference	Hold-out validation loss*	Hold-out empirical KLD**
1	All	All	Empirical	Zimbabwe	[30]	47.23	0.009
2	Physical only	All	Empirical	Zimbabwe	[30]	47.18	0.009
3	All	Home	Empirical	Zimbabwe	[30]	44.37	0.006
4	Physical only	Home	Empirical	Zimbabwe	[30]	46.40	0.008
5	All	All	Synthetic	DRC	[71]	79.60	0.04
6	All	Home	Synthetic	DRC	[71]	62.68	0.03
7	All	Home	Synthetic	Burundi	[71]	NA***	NA***
8	All	All	Synthetic	Burundi	[71]	NA***	NA***

464 * Mean negative log-likelihood times two for the validation dataset (Dataset 3, Extended Data Table 1) not
465 used in training
466 ** Kullback-Leibler divergence[74] from the empirical distribution of the validation dataset to the mean
467 modelled distribution

468   *** Not used for validation; only for the sensitivity analysis (Figure S6)
469

Model fitting
Our model fitting process was two-fold. We first fitted and validated the model of community
contact transmission using the age distributions of cases from the historical and current clade
Ia outbreaks in DRC. We used this model to estimate age-dependent susceptibility parameters
σ_a for the community contact matrix. We then constructed an expanded contact matrix
including sexual contacts and calibrated it to the age-sex distributions of cases from clade Ib
outbreaks in DRC and Burundi in order to estimate additional parameters that describe
transmission dynamics of clade Ib through community and sexual contact routes.
478

(i) Analysis of clade Ia outbreaks
As in the previous section, we defined the next generation matrix $K=\{k_{ab}\}=\beta\sigma_a c_{ab}$. The
contact matrix c_{ab} was constructed from existing datasets (Extended Data Table 2), with
adjustment for age structures between the source and modelled populations where different
(using the density correction method described by Arregui et al.[75]). Based on data
availability, we used 6 age groups (0-4, 5-10, 10-19, 20-29, 30-39 and 40+) for the 2011–
2015 Tshuapa data and 8 age groups (0-4, 5-10, 10-14, 15-19, 20-29, 30-39, 40-49 and 50+)
for the 2024 DRC endemic provinces data. When the age groups in the case data were coarser
than the contact matrix, the elements were merged accordingly[76]. Transmissibility β is a
nuisance scaling parameter and was not estimated. We assumed that children aged 0-4 years
may have potentially higher susceptibility to infection and that adults born before the
cessation of the smallpox vaccine in 1980[14] may have lower susceptibility. To account for
this, we parameterised σ_a as
492

493
$$\sigma_a = \begin{cases} \sigma_0 & (\text{aged } 0-4) \\ 1-\varepsilon & (\text{smallpox immunised}) \\ 1 & (\text{otherwise}) \end{cases}. \tag{5}$$

Here, ε represents the net vaccine protection among the vaccinated cohorts, i.e. the product of
the vaccine effectiveness and coverage in the historical campaign (also referred to as the
effective vaccine coverage). The susceptibility for the age group that contains both
vaccinated and unvaccinated cohorts (ages 30-39 in the historical Tshuapa dataset and the
ages 40-49 in the 2024 outbreak datasets) was specified as the weighted average between 1-ε
and 1, based on the proportion of individuals born in or before 1980 in those age groups[70] as
of 2013 (the midpoint of the Tshuapa dataset period) or 2024. Given the high historical
vaccination coverage in DRC (89–97%)[35], the estimated vaccine protection ε would provide
an approximation (or at least a reasonable lower bound) for the effectiveness against clade I
mpox. We estimated σ_0 and ε by the Bayesian importance sampling method[77] based on the
multinomial likelihood in Equation 1 and obtained median estimates and 95% credible
intervals (CrIs).
507

The exploratory model development was done using the historical Tshuapa dataset. The
resulting model, described above, was then validated using the 2024 DRC endemic provinces
dataset. We estimated two parameters σ_0 and ε from the historical Tshuapa dataset alone and
compared the model outputs (after accounting for a decade's shift in the age of vaccinated
cohort) with the age distribution from the (hold-out) 2024 DRC endemic provinces dataset
(Figure S1). The model outputs showed good overall concordance with the validation data.
We obtained final parameter estimates from the joint estimation using both datasets for the
subsequent analyses.

(ii) Analysis of clade Ib outbreaks

We parameterised the block components representing sexual contacts among high-activity

individuals Σ_{MF}, Σ_{FM}, S_{MF} and S_{FM} in Equation 3 as follows. We assumed age groups of 15-

19, 20-29, 30-39 and 40-49 years potentially contain high-activity populations, i.e. $p_a = q_a = 0$

outside those age groups. We assumed a simple proportionate mixing assumption (i.e. no age

assortativity) where each high-activity individual makes sexual contacts at a specified rate

over the infectious period of MPXV, which are randomly assigned to other high-activity

individuals of the opposite sex irrespective of age. We denote the mean sexual contact rates

among high-activity females and males infected through community contact as v_F and v_M;

similarly, we denote the mean sexual contact rates among those infected through sexual

contact as w_F and w_M. Note that v_F, v_M, w_F and w_M are defined to have the same scale as the

community contact matrix in relation to onward transmission, i.e. one unit of v_F, v_M, w_F and

w_M represent the amount of sexual contact that contributes to transmission equivalent of one

daily community contact. The entries of Σ_{MF}, Σ_{FM}, S_{MF} and S_{FM} are then modelled as:

531
$$(\Sigma_{MF})_{ab} = v_F \frac{n_a p_a}{\sum_a n_a p_a},$$

532
$$(\Sigma_{MF})_{ab} = v_M \frac{m_a q_a}{\sum_a m_a q_a} \tag{6}$$

533
$$(S_{MF})_{ab} = w_F \frac{n_a p_a}{\sum_a n_a p_a}$$

534
$$(S_{FM})_{ab} = w_M \frac{m_a q_a}{\sum_a m_a q_a},$$

where n_a and m_a are the relative population size for males and females by age group (we

assumed even sex distribution[78,79], i.e. $\sum_a n_a = \sum_a m_a = 1/2$). Note that S_X and Σ_X are

proportional with a ratio of $w_X : v_X$. We assumed that infection through community contact

has no association with one's sexual contact rate, i.e. v_F and v_M represent the mean sexual

contact rates among the high-activity groups. Reciprocity of heterosexual contacts then

requires:

541
$$v_F \sum_a m_a q_a = v_M \sum_a n_a p_a. \tag{7}$$

We estimated the ratio between w_F and v_F assuming that the sexual network among high-

activity groups is represented by a configuration network (i.e. no degree assortativity; an

alternative scenario is discussed in the Supplementary Information), in which v_X corresponds

to the mean degree and w_X to the mean neighbour degree (or degree-weighted average of

degree)[37]. We assumed female sex workers (FSWs) represent the behaviour of the majority of

high-activity female individuals and used an FSW survey data from DRC[80]. Since both the

mean degree and mean neighbour degree can be characterised by the mean μ and standard

deviation (SD) σ of the sexual contact degree distribution $\pi(x)$, their ratio is estimated as:

551
$$\frac{w_F}{v_F} = \frac{\int_0^\infty x^2 \pi(x) dx}{\left(\int_0^\infty x \pi(x) dx\right)^2} = 1 + \frac{\sigma^2}{\mu^2} \tag{8}$$

We used the reported mean and SD for the weekly number of clients (mean: 17.61; SD: 12.0)

and non-paying partners (mean: 5.51; SD: 22.0) per FSW[80] to derive $w_F/v_F = 2.17$. Here we

assumed that the numbers of clients and non-paying partners are uncorrelated (an alternative

scenario discussed in Supplementary Information); the sexual contact degree as a sum of

these two numbers would then have a mean $17.61 + 5.51 = 23.12$ and a variance of $12.0^2 +$

$22.0^2 = 628$, which were supplied to Equation 8. This and the reciprocity requirement in

Equation 7 assure that v_M and v_F are derived from w_F, $\{p_a\}$ and $\{q_a\}$. We also set boundary
conditions for the total number of high-activity individuals, i.e. $\sum_a n_a p_a$ and $\sum_a m_a q_a$.
Namely, we assumed that high-activity males account for 10% of the male population among
the sexually-active age groups and high-activity females 1% of the total female population,
based on engagement data in commercial sex in Africa (proportion of reporting paid sex in
the last 12 months among males aged 15-59 in Mozambique[81] and the estimated number of
FSWs per capita in DRC[43]).
566
Assuming the community transmission patterns are similar between clades Ia and Ib, we used
the same σ_a estimated for clade Ia throughout the clade Ib analysis. This implicitly assumes
uniform smallpox vaccine coverage across geographical settings (see Figure S7 for sensitivity
analysis). The rest of the parameters, $\{p_a\}$, $\{q_a\}$, w_F, w_M were estimated by the Markov-chain
Monte Carlo (MCMC) method (No-U-Turn Sampler[82]). We obtained 2,000 MCMC samples,
while ensuring the effective sample sizes (ESS) of at least 500 for every parameter; most
parameters achieved an ESS of ~1,000 or larger. Using these posterior samples, we estimated
the proportion of R_{eff} attributable to sexual transmission as a measure of relative contribution
of sexual contacts to mpox transmission. We defined it as the relative reduction in R_{eff} by
excluding sexual contacts from the next generation matrix. Namely, this proportion is given
as $1 - \lambda_C/\lambda_{SC}$, where λ_{SC} and λ_C are the dominant eigenvalue of the next generation matrix (see
Equation 3) with and without the sexual contact blocks (S_X and Σ_X), respectively.
579
580   **Extended Data Table 3. Model parameters.**

Notation	Dimension*	Description	Reference
σ_0	1	Relative susceptibility of children aged 0–4	Estimated Prior: Unif(0,1)
ε	1	Net vaccine protection among immunised cohorts (born in or before 1980)	Estimated Prior: Unif(0,1)
C	6×6 or 8×8	Community contact matrix	[83]
β	1	Transmissibility	Nuisance parameter (not explicitly estimated)
p_a	4	Proportion of high-activity individuals among males of age group a: (aged 15–19, 20–29, 30–39, and 40–49 years)	Estimated Prior: LogNorm(0, 5) truncated on [0, 1]
q_a	4	Proportion of high-activity individuals among females of age group a: (aged 15–19, 20–29, 30–39, and 40–49 years)	Estimated Prior: LogNorm(0, 5) truncated on [0, 1]
v_F	1	Mean sexual contact rate of high-activity females (mean degree)	Derived
v_M	1	Mean sexual contact rate of high-activity males (mean degree)	Derived

w_F	1	Mean sexual contact rate of high-activity females with sexually-acquired infection (mean neighbour degree)	Estimated Prior: LogNorm(0, 2)
w_M	1	Mean sexual contact rate of high-activity males with sexually-acquired infection (mean neighbour degree)	Estimated Prior: LogNorm(0, 2)
n_a	8	Relative population size for males of age group a	[69,70]
m_a	8	Relative population size for females of age group a	[69,70]

581   * The number of free parameters represented by each notation
582
Model averaging and outputs
We used model averaging over candidate models using different contact matrix data
(Extended Data Table 2) with Watanabe-Akaike weights[36] based on the combined likelihood
for datasets 2, 3 and 4 in Extended Data Table 1. We did not include the likelihood for
subprovincial datasets (datasets 5 and 6) or Burundi (dataset 7) in the weights for the main
analysis due to overlapping cases and potentially different data collection and contexts,
respectively. Alternatively, weights also using these likelihoods were included in our
sensitivity analysis (see Supplementary Information). Models 5 and 6 using synthetic contact
matrices had substantially worse likelihood than other models resulting in effectively zero
weights and thus were not included in the main analysis.
593
594   *(i) Frequency of transmission between age groups*
595   We estimated the relative frequency of infector-infectee age pairs at equilibrium using the
596   next generation matrix and the normalised dominant eigenvector:
597

$$M = B_\sigma \begin{bmatrix} O & O & S_{MF} & \Sigma_{MF}Q \\ C_{MM} & C_{MM} & C_{MF} & C_{MF} \\ S_{FM} & \Sigma_{FM}P & O & O \\ C_{FM} & C_{FM} & C_{FF} & C_{FF} \end{bmatrix} \mathrm{diag}\left(\begin{bmatrix} \begin{bmatrix} i^t_{M_H} \end{bmatrix} \\ i^t_m \\ i^t_{F_H} \\ i^t_f \end{bmatrix} \right). \tag{9}$$

599
600   We then aggregated the block elements of M to obtain a sex- and transmission-route
601   aggregated frequency matrix.
602
(ii) Projection of time evolution of the reproduction numbers
We projected the effective reproduction number (R_y) of mpox in DRC and Burundi by
shifting the age of the smallpox-immunised cohorts and modifying the contact matrices[75]
according to the population projection[70]. Referring to the previous estimate[21], we assumed the
effective reproduction number of clade Ia in the Tshuapa province was 0.82 in 2015
(midpoint of the estimation period 2013–2017). We then projected the effective reproduction
number for clade Ia between 2010 and 2030 as the dominant eigenvalue of the next
generation matrix for each year, accounting for the following two factors. First, the relative
susceptibility of age groups that include the immunised cohorts was adjusted according to the
estimated net protection ε (model averaging weights from Table 1) and the proportion of
immunised cohorts (those born in or before 1980 as per the population projection[70]) within
the age groups. Second, the community contact matrices are also adjusted to the projected

population age distribution by the density correction method[75]. For clade Ib, we projected the effective reproduction number in 2024 onward assuming the patterns of community contact transmission are shared between clades Ia and Ib. Namely, the R_y for clade Ib was projected by incorporating additional sexual contact contributions described in Equation 3 (estimated for the Kivus and Burundi) into the projected next generation matrices for clade Ia, assuming time-invariant sexual contact patterns. Here we assumed a hypothetical 'pre-outbreak' state and neglected potential infection- or mpox vaccination-derived immunity as well as any future interventions.

(iii) The effective reproduction numbers under focused and mass vaccination scenarios
We modelled the change in the effective reproduction number of clade Ib under a combination of FSW-focused and mass vaccination scenarios. We used an mpox vaccine effectiveness estimate of 86%[84], which we assumed to protect against infection but not against onward transmission in the case of breakthrough infections. FSW-focused vaccination was assumed to be offered to female high-activity individuals of age 20–39. For mass vaccination, all individuals aged 20–39 years, regardless of sexual activity levels, were included. The baseline effective reproduction numbers without vaccines were matched to the values projected in (ii) for each population setting.

Data and code availability
All data used in this study are publicly available. All the analysis was conducted in Julia v.1.9.3 and R v.4.3.1. Source codes are available on a GitHub repository:
https://github.com/akira-endo/mpoxclade1_eigen.

Acknowledgement

H.M., T.R.A., S-m.J. and A.E. are supported by the Japan Science and Technology Agency (JST) (JPMJPR22R3, to A.E.) and Japan Agency for Medical Research and Development (JP223fa627004). H.M. and A.E. are supported by the Japan Society for the Promotion of Science (JSPS) (JP22K17329, to A.E.). H.M., J.H.F. and A.E. are supported by Cabinet Agency for Infectious Disease Crisis Management, Cabinet Secretariat of Japan. T.R.A. is also supported by the Rotary Foundation (GG2350294), the Nagasaki University World-leading Innovative & Smart Education (WISE) Program of the Japanese Ministry of Education, Culture, Sports, Science and Technology (MEXT) and the JSPS (JP24KJ1827). KE is supported by the Ministry of Education, Singapore, under the Academic Research Fund Tier 1 Seed Award (RLMOE100201900000001), a Singapore Ministry of Education startup grant (LKCMedicine-SUG, #022487-00001) and JST (JPMJPR23R3). S-m.J. is supported by the Centers for Disease Control and Prevention (CDC) Safety and Healthcare Epidemiology Prevention Research Development programme (200-2016-91781). A.E. is also supported by National University of Singapore Start-Up Grant. A.N. and A.E. are supported by the National Institutes of Health (K01AI166347, to A.N.), the National Science Foundation (NSF #2230125, to A.N.). A.N. is also supported by JST (JPMJPR21R8). S.J. and B.L.D. are supported by the Ministry of Education Reimagine Research Grant and Programme for Research in Epidemic Preparedness and REsponse (PREPARE), Ministry of Health, Singapore. J.H.F is supported by PREPARE.

Competing interests
None declared.

Author contribution
Conceptualisation: A.E. Methodology: H.M., T.R.A, K.P. and A.E. Investigation: H.M.,
T.R.A and A.E. Visualisation: H.M., T.R.A and A.E. Data curation: T.R.A., S.J., K.P. and
A.E. Funding acquisition: A.E. and A.N. Writing – original draft: H.M., T.R.A and A.E.
Writing – review and editing: B.L.D., D.B., J.H.F., S.J., P.K.M., K.E., S-m.J., A.N. K.P.,
A.M.W., D.S.N. and D.N.

References
1. Vakaniaki, E. H. *et al.* Sustained human outbreak of a new MPXV clade I lineage in

eastern Democratic Republic of the Congo. *Nat. Med.* 1–5 (2024) doi:10.1038/s41591-

024-03130-3.

2. Laurenson-Schafer, H. *et al.* Description of the first global outbreak of mpox: an

analysis of global surveillance data. *Lancet Glob. Health* **11**, e1012–e1023 (2023).

3. World Health Organization. *Multi-Country Outbreak of Mpox, External Situation*

Report #39 - 6 October 2024. https://www.who.int/publications/m/item/multi-country-

outbreak-of-mpox--external-situation-report--39---6-october-2024 (2024).

4. World Health Organization. *Multi-Country Outbreak of Mpox, External Situation*

Report#36- 14 September 2024. https://www.who.int/publications/m/item/multi-

country-outbreak-of-mpox--external-situation-report-36--14-september-2024 (2024).

5. Vakaniaki, E. H. *et al.* Sustained human outbreak of a new MPXV clade I lineage in

eastern Democratic Republic of the Congo. *Nat. Med.* 1–5 (2024) doi:10.1038/s41591-

024-03130-3.

6. Wawina-Bokalanga, T. *et al.* Co-circulation of monkeypox virus subclades Ia and Ib in

Kinshasa Province, Democratic Republic of the Congo, July to August 2024.

Eurosurveillance **29**, 2400592 (2024).

7. Multi-country outbreak of mpox, External situation report #42- 9 November 2024.

https://www.who.int/publications/m/item/multi-country-outbreak-of-mpox--external-

situation-report--42--9-november-2024.

8.   Public Health Emergency Operations Center, Ministry of Public Health and the Fight

against AIDS in Burundi. Rapport de situation sur l'épidémie de la variole simienne

sitrep n° 094 [Situation report of the mpox epidemic No. 94] 2024.

9.   Multi-country outbreak of mpox, External situation report #53 - 29 May 2025.

https://www.who.int/publications/m/item/multi-country-outbreak-of-mpox--external-

situation-report--53---29-may-2025.

10.   Mitjà, O. *et al.* Monkeypox. *The Lancet* **401**, 60–74 (2023).

11.   Nolen, L. D. *et al.* Introduction of Monkeypox into a Community and Household: Risk

Factors and Zoonotic Reservoirs in the Democratic Republic of the Congo. (2015)

doi:10.4269/ajtmh.15-0168.

12.   Hammarlund, E. *et al.* Multiple diagnostic techniques identify previously vaccinated

individuals with protective immunity against monkeypox. *Nat. Med.* **11**, 1005–1011

(2005).

13.   Karem, K. L. *et al.* Monkeypox-Induced Immunity and Failure of Childhood Smallpox

Vaccination To Provide Complete Protection. *Clin. Vaccine Immunol.* **14**, 1318–1327

(2007).

14.   Rimoin, A. W. *et al.* Major increase in human monkeypox incidence 30 years after

smallpox vaccination campaigns cease in the Democratic Republic of Congo. *Proc.*

Natl. Acad. Sci. **107**, 16262–16267 (2010).

15.   Christodoulidou, M. M. & Mabbott, N. A. Efficacy of smallpox vaccines against Mpox

infections in humans. *Immunother. Adv.* **3**, ltad020 (2023).

16.   Li, E. *et al.* Duration of humoral immunity from smallpox vaccination and its cross-

reaction with Mpox virus. *Signal Transduct. Target. Ther.* **8**, 1–7 (2023).

17.   Colombe, S. *et al.* Effectiveness of historical smallpox vaccination against mpox clade

II in men in Denmark, France, the Netherlands and Spain, 2022. *Eurosurveillance* **29**,

2400139 (2024).

18.   Hoff, N. A. *et al.* Evolution of a Disease Surveillance System: An Increase in Reporting

of Human Monkeypox Disease in the Democratic Republic of the Congo, 2001-2013.

Int. J. Trop. Dis. Health 1–10 (2017) doi:10.9734/IJTDH/2017/35885.

19.   Bunge, E. M. *et al.* The changing epidemiology of human monkeypox—A potential

threat? A systematic review. *PLoS Negl. Trop. Dis.* **16**, e0010141 (2022).

20.   Subissi, L. & Mitjà, O. Rising mpox trends in DR Congo: the neglected spread of an

epidemic. *The Lancet* **405**, 358–360 (2025).

21.   Charniga, K. *et al.* Updating Reproduction Number Estimates for Mpox in the

Democratic Republic of Congo Using Surveillance Data. (2024) doi:10.4269/ajtmh.23-

0215.

22.   Kinganda-Lusamaki, E. *et al.* Clade I Mpox virus genomic diversity in the Democratic

Republic of the Congo, 2018 - 2024: Predominance of Zoonotic Transmission.

2024.08.13.24311951 Preprint at https://doi.org/10.1101/2024.08.13.24311951 (2024).

23.   Government of the Democratic Republic of the Congo & World Health Organization.

Rapport de La Situation Epidemiologique de La Variole Simienne (Mpox) En RDC

Sitrep No 020 (17 Au 23 Juin 2024) - Democratic Republic of the Congo | ReliefWeb.

https://reliefweb.int/report/democratic-republic-congo/rapport-de-la-situation-

epidemiologique-de-la-variole-simienne-mpox-en-rdc-sitrep-no-020-17-au-23-juin-2024

(2024).

24.   Government of the Democratic Republic of the Congo & World Health Organization.

Rapport de La Situation Épidémiologique de La Variole Simienne (Mpox) En RDC

Sitrep No 025 (29 Août 2024) - Democratic Republic of the Congo | ReliefWeb.

https://reliefweb.int/report/democratic-republic-congo/rapport-de-la-situation-

epidemiologique-de-la-variole-simienne-mpox-en-rdc-sitrep-no-025-29-aout-2024

(2024).

25. World Health Organization. *Mpox - Democratic Republic of the Congo*.

https://www.who.int/emergencies/disease-outbreak-news/item/2024-DON522.

26. Endo, A. *et al.* Heavy-tailed sexual contact networks and monkeypox epidemiology in

the global outbreak, 2022. *Science* **378**, 90–94 (2022).

27. Murayama, H. *et al.* Accumulation of Immunity in Heavy-Tailed Sexual Contact

Networks Shapes Mpox Outbreak Sizes. *J. Infect. Dis.* **229**, 59–63 (2024).

28. World Health Organization. *Joint ECDC-WHO Regional Office for Europe Mpox*

Surveillance Bulletin: 09 October 2023.

https://www.who.int/andorra/publications/m/item/joint-ecdc-who-regional-office-for-

europe-mpox-surveillance-bulletin--09-october-2023 (2023).

29. Diekmann, O., Heesterbeek, J. A. P. & Roberts, M. G. The construction of next-

generation matrices for compartmental epidemic models. *J. R. Soc. Interface* (2010)

doi:10.1098/rsif.2009.0386.

30. Melegaro, A. *et al.* Social Contact Structures and Time Use Patterns in the Manicaland

Province of Zimbabwe. *PLOS ONE* **12**, e0170459 (2017).

31. Ježek, Z., Grab, B., Szczeniowski, M. V., Paluku, K. M. & Mutombo, M. Human

monkeypox: secondary attack rates. *Bull. World Health Organ.* **66**, 465–470 (1988).

32. Whitehouse, E. R. *et al.* Clinical and Epidemiological Findings from Enhanced

Monkeypox Surveillance in Tshuapa Province, Democratic Republic of the Congo

During 2011–2015. *J. Infect. Dis.* **223**, 1870–1878 (2021).

33. Buckland, S. T., Burnham, K. P. & Augustin, N. H. Model Selection: An Integral Part

of Inference. *Biometrics* **53**, 603–618 (1997).

34.   Titanji, B. K. *et al.* Effectiveness of Smallpox Vaccination to Prevent Mpox in Military

Personnel. *N. Engl. J. Med.* **389**, 1147–1148 (2023).

35.   Taube, J. C., Rest, E. C., Lloyd-Smith, J. O. & Bansal, S. The global landscape of

smallpox vaccination history and implications for current and future orthopoxvirus

susceptibility: a modelling study. *Lancet Infect. Dis.* **23**, 454–462 (2023).

36.   Watanabe, S. Asymptotic Equivalence of Bayes Cross Validation and Widely

Applicable Information Criterion in Singular Learning Theory. *J. Mach. Learn. Res.* **11**,

3571–3594 (2010).

37.   Catanzaro, M., Boguñá, M. & Pastor-Satorras, R. Generation of uncorrelated random

scale-free networks. *Phys. Rev. E* **71**, 027103 (2005).

38.   Masirika, L. M. *et al.* Mapping and sequencing of cases from an ongoing outbreak of

Clade Ib monkeypox virus in South Kivu, Eastern Democratic Republic of the Congo

between September 2023 to June 2024. 2024.09.18.24313835 Preprint at

https://doi.org/10.1101/2024.09.18.24313835 (2024).

39.   Sex workers find themselves at the center of Congo's mpox outbreak | AP News.

https://apnews.com/article/mpox-outbreak-sex-workers-miners-kamituga-congo-

5f6f3c7e9f0abd1917f8daf41678ea2b.

40.   De Serres, G., Gay, N. J. & Farrington, C. P. Epidemiology of Transmissible Diseases

after Elimination. *Am. J. Epidemiol.* **151**, 1039–1048 (2000).

41.   Beer, E. M. & Rao, V. B. A systematic review of the epidemiology of human

monkeypox outbreaks and implications for outbreak strategy. *PLoS Negl. Trop. Dis.* **13**,

e0007791 (2019).

42.   CDC. ACIP Recommendations. *Advisory Committee on Immunization Practices (ACIP)*

https://www.cdc.gov/acip/vaccine-recommendations/index.html (2024).

43.   UNAIDS. *Democratic Republic of the Congo*.

https://www.unaids.org/en/regionscountries/countries/democraticrepublicofthecongo.

44.   Mpox and stigma. *Bull. World Health Organ.* **102**, 848–849 (2024).

45.   Adebisi, Y. A., Ezema, S. M., Bolarinwa, O., Bassey, A. E. & Ogunkola, I. O. Sex

Workers and the Mpox Response in Africa. *J. Infect. Dis.* jiae435 (2024)

doi:10.1093/infdis/jiae435.

46.   Protecting at-risk populations from mpox in the Democratic Republic of the Congo |

WHO | Regional Office for Africa. https://www.afro.who.int/countries/democratic-

republic-of-congo/news/protecting-risk-populations-mpox-democratic-republic-congo

(2024).

47.   World Health Organization. *Multi-Country Outbreak of Monkeypox, External Situation*

Report #1 - 6 July 2022. https://www.who.int/publications/m/item/multi-country-

outbreak-of-monkeypox--external-situation-report--1---6-july-2022 (2022).

48.   Ogoina, D. A brief history of monkeypox in Nigeria. https://www.idsociety.org/science-

speaks-blog/2022/a-brief-history-of-monkeypox-in-nigeria/ (2022).

49.   Ogoina, D. & James, H. I. Mpox among Linked Heterosexual Casual Partners in

Bayelsa, Nigeria. *N. Engl. J. Med.* **388**, 2101–2104 (2023).

50.   Strathdee, S. A., Crago, A.-L. & Shannon, K. Harm reduction and rights-based

approaches to reduce monkeypox transmission among sex workers. *Lancet Infect. Dis.*

23, e43–e46 (2023).

51.   Wawina-Bokalanga, T. *et al.* Epidemiology and phylogenomic characterisation of two

distinct mpox outbreaks in Kinshasa, DR Congo, involving a new subclade Ia lineage: a

retrospective, observational study. *The Lancet* **406**, 63–75 (2025).

52. Kangbai, J. B. *et al.* Clinical and epidemiological characteristics among probable and

confirmed patients with mpox in Sierra Leone reported from January to May 2025.

2025.05.30.25328691 Preprint at https://doi.org/10.1101/2025.05.30.25328691 (2025).

53. Blackburn, D. Epidemiologic and Clinical Features of Mpox in Transgender and

Gender-Diverse Adults — United States, May–November 2022. *MMWR Morb. Mortal.*

Wkly. Rep. **71**, (2022).

54. Pembi, E., Omoleke, S., Paul, H., Augustine, T. & Cuevas, L. E. Monkeypox outbreak

in a correctional center in North Eastern Nigeria. *J. Infect.* **85**, 702–769 (2022).

55. Katoto, P. D. *et al.* Shifting transmission patterns of human mpox in South Kivu, DR

Congo. *Lancet Infect. Dis.* **24**, e354–e355 (2024).

56. Mpox – African Region. https://www.who.int/emergencies/disease-outbreak-

news/item/2024-DON528.

57. Hazra, A. *et al.* Mpox in people with past infection or a complete vaccination course: a

global case series. *Lancet Infect. Dis.* **24**, 57–64 (2024).

58. Cowen, E. W., Tkaczyk, E. R., Norton, S. A. & Leslie, K. S. Mpox: A rapidly evolving

disease. *JAMA Dermatol.* **159**, 424–431 (2023).

59. Mitjà, O. *et al.* Mpox in people with advanced HIV infection: a global case series. *The*

Lancet **401**, 939–949 (2023).

60. Conflict in DR Congo: What's happening & how to help - Democratic Republic of the

Congo | ReliefWeb. https://reliefweb.int/report/democratic-republic-congo/conflict-dr-

congo-whats-happening-how-help (2024).

61. *UNICEF Burundi Humanitarian Situation Report No. 1 (Level 3 Mpox): 25 July to 15*

September 2024 - Burundi | ReliefWeb. https://reliefweb.int/report/burundi/unicef-

burundi-humanitarian-situation-report-no-1-level-3-mpox-25-july-15-september-2024

(2024).

62.   Exploitation, Sexual and Gender-Based Violence Increases in DRC - Democratic

837   Republic of the Congo | ReliefWeb. https://reliefweb.int/report/democratic-republic-

838   congo/exploitation-sexual-and-gender-based-violence-increases-drc (2024).

839   63.   Marziano, V., Guzzetta, G., Longini, I. & Merler, S. Epidemiologic Quantities for

840   Monkeypox Virus Clade I from Historical Data with Implications for Current

841   Outbreaks, Democratic Republic of the Congo - Volume 30, Number 10—October 2024

842   - Emerging Infectious Diseases journal - CDC. doi:10.3201/eid3010.240665.

843   64.   Mpox research and innovation - aligning research response with outbreak goals.

844   https://www.who.int/news-room/events/detail/2024/08/29/default-calendar/mpox-

845   research-and-innovation---aligning-research-response-with-outbreak-goals (2024).

846   65.   Public Health Emergency Operations Center, Ministry of Public Health and the Fight

847   against AIDS in Burundi. Rapport de situation sur l'épidémie de la variole simienne

848   sitrep n° 053 [Situation report of the mpox epidemic No. 53] 2024.

849   66.   World Health Organization. 2022-24 Mpox (Monkeypox) Outbreak: Global Trends.

850   https://worldhealthorg.shinyapps.io/mpx_global/.

851   67.   Funk, S. *et al.* socialmixr: Social Mixing Matrices for Infectious Disease Modelling.

852   (2024).

853   68.   Population Census Office, Zimbabwe National Statistics Agency. Zimbabwe Population

854   Census 2012.

855   https://web.archive.org/web/20240712073454/https://www.zimstat.co.zw/wp-

856   content/uploads/publications/Population/population/Manicaland.pdf (2024).

857   69.   DECRET N0 100/032 DU 27 MARS 2025 PORTANT PUBLICATION DES

858   RESULTATS PRELIMINAIRES DU RECENSEMENT GENERAL DE LA

859   POPULATION, DE L'HABITAT, DE L'AGRICULTURE ET DE L'ELEVAGE AU

860   BURUNDI, EDITION 2024 (RGPHAE, 2024) – Présidence de la République du

Burundi. https://presidence.gov.bi/2025/04/03/decret-n0-100-032-du-27-mars-2025-

portant-publication-des-resultats-preliminaires-du-recensement-general-de-la-

population-de-lhabitat-de-lagriculture-et-de-lelevage-au-burundi-edition-2024-r/ (2025).

70. United Nations, Department of Economic and Social Affairs, Population Division.

World Population Prospects 2024, Online Edition. https://population.un.org/wpp/.

71. Prem, K. *et al.* Projecting contact matrices in 177 geographical regions: An update and

comparison with empirical data for the COVID-19 era. *PLOS Comput. Biol.* **17**,

e1009098 (2021).

72. Anderson, R. M., Medley, G. F., May, R. M. & JOHNSON, A. M. A Preliminary Study

of the Transmission Dynamics of the Human Immunodeficiency Virus (HIV), the

Causative Agent of AIDS. *Math. Med. Biol. J. IMA* **3**, 229–263 (1986).

73. Gomes, M. G. M. *et al.* Individual variation in susceptibility or exposure to SARS-CoV-

2 lowers the herd immunity threshold. *J. Theor. Biol.* **540**, 111063 (2022).

74. Kullback, S. & Leibler, R. A. On Information and Sufficiency. *Ann. Math. Stat.* **22**, 79–

86 (1951).

75. Arregui, S., Aleta, A., Sanz, J. & Moreno, Y. Projecting social contact matrices to

different demographic structures. *PLOS Comput. Biol.* **14**, e1006638 (2018).

76. Sebastian Meyer. R: Aggregate a Contact Matrix. (2024).

77. Kloek, T. & Van Dijk, H. K. Bayesian Estimates of Equation System Parameters: An

Application of Integration by Monte Carlo. *Econometrica* **46**, 1 (1978).

78. UNSD — Demographic and Social Statistics. *Demographic Yearbook – 2020*.

https://unstats.un.org/unsd/demographic-social/products/dyb/dyb_2020/ (2020).

79. Democratic Republic of the Congo - Subnational Population Statistics -

cod_admpop_adm2_2020.csv - Humanitarian Data Exchange.

https://data.humdata.org/dataset/cod-ps-cod/resource/bf8edc01-c28a-470d-9720-

cb2d9435dfdf.

80. Inungu, J. N., Kandala, B. F. N. & Atungale, A. M. Prevalence of HIV and Syphilis and

Related Risk Behaviors Among Female Sex Workers in the Democratic Republic of the

Congo. *Eur. J. Environ. Public Health* **6**, em0114 (2022).

81. Semá Baltazar, C. *et al.* Who Are the Men Who Pay for Sex in Mozambique? Results

from the National HIV/AIDS Indicator Survey 2015. *Arch. Sex. Behav.* **50**, 2057–2065

(2021).

82. Hoffman, M. D. & Gelman, A. The No-U-Turn sampler: adaptively setting path lengths

in Hamiltonian Monte Carlo. *J Mach Learn Res* **15**, 1593–1623 (2014).

83. Melegaro, A. *et al.* Social Contact Structures and Time Use Patterns in the Manicaland

Province of Zimbabwe. *PLOS ONE* **12**, e0170459 (2017).

84. Wolff Sagy, Y. *et al.* Real-world effectiveness of a single dose of mpox vaccine in

males. *Nat. Med.* **29**, 748–752 (2023).

900
901