Modelling the potential spread of Clade Ib MPXV in Asian cities

Authors: Shihui Jin[1], Gregory Gan[1], Akira Endo[1,2,3], Kiesha Prem[1,2], Rayner Kay Jin Tan[1], Jue Tao Lim[4], Keisuke

Ejima[4], Borame L Dickens[1]

1. Saw Swee Hock School of Public Health, National University of Singapore and National University Health

System, Singapore

2. Department of Infectious Disease Epidemiology, London School of Hygiene and Tropical Medicine, London, UK

3. School of Tropical Medicine and Global Health, Nagasaki University, Nagasaki, Japan

4. Lee Kong Chian School of Medicine, Nanyang Technological University, Singapore

Correspondence to: Borame L Dickens; ephdbsl@nus.edu.sg; National University of Singapore, 117549,

Singapore

Abstract

Background

The ongoing 2023–2024 mpox outbreak in several African countries, driven by the novel Clade Ib strain, has resulted in imported cases reported in Sweden, Thailand, and India. The potentially high transmissibility of this new strain and shifts in transmission modes may make territories in Asia, which were minimally affected by previous mpox waves, vulnerable to community-wide transmission following importation. While this highlights the importance of early preparedness, current knowledge of the virus's transmission dynamics remains too limited to effectively inform policy-making and resource planning.

Methods

A compartmental model was constructed to characterise the potential mpox transmission dynamics. Importation-triggered outbreaks were simulated in 37 Asian cities with one, three, and five initial local exposures. The impacts of various non-pharmaceutical interventions (NPIs), including isolation and quarantine, were projected and compared.

Findings

Simulated outbreak sizes varied substantially among the cities, with large-scale outbreaks expected in territories with high proportions of sexually active individuals at risk or low immunity from smallpox vaccines. Case count would increase linearly with initial exposures. In the scenario with three initial exposures, up to 58 cases per million residents were expected without interventions. Isolation for diagnosed cases would lower the outbreak size by up to 49.4%. Quarantining close contacts would further decrease cases by 30 percentage points over one year.

Interpretation

The potential mpox outbreak in Asia could be alleviated through strong surveillance and timely response. NPIs are recommended for outbreak management due to their demonstrated effectiveness and practicability.

Keywords: Infectious disease, Mpox, MPXV, NPI, Vaccination

1. Introduction

An outbreak caused by Clade I monkeypox virus (MPXV) was estimated to have emerged in September 2023 in South Kivu province (1), Democratic Republic of the Congo (DRC), with transmission likely driven primarily by sexual contact. The novel strain of the outbreak, later designated as Clade Ib, resulted in surging cases reported in the DRC, alongside the endemic Clade Ia already circulating in other provinces, and quickly spread to neighbouring African countries, including Burundi, Rwanda, Uganda, and Kenya. As of September 2024, importation has also been detected outside the African continent, in Sweden, Thailand, and India (2,3). In response to the rapid spread of Clade Ib in Africa, the World Health Organisation re-declared mpox as a public health emergency of international concern (PHEIC), aiming to raise the global awareness and prevent a repeat of the 2022 mpox outbreak, whose global circulation could potentially be attributed to initial negligence in surveillance and prevention (4).

During the 2022 wave caused by Clade IIb MPXV, over 90% of the cases reported were from Europe and the Americas. In contrast, the Western Pacific, Southeast Asia, and Eastern Mediterranean regions—encompassing most Asian countries—collectively accounted for fewer than 5% of all Clade IIb cases (5). Whilst the possibility of high under-ascertainment exists due to the limited surveillance efforts and reluctance in seeking medical help due to stigmas associated with the infection (6), the proportion of the population in Asia with active infection-induced immunity against the virus is still likely to be much lower compared to other regions of the world with higher cumulative prevalence.

While Clade IIb MPXV was predominantly spread through sexual transmission within the gay, bisexual and other men-who-have-sex-with-men (GBMSM) community (7), heterosexual contact appears to play a more critical role in the ongoing 2023–2024 mpox outbreak driven by Clade Ib in the DRC and neighbouring countries, as female cases, especially sexual workers, comprised approximately half of all the confirmed infections (1,8). Apart from the well-documented sexual mode of transmission, evidence from the ongoing outbreak, including the non-trivial contribution of children to reported case counts, has suggested potential higher prevalence of non-sexual transmission routes compared to the previous 2022 outbreak (8–10). These two possible changes in transmission modes, together with the high global connectivity and international travel, raise the risk of an importation-triggered

mpox outbreak affecting the general population, particularly in an Asia setting with minimal prior exposure to the

virus.

Throughout the COVID-19 pandemic, territories in Asia gained valuable experience in implementing non-

pharmaceutical interventions (NPIs) (11,12). Social distancing measures like quarantine and isolation, which

reduced interpersonal contact, effectively contained the disease transmission (13). These successful stories offer

hope for the authorities in this area to leverage NPIs to curb a potential mpox outbreak, should it be widespread in

the general community. Furthermore, the absence of definitive evidence for respiratory transmission (14) for MPXV

implies that effective outbreak management might be achieved through less stringent NPIs, such as isolating

infected individuals, compared to those implemented during the COVID-19 pandemic.

Nevertheless, few NPIs were applied to manage the 2022 mpox outbreak, which affected only a limited

subpopulation. This creates great uncertainty about their true effectiveness and sufficiency in averting a more

widespread mpox outbreak once it establishes itself in the general population. In addition, the distinct transmission

patterns of Clade Ib MPXV, especially regarding the prolonged infectious duration and the combination of

transmission through sexual and non-sexual routes (10,15), make it challenging to directly extrapolate the impacts of

these containment measures from COVID-19 surveillance data.

To address these gaps, we performed this study to model potential mpox outbreaks in 37 Asian cities, covering all

key territories in this region. We collected territory-specific data on the immunity levels from smallpox vaccination,

age structure, and sexually active populations which were assumed of higher transmission risk. These statistics

enabled us to simulate importation-triggered outbreaks utilising a modified deterministic susceptible-exposed-

infectious-recovered (SEIR) model and the transmissibility parameters estimated from the surveillance data in DRC.

We further projected the impacts of NPIs on curbing outbreaks and reducing disease burden. These results

collectively provide insights into factors shaping the outbreak dynamics, informing policy-makers of strategies to

pre-empt or mitigate mpox transmission across the continent.

2. Methods

2.1 SEIR-style transmission model

We adapted a deterministic SEIR compartmental model to characterise potential disease transmission patterns, as shown in Figure 1. Additional compartments, including Quarantined (, Diagnosed ( ), and Dying ( ), were incorporated into the model to facilitate a more accurate reflection of a real-world outbreak, with disease surveillance, variations in disease burden among individuals, and the probable presence of intervention measures.

Figure 1. Model schematic.   and refer to the effective infectious high-risk active and low-risk subpopulations, respectively, where   is the reduced capability of disease spreading due to quarantine or isolation and equals 1 in the absence of them.   is the number of traced close contacts, whose value depends on contact tracing capacity, number of newly diagnosed cases and existing exposed population. See Table S1 for more details for these parameters.

Due to the possible disparity in transmission potential of mpox via sexual and non-sexual activities, we segregated the affected population into two cohorts: high-risk ( ) and low-risk ( ) individuals, assuming the latter could only contract or spread the infection through non-sexual means. Diagnostic rate and disease severity were not distinguished between these two groups.

Infections who underwent clinical testing would be diagnosed and reported, while those untested were assumed to have mild symptoms and would recover without medical assistance. We further stratified the diagnosed cases by

severity. The majority (97%) were expected to have mild symptoms and be able to recover, while the remaining 3%

were at high risk of death,(2) with their survival status depending on the accessibility of medical resources.

Intervention measures built into the model include isolating diagnosed cases and quarantining the close contacts of

confirmed cases. Both groups were presumed to have limited daily interactions with others. Please refer to section

2.3 and the Supplementary Information for more details.

2.2 Outbreak simulations for Asian cities

Utilising the established compartmental model, we simulated potential localised mpox outbreaks in 37 major

municipals across Asia, each being either the capital or with the largest population size in its respective territory. We

assumed the average number of secondary infections generated by an infector through sexual and non-sexual routes

in a fully susceptible population to be 1.62 (R_s) and 1.02 (R_c), respectively. These values were derived from the

estimated overall effective reproduction number and contributions of the two transmission routes during the 2024

mpox outbreak in South Kivu (16). We validated this combination of R_s and R_c combination by fitting the

confirmed death counts (17) to our compartmental model (Figure S1), which traced the index animal-to-human

transmission event back to mid-September 2023, consistent with findings in literature (1).

We initiated the outbreaks by modelling the outcome of a one-time importation event, in which one or more

susceptible individuals in the city were converted to exposed status. We assumed no pre-existing exposure or active

infections in the population prior to the importation. Neither did we account for cross protection from previous

mpox outbreaks, due to the limited exposure to previous waves among individuals outside the GBMSM community

and the few documented mpox cases in the region.(5) However, protection from smallpox vaccines were considered

given their high efficacy and proven ability to prevent mpox infections, particularly among individuals aged over 45

years.(18,19) We obtained the age-specific vaccination rates in 2024 using the model developed by Taube et al.,(20)

and aggregated the overall population immunity level using United Nations age distribution data(21) and an 80.7%

vaccine effectiveness.(20) The corresponding proportion of individuals were assigned to the recovered compartment,

assuming lifelong immunity from vaccination. These statistics were assumed invariant across cities within the same

territory.

Following the approach of Murayama et al. (16), we approximated the size of the high-risk subpopulation by

considering the number of sex workers and their male clients, as they were more likely to engage in higher

frequencies of sexual activity compared to the general population, thus more likely to contract the disease or infect

others. Territory-specific sex worker statistics were collected from UNAIDS and the International Union of Sex

Workers (IUSW) (22,23). while the estimated proportions of clients among males aged 15–49 years were obtained

from Carael et al. (24). We further assumed that males comprised half of the population in each city and that all

high-risk individuals fell in the age group of 15–49 years. Based on these assumptions, we calculated the proportions

of sexually active individuals at high risk in each territory and their corresponding immunity level. To account for

the potential bias in the territory-specific estimates due to limited data and potential temporal changes in the

statistics (24), we additionally conducted a sensitivity analysis to explore how variations in the high-risk population

could shape the outbreak dynamics in each city (Figure S6–S8).

We first simulated the outbreaks initiated by a single local exposure resulting from a one-time importation event,

modelling the scenarios in which the index exposure occurred in either the low- or high-risk subpopulation. To

assess the impact of importation size on disease transmission, we generated epidemic curves for the initial exposure

sizes of one, three, and five, with the proportion of high-risk exposures matching that of the high-risk individuals

among the overall susceptible population to reflect the average spread patterns. We did not assume any subsequent

importation events or the implementation of NPIs (e.g., isolation or quarantine) in these scenarios.

2.3 Intervention effect projection

We evaluated two community-level NPIs: mandatory stay-at-home requirement for diagnosed cases (henceforth

referred to as 'isolation') and quarantine of their close contacts captured by contact tracing (henceforth referred to as

'quarantine'). These measures aimed at segregating the susceptible population from the detected infections and

individuals exposed but not yet infectious, thereby reducing the risk of secondary infection.

Two intervention strategies, including

1) isolation alone, and

174   2) isolation combined with quarantine

were proposed and assessed under the scenario of three initial exposures due to a one-time importation event. For

each strategy, four levels of quarantine or isolation effectiveness were assessed, in which the segregation was

expected to reduce contacts between the susceptible and quarantined or isolated individuals by 25%, 50%, 75%, or

100%. The varying isolation effectiveness levels would allow for potential secondary infections of close contacts

due to incomplete adherence, particularly in the case when non-sexual transmission be an established pathway of

transmission (10). Further details regarding these NPIs are elaborated in the Supplementary Information.

We simulated epidemic trajectories over an 18-month (i.e., 540-day) period for all the aforementioned scenarios,

from which we derived statistics to reflect epidemic growth and outbreak size, including time from the initial

exposure to reaching specific infection or confirmed case counts, and cumulative cases or potential deaths within

one year following the initial exposure. All the analyses and visualisation were performed using the R software

(25).

3. Results

3.1 Susceptibility profiles of the 37 territories

The size of the susceptible population varied substantially across the 37 territories. Under the assumption of a 80.7% vaccine efficacy, 34 territories had higher immunity levels due to smallpox vaccines compared to DRC, with 27 having 75%–90% of their population remaining susceptible to MPXV. Territories in East Asia, including China, Hong Kong SAR, Japan, South Korea, and Taiwan, had the lowest percentages of susceptible population, at 60%–70%. Nevertheless, the proportion of high-risk susceptible individuals was relatively larger in theses territories, with at least 1.5% belonging to this subpopulation. Several Southeast Asian territories, such as Thailand, Cambodia, Laos, and Singapore, also had relatively high proportions of high-risk susceptible subpopulations, accounting for over 2% of the total population. This was particular in Thailand, where the proportion reached a high value of 2.9%. In contrast, the high-risk susceptible subpopulation was relatively small in many territories in Central and West Asia, hovering around or even below 1% (Figure 2).

Figure 2. Proportion of total susceptible population (x-axis) and that of high-risk susceptible individuals (y-axis) in each territory. A vaccine efficacy of 80.7% was assumed for smallpox vaccines. Territories within the same subregion (Central, East, Southeast, Southern, or West Asia) are represented by similar colours. The grey horizontal and vertical lines indicate the respective statistics in DRC.

3.2 Virus spread rate under the scenario involving one initial exposure

When the index local exposure to Clade Ib MPXV occurred in the low-risk group, it would take at least 4.3 months

to reach five cumulative infections in the local population. Five cities—including Phnom Penh (KHM), Dubai

(ARE), Sanaa (YEM), Vientiane (LAO), and Kabul (AFG)—would expect five or more infections within the first

five months, while Tokyo would expect no more than 4 infections by the end of 18 months following the index local

exposure (Figure 3).

Figure 3. Geospatial distribution of simulated virus spread rate. Time in months from the initial local exposure

to five local infections where the initial exposure occurred in either the (a) low- or (b) high-risk group, across the 37

cities. The localized outbreak was modelled with only one initial exposure.

However, patterns would be significantly different if the index exposure took place in the high-risk group. Bangkok,

the capital of Thailand, would be the first to reach five local infections within 1.5 months, due to its largest size of

high-risk susceptible population among the 37 cities. The time it took to reach five infections would also be

substantially shortened to under two months for Hong Kong (HKG), Kuala Lumpur (MYS), and Singapore (SGP),

which would otherwise expect a much slower outbreak, with fewer than five infections in six months. By

comparison, places like Ho Chi Minh City (VNM), Jakarta (IDN), and Almaty (KAZ) would remain at relatively

low outbreak risk. Nonetheless, all the 37 cities would experience a localised outbreak with at least five infections

by the end of the sixth month after the initial local exposure (Figure 3).

3.3 Impact of importation size on local transmission

For all the 37 cities, the time from initial local exposures to first observed case decreased significantly as the size of initial exposures increased. With one initial exposure, the duration varied greatly across cities, ranging from 3.4 months to over 18 months. This period was reduced to approximately one month (29–34 days) for all the cities with three initial exposures and further shortened to around half a month (16–17 days) when five initial exposures were modelled (Figure 4).

Figure 4. Summary statistics for simulated outbreaks in 37 Asian cities. The three statistics reported are: time from initial exposure to first confirmed case (Column 1), number of confirmed cases per million residents (Column 2), and total number of individuals at risk of dying (Column 3) within one year following the initial exposures. The scenarios considered for each city include one, three, and five initial exposures. Cities within each subregion were

ordered by potential death counts in the scenario with five initial exposures. In the 540-day simulation for Japan, no

confirmed cases were observed in the scenario involving only one initial exposure although infections were present.

The top three cities with the highest risks of large, localized outbreaks were all located in Southeast Asia, including

Bangkok, Vientiane, and Phnom Penh. The expected number of cases per million residents in one year in the three

cities were 19, seven, and six, respectively, in the scenario assuming one initial local exposure. These figures would

raise to 96, 35, and 31 when the initial exposure size increased to five. In Central, West, South, and East Asia,

Dushanbe, Muscat, Colombo, and Hong Kong were projected to experience the largest outbreaks, with the initial

local exposure leading to 2.7, 2.5, 1.8, and 0.5 local cases per million residents in one year. Nevertheless, it should

be noted that the ranking of outbreak risks would shift if absolute case counts were used as the metric, with higher

number of local cases expected in densely populated cities. For instance, approximately four, 11, and 19 cases would

be detected in Hong Kong within one year under the scenarios involving one, three, and five initial local exposures,

respectively, while Dubai would expect around three, nine, and 14 cases under the same conditions. The number of

individuals at risk of dying, however, would be substantial only in Bangkok, with up to 11 potential deaths projected

in the scenario involving five initial exposures. In contrast, other cities would expect at most one death in this

hypothetical worst-case scenario. The overall impacts on outbreak size and disease burden over the one-year period

remained consistent across the two increments in the initial exposure size.

3.4 Effectiveness of NPIs on outbreak control

Both isolation and quarantine would contribute to significant reductions in confirmed cases for the 37 cities. In the

scenario with three initial exposures, compared to the baseline without NPIs, isolation alone would bring a 17.9%

(interquartile range [IQR]: 14.8%–22.0%), 31.4% (IQR: 26.4%–37.6%), 41.5% (IQR: 35.5%–49.0%), and 49.4 %

(IQR: 42.7%–57.4%) reduction to the cumulative cases within one year following the initial local exposures, when it

decreased the interpersonal contacts by 25%, 50%, 75%, and 100%, respectively. Adding quarantine would further

lower the cumulative cases by 29.8 (IQR: 26.6–32.8), 34.4 (IQR: 32.7–36.7), 34.0 (IQR: 30.7–34.7), and 30.8

(27.5–32.5) percentage points for the four levels of isolation effectiveness (Figure 5).

Figure 5. Projected reduction in confirmed cases in one year following the initial three exposures due to (a) isolation or (b) isolation combined with quarantine. The reduction proportions were calculated for each level of isolation effectiveness (25%, 50%, 75%, and 100% reduction in interpersonal contacts). compared to the baseline scenario with no NPIs and three initial local exposures per city. Grey box plots display the median, interquartile ranges, and ranges of reductions across the 37 cities. In the legend bar on the right, these cities were ordered by percentage reduction in cases for the scenario in which isolation would eliminate all interpersonal contacts, with a darker colour indicating a greater decrease.

The impacts of NPIs also varied greatly across the 37 cities, with a distribution showing a long right tail. Cities experiencing larger outbreaks tended to benefit more from NPIs. For example, in Bangkok, isolation alone could decrease 47.8%, 72.8%, 85.9%, and 92.5% of confirmed cases at the four effectiveness levels, respectively. In contrast, cities like Ho Chi Minh City and Tokyo, where fewer than four cases were projected over one year, would only see a moderate reduction in case counts of less than 40%, even in an ideal setting when isolation eliminated contacts between the susceptible and infectious population (Figure 5, Figure S4). Nevertheless, in all the cities except Bangkok, local case counts would be minimal with highly effective isolation and quarantine. With isolation alone, the maximum number of cases in a city would be 19, 12, eight and six at 25%, 50%, 75%, and 100% effectiveness, respectively. These figures would drop to nine, three, two, and one, respectively, with the further introduction of quarantine (Figure S4).

4. Discussion

In this study, we simulated localized mpox outbreaks triggered by a single importation event to project the potential spread of Clade Ib MPXV in 37 Asian cities with varying population sizes, levels of immunity from smallpox vaccination, and proportions of sexually active population at higher transmission risk. In the scenario involving three initial exposures, an outbreak of up to 58 cases per million residents was anticipated within one year in the absence of interventions, while the implementation of potent isolation and quarantine measures could lower the incidence to fewer than two per million.

Despite this, a delay of at least 16 days was observed between the initial local viral exposure due to importation events and the first reported local case, with at least two additional months before exponential growth in infections across various scenarios assessed (Figure S3). This delay could provide decision-makers with sufficient time to set up emergency infrastructure, testing centres and create treatment guidelines, and to also then begin containing any potential spread before the number of infections become a burden on the healthcare system.

Our simulation results revealed positive correlations between outbreak sizes and both the proportion of sexually active individuals at high transmission risk and the level of immunity due to smallpox vaccination in each city. However, the impact of the high-risk population size on outbreak scales outweighed that of vaccine-induced immunity, as suggested by the significantly higher number of cases and deaths projected in Bangkok, Thailand, compared to other cities across scenarios with varying initial exposure sizes. Despite 31% of the Thailand's population being protected by smallpox vaccines and presumed immune to MPXV, over 10% of males aged 15–49 years were documented to engage with sex workers (24), leading to an estimated 2.9% of the population being unvaccinated and categorised as high-risk in our study (Figure 2). By comparison, Dubai was projected to have fewer than 5% of the cases and potential deaths seen in Thailand, although the territory had only 7% vaccine-induced immunity (Figure 2, Figure 4). This discrepancy underscores the critical role of sexual transmission in driving localized outbreaks, aligning with the results in our sensitivity analysis, in which Bangkok was predicted to be among cities with the lowest outbreak risks under the assumption of uniform high-risk proportion among individuals aged 15–49 years (Figure S6–S8).

Both NPIs assessed in our analysis—case isolation, and quarantining of close contacts—demonstrated promising

potential in mitigating the outbreaks across all Asian cities. While median reduction in the total case counts over one

year was expected to be around 50% in the ideal scenario where isolation was 100% effective in preventing contacts

between susceptible and isolated infectious individuals, the rate of case reduction would diminish per 25%

increment in isolation effectiveness. The actual effectiveness would depend on stringency of isolation policies, the

isolation facilities required (e.g., home or designated healthcare centres), and compliance of affected individuals,

which may vary between countries. Nevertheless, the moderate simulated outbreak sizes in most cities suggest the

feasibility of less restrictive isolation measures, such as home isolation for confirmed cases, since even with 50%

isolation effectiveness, our model projected no more than 12 confirmed cases in any Asian city except Bangkok

(Figure S3). Furthermore, given the significantly longer infectious period of mpox compared to COVID-19 (15),

extended isolation in healthcare facilities could easily strain pre-existing infrastructure with the surge in case counts.

The added benefits of quarantining close contacts, beyond isolation alone, was consistently projected to slowing the

disease spread, contributing to an additional reduction of around 30 percentage points in the cases counts over one

year. This aligns with its extensive use during the COVID-19 pandemic (26,27). However, it is important to note

that effective contact tracing would require substantial human resources for identifying close contacts, which may

not always be readily available. As such, isolation is recommended as the primary intervention strategy for

containing all potential outbreaks, with contact tracing and quarantining serving as supplementary measures in

scenarios involving relatively large outbreak scales.

It is worth highlighting that our estimates of outbreak sizes and disease burden were based on simulated epidemic

trajectories initiated by a one-time importation, whose outcome were modelled through a fixed size of initial

exposures in the fully susceptible local population for each city. We did not account for recurring importation events

due to the observation of single imported cases in individual affected countries outside the African continent by

September 2024 and the challenges in quantifying the infectivity of incoming infectious travellers on local residents,

as MPXV is mainly transmitted through close contacts (10) and contact patterns between travellers and the local

population might differ significantly from those within the local community. Nevertheless, given the high disease

transmissibility, especially through sexual transmission, the role of importation would likely be minimal in

influencing the epidemic curve once the virus begins to circulate within the city.

In addition, it should be noted that our approximation of high-risk population sizes was based on reported numbers

of sex workers and their clients (22–24). Although consistent data sources were utilized for most territories and the

same calculation method was applied to enhance the comparability across territories, relevant statistics were

unavailable for certain territories, especially those in the Middle East. Consequently we extrapolated the figures

using subregional average proportions, which were lower than the reported values for other parts of Asia.

Possibilities also exist that the number of individuals engaging with sex-workers was under-reported in surveys due

to strict cultural norm in some Asian territories and the stigma associated with acknowledging commercial sex

(24,28). Both factors could lead to an underestimation of the high-risk population sizes and hence an

underestimation of outbreak size, as sexual transmission was a key driver of disease spread in our simulations.

Another major limitation of our study pertains to the large uncertainty surrounding transmission-related model

parameters. The co-circulation of multiple MPXV clades in Africa (29), along with a lack of comprehensive

sequencing data, complicates efforts to pinpoint the transmissibility of MPVX Clade Ib. The disparities in

transmission dynamics across regions with reported MPXV Clade Ib introduced additional unsureness regarding the

contribution of sexual transmission to the overall outbreak (16). In this study, we based our estimates on disease

surveillance data from South Kivu, assuming that transmission in the province was predominately driven by MPXV

Clade Ib with minimal spillover from neighbouring areas. Since the cases identified outside the African continent

primarily consisted of importations and local transmission statistics remained scarce by September 2024, we

extrapolated the transmissibility estimates to the Asian urban setting. However, the contact patterns, a driving factor

of the transmission potential, might vary significantly across cities. Furthermore, the reliance on surveillance data

sourcing from Africa, where healthcare and monitoring systems may face constraints, might have led to an

underestimation of infection sizes and an overestimation of disease severity. Consequently, the projected potential

death counts based on African data in our study may far exceed reality in many Asian cities. With the availability of

more case data and epidemiological parameters, our model may require updating.

Despite these limitations, the transmission model proposed in this study is sufficiently flexible to accommodate for a

range of scenarios with diverse disease transmission dynamics, enabling decisionmakers to plan for different

potential outbreak possibilities. This study explores the impact of NPI impacts on mpox outbreaks, an area much

less explored compared to the virus's transmission patterns (1,30). This has provided evidence for adapting the

containment measures widely applied during the COVID-19 pandemic to effectively managing new outbreaks with

probably high community transmission potential.

Our simulations suggest that while mpox outbreaks triggered by importation could lead to substantial morbidity and

mortality in an Asian city with large populations of sexually active individuals at risk, they would still remain

controllable provided adequate preparation and a timely response from decision-makers, underscoring the

significance of robust surveillance systems. In addition, the NPIs evaluated in this study have the potential to

suppress outbreak sizes and curb disease spread by up to 93% provided proper implementation during a large-scale

outbreak. Based on these findings, there exists an urgent need for strong surveillance systems, efficient contact

tracing, quarantining of close contacts, and institutional isolation of cases.

Declaration of interests

The authors declare no conflict of interests

Author contributions

SJ, GG, and BLD conceived and designed the study. SJ implemented the statistical analysis and created the figures and tables. SJ wrote the original draft of the manuscript. GG, AE, KP, RKJT, JTL, and BLD reviewed and edited the manuscript.

Data availability

Data utilized for this study area available upon request.

Funding

This work was supported by Ministry of Education Reimagine Research Grant; and PREPARE, Ministry of Health. AE is supported by the Japan Science and Technology Agency (JST) (JPMJPR22R3), JSPS Grants-in-Aid KAKENHI (JP22K17329) and National University of Singapore Start-Up Grant.

References

1. Vakaniaki EH, Kacita C, Kinganda-Lusamaki E, O'Toole Á, Wawina-Bokalanga T, Mukadi-Bamuleka D, et al.

Sustained human outbreak of a new MPXV clade I lineage in eastern Democratic Republic of the Congo. Nat

Med [Internet]. 2024 Jun 13 [cited 2024 Aug 29];1–5. Available from: https://www.nature.com/articles/s41591-

024-03130-3

2. European Centre for Disease Prevention and Control. Epidemiological update: Mpox due to monkeypox virus

clade I [Internet]. 2024 [cited 2024 Sep 6]. Available from: https://www.ecdc.europa.eu/en/news-

events/epidemiological-update-mpox-due-monkeypox-virus-clade-i

3. CIDRAP. India reports imported clade 1b mpox case [Internet]. 2024 [cited 2024 Sep 30]. Available from:

https://www.cidrap.umn.edu/mpox/india-reports-imported-clade-1b-mpox-case

4. WHO Director-General declares mpox outbreak a public health emergency of international concern [Internet].

[cited 2024 Aug 29]. Available from: https://www.who.int/news/item/14-08-2024-who-director-general-

declares-mpox-outbreak-a-public-health-emergency-of-international-concern

5. Multi-country outbreak of mpox, External situation report#35- 12 August 2024 [Internet]. [cited 2024 Aug 29].

Available from: https://www.who.int/publications/m/item/multi-country-outbreak-of-mpox--external-situation-

report-35--12-august-2024

6. Chan ZYS, Chong SY, Niaupari S, Harrison-Quintana J, Lim JT, Dickens B, et al. Receptiveness to monkeypox

vaccines and public health communication strategies among gay, bisexual and other men who have sex with

men in Singapore: cross-sectional quantitative and qualitative insights. Sex Transm Infect [Internet]. 2024 Sep 1

[cited 2024 Aug 29];100(6):362–7. Available from: https://sti.bmj.com/content/100/6/362

7. Endo A, Murayama H, Abbott S, Ratnayake R, Pearson CAB, Edmunds WJ, et al. Heavy-tailed sexual contact

networks and monkeypox epidemiology in the global outbreak, 2022. Science [Internet]. 2022 Oct 7 [cited

2024 Aug 29];378(6615):90–4. Available from: https://www.science.org/doi/10.1126/science.add4507

8. Rivers C, Watson C, Phelan AL. The Resurgence of Mpox in Africa. JAMA. 2024 Oct 1;332(13):1045–6.

9. Risk assessment for the EU/EEA of the mpox epidemic caused by monkeypox virus clade I in affected African
countries [Internet]. 2024 [cited 2024 Sep 1]. Available from: https://www.ecdc.europa.eu/en/publications-
data/risk-assessment-mpox-epidemic-monkeypox-virus-clade-i-africa

10. Mpox [Internet]. [cited 2024 Aug 29]. Available from: https://www.who.int/news-room/fact-sheets/detail/mpox

11. Yeoh EK, Chong KC, Chiew CJ, Lee VJ, Ng CW, Hashimoto H, et al. Assessing the impact of non-
pharmaceutical interventions on the transmissibility and severity of COVID-19 during the first five months in
the Western Pacific Region. One Health [Internet]. 2021 Jun 1 [cited 2024 Aug 29];12:100213. Available from:
https://www.sciencedirect.com/science/article/pii/S2352771421000033

12. El Guerche-Séblain C, Chakir L, Nageshwaran G, Harris RC, Sevoz-Couche C, Vitoux O, et al. Experience
from five Asia-Pacific countries during the first wave of the COVID-19 pandemic: Mitigation strategies and
epidemiology outcomes. Travel Medicine and Infectious Disease [Internet]. 2021 Nov 1 [cited 2024 Aug
29];44:102171. Available from: https://www.sciencedirect.com/science/article/pii/S147789392100212X

13. Fitzgerald DA, Wong GWK. COVID-19: A tale of two pandemics across the Asia Pacific region. Paediatric
Respiratory Reviews [Internet]. 2020 Sep 1 [cited 2024 Sep 3];35:75–80. Available from:
https://www.sciencedirect.com/science/article/pii/S1526054220301032

14. Beeson A, Styczynski A, Hutson CL, Whitehill F, Angelo KM, Minhaj FS, et al. Mpox respiratory transmission:
the state of the evidence. The Lancet Microbe [Internet]. 2023 Apr 1 [cited 2024 Aug 29];4(4):e277–83.
Available from: https://www.thelancet.com/journals/lanmic/article/PIIS2666-5247(23)00034-4/fulltext

15. Byrne AW, McEvoy D, Collins AB, Hunt K, Casey M, Barber A, et al. Inferred duration of infectious period of
SARS-CoV-2: rapid scoping review and analysis of available evidence for asymptomatic and symptomatic
COVID-19 cases. BMJ Open [Internet]. 2020 Aug 1 [cited 2024 Sep 3];10(8):e039856. Available from:
https://bmjopen.bmj.com/content/10/8/e039856

16. Murayama H, Asakura TR, Dickens BL, Foo JH, Jin S, Mukadi PK, et al. Roles of community and sexual
contacts as drivers of clade I mpox outbreaks [Internet]. medRxiv; 2024 [cited 2024 Oct 17]. p.
2024.10.15.24315554. Available from: https://www.medrxiv.org/content/10.1101/2024.10.15.24315554v1

17.   Francesco Branda. African Disease Analysis and Monitoring System (ADAMS) [Internet]. 2024 [cited 2024

Oct 6]. Available from: https://github.com/fbranda/ADAMS/tree/main

18.   Matusali G, Petruccioli E, Cimini E, Colavita F, Bettini A, Tartaglia E, et al. Evaluation of Cross-Immunity to

the Mpox Virus Due to Historic Smallpox Vaccination. Vaccines (Basel) [Internet]. 2023 Sep 28 [cited 2024

Sep 1];11(10):1541. Available from: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC10610801/

19.   Population and Population Structure [Internet]. [cited 2024 Sep 1]. Population and Population Structure.

Available from: http://www.singstat.gov.sg/find-data/search-by-theme/population/population-and-population-

structure/latest-data

20.   Taube JC, Rest EC, Lloyd-Smith JO, Bansal S. The global landscape of smallpox vaccination history and

implications for current and future orthopoxvirus susceptibility: a modelling study. The Lancet Infectious

Diseases. 2023 Apr 1;23(4):454–62.

21.   United Nations. World Population Prospects - Population Division [Internet]. 2024 [cited 2024 Oct 6].

Available from: https://population.un.org/wpp/Download/Standard/Population/

22.   UNAIDS. AIDSinfo [Internet]. 2024 [cited 2024 Oct 6]. Available from: https://aidsinfo.unaids.org/

23.   International Union of Sex Workers. Sex Worker Statistics by Country [Internet]. 2024 [cited 2024 Oct 6].

Available from: https://iusw.org/sex-worker-statistics/

24.   Carael M, Slaymaker E, Lyerla R, Sarkar S. Clients of sex workers in different regions of the world: hard to

count. Sex Transm Infect. 2006 Jun;82(Suppl 3):iii26–33.

25.   R Core Team. R: A Language and Environment for Statistical Computing [Internet]. Vienna, Austria: R

Foundation for Statistical Computing; 2024. Available from: https://www.R-project.org/

26.   Shen M, Peng Z, Guo Y, Rong L, Li Y, Xiao Y, et al. Assessing the effects of metropolitan-wide quarantine on

the spread of COVID-19 in public space and households. International Journal of Infectious Diseases [Internet].

2020 Jul 1 [cited 2024 Aug 29];96:503–5. Available from:

https://www.sciencedirect.com/science/article/pii/S120197122030326X

27. Girum T, Lentiro K, Geremew M, Migora B, Shewamare S. Global strategies and effectiveness for COVID-19

prevention through contact tracing, screening, quarantine, and isolation: a systematic review. Trop Med Health

[Internet]. 2020 Nov 23 [cited 2024 Aug 29];48(1):91. Available from: https://doi.org/10.1186/s41182-020-

00285-w

28. Husson L. Who Are the Clients and What They Say about Prostitution in South-East Asia? Moussons

Recherche en sciences humaines sur l'Asie du Sud-Est. 2017 May 4;(29):209–62.

29. Kuppalli K, Dunning J, Damon I, Mukadi-Bamuleka D, Mbala P, Ogoina D. The worsening mpox outbreak in

Africa: a call to action. The Lancet Infectious Diseases [Internet]. 2024 Sep 10 [cited 2024 Oct 8];0(0).

Available from: https://www.thelancet.com/journals/laninf/article/PIIS1473-3099(24)00577-2/fulltext

30. Kibungu EM, Vakaniaki EH, Kinganda-Lusamaki E, Kalonji-Mukendi T, Pukuta E, Hoff NA, et al. Clade I–

Associated Mpox Cases Associated with Sexual Contact, the Democratic Republic of the Congo - Volume 30,

Number 1—January 2024 - Emerging Infectious Diseases journal - CDC. Emerg Infect Dis [Internet]. 2024

[cited 2024 Aug 29];30(1):172–6. Available from: https://wwwnc.cdc.gov/eid/article/30/1/23-1164_article