Title: Study Research Protocol for Phenome India-CSIR Health Cohort Knowledgebase (PI-CHeCK): A Prospective multi-modal follow-up study on a nationwide employee cohort

Running Title: Protocol paper for Phenome India project

Authors: Phenome India Consortium

Correspondence:
Dr. Shantanu Sengupta
Office: Room No. 201,
CSIR- Institute of Genomics and Integrative Biology (IGIB),
Sukhdev Vihar, Mathura Road,
New Delhi - 110025
Tel: 91-11-29879 201
Fax: 91-11-27667 471
e-mail: shantanus@igib.res.in

Abstract

Predicting individual health trajectories based on risk scores can help formulate effective preventive strategies for diseases and their complications. Currently, most risk prediction algorithms rely on epidemiological data from the Caucasian population, which often do not translate well to the Indian population due to ethnic diversity, differing dietary and lifestyle habits, and unique risk profiles. In this multi-center prospective longitudinal study conducted across India, we aim to address these challenges by developing clinically relevant risk prediction scores for cardio-metabolic diseases specifically tailored to the Indian population. India, which accounts for nearly 18% of the global population, also has a significant diaspora worldwide. This program targets longitudinal collection and bio-banking of samples from over 10,000 employees both working and retirees of the Council of Scientific and Industrial Research and their spouses, with baseline sample collection already completed. During the baseline collection, we gathered multi-parametric data including clinical questionnaires, lifestyle and dietary habits, anthropometric parameters, lung function assessments, liver elastography by Fibroscan, electrocardiogram readings, biochemical data, and molecular assays, including but not limited to genomics, plasma proteomics, metabolomics, and fecal microbiome analysis. In addition to exploring associations between these parameters and their cardio-metabolic outcomes, we plan to employ artificial intelligence algorithms to develop predictive models for phenotypic conditions. This study could pave the way for precision medicine tailored to the Indian population, particularly for the middle-income strata, and help refine the normative values for health and disease indicators in India.

Keywords: Phenome, Cardio-metabolic, multi-omics, cohort, risk scores

Introduction

Non-communicable diseases (NCDs) contribute significantly to human morbidity and mortality, accounting for 71% of the total 55 million deaths that occur annually worldwide (1). Of these, approximately 17 million deaths occur in individuals younger than 70 years, with 86% of these early fatalities happening in low and middle-income countries (LMICs) (1). India alone accounts for 5.8 million of these deaths, nearly 10% of the global mortality due to NCDs. Notably, the onset of cardiovascular diseases (CVD) in this population occurs nearly a decade earlier (2, 3). Among NCDs, cardiovascular diseases are the leading cause of death, with 17.9 million fatalities, followed by cancer (9.3 million), respiratory disorders (4.1 million), and diabetes (2 million). Conditions like Nonalcoholic fatty liver disease (NAFLD) and nonalcoholic steato-hepatitis (NASH), which also increase cardiovascular risks, are expected to see a rise in global prevalence from 24%-40% to over 55% by 2040 (4-6).

Given India's diverse ethnicities and geographic variations, understanding the specific risk factors for NCDs in India is crucial. This is particularly important because India is home to one-sixth of the world's population and is considered a global genetic melting pot (7). Additionally, a significant portion of the Indian population resides in the diaspora across other countries. Therefore, comprehensive longitudinal studies across India are essential to identify unique risk factors that can guide the development of targeted preventive and personalized health strategies for both the Indian population and its global diaspora. Identifying these factors is crucial for formulating effective policies and guidelines for precision medicine, which enables the prediction of individual health trajectories and allows for early intervention to prevent disease onset or complications. Risk scores have traditionally been used to predict health outcomes based on data from prospective cohort studies. With the advent of multi-omics data and artificial intelligence (AI)-based analytical tools, there are unprecedented opportunities for developing novel personalized risk metrics for predicting health outcomes (8, 9). A longitudinal population study involving baseline phenotypic measurements and long-term follow-up can help elucidate the complex interrelationships between environmental, genetic, and molecular factors and the risk of subsequent diseases. This approach allows for the measurement

of parameters well before the disease onset, minimizing selection bias and avoiding issues related to reverse causality. Such epidemiological data are vital for developing risk prediction tools and advancing precision health. Currently, most risk prediction algorithms are based on epidemiological data from Caucasian populations, which are often inaccurate when applied to the Indian context due to ethnic diversity, differing dietary and lifestyle patterns and distinct risk profiles (10-12). Importantly, existing studies on the Indian population are limited in both phenotypic information and diversity.

The Indian population is known to exhibit a predisposition to central obesity, increasing the risk of cardio-metabolic diseases (13, 14). Detailed phenotyping and understanding of these risks' genetic and metabolic underpinnings is essential. A longitudinal approach that collects phenotypic data across multiple time points (including physiological function, multi-omics data, and microbiome diversity) will enhance understanding of these risks. Integrating various omics data, such as phenome, metabolome, proteome, and genome, is crucial for developing preventive protocols for various NCDs. Phenomics can help discover new risk factors, diagnostic biomarkers and therapeutic targets for precision medicine (15).

To address the risk factors of cardio-metabolic diseases, particularly diabetes and NAFLD, the Council of Scientific and Industrial Research (CSIR) has initiated the multi-center, pan-India study "Phenome India-CSIR Health Cohort Knowledgebase" (PI-CheCK). The study encompasses its 37 constituent laboratories and associated centers across 17 states and 2 union territories (UT) (16) (Figure 1).

Figure 1: Sampling zones, sites in Phenome India Project, and coordinating labs. Zones are colored and defined for logistics purposes, not in any conventional manner.

CSIR encompasses a diverse range of ethnicities, social-ecological regions and occupational exposures, primarily situated in urban and semi-urban areas which may contribute key factors relevant to this study. At the same time, CSIR staff predominantly come from urban and semi-urban middle- and higher-middle-income groups, resulting in a relatively homogenous socio-economic profile. This planned cohort, comprising approximately 10,000 individuals, includes well-educated working and retired staff as well as their spouses. While India has some notable cohort studies, these are often limited by geographic scope, or the parameters being monitored (6, 17-21). To the best of our knowledge, no nationwide longitudinal cohort of this scale and characterization has been reported before. The study will involve prospective data collection initially over five years at three-time points, gathering extensive health information. This will include but is not limited to:- physiological assessment of lung function via spirometry and oscillometry, cardiac function and heart rate variability (HRV) analysis using echocardiogram (ECG), body composition analysis (BCA), liver function tests through transient elastography (Fibroscan), dermatological assessment of skin moisture, sebum and hydration, as well as anthropometry including muscle strength through dynamometer. In addition, molecular

and biochemical assays of blood samples will be conducted, and a comprehensive sample and data repository will be established for multiple omics and microbiome analyses.

The goal is to generate a first-of-its-kind Indian dataset encompassing various health-related and biomedical parameters from a nationwide employee cohort, tracked over several years. This dataset will have significant public health implications, with the potential to identify diagnostic and prognostic biomarkers. An employee cohort is expected to have a lower dropout rate, as participants are educated and literate, which enhances the study's effectiveness. Moreover, this will promote discussions of clinically relevant findings, ultimately contributing to the development of a healthy and productive scientific workforce at CSIR.

Study Objectives

The primary objectives of the PI-CHeCK study are:

a) To develop a prospective longitudinal cohort of consenting CSIR employees, retirees and their spouses.

b) To develop clinically useful personalized risk prediction scores for cardio-metabolic disorders for pre-diabetes, diabetes, and fatty liver disease.

The long-term objectives for the listed disorders of the study are:

1. To develop risk prediction models for diabetes and liver fibrosis:

(i) Glycemic control: Development of a binary prediction model for the normo-glycemic population to develop:

 (a) Diabetes (prior glycated hemoglobin (HbA1c) <5.7 and fasting blood sugar (FBS) <100 to HbA1c ≥6.5 and FBS ≥126 mg/dL (as per American Diabetes Association (ADA) criteria)

(b) Pre-Diabetes (with prior HbA1c <5.7 and FBS <100 to HbA1c between ≥5.7 and <6.5 and FBS ≥100 but less than <126 mg/dL as per ADA criteria)

(ii) NAFLD/Liver Fibrosis: Development of a binary prediction model for F0/F1 (no fibrosis) to F2/F3/F4 (moderate to severe fibrosis/cirrhosis) on liver Fibroscan (Transient Elastography)

2. To create and maintain a database and linked bio-repository/biobank.

Methods

Design of the study

The study design is primarily longitudinal with the baseline sample collection phase already being completed and two subsequent follow-up phases planned.

Sampling sites and sample size determination

All 37 CSIR labs and associated centers were organized into four zones to streamline logistics management. These zones were defined based on the geographical locations of the labs. The baseline number of regular employees at each lab was used as a reference, and this number was multiplied by four to account for the spouses of current employees, an equal number of pensioners, and the spouses of pensioners. The assumptions made were a) the number of pensioners is at least equal to the number of employees in each lab, and b) both employees and pensioners have spouses. This approach provided a total count for each lab, which was then adjusted to ensure that each zone contained at least 2,500 samples. Information on the four zones and their respective labs is provided in Supplementary Table S1. Four zonal nodal Institutes were designated to coordinate this effort: the Indian Institute of Chemical Biology (CSIR-IICB) for the East zone; the National Chemical Laboratory (CSIR-NCL) for the West zone; the Institute of Genomics and Integrative Biology (CSIR-IGIB) for the North zone and the Centre for Cellular and Molecular Biology (CSIR-CCMB) for the South zone (Figure 1).

Inclusion/Exclusion criteria

All CSIR permanent staff, including employees, retirees, and their spouses, were eligible for enrollment in this study. The exclusion criteria included pregnancy and being under 18 years of age. Additionally, participants were excluded from a specific scanning modality if they reported a history of medical surgery or complications related to that modality, as detailed in Supplementary Table S2. Participants were instructed to fast for at least 10-12 hours before arriving at the phlebotomy sampling site.

Tests and scans conducted

Three main categories of assays were planned: blood-based, scanning-based and stool sample-based. Statistically with the objectives defined, we required 3300 subjects as listed in Table 1. The sample size for risk calculation was planned for a regression model with nearly 20 parameters and a c-statistic of 0.85, which was determined with STATA 15 (22-25). Transparent reporting of a multivariable prediction model for individual prognosis or diagnosis (TRIPOD): the TRIPOD Statement and The Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) guidelines for cohort studies will be adhered to, as and when to be reported (26, 27).

	Diabetes		Fatty Liver
	Q1	Q2	Q1
Definition of End Point	What percentage of normoglycaemic population turned diabetic	What percentage of Normoglycaemic population turned pre-diabetic	What percentage of people with fo or f1 fibrosis grading went to f3 or f4
Criteria 1	HbA1c >= 6.5 from <5.7	Hba1c btwn 5.7 and 6.5 from <5.7	Transient elastography reading <6.5 to >=8.6 or 10
Study Design	Longitudinal	Longitudinal	Longitudinal
Current Prevalence	10	10.3	15
C Statistics	0.85	0.85	0.85
Dropout	30	30	30
Prediction Parameters	30	20	20
Sample Size	3227	1446	1074
Events Needed	323	188	161
Incidence	20	13	
Years of Follow up	3-4 years	2-3 years	

Table 1 Calculation of Statistical Sample size

In response to long-term objectives, we could observe that we need ~3300 samples for events to occur. However, to account for sample availability, considering geographical region, age, sex, etc, and dropout with available resources, we will do phenotyping of 10000 and deep phenotyping of 5000 subjects. It may be noted that the samples of all 10000 individuals will be bio-banked, and if required, based on outcome data, we can do the deep phenotyping on these as well.

A brief description of the test and scans performed and planned is given in Table 2. Biochemical tests and molecular assays, including metabolomics, proteomics, GSA, Cytokine, and Immuno-phenotyping, would be done in a restricted number of samples and not all as described above

Blood-based	
1. Biochemical Analysis	5. Genomics-Global Screening Array (GSA)[*], Whole Genome Sequencing[$] and Whole Exome sequencing[$]
2. Proteomics and Metabolomics	6. Metal Content Analysis
3. Cytokine[*] and Immuno-phenotyping[*#]	7. Telomere Length and Activity[*#]
4. RNA-Micro RNA profiling and RNA sequencing[#]	
Scanning-based	
1. Transient Elastography (TE) for Liver-Fibroscan based	5. Anthropometry-Height, Dynamometer for grip strength, and anthropometry tape-based measurements of Chest, Waist, and Hip Circumference
2. Spirometry for Lung	6. ECG
3. Oscillometry for Lung	7. Skin Health Analysis[*$]
4. Body Composition Analysis (BCA)	
Stool Sample-based	
1. Gut Microbiome sequencing	

[*]Baseline only, [#]Subset of samples only,
[$]Exploratory
Table 2: Different assays planned in the Phenome India Project.

Cohort enrolment and sample collection

The Institutional Human Ethics Committee (IHEC) approved the study at CSIR-IGIB (reference number: CSIR-IGIB/IHEC/2023-24/16). The study was registered vide CTRI number-CTRI/2024/01/061807. The baseline sample collection was planned for seven months, from December 2023 to June 2024. In each lab, on average, a week was spent sampling nearly 50 participants per day. The two follow-up samplings are planned to be carried out in 2025-26 and 2026-27.

To maximize participation, awareness of the program's significance and its participatory nature was promoted at each of the constituent laboratories of CSIR through brochures, web-based communications and personal interactions. A signed physical informed consent form was obtained from the participants during sample collection.

Registration & Survey Questionnaire, Slot Booking

Once awareness of the project was raised within a laboratory, participants were registered through a secure online portal-the CSIR Cohort portal (available at https://www.csircohort.org/) inspired by HIPAA guidelines, the portal implements security measures like two-factor authentication, data encryption, and user timeouts, role-based access, data integrity controls and logging privileges. It also provides supporting documents to inform participants about the study's objectives and protocols. Using two-factor authentication, participants can access their recorded health parameters and reports anytime through the portal.

Participants completed a baseline questionnaire, which included questions about their family and residence. This information was recorded solely for authentication purposes and stored securely in a separate, encrypted database. Each entry is indexed with a unique Registration ID to ensure participant anonymity.

Each laboratory had a coordinator and a co-coordinator, who were and will be responsible for verifying the authenticity of participants. Once the respective coordinator approves a participant's registration, they can proceed to fill out the online questionnaire, which is integrated with a secure Research Electronic Data Capture (REDCap) web application hosted in-house for easy self-reporting (28, 29). REDCap is designed to support de-identified data capture for research studies, providing an intuitive interface for validated data entry. Details of the questionnaire are provided in Supplementary Tables S3 and S4.

Participants could only book a slot for sample collection after successfully submitting the questionnaire. They were then able to choose a date and time for their sample collection. Registered participants received notifications about available sample collection dates at their respective labs, allowing them to schedule their visit at their convenience. For microbiome analysis, participants were required to collect the sampling kit within 48 hours of their chosen sampling time. Reminders were sent via email and mobile through the portal. Reports of the anthropometric parameters, scanning-based tests and basic biochemistry were sent to the participants through the portal, which also requires two-factor authentication for access. On the day of the sampling, the participants were provided with a lab ID to maintain confidentiality. The lab ID was generated by the computer against the registration ID provided by the participant.

Data Acquisition

Each of the activity stations was staffed by trained personnel. All the labs and center stations were equipped with the same equipment except Oscillometry. Details of the various scanning modalities are provided in Supplementary Table S5, and a brief description is provided below. Hard copies of the health card after recording their anthropometric parameters, spirometry, oscillometry, liver elastography results, body composition analysis and ECG were given to the participants. At the end of each day, the station administrator uploads all data in PDF and Excel formats to the portal.

Anthropometry

Anthropometry was carried out to measure height, weight, chest (CC), waist (WC), abdominal (AC), and hip circumferences (HC). Height (in cm) – As recommended by Integrated Child Development Services (ICDS) guidelines, a standard stadiometer was utilized at all centers. Weight (in kg) - Weight was measured using the Accuniq Body Composition Analyzer (Manufacturer- SELVAS Healthcare, Korea Model BC 380). Body Circumferences (in cm) - CESCORF measuring tape (Manufacturer – Cescorf, Brazil) was utilized to measure CC, WC, AC, and HC. The following definitions were adopted to measure circumferences thrice at each site (30, 31):

CC-"Measure the chest circumference at the most significant part of the chest, which is usually across the level of the nipple line in males and just above the breast tissue in females. Measurements were taken at the end of a normal expiration".

WC- "A horizontal measure is taken at the midpoint between the lower margin of the last palpable rib and the top of the iliac crest".

AC- "The tape is held behind the participant with one edge at the horizontal plane through the center of the umbilicus".

HC- "The participant is standing erect, and the feet close together. A horizontal measure is taken around the widest portion of the hips and buttocks".

Dynamometer

It was carried out to record grip strength and the participant was required to sit upright in a chair with feet flat on the floor and legs uncrossed. 200 lbs/90 kg capacity Model 12-0241 Brand-Baseline, USA dynamometer was utilized in the study and was ensured for factory calibration. After positioning the participant comfortably, they would hold the hand dynamometer

with their fingers wrapped around one side of the handle and thumb around the other side with the dynamometer set in the second position. The dynamometer should face forward and the forearm should be held parallel to the floor. Participants' position should be seated with their shoulders adducted, sitting upright with their elbows bent at a 90-degree angle and their wrists set between 0-30 degrees extension. The participant was told to squeeze the hand dynamometer with maximum isometric effort (as hard as possible) for at least 3 seconds and relax. Starting with the right hand, the test was repeated with the opposite left hand and vice versa if the dominant hand was left three times in each hand. It was reset in each repeat. Dynamometer data can be influenced by the duration each participant gets to press it and familiarization with the device. To address these factors, each respondent was given a brief opportunity to acquaint themselves with the dynamometer through a short hands-on practice session. During the test, the dynamometer shouldn't touch any body part or any other object (32-34).

Body Composition Analysis

Body Composition Analysis (BCA) was done on the Accuniq Body Composition Analyzer (manufacturer- SELVAS Healthcare, Korea Model BC 380), a three-frequency system and was carried out as per the standard protocol specified by the manufacturer.

Transient Elastography of Liver

The FS Mini 430 Plus model (Manufacture-Echosens, France) with M or XL probes was used to acquire the scan and data. Fibroscan is a simple and safe technique to assess liver stiffness and fat content, and it takes 5-10 minutes on average per examination. Preliminary preparation required the subject to remain fasting for at least 6 hours before the procedure, which was met through the fasting condition specified. The subject is then asked to lie supine with the right arm in full abduction.

The examiner places the probe along the 5th to 7th intercostal space in the mid or anterior axillary line to obtain a view of the right lobe of the liver. Once the area is identified, measurements are obtained using the appropriate fibro scan probe.

Liver Stiffness Measurement (LSM) and Continuous Attenuation Parameter (CAP) measurements were captured by trained personnel blinded to participants' clinical data. Although the literature points out the use of an XL probe for body mass index (BMI) greater than 30, we used the recommender system of the device for probe selection. Failure of transient elastography is defined as the inability to obtain valid LSM or CAP value (35-37). Manufacturer guidelines were followed to acquire 10 valid measurements.

Spirometry

EasyOne Air hand-held spirometer (Manufacturer-NDD, Switzerland), a widely utilized handheld spirometer based on ultrasound sensing, was used in the study to capture spirometry data. American Thoracic Society (ATS)/European Respiratory Society (ERS) guidelines and interpretation were followed to capture spirometry data (38-41). Forced expiratory volume in first second (FEV1) and forced vital capacity (FVC) are generally used to diagnose lung function abnormalities, with an FEV1/FVC ratio of <0.7 indicative of obstructive airway disease (OAD). If the ratio is normal, i.e. ≥0.7, FEV1 and/or FVC of <80% predicted indicate low lung function with restricted lung function measured through FVC and preserved ratio through FEV1 (42).

Oscillometry

Oscillometry was performed before spirometry using the Tremoflo C-100 Airwave Oscillometry System (Thorasys, Canada) or Resmon Pro V3 System. Oscillometry is used to measure perturbations to sound signals of different frequencies superimposed onto normal tidal breathing, which helps characterize the resistance of small airways and airway distensibility

(43). The signal was recorded for at least three artifact-free breaths, which sets the QC before capturing acceptable maneuvers. Resistance and Reactance at 5 Hz (R5 and X5), Resistance at 20 Hz or 19 Hz (R20/R19), Resistance of small airways, 5 to 20 Hz (R5-20 or R5-R19) and frequency response (Fres) were primarily monitored.

ECG

ECG was performed at a 500 Hz sampling rate using Pisces 1012 model 12 lead ECG, (Manufacturer Allengers, India). The ECG uses a 12-lead setup with a high-pass frequency filter at 0.05 Hz, a low-pass filter at 35 Hz, and a 50 Hz notch filter. The sweep speed was 25 mm/s, and the amplitude was 1.00 cm/mV. A two-minute recording was taken while the participant remained motionless. The data is saved automatically, generating a PDF uploaded to the participant's dashboard.

Skin analysis and phenotyping

To develop standards for objective assessment of skin barrier in Indian skin types, skin barrier function was measured by utilizing Courage Khazaka (Manufacturer Courage Khazaka Electronic GmBH, Germany) in terms of four biophysical parameters, employing distinct probes for measurement: Skin sebum content (Sebumeter): The measurement process involves using grease spot photometry with the Sebumeter® SM 815. The matte tape contacts the skin or hair, becoming transparent based on surface sebum. A photocell measures transparency when the tape is inserted into the device aperture, indicating sebum content by light transmission. The Sebumeter measures sebum levels on the forehead and left cheek, with readings taken over 30 seconds at each site. Transepidermal Water Loss (TEWL) (Tewameter): The Tewameter® probe measures the density gradient of the water evaporation from the skin (Δc) proportional to the TEWL. The probe takes three 10-second measurements on the forehead, left cheek, and left volar forearm to assess TEWL. Stratum Corneum Hydration (Corneometer®): The measurement is based on capacitance measurement of a dielectric medium, here the

stratum corneum, the uppermost layer of the skin. With increasing hydration, its dielectric properties change. The measurement is based on the fact that water has a higher dielectric constant than most other substances (mainly <7) and is done on the forehead, left cheek, and left volar forearm. Skin Elasticity (Cutometer): The measuring principle of the Cutometer® is based on the suction method, where negative pressure deforms the skin mechanically. The device creates pressure, drawing the skin into the probe's aperture. After a defined time, it is released again.

Blood Collection, Testing and Storage

The protocol for blood sample collection, preparation and management was conducted in collaboration with a pathology laboratory that ensured the collection of high-quality samples for subsequent research purposes. The routine blood tests described in Supplementary Table S6 were performed by the diagnostic laboratory, which was accredited by the National Accreditation Board for Testing and Calibration Laboratories (NABL) and the College of American Pathologists (CAP). Other biochemical analyses are being performed at CSIR-IGIB (Supplementary Table S7). The results are uploaded to the central database, and the participants are provided passcode-enabled access to their results. The data is stored in a data repository at CSIR-IGIB with the provision of a backup at CSIR-4PI. The sample flow and processing work plan is depicted in Figure 2.

Blood Sample Collection

Participants came to the collection centers with prior fasting for 10-12 hours and 20 mL of blood was drawn at the phlebotomy station into specific blood tubes such as Ethylenediamine tetraacetic acid (EDTA) (4 tubes), serum-separating tube (SST; 2 tubes), and Fluoride (1 tube). (One EDTA, One-Serum and One Fluoride tube were transported, maintaining a proper cold chain to the pathology laboratory for basic biochemistry analysis. Plasma and serum samples separated in

the sample collection site were transported using a cold chain to CSIR-IGIB, where they were aliquoted, provided a test ID and stored at -80°C for future analysis. The sample flow and processing work plan is shown in Figure 2.

Figure 2: Sample flow and processing in Phenome India Project

Bio-Banking

Each aliquot was assigned a unique 4-character Test ID and frozen using a Cryomed Controlled-Rate Freezer (CRF) for approximately 14 minutes before being stored in an ultra-low deep freezer. The samples are stored in 0.5 ml 2D-tubes marked with unique quick response (QR) codes and tube numbers for precise traceability and data confidentiality. A robust sample management system is maintained at the laboratory that can track samples and facilitate their future usage. A meticulous biobank system will ensure the expansion of global biobank networks and comprehensive data-sharing frameworks to enhance the scope and impact of research. It will further act as a foundation for reliable research by accurately correlating samples with participants and health outcomes.

Assays to be performed and planned

The different experimental assays to be performed and planned are summarized in Table 3.

Study Area	Assay Sample Number	Molecular Study	Instrument

Metabolomics	Metabolomics using the MxP Quant 500 Kit (Biocrates) 5000	Quantification of plasma metabolites	Triple quadrupole mass spectrometer (Sciex, QTRAP 6500 plus, Web IDQ Software
Proteomics	MRM (Multiple Reaction Monitoring) for protein quantification 5000	Targeted quantification of proteins related to cardio-metabolic diseases	Triple quadrupole mass spectrometer (Sciex, QTRAP 6500 plus)
Metallomics	Measurement of metal ions using Matrix-matched serum calibrators (Recipe) used for calibration 10000	Determining the concentration of 30 essential, trace, and toxic metals in serum.	ICP-MS (Inductively Coupled Plasma Mass Spectrometer)
RNA-seq Assay	RNA sequencing using Illumina TruSeq Stranded Total RNA Library Prep Gold Illumina sequencing using 2 x 151 cycles of SBS (Sequence by Synthesis) 500	Differential Gene expression analysis from RNA samples	Illumina TruSeq, NextSeq 2000, Agilent 2100 Bioanalyzer, FastQC, Trimmomatic, Salmon, DESeq2

Genomic Studies (GSA)	Genotyping using Global Screening Array (GSA) 3.0 (Illumina) 5000	Identification of genetic markers and polygenic risk scores	iScan (Illumina)
miRNA Profiling	Serum miRNA profiling using RT-PCR with universal and miRNA-specific primers 5000	miRNA profiling in human serum to identify biomarkers for various diseases	QuantStudio™ 6-Flex (Applied Biosystems), Roche SYBR Master Mix
Telomere	Quantitative PCR for telomere length analysis 5000	Telomere analysis as a molecular marker for health and disease	qPCR (quantitative polymerase chain reaction)
Gut Microbiome	Metagenomic sequencing 6250	Profiling taxonomic composition of the gut microbiome	Novaseq™6000 Platform (Illumina)
Cytokine Analysis	Multiplexing assay using Bio-Plex Pro Human Cytokine Assay 5000	Cytokine profiling in plasma to study the immune response	Bio-Plex 200 (Biorad),

Table 3: Summary of the different assays proposed in the Phenome India Project

Data acquisition and storage

Data Management

The backend application programming interface (API) and frontend application were integrated and deployed primarily on a virtual machine at CSIR-IGIB. Data was stored in a relational database management system (RDBMS)-based architecture on the back end. The front end was developed using server-side scripting language-based architecture and framework with modular supports from JavaScript, CSS (Cascading Style Sheets), and web server-related programming widgets. Robust data security and access management protocols were deployed, and development using non-standard protocols was avoided to prevent potential SQL (Structured Query Language) Injection attacks. Data backup, recovery, and mirrored deployment (disaster management protocols) are proposed (Figure 3).

Figure 3: Overall flow of data in the Phenome India Project.

Data acquisition

The data acquisition process for the PI-CHeCK study involves collecting data from 10,000 participants of diverse backgrounds initially over a five-year period. The study includes surveys, laboratory tests, clinical tests, and scanning modalities. Data is uploaded via a secure protocol involving restricted access to the station admins of the respective modality. To protect data security and privacy, participant information is de-identified, and Base64 encoded, with one-time password-enabled restricted access granted only to authorized personnel. The data is stored in the in-house server, established at CSIR-IGIB, where the PI-CHeCK portal is hosted. The data backup is planned to be managed by a scheduled task that automatically stores backups on the same server and a remote server (through file transfer protocol) at regular intervals. Researchers may need access to participants' data stored in a centralized database for their studies. However, access to this data is controlled to ensure participant privacy and compliance with regulations inspired by health insurance portability and accountability act (HIPAA) guidelines.

Data security and privacy

Apart from basic security measures of password hashing, one-time password-enabled access and security headers, the portal is equipped with security measures to counter cross-site request forgery and handle injection attacks by implementing security cross-site scripting. Arrangements are made to encrypt data in transit and rest in a secure server. Further, the management of sessions strategically prevents interception and unauthorized access.

Utmost care has been taken to maintain data privacy, i.e., data is secured at rest, during transit, and editing. Collected multi-omics/clinical/lifestyle data from different CSIR institutes is stored in a relational database management system. The database can be queried using a secured graphical user interface (GUI) application. Individuals (who provided samples) were treated as 'normal users' of the application and allowed only to view the submitted data. Power users with 'Lab' privileges can generate sample IDs (multiple samples for each 'normal user'), add/link protocols, and upload data files (raw and processed) against anonymized sample IDs only.

A centralized data platform capable of encompassing about 500 terabytes (TB) and appropriate data security, including data protection and database development, was set up at CSIR-IGIB. CSIR-4PI has been recognized as a data recovery site. To enable collaboration amongst the participating CSIR labs, curated de-identified data will be made available for in-depth analysis.

Data Sharing

Data accessibility is enabled for authorized users or systems to access specific datasets. It involves establishing protocols and permissions to ensure that only authorized users can access the necessary data. Access to participants' data is strictly controlled to ensure privacy and regulatory compliance. Authorized researchers with approved projects must submit a Data Access Request Form, reviewed by a Data Access Committee to avoid overlapping objectives. Access is granted for a defined period, with data sharing restricted to individuals listed in the request. Role-based access control (RBAC), encryption, and audit logs are implemented to prevent unauthorized access and track data usage. Secure data sharing using secure file transfer protocol (SFTP) is essential for maintaining sensitive research data's confidentiality, integrity, and availability, ensuring compliance with ethical and regulatory standards. SFTP provides a secure alternative to traditional methods such as email and FTP by encrypting data during transmission and implementing robust authentication

mechanisms. Our protocol includes configuring the SFTP server, establishing user groups with appropriate permissions, and setting up secure directories, with additional measures like virtual private network (VPN) configuration and tools such as graphical user interface or command-line interfaces (CLI) to enhance security and efficiency.

Data backup and mirroring

Data backup and mirroring efforts will be implemented at the CSIR-4PI facility situated in the south zone, which is geographically distinct from the primary data storage site in the north zone.

Data Analysis

In addition to data visualization, we plan to analyze inter-parameter linear/nonlinear correlation keeping given age/gender covariates, use open source tools to identify patterns within the data, develop predictive models using machine learning (ML) algorithms to predict certain phenotypic conditions, create an integrative network of various genomics, proteomics, and metabolomics parameters to understand the systemic regulation (Figure 4).

Sample and indexed participant similarity can be achieved through unsupervised approaches including, but not limited to, clustering and Euclidean/Mahalanobis distance-based methods, which can be used to quantify these similarities. These similarity scores can then be fed and used to train personalized models for risk score assessment.

Using collaborative filtering techniques, individuals having similar phenotypes can be clubbed together. Employing cohort participants' similarities will help identify a subset proximal to an index patient, which will then be used to train a personalized model in a disease-specific manner. The approach for the personalized model will depend on the type of disease and ground truth label criteria. Recent studies have used various algorithms ranging from recurrent neural networks and ensemble

techniques as personalized models for disease state prediction (8, 9). Multi-view datasets will allow us to explore the quantitative association of collected risk factors with the comorbidity- and disease-specific outcomes. The developed framework will be based on explainable AI, rendering the weight of each contributing factor to the predicted outcome.

Figure 4: Schematics depicting multi-modal data integration for risk score development.

Discussion

The incidences of metabolic and other non-communicable complex diseases are increasing at an alarming rate in India which is now considered to be the hub of cardio-metabolic diseases and diabetes in the world (19, 44). From the public health perspective, it is of prime importance to understand the risk factors driving this rise in disease prevalence and develop effective risk stratification, prevention and management strategies.

Large-scale longitudinal cohort studies are essential for investigating disease causation and linking risk factors to health outcomes. For example, the Framingham Heart Study, an ongoing study initiated in 1948, has provided valuable insights into risk factors for heart disease, such as lipid levels, diet, exercise, hypertension, cigarette smoking, etc. This research led to the development of the Framingham Risk Score (FRS), a 10-year cardiovascular risk prediction tool (45-49). However, the FRS may be less applicable to Indian patients due to diverse lifestyles and risk profiles, highlighting the need for a tailored risk matrix based on long-term data specific to the Indian population (50, 51). Over the last few decades, several studies have aimed to identify diagnostic and prognostic markers in India. Most of these were case-control studies, focused on a limited number of biochemical markers and lacked the capacity to analyze critical disease trajectories (52-56). Therefore, a comprehensive longitudinal study across India is necessary to explore the complex interactions between environmental, genetic, and molecular factors influencing disease risk.

The Council of Scientific and Industrial Research (CSIR), with its extensive network of over 37 laboratories and centers nationwide, is ideally positioned to address these gaps. This capability was demonstrated during the COVID-19 pandemic, where a longitudinal cohort provided vital insights into infection dynamics, antibody response dynamics, herd immunity

trends across India, and the likelihood of significant outbreaks in a population (16, 57-59). The study helped obtain sentinel data about the spread pattern and different characteristics of SARS-CoV-2 infection across India.

Recent advances in molecular techniques in genomics, transcriptomics, proteomics, metabolomics, and metagenomics have significantly enhanced our understanding of several facets of human biology. Through the Phenome India project, CSIR aims to leverage these advancements alongside the prospective collection of biological samples from a diverse population, enabling long-term follow-up to establish causal relationships and develop diagnostic and prognostic biomarkers for chronic non-communicable diseases. The extensive phenotyping of participants will make this cohort unique and invaluable. To the best of our knowledge, tests like fibroscan, oscillometry, and skin barrier assessments are being conducted in such a large and diverse population for the first time in India. The prospective collection and analysis of fecal microbiomes adds a new dimension to understanding cardio-metabolic disease risk. Additionally, evaluating metal content in participant's serum samples together with metabolomics, proteomics and genomics will help illuminate the complex interactions between genetic, dietary, and environmental factors impacting metabolic diseases. This deep phenotyping is also expected to unravel several novel genotypic and phenotypic associations in the Indian population.

Employee-based cohorts and restricting a longitudinal cohort study to urban and semi-urban areas introduce both limitations and advantages that can impact the study's generalizability and data quality and feasibility. Employees, especially those in formal jobs, may not represent the general population due to differences in socioeconomic status, education, and access to healthcare. Urban and semi-urban populations differ significantly from rural populations in terms of lifestyle, healthcare access, and environmental exposures. Urban and semi-urban populations may have better access to healthcare and technology than the rural populations. The significant socio-demographic differences between urban and rural populations and gene-environment correlations in rural and urban populations may differ in their genetic risks for certain health outcomes

and it could bias the study's findings. This could skew the findings and limit the findings of health trends across different geographic regions.

However, there are advantages to an employee-based cohort focused on urban and semi-urban areas, particularly in terms of recruitment and follow-up feasibility. Educated participants are more likely to understand the study's importance and comply with data collection and follow-up protocols. Employees are more likely to come from middle or upper socioeconomic backgrounds, which may be seen as potentially skewing the health profile of the cohort and limiting insights into diseases associated with poverty; however, it makes it a uniform cohort and homogenous in terms of socioeconomic strata, education and access to healthcare. The current cohort while it may miss conditions prevalent in rural areas, such as certain infectious diseases, malnutrition, or specific chronic conditions, it would stand as one of a kind for a homogenous population being screened at this large scale and specifically in the context of non-communicable diseases

In conclusion, this multicenter, multi-modal prospective cohort study represents a significant step towards precision medicine, in the Indian context. This level of comprehensive phenotyping has not been attempted before and when combined with data analytics and AI/ML approaches, it aims to produce reliable risk prediction tools for cardio-metabolic disorders.

Ethics

The study was approved by IHEC at CSIR-IGIB (reference no.: CSIR-IGIB/IHEC/2023-24/16). The study was registered vide Clinical Trials Registry- India (CTRI) number- CTRI/2024/01/061807.

Funding

The study is funded by CSIR through grant HCP47.

Acknowledgments

We acknowledge the support from CSIR, participants, and volunteers. Further, following names are acknowledged:

Arpana Parihar, Shalu Yadav, Mohd Abubakar Sadique, Akshay Singh Tomar, Varsha Agrawal, Sakshi Rajput, Aabha Kushwah, Shilpee Chauhan, Varsha Parmar, Pushpesh Ranjan, Neeraj Kumar, Abhijeet Bamoria, Abhisek Giri, Navnidhi Tripathi, Jyoti Lodhi, Parul Shrivastava, Prachi Shrivastava, Komal Sharma (**AMPRI**), Vikas Pathak (**CBRI**), Surya Narayan Mishra, Sathya Swetha, Gururaj Kalshetti, Rakesh Pal Bhagat, Chandreshwar Raju Kataru, Hari Charan Goud Nerella, Amit Chakraborty, Prangya Paramita Sahoo, Alagu Sankareswaran, Sofia Banu, Priyadarshini Sanyal, Sourav Ganguly, Manash Kumar Behera, Pallavi Rao T, Akila Ramesh, Niharika Tiwary, Devika Mahimakar, Abhishek Saha, Shreya, Pramoda, Ruqaiya Tasneem (**CCMB**), Khushbu Singh Raghav, Shivam Tiwari (**CEERI**), N Vinodkumar, Rani C Avilash Sumithra , Pramila Epparti, M Archana (**CFTRI**), Swachchha Majumdar, Sourja Ghosh (**CGCRI**), Debabrata Chanda, Dnyaneshwar Umrao Bawankule, Pankhuri Singh, Pankaj Kumar Shukla, Parmanand Kumar, Sumati Sen, Chandra Kant, Sarita Upadhyay, Sumit Kushwaha, Sukriti Srivastava, Daneshvar Prasad, Kavita Singh, Om Prakash Choudhary Kar, Debasmita Sahoo, Anant Kumar Chaudhary (**CIMAP**), Srinivetha Pathmanapan, Syed Nasar Rahaman, Vikash Negi, Jegan K, Sankar

Palanivel, Lishadevi M, Pranathy K, Vandhana Anumaiya, Kirubakaran BJ, Thiyaneshwar (**CLRI**), Sujata Pachhal, Sukanta Maji, Sonali Garai (**CMERI**),Mukti Advani, Rina Singh (**CRRI**), Sandeep Kumar, Rajni, Preetismita Borah, Manish Kumar, Goraj Singh, Suraj Prakash, Ajay Kumar, Inderjit, Shubham Patial, Shavita, Amandeep, Jyoti Jangra, Aman Grewal, Shivalika, Rajrani, Swapna, Dinesh Kumar, Dhiru, Munish, Akriti, Deepak Kumar, Gourav, Komal, Prashant Kumar, Amrit Marthu, Rishav Rai, Amir Khan, Satinder Kaur, Netra, Pankaj, Anuj Sharma, Parneet Kaur, Krishna, Pallavi Suman, Abhishek, Yash (**CSIO**), Haresh Khasiya, Vishal Dumraliya, Chinmay Chauhan, Priyanka Pandya, Disha Rohit, Neha Sharma, Aadya Yadav, Riya Soni, Dhaval Nathani, Mujeer Habsi, Chand Kothadia, Parth Nathani, Keyur Patel, Jigar Aal, Muskan Kalla, Jayesh Rathod, Saif Syed, Abhit Shah, Chirag Khambhaliya, Mrunalini Gohil, Aditya Vaja, Asmita Dhimmar, Rupal Dubey, Jalak Maniar, Kashish Sharma, Dhruvil Chavda, Charu Waghmare, Anushka Tiwari, Shravi Jain, Arun Rathod, Ankita Shandul, Bharti Rana, Deepesh Khandwal, Tulika Mairal, Rajveer Parmar, Mohit Dangariya (**CSMCRI**), Ajeet Singh (**HRDC**), Salwa Naushin, Subhash Gurjar, Rajesh Kumar (**IGIB**), Anuj, Aditya Ranout, Trilok Saini, Sahdev Choudhary, Amit Kumar, Shweta Sharma, Neha Bhardwaj, Kajal Kalia, Suresh Kumar, Abhishek Goel, Ravi Kumar, Surbhi Mali, Komal Goel, Swati Katoch, Shagun Dogra, Vinesh Sharma, Rajneesh Kumar, Poonam Dhiman, Pardeep Poonia, Sahiba Chahal, Rahul Kumar, Athrinandan s Hegde, Rupinder Kaur, Vivek Kumar, Prakriti Sharma, Avisha Sharma, Ankita, Mohak Mali, Priyanka Koundal, Ashruti (**IHBT**), Rajendra Prasad Punna, Kiran Kumar A, Taslim B Shaikh, Nidhi Sharma, Anjali Veeram, Komal walvekar, Mani Sharma, Suriya panneer, Rajwinder Kaur, Ramasatyasri Kotipalli, Hari Priya Sripadi, Vaishnavi Kambhampati, Pinkal Sondarva, Divya Vani kundurthi, Maneesha Yeddula, Sindoora Dhoolipala (**IICT**), Abhilek K Nautiyal, Harleen Kaur, Ayan Banerjee, Rahul Gautam, Mirnalini Juyal, Mridul Budakoti, Preetanshika Tracy, Pratima Ashoke Patel, Ritu Mourya, Harish Panwar (**IIP**), Sneha Mohanty, Mohd Tauseef Khantwal, Vaibhavi Lahane, Shreya Tripathi, Sunita Devi, Gulafsha Siddiqui, Km Priya, Priyanka Goswami, Sandip Chatterjee, Sakshi Singh, Sachin Mishra, Abhineet Singh, Vishal Narayan Soni, Devendri Khantwal, Murk Kumari, Sanjivani Naik, Apoorva Dewangan, Ananta Joshi, Akansha Chaurashiya, Akansha Chauhan, Sheetal Agarwal, Joel Saji, Nikita Shraogi (**IITR**), Ankita Das, Mayuri Kumari,

Amit Kumar Swain, Deepak Kumar Ahirwar, Nilotpal Kapri, Ghrutanjali Sahu, Santosh Kumar Behera, Suchismita Senapati, Subhashree Nayak, Yogesh Chaudhary, Anagonou Bertrand, Bibekananda Nayak, Suraj Kumar Sahu, Saswati Suryasnata, Champakeswara Mahanta, Annapurna Behera, Satyabrata Sahoo, Akshyurna Pattnaik, Rakhi Rani Biswas, Chandrama Dipa, Adyasha Bijay Mishra, Dinesh Kumar Panda, Sudipta Nayak, Avishek Kar, Kajal Sundaray, Swagatika Dash, Alok Pattanaik, Sonali Swapanjali Bhoi, Swikruti Mishra, Shalini Das, Parinita Mishra, Debidatta Barik, Tejaswini Das, Subhashree Pattnaik (**IMMT**), Kailash T Bhamare, Chander Shekhar Sharma, Amit Kumar, Vineet Kumar, Deepak Bhatt, Surjeet Singh, Harminder Singh, Paramjit Lal, Sandeep Kumar, Nitin Sharma, Jaideep Mehta, Renu, Rohtas Ranga, RK Dhiman, Anjali Koundal, Bhumika Vaidya, Amit Kumar, Shweta Pandey, Anunay Sinha, Joyshree Das, Shubham, Bulbul Roy, Pompi Bhadra, Meena Sharma, Sana Khatun, Payal Thakur, Bhagyashree Rabha, Meenal Rastogi, Mayur Sudhakar Zarkar, Somnath Chindhe, Shreya Singh, Neelam (**IMTECH**), Kiran Raj S, Ravi Kumar Sharma Kumar (**IPU**), Nataraj D, Dr. Swetha C Desai, Shivanna TB, Murthy BYK, Raghu G, Sunil A, Shashidhara KN, Raghavendra Swamy M, Aravind Kumar Joseph, Purushotham HM, Shiva Kumar SR, Muniprakash M, Shekhar Gogeri, Siva Kumar R, Jaikesh S, Ranganatha S, SatishKumar Shri, Shashikala U, Praveen Kumar JD, Jitendra Ping (**NAL**), Koteswara K Rao, Parveen Goyal, NMR Ashwin, Rakesh S Joshi, Rupali Waichal, Shashikala Ranjane, Abujunaid Khan, Priyanka M Bankar, Rachel Samson, Apurva R Barge, Ashish S Jagtap, Amreen Sheikh, Shahaji Palaskar, Jugal Kanerkar, Arvind Chourasiya, Bhagyashree Likhitkar, Pradeep S, Deepak Jadhav, Pranay Awathare, Swaraj Jathar, Pooja Deshmukh, Vaishnavi Salunkhe, Bharat Pande, Aruna Choughule, Yugendra Patil, Rohit S Dashpute, Minal R Bhalerao, Vinod Kamble, Yashodhara Shinde, Vineetkumar S Nair, Prathmesh Ghongade, Prashant Kalaskar, Ganesh Jadhav, Sakshi Pingle, Preshitta Bhat, Shubham Choure, Tanaji B Devkate, Vikram Nichit, Shrutika M Shewale, Shyam K Gawari, Sonali Gaikwad, Abhishek Tripathi, Sayli Jamdade, Ashtamy MG, Shubham Kumar, Shivani V Palkar, Rutuja Ugle, Kaumudi Joshi, Khushboo Bavishi, Priya Bhavsar, Nandini Rangari, Ayushi Rushesari, Maria Kuwazerwala, Taha Farhan Siddiqui, Anand Kumar Shukla, Prakashini Saroj Nilgirwar, Sakshi R Mangate, Sindhu Bali, Prabha Oli (**NCL**), Pankaj Pardhi (**NEERI**), Jayashree Chiring Phukon, Shridhar Hiremath, Prachurjya

Dutta, Parishmita Borgohain, Moirangthem Goutam Singh, Devpratim Koch, Dipanneeta Das Gupta, Pankaj Barman, Sahana SK, Monojit Kumar Roy, Aditya Sarkar, Bhaben Sharma, Shyamalima Mech, Masum Saikia, Trishna Rani Borah, Gayatri Gogoi, Anupriya Borah, Udeshna Changmai, Himadri Das, Anshuman Goswami, Rocktotpal Mahanta, Rina Yumnam, Sukanya Borthakur, Manabendra Borah, Sarangthem Dinamani Singh, Pronami Gogoi, Priyanka Saikia, Umakanta Tanti, Mohan Kurmi, Ravi Kumar Sahu, Nayan Jyoti Borah, Babli Borah, Sindhu Sharma, Trishna Dutta, Sarifa Mafuz Ahmed, Bidyut Prakash Deka, Darshana Tamuli, Meghna Kakoty, Bhaswat V. Borah, Yashodhara Goswami, Priyanka Boro (**NEIST**), Inshamol KP, Athulya, Akhila, Amrutha M, Anaga, Anaswara P A, Anirudh, Anjitha, Anupama, Anusha, Anusree P, Aparna, Aparna Dinil, Arathy, Archa R S, Archana, Arnold, Athira V C, Biji Raphy, Dileep R Nair, Evan, Gayathri, Gopika R, Greeshma, Greeshma Jayan, Karthika N, Karthika Nath, Kavya Mohan, Krishan Unni, Lakshmi M Nair, Lakshmi Shaji, Nandana A B, Neeraja, Neeraja M, Neetha P Sobandas, Parvathy, Reena R, Saranyadevi, Shahansha, Shamna Fathima, Shehbas C, Shilpa, Sruthi, Swapna B, Varsha, Vishnu, Valan Rebinro, Adarsh VP, Jedy Jose (**NIIST**), Govind Ranade, AS Unnikrishnan, Sohan Pal Meena, Karishma Pradeep Chari, Vitasta Jad, Vikash Kumar, Nishamol M, Helen Agnes, Natasha Maria Barnes, Bhumika Rohidas Shirodkar, Shania Waluscha Moeres, Yuvrani Halarnkar, Ramila Ram Gaonkar, Mrinalini Chandra Mohan, Pratibha Bachhley (**NIO**), Himani Meena, Abhinav Banait, Shekar Sharma (**NIScPR**), Ashok Kr Shaw, T Kr Minz, Rahul Tirkey, Mousumi Sarkar, Chandan Kumar Chowdhury, Raushan Kumar, Amit Kumar, Sudip Kumar, Surajit Kundu, Mahesh Pramanik, Deepak Dhivar, Aman Beldar, Tapan Kumar Shaw, Premlal, Akul Gope, Nisha Gupta, Alka Kumari, Vinod, Shikha Kumari, Abhishek Kumar Machhua, Uttam Mukhi, Rahul Mukhi, Manju, Jaydev Sahu, Ravi Ranjan Kumar (**NML**), Rajesh, Sumana Gajjala, Sudesh Yadav, Vinod Kumar Tanwar, Vishesh Garg, Manoj Kumar Pandey, Vikash Sharma (**NPL**)

Author Contributions

Phenome India Consortium Authors

Project conceptualization

Shantanu Sengupta, Debasis Dash, Viren Sardana, Kumardeep Chaudhary (**IGIB**), Giriraj Ratan Chandak, Swasti Raychaudhuri, Karthik Bharadwaj Tallapaka (**CCMB**), Mahesh J Kulkarni (**NCL**), Partha Chakraborty, Dipyaman Ganguly (**IICB**), Umakanta Subudhi (**IMMT**).

Project planning/Collection management

Shantanu Sengupta, Debasis Dash, Viren Sardana, Kumardeep Chaudhary, Vamsi K. Yenamandra, Rakesh Sharma, Aastha Mishra, Ajay Pratap Singh, Swarnendu Bag, Beena Pillai, Vivek Rao, Sheetal Gandotra, Shantanu Chowdhury, Rajesh Pandey, Bhavana Prasher, Pankaj Pandey, Ankita Sahu, Sudhir Rohilla, Komal Jindal, Vivek Junghare, Tarani Mathur, Meghana Arvind, Satyartha Prakash, Yogesh Kumar, Vikas M Hiremath, Vignesh S Kumar, Deeksha Yadav, Deepak (**IGIB**), Swasti Raychaudhuri, Giriraj Ratan Chandak, Karthik Bharadwaj Tallapaka, Rakhesh K V (**CCMB**), Ashok P Giri, Narendra Y Kadoo, Mahesh J Kulkarni, Dhanasekaran Shanmugam, Mahesh S Dharne, Syed G Dastager, Chiranjit Chowdhury (**NCL**), Partha Chakraborty, Dipyaman Ganguly, Arun Bandyopadhyay, Shilpak Chatterjee (**IICB**), Umakanta Subudhi, Bhavani S Jena, Trupti Das, Boopathy Ramasamy, Pavan Kumar (**IMMT**), Ramakrishna Sistla, Prabhakar Sripadi, Jagadeshwar Reddy Thota, Ramesh Ummanni, Srinivasa Rao M, Sai Balaji Andugulapati (**IICT**), Amit Lahiri, Mrigank Srivastava, Vivek Bhosale (**CDRI**), Iranna Gogeri (**4PI**), Raju Khan, Narendra Singh (**AMPRI**), Rajesh Kumar Verma, Neeraj Jain (**CBRI**), Ganesh Venkatachalam, Murugan Veerapandian (**CECRI**), Deepak Bansal, Amit Kumar, Dinesh Gupta(**CEERI**), Prakash M Halami, S P Muthukumar (**CFTRI**), Anirban Pal, Anil Kumar Maurya (**CIMAP**), Jai Krishna Pandey, Bhanu Pandey, A K Raman (**CIMFR**), Suresh Kumar Anandasadagopan (**CLRI**), Swati Saha, Vishal Anand (**CMERI**), Suman Singh, Anamika Kothari (**CSIO**), Avinash Mishra, Mangal S Rathore (**CSMCRI**), Preeti Srivastava, Pooja Aggarwal (**HQ**), Shreedhar Kanagarjan (**HRDG**), Yogendra Padwad, Vikram Patial (**IHBT**), Sumit G Gandhi (**IIIM, Jammu**), Fayaz Malik (**IIIM, Sinagar**), Debashish Ghosh, Jyoti Porwal (**IIP**), Vikas Srivastava, Prabhanshu Tripathi (**IITR**), Anshu Bhardwaj, Srinivasan Krishnamurthi, Rashmi Kumar , Deepak Sharma, Amit Tuli (**IMTECH**), Indrani Ghosh (**IPU**), Prakash L, Satisha Shri (**NAL**), Chandana Venkateswara Rao, Sanjeev Kumar Ojha, Brahma Nanda Singh, Vijayanandraj Selvaraj (**NBRI**), Shilpa Paranjape, Prashanti Niwant (**NEERI**), Jatin Kalita, Prasenjit Manna, Romi Wahengbam, Tridip Phukan,

Pankaj Bharali (**NEIST**), E V S S K Babu, Biswajit Mandal, T Vijaya Kumar (**NGRI**), Rajeev K Sukumaran, Rameshkumar N (**NIIST**), Samir Ravikant Damare, Sameer Damare,Kalpana Sandesh Chodankar (**NIO**), Arvind Meena, Arun Uniyal (**NIScPR**), Ansu J Kailath, Krishna Kumar, Roshan Kumar, Nikhil Kumar, Kuldeep Singh Gour (**NML**), Ved Varun Agrawal, Arun Kant Singh (**NPL**), Maheswaran Srinivasan, Vasudevan Pandurangan (**SERC**), Lalita Goyal (**TKDL**), Rashmi Arya, Manisha Sakpal (**URDIP**).

Portal development and management
Kumardeep Chaudhary, Debasis Dash, Viren Sardana, Ajay Pratap Singh, Shantanu Sengupta, Satyartha Prakash, Vignesh S Kumar, Anshul Verma, Safeer Khan, Anshul Bhardwaj, Sudhir Rohilla, Prateek Singh (**IGIB**), Gopal Krishna Patra (**4PI**), Nikhilesh Yadav (**NCL**), Abbani Rakesh (**NAL**)

Sample Collection / Scanning
Pankaj Pandey, Sudhir Rohilla, Vivek Junghare, Komal Jindal, Ankita Sahu, Satyartha Prakash, Meghana Arvind, Rajat Ujjainiya, Tarani Mathur, Shilpa Ray, Ayushi Narayan, Harleen Kaur, Vishu Gupta, Mohit Kumar Divakar, Mamta Rathore, Prateek Singh, Vikas M Hiremath, Saraswati Awasthi, Kanika Singh, Divya Bhalla, Anubhuti Bansal, Yogesh Kumar, Vignesh S Kumar, Shivani Chitkara, Charvy Rana, Amit Maurya, Sreeshma Raj K, Anshul Verma, Pulkit Hasmukhbhai Leuva, Pratik Pathade, Safeer Khan, Anshul Bhardwaj, Bharti, Shail Kumari, Bharti sharma, Ruchi, Shubham Kumar , Tanmay pawaskar, Sumant Kumar, Mohit, Shyam Singh Bisht, Deeksha Yadav, Prajwali Sawant, Rohan Bhardwaj, Shailja kant Upadhyay, Rushikesh joshi, Shaivyanand Singh, Shivam Dhawan, Vanshika Srivastava, Dikkshita Baruah, Rohit Kumar, Praveena Mishra, Ankit Basnal, Md. Intyaz Ali, Pranjal Tewari, Swati, Abhishek Kumar, Nancy Rawat , Azhar uddin, Kamal singh Pindari, Deepak (**IGIB**), MK Kanakavalli, Rakhesh K V, Neha Kumari, Jukanti Akshitha, Deepshikha Esari, Mohammed Osed Annan, Mahfuj Hasan, Dibya Rana Saha Roy (**CCMB**), Ajit A Sutar, Sagar Baulia, Ameya A Pawar, Monika Sharma, Milind Kale, Ankita Namdeo, Rajesh S, Rashdajabeen Q Shaikh, Vaishnavi N Mahajan (**NCL**), Ankita Mridha, Saheli Chowdhury, Pratitusti Basu, Dipamoy Dutta, Dwaipayan Saha, Ruby Banerjee (**IICB**), Sai Adarsh Sahu, Sk Rameej Raja (**IMMT**), Abhisheik Eedara, Vishwa Priya, Navya Sahithi Pelimelli, srilekha Anumulapuri (**IICT**), Shail Singh, Lakra Promila, Swarnali Basu, Rahul Roy, Shweta Tiwari, Shikha Yadav, Amit Kumar Shahravat, Kabita Sarkar, Adrija Rakshit, Deepanshu Sindhwani, Kajal KM, Smita Pandey (**CDRI**), Sangavi Pakkyam (**CECRI**), Vipul Sharma (**CEERI**),

Dikchha Singh (**CIMFR**), Parimala karupannan (**CLRI**), Ravi Raj, Ankita Kumari (**IHBT**), Nancy Sharma, Sahaurti Sharma, Sakshi Nagial (**IIIM,Jammu**), Mir Shahid Maqbool, Kaneez Fatima (**IIIM,Srinagar**), Suchismita Benjwal, Pramod Chauhan (**IIP**), Neha Mehrotra (**IITR**), Parvez Ahmad, Priyadarshan Kinatukara, Bhupender Singh, Shiva Sundharam S, Pranavathiyani G, Kuldeep Singh, Rakesh Kumar, Pradip Sen, Siddhakam Palmal, Ritu Jatav, Lalit Kumar, Pravin Kumar, Priyanshu Singh Raikwar, Simran Gambhir (**IMTECH**), Madan Mohan Pandey (**NBRI**), Manuj Kr Das, Borsha Rajkumari, Sukanya Borkakoti, Mamta Thapa, Ashique Hussain, Ishant Jyoti Nath (**NEIST**), Rahul R Menon, Sanoj Mohan, Arnold Moses (**NIIST**), Akshika (**NIScPR**), Navneet Singh Randhawa, Priyanka Singh, K Sudhakara Rao (**NML**).

Sample aliquoting

Aastha Mishra, Ankita Sahu, Meghana Arvind, Shivani Chitkara, Deeksha Yadav, Mohit, Shyam Singh Bisht, Shail Kumari, Ankit Basnal, Ansuman Sahu, Ruchi, Tanmay Pawaskar, Nancy Rawat, Pranjal Tewari, Vignesh S Kumar, Safeer Khan, Anshul Bhardwaj (**IGIB**)

Assays

Shantanu Sengupta, Viren Sardana, Kumardeep Chaudhary, Vamsi K. Yenamandra, Rakesh Sharma, Aastha Mishra, Vivek Rao, Shantanu Chowdhury, Beena Pillai, Sheetal Gandotra, Rajesh Pandey Ankita Sahu, Satyartha Prakash, Vignesh S Kumar, Safeer Khan, Shilpa Ray, Tarani Mathur, Rajat Ujjainiya, Mamta Rathore, Yogesh Kumar, Ankur Halder, Azhar uddin, Shivani Chitkara, Md Quasid Akhter, Bharti, Shubham Kumar, Sumant Kumar, Shail Kumari, Anshul Bhardwaj, Ruchi, Deeksha Yadav, Jitendra Kumar, Abhishek Kumar, Swati (**IGIB**), Partha Chakraborty, Dipyaman Ganguly (**IICB**). Giriraj Ratan Chandak, Swasti Raychaudhuri, Karthik Bharadwaj Tallapaka, Jukanti Akshitha, Dimple Lavanuru, Sumati Swain (**CCMB**), Mahesh J Kulkarni, Chiranjit Chowdhury, Rajesh S, Ajit A Sutar, Ameya A Pawar, Milind Kale (**NCL**), Dipyaman Ganguly, Partha Chakraborty, Saikat Majumder, Shilpak Chatterjee, Jahangir Alam (**IICB**), Umakanta Subudhi, Sai Adarsh Sahu, Sk Rameej Raja (**IMMT**) Ramakrishna Sistla, Sai Balaji Andugulapati (**IICT**), Amit Lahiri, Mrigank Srivastava, Shail Singh, Lakra Promila, Swarnali Basu, Shweta Tiwari, Shikha Yadav, Kabita Sarkar, Adrija Rakshit, Deepanshu Sindhwani (**CDRI**)

Genomic DNA Isolation

Aastha Mishra, Ankita Sahu, Shubham Kumar, Shyam Singh Bisht, Safeer Khan, Bharti, Ruchi, Tanmay Pawaskar, Rohit Kumar, Nancy Rawat, Swati, Pranjal Tewari (**IGIB**).

Data management and curation

Kumardeep Chaudhary, Viren Sardana, Shantanu Sengupta, Ajay Pratap Singh, Ankita Sahu, Meghana Arvind, Satyartha Prakash, Vignesh S Kumar, Pulkit Hasmukhbhai Leuva, Anshul Verma, Sreeshma Raj K, Pratik Pathade, Tarani Mathur, Kalyani Verma, Yogesh Kumar, Md. Intyaz Ali, Sudhir Rohilla (**IGIB**)

Manuscript writing, review and editing

Shantanu Sengupta, Viren Sardana, Kumardeep Chaudhary, Vamsi K. Yenamandra, Satyartha Prakash, Vignesh S Kumar, Anshul Verma, Sreeshma Raj K, Pulkit Hasmukhbhai Leuva (**IGIB**),Giriraj Ratan Chandak, Swasti Raychaudhuri, Karthik Bharadwaj Tallapaka (**CCMB**), Mahesh J Kulkarni (**NCL**), Partha Chakraborty, Dipyaman Ganguly (**IICB**), Umakanta Subudhi (**IMMT**).

Conflicts of Interest

All the authors declare no conflict of interest.

Data Availability

Anonymized data for public use may be made available after three years from completion of baseline phase of study or as per advisory from Monitoring Committee of the project, if any revisions thereof.

List of Abbreviations

AC	:	Abdominal Circumferences
ADA	:	American Diabetes Association
AI	:	Artificial Intelligence
API	:	Application Programming Interface
ATS	:	American Thoracic Society
BCA	:	Body Composition Analysis
BMI	:	Body Mass Index
CAP	:	Continuous Attenuation Parameter
CAP	:	College of American Pathologists
CC	:	Chest Circumferences
CLI	:	Command-Line Interfaces
CRF	:	Cryomed Controlled-Rate Freezer
CSIR	:	Council of Scientific and Industrial Research
CSS	:	Cascading Style Sheets
CTRI	:	Clinical Trials Registry- India
CVD	:	Cardiovascular Diseases
ECG	:	Echocardiogram
EDTA	:	Ethylenediamine Tetraacetic Acid
ERS	:	European Respiratory Society
FBS	:	Fasting Blood Sugar
FEV1	:	Forced Expiratory Volume
FRS	:	Framingham Risk Score
FVC	:	Forced Vital Capacity
GSA	:	Global Screening Array
GUI	:	Graphical User Interface
HbA1c	:	Glycated Hemoglobin
HC	:	Hip Circumferences
HRV	:	Heart Rate Variability
ICDS	:	Integrated Child Development Services

ICP-MS	:	Inductively Coupled Plasma Mass Spectrometer
IHEC	:	Institutional Human Ethics Committee
LMICs	:	Low and Middle-Income Countries
LSM	:	Liver Stiffness Measurement
ML	:	Machine Learning
MRM	:	Multiple Reaction Monitoring
NABL	:	National Accreditation Board for Testing and Calibration Laboratories
NAFLD	:	Nonalcoholic Fatty Liver Disease
NASH	:	Nonalcoholic Steato-Hepatitis
NCDs	:	Non-Communicable Diseases
OAD	:	Obstructive Airway Disease
PI-CheCK	:	Phenome India-CSIR Health Cohort Knowledgebase
qPCR	:	Quantitative Polymerase Chain Reaction
RBAC	:	Role-Based Access Control
RDBMS	:	Relational Database Management System
REDCap	:	Research Electronic Data Capture
RPM	:	Revolutions Per Minute
SBS	:	Sequence By Synthesis
SFTP	:	Secure File Transfer Protocol
SQL	:	Structured Query Language
SST	:	Serum-Separating Tube
TB	:	Terabytes
TE	:	Transient Elastography
TEWL	:	Transepidermal Water Loss
UT	:	Union Territories
VPN	:	Virtual Private Network
WC	:	Waist Circumferences

References

1.   Noncommunicable diseases 2023 [Available from: https://www.who.int/news-room/fact-sheets/detail/noncommunicable-diseases.
2.   Joshi P, Islam S, Pais P, Reddy S, Dorairaj P, Kazmi K, et al. Risk Factors for Early Myocardial Infarction in South Asians Compared With Individuals in Other Countries. Jama. 2007;297(3):286-94.
3.   Prabhakaran D, Jeemon P, Roy A. Cardiovascular Diseases in India. Circulation. 2016;133(16):1605-20.
4.   Younossi Z, Anstee QM, Marietti M, Hardy T, Henry L, Eslam M, et al. Global burden of NAFLD and NASH: trends, predictions, risk factors and prevention. Nature Reviews Gastroenterology & Hepatology. 2018;15(1):11-20.
5.   Li J, Wang Q, Ni W, Liu C, Li Z, Qi X. Global health burden of cirrhosis and other chronic liver diseases (CLDs) due to non-alcoholic fatty liver disease (NAFLD): A systematic analysis for the global burden of disease study 2019. Global Transitions. 2023;5:160-9.
6.   Prabhakar T, Prasad M, Kumar G, Kaushal K, Shenoy PS, Dubey S, et al. High prevalence of MAFLD in general population: A large cross-sectional study calls for concerted public health action. Alimentary Pharmacology & Therapeutics. 2024;59(7):843-51.
7.   Majumder PP, Basu A. A genomic view of the peopling and population structure of India. Cold Spring Harbor perspectives in biology. 2014;7(4):a008540.
8.   Sharma A, Lysenko A, Jia S, Boroevich KA, Tsunoda T. Advances in AI and machine learning for predictive medicine. Journal of Human Genetics. 2024.
9.   Ahmed Z, Wan S, Zhang F, Zhong W. Artificial intelligence for omics data analysis. BMC Methods. 2024;1(1):4.
10.   Bien SA, Wojcik GL, Hodonsky CJ, Gignoux CR, Cheng I, Matise TC, et al. The Future of Genomic Studies Must Be Globally Representative: Perspectives from PAGE. Annual review of genomics and human genetics. 2019;20:181-200.
11.   Mehta LS, Velarde GP, Lewey J, Sharma G, Bond RM, Navas-Acien A, et al. Cardiovascular Disease Risk Factors in Women: The Impact of Race and Ethnicity: A Scientific Statement From the American Heart Association. Circulation. 2023;147(19):1471-87.
12.   Mishra A, Malik R, Hachiya T, Jürgenson T, Namba S, Posner DC, et al. Stroke genetics informs drug discovery and risk prediction across ancestries. Nature. 2022;611(7934):115-23.
13.   Wells JC. Commentary: Why are South Asians susceptible to central obesity? — the El Niño hypothesis. International journal of epidemiology. 2007;36(1):226-7.
14.   Chaudhary M, Sharma P. Abdominal obesity in India: analysis of the National Family Health Survey-5 (2019–2021) data. The Lancet Regional Health - Southeast Asia. 2023;14.
15.   Ying W. Phenomic Studies on Diseases: Potential and Challenges. Phenomics (Cham, Switzerland). 2023;3(3):285-99.
16.   Naushin S, Sardana V, Ujjainiya R, Bhatheja N, Kutum R, Bhaskar AK, et al. Insights from a Pan India Sero-Epidemiological survey (Phenome-India Cohort) for SARS-CoV2. eLife. 2021;10:e66537.
17.   Kondal D, Patel SA, Ali MK, Mohan D, Rautela G, Gujral UP, et al. Cohort Profile: The Center for cArdiometabolic Risk Reduction in South Asia (CARRS). International journal of epidemiology. 2022;51(6):e358-e71.

18.   Aggarwal M, Garg NM, Agrawal A, Sardana V. Lung Function Reference Equations for Indian Ethnic Groups Based on a Handheld Forced Oscillation Device for Age 9-19 Years. Indian journal of pediatrics. 2023;90(1):61-8.

19.   Anjana RM, Unnikrishnan R, Deepa M, Pradeepa R, Tandon N, Das AK, et al. Metabolic non-communicable disease health report of India: the ICMR-INDIAB national cross-sectional study (ICMR-INDIAB-17). The Lancet Diabetes & Endocrinology. 2023;11(7):474-89.

20.   Huffman MD, Prabhakaran D, Osmond C, Fall CH, Tandon N, Lakshmy R, et al. Incidence of cardiovascular risk factors in an Indian urban cohort results from the New Delhi birth cohort. Journal of the American College of Cardiology. 2011;57(17):1765-74.

21.   Rastogi A, Rath I, Varadarajan A, Ramakrishna G, Bihari C, Maiwall R. Non-alcoholic fatty liver disease (NAFLD) in lean individuals – Single centre large cohort clinicopathologic and immunophenotypic study. Pathology - Research and Practice. 2022;238:154112.

22.   Pradeepa R, Mohan V. Epidemiology of type 2 diabetes in India. Indian journal of ophthalmology. 2021;69(11):2932-8.

23.   Ramakrishnan A, Velmurugan G, Somasundaram A, Mohanraj S, Vasudevan D, Vijayaragavan P, et al. Prevalence of abnormal liver tests and liver fibrosis among rural adults in low and middle-income country: A cross-sectional study. EClinicalMedicine. 2022;51:101553.

24.   Sreeniwas Kumar A, Sinha N. Cardiovascular disease in India: A 360 degree overview. Medical journal, Armed Forces India. 2020;76(1):1-3.

25.   Pourhoseingholi MA, Vahedi M, Rahimzadeh M. Sample size calculation in medical studies. Gastroenterology and hepatology from bed to bench. 2013;6(1):14-7.

26.   Collins GS, Reitsma JB, Altman DG, Moons KGM. Transparent reporting of a multivariable prediction model for individual prognosis or diagnosis (TRIPOD): the TRIPOD Statement. BMC Medicine. 2015;13(1):1.

27.   von Elm E, Altman DG, Egger M, Pocock SJ, Gøtzsche PC, Vandenbroucke JP. The Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) statement: guidelines for reporting observational studies. Lancet (London, England). 2007;370(9596):1453-7.

28.   Harris PA, Taylor R, Thielke R, Payne J, Gonzalez N, Conde JG. Research electronic data capture (REDCap)—A metadata-driven methodology and workflow process for providing translational research informatics support. Journal of Biomedical Informatics. 2009;42(2):377-81.

29.   Harris PA, Taylor R, Minor BL, Elliott V, Fernandez M, O'Neal L, et al. The REDCap consortium: Building an international community of software platform partners. Journal of Biomedical Informatics. 2019;95:103208.

30.   Norton K, & Eston, R. (Eds.). . Kinanthropometry and Exercise Physiology (4th ed.).2018.

31.   Peterson DD. Periodic Fitness Testing: Not Just for Athletes Anymore. Strength & Conditioning Journal. 2018;40(5).

32.   Beyer SE, Sanghvi MM, Aung N, Hosking A, Cooper JA, Paiva JM, et al. Prospective association between handgrip strength and cardiac structure and function in UK adults. PloS one. 2018;13(3):e0193124.

33.   Fess E. Clinical assessment recommendations. American society of hand therapists. 1981:6-8.

34.   Jaafar MH, Ismail R, Ismail NH, Md Isa Z, Mohd Tamil A, Mat Nasir N, et al. Normative reference values and predicting factors of handgrip strength for dominant and non-dominant hands among healthy Malay adults in Malaysia. BMC Musculoskeletal Disorders. 2023;24(1):74.

35.   Rout G, Kedia S, Nayak B, Yadav R, Das P, Acharya SK, et al. Controlled Attenuation Parameter for Assessment of Hepatic Steatosis in Indian Patients. Journal of Clinical and Experimental Hepatology. 2019;9(1):13-21.

36.   Shalimar, Kumar R, Rout G, Kumar R, Yadav R, Das P, et al. Body mass index-based controlled attenuation parameter cut-offs for assessment of hepatic steatosis in non-alcoholic fatty liver disease. Indian journal of gastroenterology : official journal of the Indian Society of Gastroenterology. 2020;39(1):32-41.

37.   Eddowes PJ, Sasso M, Allison M, Tsochatzis E, Anstee QM, Sheridan D, et al. Accuracy of FibroScan Controlled Attenuation Parameter and Liver Stiffness Measurement in Assessing Steatosis and Fibrosis in Patients With Nonalcoholic Fatty Liver Disease. Gastroenterology. 2019;156(6):1717-30.

38.   Miller MR, Hankinson J, Brusasco V, Burgos F, Casaburi R, Coates A, et al. Standardisation of spirometry. European Respiratory Journal. 2005;26(2):319-38.

39.   Moore VC. Spirometry: step by step. Breathe. 2012;8(3):232-40.

40.   Graham BL, Steenbruggen I, Miller MR, Barjaktarevic IZ, Cooper BG, Hall GL, et al. Standardization of Spirometry 2019 Update. An Official American Thoracic Society and European Respiratory Society Technical Statement. American Journal of Respiratory and Critical Care Medicine. 2019;200(8):e70-e88.

41.   Wanger J, Clausen JL, Coates A, Pedersen OF, Brusasco V, Burgos F, et al. Standardisation of the measurement of lung volumes. European Respiratory Journal. 2005;26(3):511-22.

42.   Miura S, Iwamoto H, Omori K, Yamaguchi K, Sakamoto S, Horimasu Y, et al. Preserved ratio impaired spirometry with or without restrictive spirometric abnormality. Scientific reports. 2023;13(1):2988.

43.   King GG, Bates J, Berger KI, Calverley P, de Melo PL, Dellacà RL, et al. Technical standards for respiratory oscillometry. The European respiratory journal. 2020;55(2).

44.   Jan B, Dar MI, Choudhary B, Basist P, Khan R, Alhalmi A. Cardiovascular Diseases Among Indian Older Adults: A Comprehensive Review. Cardiovascular Therapeutics. 2024;2024(1):6894693.

45.   Dawber TR, Meadors GF, Moore FE, Jr. Epidemiological approaches to heart disease: the Framingham Study. American journal of public health and the nation's health. 1951;41(3):279-81.

46.   Gordon T, Kannel WB. Predisposition to atherosclerosis in the head, heart, and legs. The Framingham study. Jama. 1972;221(7):661-6.

47.   Kannel WB, McGee D, Gordon T. A general cardiovascular risk profile: the Framingham Study. The American journal of cardiology. 1976;38(1):46-51.

48.   D'Agostino RB, Russell MW, Huse DM, Ellison RC, Silbershatz H, Wilson PW, et al. Primary and subsequent coronary risk appraisal: new results from the Framingham study. American heart journal. 2000;139(2 Pt 1):272-81.

49.   Hemann BA, Bimson WF, Taylor AJ. The Framingham Risk Score: An Appraisal of Its Benefits and Limitations. American Heart Hospital Journal. 2007;5(2):91-6.

50.   Chow CK, Joshi R, Celermajer DS, Patel A, Neal BC. Recalibration of a Framingham risk equation for a rural population in India. Journal of epidemiology and community health. 2009;63(5):379.

51.   Gupta P, Prieto-Merino D, Ajay V, Singh K, Roy A, Krishnan A, et al. Cardiovascular risk prediction in India: Comparison of the original and recalibrated Framingham prognostic models in urban populations. [version 2; peer review: 2 approved]. Wellcome Open Research. 2019;4(71).

52.   D K, Swaminathan K, Velmurugan G, Ramakrishnan A, Alexander T, Raghupathy AK, et al. Association between serum uric acid levels and cardiovascular risk factors among adults in India. Nutrition, metabolism, and cardiovascular diseases : NMCD. 2023;33(7):1330-8.

53.   Shrivastava U, Misra A, Mohan V, Unnikrishnan R, Bachani D. Obesity, Diabetes and Cardiovascular Diseases in India: Public Health Challenges. Current diabetes reviews. 2017;13(1):65-80.

54.   Unnikrishnan AG, Sahay RK, Phadke U, Sharma SK, Shah P, Shukla R, et al. Cardiovascular risk in newly diagnosed type 2 diabetes patients in India. PloS one. 2022;17(3):e0263619.

55.   Geldsetzer P, Manne-Goehler J, Theilmann M, Davies JI, Awasthi A, Danaei G, et al. Geographic and sociodemographic variation of cardiovascular disease risk in India: A cross-sectional study of 797,540 adults. PLoS medicine. 2018;15(6):e1002581.

56.   Mallinson PAC, Kulkarni B, Bhogadi S, Kinra S. Association between parents' socioeconomic conditions and nutritional status during childhood and the risk of cardiovascular disease in their adult offspring: an intergenerational study in south India. Journal of epidemiology and community health. 2021;75(11):1091-7.

57.   Ujjainiya R, Tyagi A, Sardana V, Naushin S, Bhatheja N, Kumar K, et al. High failure rate of ChAdOx1-nCoV19 immunization against asymptomatic infection in healthcare workers during a Delta variant surge. Nature Communications. 2022;13(1):1726.

58.   Yadav PD, Sardana V, Deshpande GR, Shinde PV, Thangaraj JWV, George LS, et al. Neutralizing antibody responses to SARS-CoV-2 Omicron variants: Post six months following two-dose & three-dose vaccination of ChAdOx1 nCoV-19 or BBV152. The Indian journal of medical research. 2024;159(2):223-31.

59.   Dhar MS, Marwal R, Vs R, Ponnusamy K, Jolly B, Bhoyar RC, et al. Genomic characterization and epidemiology of an emerging SARS-CoV-2 variant in Delhi, India. Science. 2021;374(6570):995-9.

Supplementary Table S1: List of CSIR Institutes serving as sample collection centers stratified by zones in PI-CHeCK study.

Institute	Institute_Name	City Name	Zone
CSIR-4PI	CSIR Fourth Paradigm Institute	Bengaluru	South
CSIR-AMPRI	Advanced Materials and Processes Research Institute	Bhopal	West
CSIR-CBRI	Central Building Research Institute	Roorkee	North
CSIR-CCMB	Centre for Cellular and Molecular Biology	Hyderabad	South
CSIR-CDRI	Central Drug Research Institute	Lucknow	North
CSIR-CECRI	Central Electrochemical Research Institute	Karaikudi	South
CSIR-CEERI	Central Electronics Engineering Research Institute	Pilani	West
CSIR-CFTRI	Central Food Technological Research Institute	Mysore	South
CSIR-CGCRI	Central Glass and Ceramic Research Institute	Kolkata	West
CSIR-CIMAP	Central Institute of Medicinal & Aromatic Plants	Lucknow	North
CSIR-CIMFR	Central Institute of Mining and Fuel Research	Dhanbad	East
CSIR-CLRI	Central Leather Research Institute	Chennai	South
CSIR-CMERI	Central Mechanical Engineering Research Institute	Durgapur	East
CSIR-CRRI	Central Road Research Institute	New Delhi	North
CSIR-CSIO	Central Scientific Instruments Organisation	Chandigarh	North
CSIR-CSMCRI	Central Salt & Marine Chemicals Research Institute	Bhavnagar	West
CSIR-IGIB	Institute of Genomics & Integrative Biology	Delhi	North
CSIR-IHBT	Institute of Himalayan Bioresource Technology	Palampur	North
CSIR-IICB	Indian Institute of Chemical Biology	Kolkata	East
CSIR-IICT	Indian Institute of Chemical Technology	Hyderabad	South
CSIR-IIIM	Indian Institute of Integrative Medicine	Jammu	North
CSIR-IIP	Indian Institute of Petroleum	Dehradun	North
CSIR-IITR	Indian Institute of Toxicology Research	Lucknow	North
CSIR-IMMT	Institute of Minerals and Materials Technology	Bhubnaneswar	East
CSIR-IMTECH	Institute of Microbial Technology	Chandigarh	North
CSIR-NAL	National Aerospace Laboratories	Bengaluru	South

CSIR-NBRI	National Botanical Research Institute	Lucknow	North
CSIR-NCL	National Chemical Laboratory	Pune	West
CSIR-NEERI	National Environmental Engineering Research Institute	Nagpur	West
CSIR-NEIST	North-East Institute of Science & Technology	Jorhat	East
CSIR-NGRI	National Geophysical Research Institute	Hyderabad	South
CSIR-NIIST	National Institute for Inter-disciplinary Science and Technology	Thiruvananthapuram	South
CSIR-NIO	National Institute of Oceanography	Goa	West
CSIR-NIScPR	National Institute of Science Communication and Policy Research	New Delhi	North
CSIR-NML	National Metallurgical Laboratory	Jamshedpur	East
CSIR-NPL	National Physical Laboratory	New Delhi	North
CSIR-SERC	Structural Engineering Research Centre	Chennai	South
CSIR-URDIP	CSIR-Unit for Research and Development of Information Products	Pune	West

Supplementary Table S2: Details of exclusion (E) criteria for different scanning modalities.

	BCA	Liver Elastography	Skin	Oscillometry	Spirometry	ECG	Dynamometer
Pacemakers, intra-cardiac defibrillators/Loop recorders, or any implantable electronic device	E	E					
Contagious disease/skin disease	E	E	E	E	E	E	E
Suspicion or ongoing treatment of active pulmonary tuberculosis	E	E	E	E	E	E	E
Active injury or an active bleed in measurement area	E		E				
Metal implant, exclude dental implants (Specific Query)	E						
History of myocardial infarction/heart attack/ recent thoracic (chest) or abdominal surgery/Pneumothorax/Hemoptysis, recent eye or ear surgery/sinus surgery or any major surgery/ies (within the previous eight weeks)					E		E
Recent history of bleeding from the mouth or vomiting blood				E	E		
Loose teeth or have facial pain/TM joint issues				E	E		
Respiratory infection in the past month				E	E		
History of Thoracic (chest), abdominal, or cerebral aneurysm/stroke					E		E
Untreated/uncontrolled/ pulmonary hypertension					E		E

Supplementary Table S3: Description of self-reported Main questionnaire.

Domain	Variables
Baseline information	Consent, Gender, Date of Birth
Lifestyle	Brushing habit, Nail biting habit, Delivery mode, Exercise, Stress, Sleep, Bowel, Diet
Female-specific	Menstruation, Menopause, Pregnancies, Live birth, Abortion/stillbirths, Gynecological issues
Addiction	Smoking, Alcohol, Chewing, Other addictions
Health issues	High blood pressure (HBP), Diabetes, High blood cholesterol (HBC), Thyroid, Neurological problems, Heart problems, Respiratory problems, Liver problems, Bone/joint problems, Kidney problems, Eye problems, Gastro-intestinal problems, Hearing problems, Cancer, Depression, Anxiety, Allergies, Recurrent infections, Surgery
Family medical history	High blood pressure (Family), Diabetes, High blood cholesterol, Thyroid problems, Neurological Problems, Heart problems, Respiratory problems, Kidney problems, Gastro-intestinal problems, Liver problems, Bone/joint problems, Eye problems, Hearing problems, Cancer, Depression, Anxiety, Allergies, Recurrent infections
COVID-19	COVID-19 Infection, COVID-19 vaccination, Long term COVID-19 complications
Transportation	Type of vehicle family own, Metro travel, Car travel, Two-wheeler travel, Bus travel, Cycle Travel, Walking.

Supplementary Table S4: Description of the self-reported Gut questionnaire.

Domain	Variables
Food item consumption	Rice, Wheat (Paratha, chapati, roti, etc), Other cereals (Bajra, jowar, ragi etc), Bread(Yeast), Pulses (Arhar, tur, moong, urd daal etc), Sprouts, Eggs, Poultry (Chicken), Meat (Mutton, pork etc), Fish and other seafoods, Curd, Paneer, Ghee, Butter, Ice cream, Cake and pastries, Chips, namkeen, Pickles, Biscuits, Chocolate, Carbonated beverages (Coke, sprite etc), Fruit juice and shakes, Tubers and starchy vegetables (Potato, corn etc), Green leafy vegetables, Fruits (Banana, apple etc), Dry fruits (Resins, almonds, dates etc), Fermented foods (Idli, dosa, dhokla etc), Fermented drinks, Vitamin supplements, Protein supplements, Probiotics
Alcohol	Beer consumption within 2 days, Alcohol consumption within two days
Travel & eating	Types of green leafy vegetables consumed last week, Travel out of city or country in the last 15 days.
Antibiotic consumption	Last antibiotic consumption, Antibiotics consumption within two weeks
Hospitalization	Surgery in last one month, Hospitalized in last one month
Stool type	Bristol stool type

Supplementary Table S5: Summary of different sampling/scanning stations.

Station (Modality)	Machine	Information
Anthropometry	Dynamometer- Baseline	Muscle strength
BCA	BIA Analyzer-3 Frequency ACCUNIQ BC380	Body Composition
Phlebotomy	Details in Supplementary Table S6	Blood parameters
Transient Elastography	Fibroscan Mini 430Plus	LSM and CAP score
ECG	Allengers Pisces 1012	Heart condition
Spirometry	NDD Easy One Air	Lung Function and Volume
Oscillometry	Tremoflo C100, Resmon Pro V3	Small Airway Assessment
Skin	Courage Khazaka	Skin health

Supplementary Table S6: Details of the phlebotomy parameters, including biochemistry assays.

Diagnostic Lab-Outsourced					
S. No.	Units	Parameter	Sample Type	Platform	Sample Size
1	10^3/uL	Absolute Basophil Count	WB	DHBC	All
2	10^3/uL	Absolute Eosinophil Count (AEC)	WB	DHBC	All
3	10^3/uL	Absolute Lymphocyte Count (ALC)	WB	DHBC	All
4	10^3/uL	Absolute Monocyte Count	WB	DHBC	All
5	10^3/uL	Absolute Neutrophil Count (ANC)	WB	DHBC	All
6	U/L	Alanine Aminotransferase (ALT/SGPT)	Serum	AUBC	All
7	Ratio	Albumin/Globulin Ratio	Serum	AUBC	All
8	U/L	Alkaline Phosphatase (ALP)	Serum	AUBC	All
9	mg/dL	Apolipoprotein A1 (APO-A1)	Serum	AUBC	All
10	mg/dL	Apolipoprotein B (APO-B)	Serum	AUBC	All
11	U/L	Aspartate Aminotransferase (AST/SGOT)	Serum	AUBC	All
12	mg/dL	Average Estimated Glucose - plasma	Calculated	DHBC	All
13	%	Basophils	Serum	DHBC	All
14	mg/dL	Blood Urea	Serum	AUBC	All
15	mg/dL	Blood Urea Nitrogen (BUN)	Serum	AUBC	All

16	Ratio	Bun/Creatinine Ratio	Calculated	AUBC	All
17	%	Eosinophils	WB	DHBC	All
18	mm/1st hour	Erythrocyte Sedimentation Rate (ESR)	WB	ALIFAX TEST-1	All
19	ng/mL	Ferritin	Serum	ATELLICA IM (SIEMENS)	All
20	U/L	Gamma Glutamyl Transferase (GGT)	Serum	AUBC	All
21	mL/min/1 73m2	estimated glomerular filtration rate (eGFR)	Calculated	AUBC	All
22	mg/dL	Glucose, Fasting	Fluoride	AUBC	All
23	Ratio	Green and king index	WB	DHBC	All
24	g/dL	Haemoglobin (HB)	WB	DHBC	All
25	%	HbA1c (Glycosylated Hemoglobin)	WB	BDB	All
26	Ratio	High-density lipoprotein (HDL)/low-density lipoprotein (LDL) Cholesterol Ratio	Serum	AUBC	All
27	%	Hematocrit (PCV)	WB	DHBC	All
28	Ratio	LDL/HDL Cholesterol Ratio	Serum	AUBC	All
29	%	Lymphocytes	WB	DHBC	All
30	pg	Mean Corp Hb (MCH)	WB	DHBC	All
31	g/dL	Mean Corp Hb Conc (MCHC)	WB	DHBC	All

32	fL	Mean Corp Volume (MCV)	WB	DHBC	All
33	Ratio	Mentzer Index	WB	DHBC	All
34	%	Monocytes	WB	DHBC	All
35	fL	MPV	WB	DHBC	All
36	%	Neutrophils	WB	DHBC	All
37	mg/dL	Non-HDL Cholesterol	Serum	AUBC	All
38	10^3/uL	Platelet Count (PLT)	WB	DHBC	All
39	%	Red Cell Distribution Width - Coefficient of Variation (RDW-CV)–	WB	DHBC	All
40	fL	Red Cell Distribution Width-Standard Deviation (RDW - SD)	WB	DHBC	All
41	Ratio	Red Cell Distribution Width Index (RDWI)	WB	DHBC	All
42	10^6/uL	Red Blood Cell Count (RBC)	WB	DHBC	All
43	g/dL	Serum Albumin	Serum	AUBC	All
44	mg/dL	Serum Bilirubin, (Direct)	Serum	AUBC	All
45	mg/dL	Serum Bilirubin, (Indirect)	Serum	AUBC	All
46	mg/dL	Serum Bilirubin, (Total)	Serum	AUBC	All
47	mg/dL	Serum Calcium	Serum	AUBC	All
48	mmol/L	Serum Chloride	Serum	AUBC	All
49	mg/dL	Serum Creatinine	Serum	AUBC	All

50	gm/dL	Serum Globulin	Serum	AUBC	All
51	mg/dL	Serum HDL Cholesterol	Serum	AUBC	All
52	ug/dL	Serum Iron	Serum	AUBC	All
53	mg/dL	Serum LDL Cholesterol	Serum	AUBC	All
54	mg/dL	Serum Phosphorus	Serum	AUBC	All
55	mmol/L	Serum Potassium	Serum	AUBC	All
56	mmol/L	Serum Sodium	Serum	AUBC	All
57	µg/dL	Serum Total Iron Binding Capacity (TIBC)	Serum	AUBC	All
58	g/dL	Serum Total Protein	Serum	AUBC	All
59	mg/dL	Serum Triglycerides	Serum	AUBC	All
60	mg/dL	Serum Uric Acid	Serum	AUBC	All
61	mg/dL	Serum very low-density lipoprotein (VLDL) Cholesterol	Serum	AUBC	All
62	Ratio	SGOT/SGPT Ratio	Calculated	AUBC	All
63	uIU/mL	Thyroid Stimulating Hormone (TSH)-Ultrasensitive	Serum	AIS	All
64	Ratio	Total CHOL/HDL Cholesterol Ratio	Calculated	AUBC	All
65	mg/dL	Total Cholesterol	Serum	AUBC	All
66	10^3/uL	Total Leucocyte Count (TLC)	WB	DHBC	All
67	%	Transferrin Saturation %	Serum	AUBC	All

68	pg/mL	Unsaturated iron binding capacity (UIBC)	Serum	AUBC	All
69	Ratio	Urea/Creatinine Ratio	Calculated	AUBC	All
70	pg/mL	Vitamin B12	Serum	AIS	All
71	ng/mL	Vitamin D (25-OH Vitamin D)	Serum	AIS	All

WB: Whole Blood; DHBC: DxH-900 Beckman Coulter; AUBC: AU5800 Beckman Coulter; BDB: BIORAD D-100 (BIORAD); AIS: ATELLICA IM (SIEMENS)

Supplementary Table S7: Details of the phlebotomy parameters, including biochemistry assays (In -House).

S No	Units	In House Parameter	Sample Type	Platform	Sample Size
		In -House			
1	ng/mL	C-Peptide	Plasma	AA	All
2	mU/mL	Insulin	Plasma	AA	All
3	pg/mL	intact parathyroid hormone (iPTH)	Plasma	AA	All
4	ng/mL	Folate	Plasma	AA	All
5	mmol/dL	Magnesium	Plasma	AA	All
6	ug/dL	Cortisol Morning	Plasma	AA	All
7	umol/L	Homocysteine	Plasma	AA	5000
8	g/L	Haptoglobin	Plasma	AA	3000
9	pg/mL	B-type natriuretic peptide (BNP)	Plasma	AA	3000
10	AU/mL	SARS-CoV2 IgG Semi Quantitative	Plasma	AA	6000
11	U/mL	anti-thyroid peroxidase (Anti-TPO)	Serum	AA	5000
12	ng/mL	prostate-specific antigen (PSA)	Serum	AA	All Males
13	ng/mL	Prolactin	Serum	AA	5000
14	pmol/L	Active B12	Serum	AA	5000

AA: Abbott ARCHITECT

IIIM
Jammu

IHBT
Palampur

CSIO, IMTECH
Chandigarh

IIP
Dehradun

CSIR HQ, IGIB, NPL, CRRI,
NIScPR, TKDL, HRDG

CBRI
Roorkee

HRDC
Ghaziabad

Delhi

NEIST
Jorhat

CEERI
Pilani

CMERI
Durgapur

CDRI, CIMAP,
IITR, NBRI
Lucknow

AMPRI
Bhopal

CGCRI, IICB
Kolkata

NML
Jamshedpur

CIMFR
Dhanbad

CSMCRI
Bhavnagar

NEERI
Nagpur

IMMT
Bhubaneswar

NCL, URDIP
Pune

NGRI, CCMB, IICT
Hyderabad

NAL, 4PI
Bengaluru

NIO
Goa

CMC, CLRI, SERC
Chennai

CFTRI
Mysore

CECRI
Karaikudi

NIIST
Thiruvananthapuram

Zonal Distribution
& Nodal labs:

North Zone - **IGIB**

West Zone - **NCL**

South Zone - **CCMB**

East Zone - **IICB**