Automated Deep Learning-Based Detection of Early Atherosclerotic Plaques in Carotid Ultrasound Imaging

Murad Omarov[1], Lanyue Zhang[1], Saman Doroodgar Jorshery[2,3], Rainer Malik[1], Barnali Das[1], Tiffany R. Bellomo[4], Ulrich Mansmann[5], Martin J. Menten[6,7], Pradeep Natarajan[8-10], Martin Dichgans[1,11-13], Marianne Kalic[14], Vineet K. Raghu[3], Klaus Berger[14], Christopher D. Anderson[2,15,16], Marios K. Georgakis[1,2,11]

1. Institute for Stroke and Dementia Research, LMU University Hospital, LMU Munich, Munich, Germany

2. Program in Medical and Population Genetics and Cardiovascular Disease Initiative, Broad Institute of MIT and Harvard, Cambridge, MA, USA

3. Cardiovascular Imaging Research Center, Massachusetts General Hospital, Harvard Medical School, Boston, MA, USA

4. Division of Vascular and Endovascular Surgery, Massachusetts General Hospital, Boston, MA, USA

5. Institute for Medical Information Processing, Biometry and Epidemiology, Medical Faculty, LMU Munich, Munich, Germany

6. BioMedIA, Department of Computing, Imperial College London, London, United Kingdom

7. Institute for AI in Healthcare and Medicine, School of Computation, Information and Technology, Technical University of Munich, Munich, Germany

8. Cardiovascular Research Center and Center for Genomic Medicine, Massachusetts General Hospital, Boston, MA, USA

9. Cardiovascular Disease Initiative, Broad Institute of Harvard and MIT, Cambridge, Boston, MA, USA

10. Department of Medicine, Harvard Medical School, Boston, MA, USA

11. Munich Cluster for Systems Neurology (SyNergy), Munich, Germany

12. German Center for Neurodegenerative Diseases, (DZNE, Munich), Munich, Germany

13. German Centre for Cardiovascular Research (DZHK, Munich), Munich, Germany

14. Institute of Epidemiology and Social Medicine, University of Münster, Münster, Germany.

15. McCance Center for Brain Health, Massachusetts General Hospital, Boston, MA, USA.

16. Department of Neurology, Brigham and Women's Hospital, Boston, MA, USA.

Corresponding author:

Marios K. Georgakis, MD, PhD

Institute for Stroke and Dementia Research

Ludwig-Maximilians-University Hospital

Feodor-Lynen-Str. 17, 81377 Munich, Germany

Tel: +49-89-4400-46127, Fax: +49-89-4400-46040

E-mail: marios.georgakis@med.uni-muenchen.de; mgeorgak@broadinstitute.org

Abstract

Background

Carotid plaque presence is associated with cardiovascular risk, even among asymptomatic individuals. While deep learning has shown promise for carotid plaque phenotyping in patients with advanced atherosclerosis, its application in population-based settings of asymptomatic individuals remains unexplored.

Methods

We developed a YOLOv8-based model for plaque detection using carotid ultrasound images from 19,499 participants of the population-based UK Biobank (UKB) and fine-tuned it for external validation in the BiDirect study (N = 2,105). Cox regression was used to estimate the impact of plaque presence and count on major cardiovascular events. To explore the genetic architecture of carotid atherosclerosis, we conducted a genome-wide association study (GWAS) meta-analysis of the UKB and CHARGE cohorts. Mendelian randomization (MR) assessed the effect of genetic predisposition to vascular risk factors on carotid atherosclerosis.

Results

Our model demonstrated high performance with accuracy, sensitivity, and specificity exceeding 85%, enabling identification of carotid plaques in 45% of the UKB population (aged 47–83 years). In the external BiDirect cohort, a fine-tuned model achieved 86% accuracy, 78% sensitivity, and 90% specificity. Plaque presence and count were associated with risk of major adverse cardiovascular events (MACE) over a follow-up of up to seven years, improving risk reclassification beyond the Pooled Cohort Equations. A GWAS meta-analysis of carotid plaques uncovered two novel genomic loci, with downstream analyses implicating targets of investigational drugs in advanced clinical development. Observational and MR analyses showed associations between smoking, LDL cholesterol, hypertension, and odds of carotid atherosclerosis.

Conclusions

Our model offers a scalable solution for early carotid plaque detection, potentially enabling automated screening in asymptomatic individuals and improving plaque phenotyping in population-based cohorts. This approach could advance large-scale atherosclerosis research.

Keywords: atherosclerosis, carotid artery, vascular ultrasound, cardiovascular disease, machine learning, genetics

CLINICAL PERSPECTIVE

Carotid ultrasound is a well-established method for assessing subclinical atherosclerosis with potential to improve cardiovascular risk assessment in asymptomatic individuals. Deep learning could automate plaque screening and enable processing of large imaging datasets, reducing the need for manual annotation. Integrating such large-scale carotid ultrasound datasets with clinical, genetic, and other relevant data can advance cardiovascular research. Prior studies applying deep learning to carotid ultrasound have focused on technical tasks–plaque classification, segmentation, and characterization–in small sample sizes of patients with advanced atherosclerosis. However, they did not assess the potential of deep learning in detecting plaques in asymptomatic individuals at the population level.

We developed an efficient deep learning model for the automated detection and quantification of early carotid plaques in ultrasound imaging, primarily in asymptomatic individuals. The model demonstrated high accuracy and external validity across population-based cohort studies. Predicted plaque prevalence aligned with known cardiovascular risk factors. Importantly, predicted plaque presence and count were associated with future cardiovascular events and improved reclassification of asymptomatic individuals into clinically meaningful risk categories. Integrating our model predictions with genetic data identified two novel loci associated with carotid plaque presence—both previously linked to cardiovascular disease—highlighting the model's potential for population-scale atherosclerosis research.

Our model provides a scalable solution for automated carotid plaque phenotyping in ultrasound images at the population level. These findings support its use for automated screening in asymptomatic individuals and for streamlining plaque phenotyping in large cohorts, thereby advancing research on subclinical atherosclerosis in the general population.

1 Introduction

Atherosclerosis, characterized by lipid accumulation and plaque formation within arterial walls[1], is the primary condition underlying cardiovascular disease (CVD), the leading cause of global mortality and morbidity[2,3]. Despite significant advancements in pharmacotherapies for lipid lowering and in the management of other vascular risk factors such as diabetes and hypertension, the alarmingly high and rising prevalence of CVD highlights the need for novel risk assessment and preventive strategies[4,5]. Atherosclerosis is a chronic disease, with subclinical atherosclerotic lesions developing silently over decades. Current CVD risk assessment tools used in clinical practice, including the Pooled Cohort Equations (PCE)[6], Framingham Risk Score[7], and Systematic Coronary Risk Evaluation (SCORE[8] and SCORE2)[9,10], rely on demographic, clinical, and biochemical factors but do not account for the presence of subclinical atherosclerosis[11,12]. Imaging studies that enable screening for subclinical atherosclerotic lesions in asymptomatic individuals suggest that atherosclerotic plaques are highly prevalent, even among individuals traditionally considered at low CVD risk[13–18]. Coronary artery calcium (CAC) scoring on computed tomography (CT) has gained traction for assessing subclinical atherosclerosis. However, it faces limitations as a screening tool due to ionizing radiation exposure and its relatively high cost for widespread application[19,20].

In contrast, carotid ultrasound offers a non-invasive, radiation-free, and widely accessible modality for assessing subclinical atherosclerotic lesions[19,21]. While traditional assessment of carotid intima-media thickness (cIMT) does not reliably predict incident CVD risk[22], the detection of carotid atherosclerotic plaques is associated with an increased risk of future events[21,23,24]. Despite promising results, it remains uncertain whether population screening for carotid plaques could reclassify asymptomatic individuals into higher-risk categories that justify the initiation of preventive pharmacotherapies[25]. Many published studies are constrained by relatively small sample sizes or insufficient follow-up[24,26–30].

Several large-scale population-based cohorts have incorporated carotid ultrasound imaging into their data collection processes,[31–33] but evaluating plaque presence across thousands of images remains a labor-intensive task. Recent advancements in deep learning have enhanced medical

imaging analysis, enabling greater precision and the automation of processing large volumes of imaging data[34,35]. Although previous deep learning models for ultrasound images have shown potential in various tasks, including carotid wall and plaque segmentation[36–38], these studies have focused on specific patient populations with advanced atherosclerotic disease, such as stroke patients or individuals with known carotid artery disease, which limits the generalizability of their findings to the broader population of asymptomatic individuals. Automating the detection of early carotid plaques in ultrasound images at a general population level could accelerate plaque screening in asymptomatic individuals and streamline plaque phenotyping in population-based cohorts. The integration of plaque phenotypes with other imaging modalities, as well as genetic, omics, and clinical data collected in the context of these studies would facilitate in-depth research into the biology of subclinical atherosclerosis, enabling explorations into the natural history of the disease and uncovering novel drug targets.

Here, we introduce a computer vision model designed to detect early atherosclerotic plaques in population-based settings. We utilized ultrasound images from 19,499 deeply phenotyped participants of the population-based UK Biobank (UKB). Our model demonstrated high performance in both detecting plaques and quantifying their counts, and showed external validity in a population-based cohort of 2,105 individuals in Germany (BiDirect). To demonstrate the potential of our model for population plaque screening and research into subclinical atherosclerosis, we leveraged its predictions to: (1) estimate age- and sex-specific prevalence rates of carotid plaques and compare them to those from other population-based cohorts applying manual plaque annotation; (2) confirm relationships with established vascular risk factors; (3) show associations of plaque presence and count with the risk of future CVD events; (4) assess whether plaque presence and count could improve CVD risk prediction; and (5) investigate whether integration with genetic data could uncover biological mechanisms underlying atherosclerotic disease using genome-wide association study (GWAS) and Mendelian Randomization (MR).

Methods

Study population

In this study, we utilized data from the UKB for the primary analysis and the BiDirect Study

(Münster, Germany) for external validation. The UKB is a large-scale prospective cohort study

that recruited 502,422 individuals aged 40 to 69 at baseline from across the United Kingdom

between 2006 and 2010[39]. Participants underwent detailed assessments, which included

comprehensive data collection through questionnaires, physical measurements, and biological

sample collections.

Following a baseline visit between 2006 and 2010, a total of 82,340 individuals returned for a

follow-up imaging visit starting in 2014, which included a carotid ultrasound. A total of 177,757

raw images from 19,768 individuals were released by the UKB and used in this study

(**Supplementary Figure 1**). Four anatomic views of the distal common carotid artery and the

bifurcation were available for each side for nearly every UKB participant who underwent a

carotid ultrasound: images along the main longitudinal axis, images along the short axis, and

images along the main longitudinal axis at two different angles for each artery, which were used

by the analysts for cIMT quantification. Our study focused on images from 19,507 UKB

participants derived along the main longitudinal axis. Eight individuals withdrew from the study

post-recruitment (field 190), resulting in a total sample size of 19,499. For participants with

repeat imaging visits, only the ultrasound data from the first visit were retained for analysis.

For external validation of the detection model, we used carotid ultrasound images from the

BiDirect Study, a twelve-year monocentric prospective cohort study established to explore the

relationship between depression and subclinical atherosclerosis[40]. The study recruited patients

with clinically diagnosed depression or cardiovascular disease, as well as a population control

group. The study participants underwent a standardized carotid ultrasound examination for

assessment of cIMT in the context of their baseline and follow-up visits. For the purposes of our

study, we utilized carotid ultrasound images along the longitudinal axis from 2,105 patients with

available ultrasound data at baseline. The ultrasound examinations in BiDirect were conducted

with a different device (Acuson X300, Siemens) than the one used in the UKB (CardioHealth Station, Panasonic) and followed a different protocol.

Pre-processing

The flowchart for extracting the carotid ultrasound imaging data for analysis from the UKB participants is summarized in **Supplementary Figure 1**. After developing an algorithm (**Supplementary Figure 2**) that automatically detects images along the longitudinal axis, we extracted 45,210 long-axis images without cIMT measurements from the left and right arteries of 19,507 individuals. The obtained images were cropped to a size of 480x448 pixels to retain only the ultrasound image while maintaining the original resolution. After excluding participants who withdrew from the study and retaining only the images from the first ultrasound visit, a total of 19,362 left and 19,370 right carotid images from 19,499 individuals remained for analysis.

In order to enhance contrast and reduce noise in the images, we applied two functions from the *OpenCV* v. 4.7.0 library[41]: median blur filtering (ksize = 5) and Contrast Limited Adaptive Histogram Equalization (clipLimit = 2.0, tileGridSize =(8, 8)), respectively, to facilitate the manual segmentation process. These processing steps were applied to the full sample of images in this study. Plaques were manually annotated by two medical doctors with postgraduate training in vascular imaging and subsequently validated by a doctor certified in carotid ultrasound imaging. *Label Studio* version *1.8.2* (https://github.com/HumanSignal/label-studio) was used to segment plaques on the ultrasound images. The edge coordinates of the segmentation masks were used to obtain the bounding boxes. Plaques were defined, according to standards, as focal protrusions in the arterial lumen with a thickness >50% of the surrounding carotid intima-media thickness[42]. If multiple longitudinal images were available for the same artery, those in which the model detected a plaque were prioritized, or a random image was used if plaques were found in more than one.

For the vast majority of participants in the UKB imaging cohort, only one image along the longitudinal axis of each distal common carotid—including the carotid bulb and bifurcation, but not focused on cIMT measurement (without measurement boxes)—was available for plaque

screening. In contrast, the were multiple images across the entire common carotid artery for

BiDirect participants. To increase comparability between the two cohorts, we selected, for each

BiDirect participant a single image from each artery most likely to depict the region closest to

the common carotid artery bifurcation. This selection was enabled by developing a lumen

segmentation model based on the U-Net architecture with a pretrained ResNet-34 encoder. For

this purpose, we manually segmented lumens in 300 randomly selected images, which were

split into training, validation, and testing sets comprising 204, 51, and 45 images, respectively.

The model was trained using a combined Dice and binary cross-entropy loss function. After

achieving strong segmentation performance in the test set (Dice score: 0.946; IoU: 0.906), we

post-processed the predicted segmentation masks to smooth borders and eliminate small

artifacts (i.e., objects with an area smaller than 10,000 pixels in images sized 480 × 544),

improving the IoU further to 0.914. The process for downstream analysis was implemented with

the *scikit-image* (v0.21.0) Python package, with functions from the *morphology* and *measure*

modules. The lumen segmentation model was then deployed on the full dataset of 17,864

images. After segmenting all images, the binary masks with the predicted lumen were split in

half, and the median height of the left and right halves was measured (representing lumen

thickness). The image with the largest difference between the median heights was selected for

downstream analysis, as this asymmetry was indicative of closeness to the carotid bifurcation,

which consistently appeared on the left side of the image. The lumen segmentation model

predictions and post-processed masks are presented in **Supplementary Figure 3.**

Model Development and Deployment

To train a deep learning model for the detection of atherosclerotic plaques, we manually

annotated plaques in 680 randomly selected carotid ultrasound images. A plaque was present

in 253 of these images. We performed transfer learning with fine-tuning, which involves

selecting a model pre-trained on a large dataset of natural images and then re-training it on a

new dataset. This approach allows for adjusting the model's weights and biases to better suit

the task related to the new dataset. Here, we employed the YOLOv8l[43] model for object

detection, pre-trained on over 330,000 images, and re-trained it on our dataset of 680 carotid

ultrasound images. The YOLOv8 object detection algorithm generates bounding boxes to

indicate the locations of objects of interest. The images with manually annotated plaques were

randomly divided into training, validation, and test sets in a 0.725/0.125/0.15 ratio, resulting in

103 images allocated to the test set. This distribution maximized the training set while ensuring

a sufficient number of images for assessing model performance in the test set. To enhance

model generalizability and predictive power, we randomly selected 50% of the training set (490

images) and applied various augmentation techniques from *Albumentations*[44] Python library:

GridDistortion (p=0.15), RandomBrightnessContrast ((0,0.5),(0,0.5)), HorizontalFlip(p=0.2),

GaussNoise(p=0.15), and RandomSizedCrop (min_max_height=(384, 384), p=0.4). These

augmentations increased the variability of the training set, making the model more invariant to

noise and other distortions. This augmented dataset, along with the rest of the images, was

processed in batches for further augmentation within the YoloV8 framework, as detailed below.

The model was trained with a batch size of 44 images. All tuned and fixed model parameters

are listed in **Supplementary Table 1**. The initial learning rate (*lr0*) was set to 0.001, which is ten

times lower than the default (0.01), as training was initiated from a pretrained model. A cosine

learning rate scheduler was applied to improve convergence. Early stopping with a patience of

five epochs was used to prevent overfitting. This means training would stop after five

consecutive epochs without improvement. The patience value was chosen based on an initial

exploratory run using a preliminary training-validation split, during which we plotted the total

training and validation loss curves. In the final run, training concluded after 14 epochs, with peak

performance observed at the ninth epoch. The following loss function parameters were selected

based on a grid search (**Supplementary Table 1**): distribution focal loss (DFL) = 2.5, box loss =

10 for bounding box regression, and binary cross-entropy loss (CLS) = 1.1. Default

augmentation techniques in the YOLOv8 framework were partially suppressed due to prior

augmentation efforts; specifically, mosaic, copy-paste, shear, close mosaic, flip up-down, and

mix-up were disabled. However, flip left-right (p=0.1), degrees (10), HSV-Saturation (hsv_s:

0.05), HSV-Value (hsv_v: 0.05), translate (0.1), and scale (0.1) were retained. Training was

conducted using an NVIDIA QUADRO RTX 5000 GPU (16 GB). *PyTorch*[45] version *1.12.1* and

the *Ultralytics*[43] framework version *8.1.16* were used for model development.

We evaluated the model's performance by training it on several subsets of our input

development dataset (training + validation) using 5-fold cross-validation. The consistent

performance metrics, with minimal variations in precision and recall, indicated no significant

signs of overfitting (**Supplementary Figure 4**).

Model fine-tuning for the BiDirect dataset

To fine-tune our model for the images of the BiDirect cohort that were derived following a

different scanning protocol, we manually annotated a set of 600 images using the same plaque

definition as for the UKB sample. Of these images, 29.8% (179) contained plaques. The dataset

was split into training (336 images), validation (144 images), and test (120 images), sets.

Fine-tuning was implemented using the previously developed model on the UK Biobank images,

with a reduced learning rate of 0.0001, 'warmup_epochs' set to 0, and the 'pretrained' option

enabled. No additional augmentation was employed beyond the one natively implemented in the

Ultralytics framework, with the same parameters used in training on the UKB images

(**Supplementary Table 1**). Loss coefficients were set to the default values (cls = 0.5; dfl = 1.5;

box = 7.5). Patience was set to 5 epochs to avoid overfitting. Batch size was set to 44 to fit

constraints, and the image size (*imgsz* parameter) was set to 480, as it was for the UKB

images.

Model's classification metrics for detecting plaque-positive images were estimated in a

confusion matrix by comparing ground truth annotations with bounding box predictions. A true

positive was recorded when both the annotation and the prediction contained a bounding box.

Demographic and Clinical Variables

In the UKB, age at carotid ultrasound assessment was derived by subtracting the participant's

date of birth (field 33) from the carotid ultrasound assessment visit date (extracted from the

manifest files linked to the ultrasound images). Ethnicity for the PCE risk score calculation was

determined by field 21000. Observations with missing data or responses of "Do not know" or

"Prefer not to answer" were encoded as "Other". Pre-existing CVD was defined based on self-

reported history (UKB field 20002, with 1075 - heart attack/myocardial infarction, 1081 - stroke,

1583 - ischemic stroke, 1491 - brain haemorrhage); general practice records (131298, 131300,

131302, 131368, 131366); hospital data records (defined using ICD-10 codes I20-I25, I60-I61,

and ICD-9 codes 410-412, 429-431, 434, 436, as well as operation codes K40-46, K49, K471,

K49, K50, K75, L294, and L295) observed before the initial ultrasound visit. SBP was quantified

using fields 4080 and 93 by averaging the observations from each field, followed by taking the

mean of the resulting values. Smoking status was categorized as "current" or "other", with

missing data (<0.5%) treated as "other" (field 20116). Total cholesterol, LDL and HDL

cholesterol levels were obtained from fields 30690, 30780 and 30760, respectively. When

available, values from the assessment closest to the ultrasound visit were extracted; otherwise,

baseline measurements were used. Missing values were imputed using the multivariate

imputation by chained equations method[46], affecting 12% of HDL-cholesterol values and 6% of

total and LDL-cholesterol values, based on other CVD risk factors. Five imputations were

conducted, and prediction results were pooled using Rubin's rules. Diabetes was defined by

self-reported cases (UKB field 20002 – codes: 1220, 1222, 1223), use of glucose-lowering

medications (field 20003), and hospital records prior to the first carotid ultrasound exam (ICD-9

codes: 250* and ICD-10 codes E10, E11). Information on the use of antihypertensive drugs,

diabetes medications, and statins was obtained from field 20003 (**Supplementary Table 2).**

Assessment of major adverse cardiovascular events

MACE endpoints were defined as follows: myocardial infarction (ICD-10 codes I21, I22 from

hospital inpatient data, UKB data-fields 131298 and 131300), stroke (ICD-10 codes I60–64 from

hospital inpatient data; UKB data-fields 131368 and 131366), or death due to any

cardiovascular cause (defined as the cause of death with an ICD-10 code starting with 'I',

extracted from the death registry). The date of the first episode observed after the carotid

ultrasound assessment was considered as the date of the event of interest.

Carotid intima-media thickness

We compared three cIMT measurements between UKB participants with and without carotid

plaque, as predicted by the model. Measurements of mean and maximum cIMT were obtained

using UKB data fields 22670–22681, as described by Strawbridge et al[47].

• cIMT mean: average of the mean values from four mean cIMT measurements (two angles

for each carotid artery: left and right).

• cIMT max**:** The maximum value of the cIMT measurements across both arteries.

• cIMT mean-max**:** The mean of the maximum cIMT values per artery.

All the obtained values were log-transformed for the analysis. For the analysis comparing the

two arteries, individuals with missing cIMT measurements for either the left or right artery were

excluded, resulting in a total of 18,497 individuals included in this analysis. The analysis was

also repeated separately for the right and left arteries. For the right artery, data were available

for 18,803 individuals with no missing measurements; for the left artery, data were available for

18,703 individuals with no missing measurements. Statistical tests were performed on the

specified number of individuals in each subset.

For the BiDirect participants, mean cIMT measurements were available for 1,974 and 1,956

individuals for the left and right arteries, respectively, out of 2,105 participants. Missing values

were excluded from the comparative analysis.

Logistic regression for plaque presence and Cox regression for future events

To estimate the associations between model-derived plaque presence and CVD risk factors, we

applied logistic regression, using plaque presence as the outcome variable. A history of

hypertension was defined by the use of antihypertensive treatment at the time of the carotid

ultrasound assessment. Definitions of other clinical and demographic variables are described

above.

For each subset of the cohort (full sample, primary events with and without individuals on statin

therapy), two separate Cox regression models were constructed: one with plaque presence as a

binary variable and another with the count of plaques as a categorical variable with three levels

(no plaques as the reference, one plaque, and two or more plaques). The time variable in the

survival analysis was calculated as the duration from the first ultrasound visit to the event of

interest or to the censoring date, which included date of death from causes other than CVD or

the date at the last observed event (2022-10-30) available at the time of the analysis. Controls

were censored at the time point of the latest observed event. The fitted Cox regression models

included all vascular risk factors included in the PCE (age, sex, SBP, smoking status, history of

diabetes, antihypertensive therapy, total cholesterol, and HDL cholesterol), as well as statin

usage. Interaction with sex was evaluated by including an interaction term in the Cox models.

Ancestry was not included due to its very low variance in our sample (**Table 1**). The proportional

hazards assumptions were tested with the scaled Schoenfeld residuals, and no violation of the

assumptions was detected.

All analyses were conducted using R software, version 4.4.0. We considered two-sided p-

values (unless otherwise specified) less than 0.05 to be statistically significant. The category-

free NRI was calculated using the *nricens* v.1.6[48] package, with confidence intervals and p-

values based on 1,000-fold bootstrap standard errors. The IDI metric was calculated using the

survIDINRI library v.1.1-2[49]. Harrell's C-statistic, along with its 95% confidence interval, was

used to evaluate the discriminative ability of the time-to-event models[50]. Comparisons between

models based on the C-statistic were conducted using the *CsChange* package[51].

PCE risk estimation

The calculation of the PCE risk score was performed using published equations[6]. PCE eligibility

included the following criteria:

1) $40 \leq Age \leq 79$,

2) $130 \leq$ Total cholesterol ≤ 320 mg/dL;

3) $20 \leq$ HDL-cholesterol ≤ 100 mg/dL;

4) $90 \leq$ Systolic blood pressure ≤ 200 mmHg

To enable fair model comparison and justify the recommended threshold usage, the original

PCE was calibrated to the UKB population. This recalibration was achieved by fitting calculated

log-hazards from the published PCE coefficients in a Cox regression stratified by sex to obtain

recalibrated probabilities. The calibration of the predicted risk values from the original and

recalibrated models was assessed using Greenwood–Nam–D'Agostino statistics and the

integrated calibration index[52,53]. Calibration plots for the PCE are presented in **Supplementary**

Fig. 5.

The plaque variables were incorporated into the PCE, with plaque presence as a binary variable

and plaque count as an ordinal variable, as previously described[54]. Briefly, the recalculated risk

is derived from the relative risk estimate for the novel risk factor, the baseline risk, and the

prevalence of the novel risk factor. As an example, the presence of plaque was incorporated

using Equation (1), adapted from Kooter et al[54]:

(1) $$[r] = p \times HR + (1 - p)$$

18 $$MF_{(+)} = HR/[r]$$

19 $$R_{(+)} = R_{bl}/[r]$$

20 $$R_{(-)} = R_{+}/RR$$

where [r] represents the weighted mean risk; p is the plaque prevalence; HR is the hazard ratio

for plaque presence; $MF_{(+)}$ is the multiplication factor; $R_{(+)}$ is recalculated risk in the presence of

plaque; R_{bl} is the baseline risk estimated from the baseline hazard and log-HRs estimated from

the fitted Cox model ; and $R_{(-)}$ is the recalculated risk in the absence of plaque.

HRs for both plaque presence and plaque count, used to enhance the recalibrated PCE risk

model with plaque information, were estimated from a Cox regression model fitted on the sub-

cohort eligible for PCE risk assessment. The model was adjusted for sex, age, HDL and total

cholesterol levels, antihypertensive drug use, smoking status, SBP, and statin use at the time of

the initial ultrasound assessment. Plaque information was incorporated separately for males and

females, with prevalence estimates calculated for each group.

Model calibration was assessed using the *survival.calib*[55] package in R. Reclassification tables

and NRI statistics were calculated using the *PredictABEL* v.1.2-4[56] library in R.

Genome-wide association study and meta-analysis

Genomic quality control in the UKB was performed as previously described[57]. For the GWAS

analysis, *Regenie*[58] v3.3 was employed. In step 1, we used directly genotyped variants with

MAF > 1%, <10% missingness, Hardy–Weinberg equilibrium test $P > 1 \times 10^{-15}$, and a minor

allele count > 100. Age at the time of the ultrasound exam, along with sex, the first 10 genetic

principal components, and the genotyping chip were used as covariates. Fixed-effect meta-

analysis was conducted with *METAL*[59], using effect size estimates and standard errors (option

SCHEME STDERR). Between-study heterogeneity was assessed using Cochran's Q statistic as

implemented in *METAL*. Significant heterogeneity was defined as a Cochran's Q-derived p-

value of less than 0.05. Results were clumped using the *clump_data* function of

TwoSampleMR[60,61] R package version 0.5.6) at an $r^2 < 0.001$ based on the European 1000

Genomes Project reference panel with 10,000 kb window[62]. Results were then visualized in a

Manhattan plot, which was constructed using the *gwaslab* Python package[63].

Mendelian randomization

23 MR aims to estimate the causal effect of an exposure on an outcome using genetic variants as

24 instrumental variables, under three assumptions: (1) the instrument is robustly associated with

25 the exposure; (2) the instrument is independent of confounders of the exposure–outcome

26 relationship; and (3) the instrument affects the outcome only through the exposure (no

27 horizontal pleiotropy). Violations of these assumptions introduce bias. Summary-level data

sources for exposures, along with their descriptions, are provided in **Supplementary Table 3**.

Two-sample MR was conducted using the *TwoSampleMR v0.6.7* package in R. The instrumental variable for each exposure was constructed by selecting SNVs from summary statistics files with a significant association ($p < 5×10^8$), followed by clumping for linkage disequilibrium at an $r^2 < 0.001$ threshold within a 10,000 kb window. R^2 and F-statistics were calculated for each SNP to assess instrument strength and evaluate potential weak instrument bias (**Supplementary Table 4**). Only SNPs with F-statistics > 10 were retained for subsequent analytical procedures to ensure adequate instrument strength. For IL-6 receptor-mediated signaling activity, the genetic instrument was constructed as previously described[64], using beta estimates derived from a cohort that excluded the UKB to minimize bias. Main association estimates were derived using random-effects IVW analysis. To account for potential bias due to horizontal pleiotropy, sensitivity analyses were performed using both MR-Egger and the weighted median estimator, as these methods are known to be more robust against pleiotropic effects[65,66]. Heterogeneity and horizontal pleiotropy of the genetic instruments were estimated using the *mr_heterogeneity* and *mr_pleiotropy_test* functions from the *TwoSampleMR* package. When MR-Egger indicated significant pleiotropy, we utilized the Mendelian Randomization Pleiotropy RESidual Sum and Outlier (MR-PRESSO) method from the *MRPRESSO*[67] *v1.0* package in R to identify and exclude significant pleiotropic instrumental variables ($P < 0.05$), which were considered as "outliers". Subsequently, IVW, weighted median, and MR-Egger analyses were carried out on the outlier-corrected models. Continuous traits (HbA1c, BMI, waist-to-hip ratio, LDL-cholesterol, HDL-cholesterol) were rank-inverse-normalized in source GWASs; inverse-variance-weighted (IVW) estimates represent effects per SD increment. Blood pressure coefficients were scaled by a factor of 10, expressing effects per 10 mmHg. Binary exposures (smoking initiation, type 2 diabetes) modeled on the log-odds scale were multiplied by 0.693, representing two-fold odds differences. IL-6 signaling was assessed using a 26-variant IL6R instrument per unit increase in natural logarithm-transformed C-reactive protein. False discovery rate (FDR) correction was applied to the IVW estimates to control for multiple testing.

We conducted the MR study in accordance with the guidelines of Strengthening the Reporting

of Observational Studies in Epidemiology using Mendelian Randomization (STROBE-MR).[68,69]

The STROBE-MR checklist is provided in **Supplementary Table 5.**

Data Availability Statement

The UKB provides an accessible research resource, available to researchers upon submitting a

research proposal at https://www.ukbiobank.ac.uk. The GWAS data on UKB and the GWAS

meta-analysis results obtained in this study are available in the GWAS Catalog

(https://ftp.ebi.ac.uk/pub/databases/gwas/summary_statistics/GCST90454001-

GCST90455000/) under accession numbers GCST90454354 and GCST90454353, respectively.

Additionally, the carotid plaque phenotypes derived from the developed model will be returned to

UKB for use in future studies. The list of GWAS used for MR analyses are presented in

Supplementary Table 3. Individual-level data of the BiDirect cohort can be obtained by

researchers from KB (https://medizin.uni-muenster.de/en/epi) following a written transfer

agreement.

The code used for this study will be available on GitHub upon publication of this manuscript.

Ethical Considerations

The UK Biobank has obtained approval from the Northwest Multi-Center Research Ethics

Committee. All participants have provided written informed consent. The Bidirect study received

ethical approval from the Ethics Committee of the University of Münster and the Westphalian

Chamber of Physicians in Münster, Germany (# 2009-391-f-S). Written informed consent was

obtained from all participants.

Results

Study population

A total of 177,757 images from 19,499 participants who underwent carotid ultrasound during the first imaging visit of the UKB were available for analysis (**Supplementary Fig. 1**). The protocol for carotid artery examination has been described previously[70]. For the current study, we used 38,732 images obtained in the longitudinal axis of the left and right distal common carotid artery and the bifurcation, allowing for the assessment of plaque presence along the vessel wall. The demographic and medical characteristics of the study participants are presented in **Table 1**. The mean age at the time of the carotid ultrasound examination was 64.6 years (SD = 7.59), and 50.8% of the participants were female. A total of 1,381 (7.1%) study participants had a baseline diagnosis of atherosclerotic CVD. A comparison between UKB participants with carotid ultrasound and the rest of the UKB cohort revealed a lower prevalence of CVD risk factors (**Supplementary Table 6**).

Plaque detection model

To train a deep learning model for the detection of atherosclerotic plaques, we manually annotated plaques in 680 randomly selected carotid ultrasound images. Plaques were defined as focal protrusions in the arterial lumen with a thickness greater than 50% of the surrounding carotid intima-media thickness[42]. A plaque was present in 253 of these images. We performed transfer learning with fine-tuning by employing a pre-trained YOLOv8[43] object detector as the foundation for developing the plaque detection model. The YOLOv8 object detection algorithm generates bounding boxes to indicate the locations of objects of interest. The images with manually annotated plaques were randomly divided into training, validation, and test sets in a 0.725/0.125/0.15 ratio. This distribution maximized the training set while ensuring a sufficient number of images for assessing model performance in the test set. We evaluated the model's performance by training it on several subsets of our input development dataset (training + validation) in 5-fold cross-validation. The consistent performance metrics, with minimal

variations in precision and recall, indicated no significant signs of overfitting (**Supplementary Fig. 4**).

After training, the performance of the model was evaluated in a blind test set of 103 images, of which 38 contained at least one plaque (53 plaques in total). The model achieved high classification metrics for plaque presence at the image level, with an accuracy, sensitivity, specificity, and Positive Predictive Value (PPV) of 89.3%, 89.5%, 89.2%, and 82.9% (**Figure 1A**), respectively, at an iteratively tuned confidence score threshold of 13%. The confidence score measures the model's certainty that a box contains an object of interest and was tuned to optimize and balance accuracy, sensitivity, and specificity. The Mean Average Precision at an Intersection over Union (IoU) threshold of 50% (mAP@50) was 68.4%, indicating the precision with which the model can localize objects with at least 50% overlap with the ground truth. The model's detection precision and recall were 70.4% and 71.7%, respectively. Prediction examples are illustrated in **Figure 1B** and **Supplementary Fig. 6**. Overall, the model correctly identified 42 of 53 plaques (79.2%) in 38 plaque-positive images in the test set (**Supplementary Fig. 7**).

To assess extent to which the augmentation parameters affect the results, we conducted a post hoc sensitivity analysis using varying augmentation probabilities for *GridDistortion* (0.05, 0.15, 0.25), *HorizontalFlip* (0.1, 0.2, 0.3), and *RandomSizedCrop* (0.3, 0.4, 0.5). The detection metrics ranged from 0.52 to 0.85 for precision, 0.45 to 0.66 for recall, and 0.50 to 0.74 for mAP@50 on the test set with a default confidence threshold of 50%. The results indicate that image augmentation may affect the outcomes, but the developed model achieved optimal performance under the selected augmentation scheme.

Prevalence and risk factors of carotid plaques in the UK Biobank

Next, we deployed the model on all available long-axis carotid ultrasound images from the UKB cohort (38,732 images, 19,499 individuals, **Supplementary Fig. 1**). This deployment allowed us to extract data on plaque phenotypes, including plaque presence and the count of plaques in

either artery for each individual. The count of plaques was determined by the number of

predicted bounding boxes in each image.

Overall, the model detected at least one plaque in 45% of UKB participants who underwent a

carotid ultrasound examination. In 14% of participants, the model detected at least two plaques,

and in 3.1%, three or more plaques across both arteries. The prevalence of plaques in the left

and right carotid arteries is presented in **Figure 2A**. As a quality control step for the model

predictions, we explored whether plaque presence was associated with cIMT, which was

quantified and documented for each individual at the time of the imaging assessment. Indeed,

cIMT was consistently higher for individuals for whom our model predicted a plaque (Wilcoxon

test $p < 10^{-60}$ for maximum, mean, and mean of maximum cIMT measurements). Similar results

were obtained for the left and right arteries separately (**Supplementary Fig. 8**).

Plaques were more common in male participants (47.5% vs. 42.6%; two-proportions z-test, p =

5.7×10^{-12}) and plaque prevalence was significantly associated with older age, increasing from

31.1% in participants aged 45–54 years to 62% in participants aged 75 years or older (Cochran-

Armitage trend test, $p = 1.12 \times 10^{-88}$, **Figure 2B**). In a multivariable logistic regression model,

male sex, older age, current smoking, higher systolic blood pressure (SBP), history of

hypertension, pre-existing CVD, use of statins, and higher Low-Density Lipoprotein (LDL)

cholesterol levels were all significantly associated with plaque presence (**Figure 2C**).

External validation of plaque detection model in the BiDirect study

To ensure external validity of our plaque detection model, we conducted validation in the

BiDirect cohort. Baseline characteristics of the BiDirect participants included in this study are

presented in **Supplementary Table 7.** After fine-tuning the developed model on the BiDirect

images, we evaluated it on a test set of 120 images, which included 40 plaque-positive images.

At a confidence threshold of 0.13, the fine-tuned model reached high image-level classification

metrics for plaque presence (accuracy 85.8%, sensitivity 77.5%, specificity 90%, and PPV

79.5%, **Figure 3A**). The mAP@50 of the fine-tuned model was 65.1%. The model detected 29

of 45 plaques in 40 plaque-positive images (**Supplementary Figure 9**). Examples of the

model's predictions are shown in **Figure 3B**. In a small subset of examinations (n = 64)

annotated by a sonographer assessing full image loops, our model, based on a single image,

showed high agreement in correctly detecting 90% of right and 86% of left carotid plaques.

In the full dataset of 2,105 patients, the model detected at least one plaque in 1,049 patients

(49.8%). In 12.9% of study participants, plaques were detected bilaterally. The prevalence of

plaques was 34.6% in the left artery and 28.3% in the right artery (**Figure 3C**). Plaque presence

was significantly associated with older age, increasing from 39% in participants younger than 45

years to 63.7% in those aged 69 years or older (Cochran–Armitage trend test, $p = 5.31 \times 10^{-16}$,

Figure 3D). Plaque prevalence was higher in the CVD group (59.0%) than in the depression

(45.7%) and population control groups (51.0%; one-sided two-proportion z-tests with Holm

adjustment: $p = 6.3 \times 10^{-5}$ and $p = 0.015$, respectively). BiDirect participants with predicted

plaques had higher mean cIMT values in both the left and right arteries (Wilcoxon p-values: 9.2

$\times 10^{-15}$ for the left and 4.6×10^{-8} for the right carotid artery; **Supplementary Figure 10**).

Associations of plaque phenotypes with the risk of future cardiovascular events

To estimate the effects of the presence and count of plaques in carotid ultrasound on the risk of

future major adverse cardiovascular events (MACE), we conducted a survival analysis.

Following the first imaging visit, UKB participants with available ultrasound images were

followed up for a median of 55 months (range 1-80 months). During this time interval, a total of

430 individuals experienced a MACE, defined as myocardial infarction, stroke or death due to

any cardiovascular cause. Of these, 335 were first-ever events among 18,110 participants

(1.8%) without a history of CVD at the time of the ultrasound examination, while 95 were

secondary events among 1,389 participants (6.8%) with an existing history of CVD.

Kaplan-Meier estimates indicated a higher incidence rate of MACE among individuals with

carotid plaques compared to those without plaques, demonstrating a dose-response pattern of

higher incidence with an increasing plaque count (log-rank test p-value for all pairwise

comparisons < 0.05; **Figure 4**). After adjusting for conventional cardiovascular risk factors in

Cox regression models, plaque presence was significantly associated with future risk of MACE

(Hazard Ratio (HR) 1.42, 95% CI: 1.16–1.73). The plaque count per individual showed a dose-

dependent association with future CVD risk (HR for 1 plaque vs. no plaque = 1.30, 95% CI:

1.04–1.63; HR for 2 or more plaques vs. no plaques = 1.62, 95% CI: 1.27–2.07). These

associations were consistent in the subgroup of individuals without an existing history of CVD,

as well as in those who had neither a history of CVD nor statin use (**Supplementary Figure**

11). There was no evidence of an interaction with sex (p = 0.139). The HRs were comparable

when analyzing individual MACE components, albeit with wider 95% CIs–likely due to lower

statistical power (195 myocardial infarction and 172 stroke cases; **Supplementary Figure 11**).

Sensitivity analysis with adjustment for competing risks was conducted using Fine-Gray

regression, where deaths from causes other than CVD (203 individuals) were treated as

competing risks. Subdistribution hazard ratios (HRs) from the Fine-Gray models were similar to

those derived from Cox regression models, showing HRs for both plaque presence

(subdistribution HR = 1.42, 95% CI: 1.16–1.73) and plaque count (subdistribution HR for 1

plaque vs. no plaque = 1.31, 95% CI: 1.04–1.64; HR for 2 or more plaques vs. no plaques =

1.61, 95% CI: 1.26–2.05).

Predictive power of plaque phenotypes

To examine whether assessing carotid plaque phenotypes could improve cardiovascular risk

prediction, we compared the fitn, reclassification, and discrimination of prediction models that

included conventional vascular risk factors with those that also considered plaque presence and

count. Both plaque presence and count significantly improved the overall goodness of fit of a

Cox regression model (p < 0.05), as assessed by the log-likelihood ratio test[71]. We observed a

reclassification improvement for plaque presence (category-free net reclassification

improvement (cfNRI): 0.331, 95% CI, 0.217–0.445) and plaque count (cfNRI: 0.369, 95% CI,

0.260–0.478, **Table 2**). Briefly, NRI quantifies model improvement by the difference between the

net proportion of cases for which the new model correctly increases predicted risks and net

proportion of controls for which the new model correctly decreases predicted risks[72,73].

Sensitivity analyses confirmed these improvements in individuals without a history of CVD and

statin use (**Table 2**). Despite the strong associations with the future risk of MACE, none of the

plaque phenotypes added to conventional risk factors significantly improved model

discrimination, as measured by the C-index. Specifically, adding plaque presence to

conventional risk factors only slightly changed C-index from 0.745 (95% CI, 0.723–0.767) to

0.747 (95% CI, 0.725–0.769), while adding plaque count changed it to 0.748 (95% CI, 0.726–

0.770). However, minor yet significant improvements were observed in the integrated

discrimination improvement (IDI): 0.002 (95%CI, $2x10^{-4}$–0.005) for plaque presence and 0.002

(95%CI, $2x10^{-5}$–0.006) for plaque count. All the models demonstrated good calibration,

indicating a strong alignment between observed outcomes and predicted risk estimates

(**Supplementary Figure 12**).

To assess whether plaque phenotyping would improve the reclassification of individuals when

complementing established clinical risk prediction models, we calculated the PCE risk scores for

UKB participants eligible for assessment according to ACC/AHA guidelines[6]. Due to the PCE's

tendency to overestimate risk in the UKB population, the model was recalibrated

(**Supplementary Figure 5**). Incorporating plaque presence and count into the PCE

demonstrated significant reclassification improvement, with a categorical NRI of 0.034 (95% CI,

0.006–0.062) and 0.04 (95% CI, 0.010–0.070), respectively, at the threshold of 7.5% 10-year

cardiovascular risk, which defines intermediate risk and justifies preventive initiation of statin

therapy according to current guidelines. Overall, adding plaque presence correctly reclassified

17 out of 318 patients who developed MACE into a higher risk category. Including plaque count

correctly reclassified 20 patients into a higher risk category (**Table 3**). In both cases–plaque

presence and plaque count–four patients were incorrectly reclassified into a lower risk category.

Genome-wide association study of carotid atherosclerotic plaque

As the next step, we investigated the genetic architecture of carotid atherosclerosis, defined by

plaque presence. To detect single nucleotide variants (SNVs) associated with the presence of

carotid atherosclerotic plaque, we conducted a genome-wide association study (GWAS) and

subsequently meta-analyzed our data with the largest available GWAS for carotid plaque from

the Cohorts for Heart and Aging Research in Genomic Epidemiology (CHARGE) consortium[74].

After quality control and excluding individuals without genetic data, the UKB GWAS included

18,203 White British individuals, comprising 8,250 cases with carotid plaque and 9,953 controls.

The pooled sample from the UKB and CHARGE cohorts included 66,637 individuals (29,790

cases; 36,847 controls). We identified seven independent genomic loci significantly associated

with the presence of carotid plaque, two of which were novel. Five of the loci (mapped to the

genes *EDNRA, LINC02577, CDKN2B-AS1, CFDP1, LDLR*) replicated known associations, as

the lead SNVs were in high linkage disequilibrium ($r^2 > 0.9$) with SNVs previously reported to be

associated with carotid plaque presence in the CHARGE study (**Figure 5**, **Supplementary**

Table 8). The sixth locus included the *LPA* gene, which encodes lipoprotein(a) (Lp(a)) and is a

known locus for atherosclerotic cardiovascular disease[75–79]. The lead variant at this locus

(rs56393506), associated with higher odds for an atherosclerotic plaque (OR for T allele: 1.12,

95%CI: 1.07–1.16), is an intronic variant in the *LPA* gene that has been previously strongly

associated with higher Lp(a) levels[80]. The lead SNV at the seventh locus is located in a non-

coding region (rs1893250, OR for A allele: 0.93, 95%CI: 0.90–0.95) and was previously

associated with angina pectoris[81]. Heterogeneity quantified by Cochran's Q statistic was not

significant for any of the genome-wide significant variants (**Supplementary Table 8**).

Mendelian randomization analyses

Finally, we performed Mendelian Randomization (MR) to explore whether genetically proxied

risk factors and biomarkers of CVD are associated with carotid atherosclerotic plaque. We used

the largest publicly available GWAS summary statistics to date for vascular risk factors to

generate genetic instruments for the risk variables under study (**Supplementary Table 3**). The

inverse-variance weighted (IVW) MR analyses revealed associations between higher genetically

proxied SBP, diastolic blood pressure (DBP), LDL cholesterol, interleukin-6 (IL-6) signaling

activity, and genetic predisposition to smoking initiation and type 2 diabetes (T2D) with the odds

of carotid plaque presence (**Figure 6**). Additionally, higher genetically proxied HDL cholesterol

was associated with lower odds of carotid plaque (**Figure 6**). The results for the significant IVW

associations were generally highly consistent in sensitivity analyses, including MR-Egger

regression and the Weighted Median estimator (**Supplementary Table 9**).

Discussion

In this study, we developed a computer vision model that accurately detects atherosclerotic

plaques in carotid ultrasound images and applied it to a population-based cohort of 19,499

participants from the UKB. The model demonstrated strong performance in detecting carotid

plaques and classifying plaque-positive carotid ultrasound images, achieving approximately

90% in accuracy, sensitivity, and specificity. Following fine-tuning, the model also achieved high

performance in the external BiDirect cohort, which followed a different scanning protocol with

another ultrasound device. Consistent with previous studies in comparable demographics

(mean age 64.6±7.6 years; 53% female)[14,82], our model identified at least one carotid plaque in

45% of the UKB participants. The presence of plaques was associated with conventional

vascular risk factors and was predictive of future adverse cardiovascular events over a follow-up

period of up to 7 years. Importantly, both plaque presence and count led to improved risk

reclassification for future adverse cardiovascular events beyond the established PCE risk

assessment tool. Leveraging the phenotypic depth of the UKB, we conducted the largest

genomic analysis of carotid atherosclerosis to date, identifying two novel loci and risk pathways,

including ones targeted by emerging cardiovascular therapeutics, such as Lp(a) and IL-6

signaling. The developed model enriches the UK Biobank with important information on carotid

plaque presence and count, and offers a promising solution for plaque phenotyping in large-

scale cohorts.

While atherosclerosis can develop long before clinical symptoms appear[83], modern risk

assessment tools do not consider subclinical atherosclerotic pathology. Many studies have

highlighted the potential of integrating imaging biomarkers of subclinical atherosclerosis into

conventional risk assessment tools[28]. CAC assessed through CT is a well-established predictor

of CVD risk[84]. However, CAC has limitations, including insensitivity to early stages of

atherosclerosis and exposure to radiation[19,20,85]. In contrast, carotid ultrasound is an

inexpensive, well-tolerated, radiation-free tool capable of detecting early-stage

atherosclerosis[14,86]. Our results, based on a population of almost 20,000 individuals, suggest

that both the presence and count of plaques are strongly associated with the risk of future acute

CVD events. Incorporating plaque information into Cox regression models led to significant

reclassification improvements, which were robust across the full cohort, individuals without a

history of CVD, and statin-naive participants. The reclassification metrics in our study indicate

that adding plaque information to conventional risk factors and directly incorporating it into the

PCE has the potential to improve patient stratification. Specifically, incorporating plaque count

reassigned 6.3% (20 out of 318) of individuals who went on to develop cardiovascular events

from a low to a higher risk category, making them eligible for preventive statin therapy.

Despite growing evidence that carotid ultrasound-derived plaque phenotypes—such as total

plaque area and vulnerability features—are independently associated with future CVD events

[26,87,88], their use in clinical practice remains underutilized. The primary challenges for wider

adoption include the time required for assessment and reliance on operator skill[23]. However, an

efficient artificial intelligence model could significantly streamline these labor-intensive tasks.

The model developed in this study demonstrates high performance in identifying individuals with

carotid atherosclerosis and localizing plaques from a single image of the carotid bifurcation.

Further model advancements could not only address the issue of labor intensity but also enable

the extraction of more detailed plaque features, thereby improving the predictive performance of

carotid ultrasound.

Importantly, our model enhances the phenotypic depth of the UKB to include subclinical

atherosclerosis, facilitating integration with the unique resources available in this population,

such as multi-omics and other imaging data[89]. Moving in this direction, we leveraged the

available genomic data to perform the largest exploration to date of the genetic architecture of

carotid plaque. We replicated five genomic loci previously associated with atherosclerosis

endophenotypes[74] and also found two new loci related to clinical cardiovascular outcomes.

Furthermore, downstream MR confirmed the effect of genetic predisposition to known vascular

risk factors, such as smoking, high blood pressure, and LDL cholesterol, on carotid plaque

presence. Importantly, the GWAS and MR results showed that genetic variation leading to

elevated Lp(a) levels and higher IL-6 signaling activity is associated with higher odds of carotid

plaque. Both pathways are believed to play key roles in atheroprogression and are the targets of

investigational drugs in advanced clinical development[75,76,90–92]. These results suggest that

drugs targeting these pathways could be promising, particularly in the preclinical stages of

atherosclerosis. Integrating carotid plaque phenotypes with additional omics layers may provide

further insights into novel drug targets for atherosclerotic CVD.

Limitations

Our study has several limitations. First, during model development, the low quality of some

images necessitated contrast enhancement and noise reduction techniques. These adjustments

may have introduced bias, especially in cases where high noise levels complicated plaque

detection. However, our model achieved high classification and detection metrics, which could

potentially be improved further by annotating more images. Moreover, the consistency of

predicted carotid plaque prevalence with previously reported plaque prevalence in similar

demographics, along with the associations of predicted plaque presence with known risk factors

and future CVD risk, supports the model's reliability[82]. Second, the UKB is a cohort of healthy

volunteers with a lower incidence rate of CVD than the general population, particularly within its

imaging subsample (**Supplementary Table 6**)[93]. This discrepancy contributes to an

overestimation of CVD risk calculated by clinical risk models, such as the PCE tool. To improve

the reliability of our reclassification evaluation, we recalibrated the model using data from the

study population to better align predicted risk with observed outcomes. Due to the low

prevalence of risk factors and events, PCE-derived absolute risk estimates remain notably low.

It is worth noting that deriving categorical NRI metrics comparable to those from other studies

with different incidence rates may be problematic[94]. We addressed this issue by calculating

continuous NRI estimates using bootstrap estimates, which are threshold-independent and less

sensitive to event rates[95]. Third, the carotid ultrasound examination took place 2 to 15 years

after the baseline visit and the assessment of cardiovascular risk factors. This gap introduces

bias into the effect estimates. To account for changes in baseline risk factors over this period,

we used, wherever available, data from the primary care records of participants collected at the

closest date to the ultrasound exam. Fourth, our study has a shorter follow-up period of 7 years

compared to most clinical risk assessment tools, including the PCE, which typically calculate 10-

year risk estimates for CVD. Fifth, we observed significant heterogeneity in the results of IVW

MR analyses for several risk factors (**Supplementary Table 9**), which could indicate the

presence of pleiotropy. To explore whether the derived estimates could be biased by directional

pleiotropy, we conducted several sensitivity analyses to test the robustness of the estimates

against different MR assumptions. Sixth, there was some population overlap between the

exposure and outcome GWAS datasets used in our MR analyses, which could introduce weak

instrument bias into the derived effect estimates. We addressed this concern by using the

largest available summary statistics for the exposure data. Given that the effective population

overlap was less than 5% for all exposure-outcome pairs, we estimated any bias in the effect

estimates to be under 5%[96]. Seventh, all GWAS used for the MR analyses were conducted in

participants of European ancestry; therefore, the estimates primarily generalize to European

populations and may not be applicable to other ancestries. Eighth, among some of the

CHARGE cohorts included in the GWAS meta-analysis, carotid plaque was defined using

criteria different from the Mannheim consensus used in this study. This variation may introduce

phenotypic heterogeneity that could affect the meta-analysis results. However, all definitions

employed are based on standard clinical criteria and reflect the presence of atherosclerotic

arterial pathology. Another limitation lies in differences between the UKB and external validation

cohort. UKB is population-based, whereas BiDirect includes a relatively high proportion of

participants with cardiovascular disease (15%; see **Supplementary Table 7**) and a high

prevalence of clinical depression (45%). The cohorts also used different ultrasound protocols,

devices, and image resolutions, necessitating model fine-tuning. Additionally, the BiDirect

sample was relatively small (N=2,105) compared with the UKB (N=19,499), which limits the

precision of the prevalence estimates. Lastly, this study analyzed samples from the UK and

Germany, which predominantly consisted of White ancestry volunteers, who are healthier than

the general population. Therefore, replicating these findings in more diverse populations and

real-world settings is crucial for improved generalizability and informed decision-making.

Conclusions

We have successfully developed, implemented, and externally validated a deep learning model

for early atherosclerotic plaque detection in carotid ultrasound imaging. Our model sets the

stage for automating subclinical carotid plaque assessment in the context of either population

screening for asymptomatic individuals or large-scale plaque phenotyping in population-based

cohorts. Our results highlight the potential of large-scale carotid plaque assessment for refining

cardiovascular risk prediction and revealing relevant biological mechanisms for atherosclerosis.

10
11
12

13 **Acknowledgements**

14 UKB data were accessed through applications 36993, 7089, and 151281. The graphical abstract

15 was created using BioRender.

16
Sources of Funding

This work was supported by the Fritz-Thyssen Foundation (grant ref. 10.22.2.024MN to MKG),

the German Research Foundation (DFG; Emmy Noether grant GZ: GE 3461/2-1, ID 512461526

to MKG; Munich Cluster for Systems Neurology EXC 2145 SyNergy, ID 390857198 to MKG), and

the Hertie Foundation (Hertie Network of Excellence in Clinical Neuroscience, ID P1230035 to

MKG). V.K.R. is supported by Norn Longevity Impetus grant. C.D.A. is supported by NIH

R01NS103924, U01NS069673, RF1NS139183, AHA 18SFRN34250007, AHA-Bugher

21SFRN812095, and the MGH McCance Center for Brain Health. The BiDirect Study was

supported by grants of the German Ministry of Research and Education (BMBF) to the University

of Muenster (KB, 01ER0816 and 01ER1506).

Disclosures

M.K.G reports consulting fees from Tourmaline bio, Inc., Pheiron GmbH, and GLG, Inc., all unrelated to this work. V.K.R has common stock in NVIDIA, Alphabet, Apple and Amazon. P.N. reports research grants from Allelica, Amgen, Apple, Boston Scientific, Genentech / Roche, and Novartis, personal fees from Allelica, Apple, AstraZeneca, Blackstone Life Sciences, Creative Education Concepts, CRISPR Therapeutics, Eli Lilly & Co, Esperion Therapeutics, Foresite Capital, Foresite Labs, Genentech / Roche, GV, HeartFlow, Magnet Biomedicine, Merck, Novartis, TenSixteen Bio, and Tourmaline Bio, equity in Bolt, Candela, Mercury, MyOme, Parameter Health, Preciseli, and TenSixteen Bio, and spousal employment at Vertex Pharmaceuticals, all unrelated to the present work. C.D.A. has received sponsored research support from Bayer AG and has consulted for ApoPharma. The other authors declare no competing interests.

References

1. Libby P, Buring JE, Badimon L, Hansson GK, Deanfield J, Bittencourt MS, Tokgözoğlu L, Lewis EF. Atherosclerosis. *Nat. Rev. Dis. Primer*. 2019;5:56.

2. Naghavi M, Ong KL, Aali A, Ababneh HS, Abate YH, Abbafati C, Abbasgholizadeh R, Abbasian M, Abbasi-Kangevari M, Abbastabar H, et al. Global burden of 288 causes of death and life expectancy decomposition in 204 countries and territories and 811 subnational locations, 1990–2021: a systematic analysis for the Global Burden of Disease Study 2021. *The Lancet*. 2024;403:2100–2132.

3. Ferrari AJ, Santomauro DF, Aali A, Abate YH, Abbafati C, Abbastabar H, Abd ElHafeez S, Abdelmasseh M, Abd-Elsalam S, Abdollahi A, et al. Global incidence, prevalence, years lived with disability (YLDs), disability-adjusted life-years (DALYs), and healthy life expectancy (HALE) for 371 diseases and injuries in 204 countries and territories and 811 subnational locations, 1990–2021: a systematic analysis for the Global Burden of Disease Study 2021. *The Lancet*. 2024;403:2133–2161.

4. Joynt Maddox KE, Elkind MSV, Aparicio HJ, Commodore-Mensah Y, De Ferranti SD, Dowd WN, Hernandez AF, Khavjou O, Michos ED, Palaniappan L, et al. Forecasting the Burden of Cardiovascular Disease and Stroke in the United States Through 2050— Prevalence of Risk Factors and Disease: A Presidential Advisory From the American Heart Association. *Circulation* [Internet]. 2024 [cited 2024 Aug 19];150. Available from: https://www.ahajournals.org/doi/10.1161/CIR.0000000000001256

5. Goh RSJ, Chong B, Jayabaskaran J, Jauhari SM, Chan SP, Kueh MTW, Shankar K, Li H, Chin YH, Kong G, et al. The burden of cardiovascular disease in Asia from 2025 to 2050: a forecast analysis for East Asia, South Asia, South-East Asia, Central Asia, and high-income Asia Pacific regions. *Lancet Reg. Health - West. Pac.* 2024;49:101138.

6. Goff DC, Lloyd-Jones DM, Bennett G, Coady S, D'Agostino RB, Gibbons R, Greenland P, Lackland DT, Levy D, O'Donnell CJ, et al. 2013 ACC/AHA Guideline on the Assessment of Cardiovascular Risk: A Report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines. *Circulation* [Internet]. 2014 [cited 2021 Oct 9];129. Available from: https://www.ahajournals.org/doi/10.1161/01.cir.0000437741.48606.98

7. D'Agostino RB, Vasan RS, Pencina MJ, Wolf PA, Cobain M, Massaro JM, Kannel WB. General Cardiovascular Risk Profile for Use in Primary Care: The Framingham Heart Study. *Circulation*. 2008;117:743–753.

8. Conroy R. Estimation of ten-year risk of fatal cardiovascular disease in Europe: the SCORE project. *Eur. Heart J.* 2003;24:987–1003.

9. SCORE2 working group and ESC Cardiovascular risk collaboration. SCORE2 risk prediction algorithms: new models to estimate 10-year risk of cardiovascular disease in Europe. *Eur. Heart J.* 2021;42:2439–2454.

10. Van Daalen KR, Zhang D, Kaptoge S, Paige E, Di Angelantonio E, Pennells L. Risk estimation for the primary prevention of cardiovascular disease: considerations for appropriate risk prediction model selection. *Lancet Glob. Health*. 2024;12:e1343–e1358.

11. Lloyd-Jones DM, Braun LT, Ndumele CE, Smith SC, Sperling LS, Virani SS, Blumenthal RS. Use of Risk Assessment Tools to Guide Decision-Making in the Primary Prevention of Atherosclerotic Cardiovascular Disease. *J. Am. Coll. Cardiol.* 2019;73:3153–3167.

12. Boi A, Jamthikar AD, Saba L, Gupta D, Sharma A, Loi B, Laird JR, Khanna NN, Suri JS. A Survey on Coronary Atherosclerotic Plaque Tissue Characterization in Intravascular Optical Coherence Tomography. *Curr. Atheroscler. Rep.* 2018;20:33.

13. Erbel R, Delaney JAC, Lehmann N, McClelland RL, Mohlenkamp S, Kronmal RA, Schmermund A, Moebus S, Dragano N, Stang A, et al. Signs of subclinical coronary atherosclerosis in relation to risk factor distribution in the Multi-Ethnic Study of Atherosclerosis (MESA) and the Heinz Nixdorf Recall Study (HNR). *Eur. Heart J.* 2008;29:2782–2791.

14. Fernández-Friera L, Peñalvo JL, Fernández-Ortiz A, Ibañez B, López-Melgar B, Laclaustra M, Oliva B, Mocoroa A, Mendiguren J, Martínez De Vega V, et al. Prevalence, Vascular Distribution, and Multiterritorial Extent of Subclinical Atherosclerosis in a Middle-Aged Cohort: The PESA (Progression of Early Subclinical Atherosclerosis) Study. *Circulation.* 2015;131:2104–2113.

15. Berry JD, Liu K, Folsom AR, Lewis CE, Carr JJ, Polak JF, Shea S, Sidney S, O'Leary DH, Chan C, et al. Prevalence and Progression of Subclinical Atherosclerosis in Younger Adults With Low Short-Term but High Lifetime Estimated Risk For Cardiovascular Disease: The Coronary Artery Risk Development in Young Adults Study and Multi-Ethnic Study of Atherosclerosis. *Circulation.* 2009;119:382–389.

16. Ibanez B, Fernández-Ortiz A, Fernández-Friera L, García-Lunar I, Andrés V, Fuster V. Progression of Early Subclinical Atherosclerosis (PESA) Study. *J. Am. Coll. Cardiol.* 2021;78:156–179.

17. Laclaustra M, Casasnovas JA, Fernández-Ortiz A, Fuster V, León-Latre M, Jiménez-Borreguero LJ, Pocovi M, Hurtado-Roca Y, Ordovas JM, Jarauta E, et al. Femoral and Carotid Subclinical Atherosclerosis Association With Risk Factors and Coronary Calcium. *J. Am. Coll. Cardiol.* 2016;67:1263–1274.

18. López-Melgar B, Fernández-Friera L, Oliva B, García-Ruiz JM, Peñalvo JL, Gómez-Talavera S, Sánchez-González J, Mendiguren JM, Ibáñez B, Fernández-Ortiz A, et al. Subclinical Atherosclerosis Burden by 3D Ultrasound in Mid-Life. *J. Am. Coll. Cardiol.* 2017;70:301–313.

19. Spence JD. Coronary calcium is not all we need: Carotid plaque burden measured by ultrasound is better. *Atherosclerosis.* 2019;287:179–180.

20. Mendieta G, Pocock S, Mass V, Moreno A, Owen R, García-Lunar I, López-Melgar B, Fuster JJ, Andres V, Pérez-Herreras C, et al. Determinants of Progression and Regression of Subclinical Atherosclerosis Over 6 Years. *J. Am. Coll. Cardiol.* 2023;82:2069–2083.

21. Stein JH, Korcarz CE, Hurst RT, Lonn E, Kendall CB, Mohler ER, Najjar SS, Rembold CM, Post WS. Use of Carotid Ultrasound to Identify Subclinical Vascular Disease and Evaluate Cardiovascular Disease Risk: A Consensus Statement from the American Society of Echocardiography Carotid Intima-Media Thickness Task Force Endorsed by the Society for Vascular Medicine. *J. Am. Soc. Echocardiogr.* 2008;21:93–111.

22. Den Ruijter HM, Peters SAE, Anderson TJ, Britton AR, Dekker JM, Eijkemans MJ, Engström G, Evans GW, de Graaf J, Grobbee DE, et al. Common carotid intima-media thickness measurements in cardiovascular risk prediction: a meta-analysis. *JAMA*. 2012;308:796–803.

23. Johri AM, Nambi V, Naqvi TZ, Feinstein SB, Kim ESH, Park MM, Becher H, Sillesen H. Recommendations for the Assessment of Carotid Arterial Plaque by Ultrasound for the Characterization of Atherosclerosis and Evaluation of Cardiovascular Risk: From the American Society of Echocardiography. *J. Am. Soc. Echocardiogr.* 2020;33:917–933.

24. Bao X, Xu B, Lind L, Engström G. Carotid ultrasound and systematic coronary risk assessment 2 in the prediction of cardiovascular events. *Eur. J. Prev. Cardiol.* 2023;30:1007–1014.

25. Inaba Y, Chen JA, Bergmann SR. Carotid plaque, compared with carotid intima-media thickness, more accurately predicts coronary artery disease events: a meta-analysis. *Atherosclerosis*. 2012;220:128–133.

26. Sillesen H, Sartori S, Sandholt B, Baber U, Mehran R, Fuster V. Carotid plaque thickness and carotid plaque burden predict future cardiovascular events in asymptomatic adult Americans. *Eur. Heart J. - Cardiovasc. Imaging*. 2018;19:1042–1050.

27. Polak JF, Szklo M, Kronmal RA, Burke GL, Shea S, Zavodni AEH, O'Leary DH. The Value of Carotid Artery Plaque and Intima-Media Thickness for Incident Cardiovascular Disease: The Multi-Ethnic Study of Atherosclerosis. *J. Am. Heart Assoc.* 2013;2:e000087.

28. Raitakari OT, Magnussen CG, Juonala M, Kartiosuo N, Pahkala K, Rovio S, Koskinen JS, Mykkänen J, Laitinen TP, Kähönen M, et al. Subclinical atherosclerosis in young adults predicting cardiovascular disease: The Cardiovascular Risk in Young Finns Study. *Atherosclerosis*. 2024;117515.

29. Matangi MF, Hétu M-F, Armstrong DWJ, Shellenberger J, Brouillard D, Baker J, Johnson A, Grubic N, Willms H, Johri AM. Carotid plaque score is associated with 10-year major adverse cardiovascular events in low–intermediate risk patients referred to a general cardiology community clinic. *Eur. Heart J. - Cardiovasc. Imaging*. 2024;jeae153.

30. Baber U, Mehran R, Sartori S, Schoos MM, Sillesen H, Muntendam P, Garcia MJ, Gregson J, Pocock S, Falk E, et al. Prevalence, Impact, and Predictive Value of Detecting Subclinical Coronary and Carotid Atherosclerosis in Asymptomatic Adults. *J. Am. Coll. Cardiol.* 2015;65:1065–1074.

31. Bild DE. Multi-Ethnic Study of Atherosclerosis: Objectives and Design. *Am. J. Epidemiol.* 2002;156:871–881.

32. Fernández-Ortiz A, Jiménez-Borreguero LJ, Peñalvo JL, Ordovás JM, Mocoroa A, Fernández-Friera L, Laclaustra M, García L, Molina J, Mendiguren JM, et al. The Progression and Early detection of Subclinical Atherosclerosis (PESA) study: Rationale and design. *Am. Heart J.* 2013;166:990–998.

33. Nasir K, Cainzos-Achirica M, Valero-Elizondo J, Ali SS, Havistin R, Lakshman S, Blaha MJ, Blankstein R, Shapiro MD, Arias L, et al. Coronary Atherosclerosis in an Asymptomatic U.S. Population. *JACC Cardiovasc. Imaging*. 2022;15:1604–1618.

34. Khera R, Oikonomou EK, Nadkarni GN, Morley JR, Wiens J, Butte AJ, Topol EJ. Transforming Cardiovascular Care With Artificial Intelligence: From Discovery to Practice. *J. Am. Coll. Cardiol.* 2024;84:97–114.

35. Lin A, Kolossváry M, Motwani M, Išgum I, Maurovich-Horvat P, Slomka PJ, Dey D. Artificial Intelligence in Cardiovascular Imaging for Risk Stratification in Coronary Artery Disease. *Radiol. Cardiothorac. Imaging.* 2021;3:e200512.

36. Jain PK, Sharma N, Giannopoulos AA, Saba L, Nicolaides A, Suri JS. Hybrid deep learning segmentation models for atherosclerotic plaque in internal carotid artery B-mode ultrasound. *Comput. Biol. Med.* 2021;136:104721.

37. Huang Q, Tian H, Jia L, Li Z, Zhou Z. A review of deep learning segmentation methods for carotid artery ultrasound images. *Neurocomputing.* 2023;545:126298.

38. Singh S, Jain PK, Sharma N, Pohit M, Roy S. Atherosclerotic plaque classification in carotid ultrasound images using machine learning and explainable deep learning. *Intell. Med.* 2024;4:83–95.

39. Bycroft C, Freeman C, Petkova D, Band G, Elliott LT, Sharp K, Motyer A, Vukcevic D, Delaneau O, O'Connell J, et al. The UK Biobank resource with deep phenotyping and genomic data. *Nature.* 2018;562:203–209.

40. Teismann H, Wersching H, Nagel M, Arolt V, Heindel W, Baune BT, Wellmann J, Hense H-W, Berger K. Establishing the bidirectional relationship between depression and subclinical arteriosclerosis--rationale, design, and characteristics of the BiDirect Study. *BMC Psychiatry.* 2014;14:174.

41. Bradski G. The opencv library. *Dr Dobbs J. Softw. Tools Prof. Program.* 2000;25:120–123.

42. Touboul P-J, Hennerici MG, Meairs S, Adams H, Amarenco P, Bornstein N, Csiba L, Desvarieux M, Ebrahim S, Hernandez Hernandez R, et al. Mannheim Carotid Intima-Media Thickness and Plaque Consensus (2004–2006–2011). *Cerebrovasc. Dis.* 2012;34:290–296.

43. Jocher, G., Chaurasia, A., Qiu, J. *YOLO* by Ultralytics [Internet]. Available from: https://github.com/ultralytics/ultralytics; 2023; Accessed: August 14, 2024.

44. Buslaev A, Iglovikov VI, Khvedchenya E, Parinov A, Druzhinin M, Kalinin AA. Albumentations: Fast and Flexible Image Augmentations. *Information.* 2020;11:125.

45. Paszke A, Gross S, Massa F, Lerer A, Bradbury J, Chanan G, Killeen T, Lin Z, Gimelshein N, Antiga L, et al. PyTorch: An Imperative Style, High-Performance Deep Learning Library [Internet]. 2019 [cited 2024 Aug 14];Available from: https://arxiv.org/abs/1912.01703

46. Buuren SV, Groothuis-Oudshoorn K. **mice** : Multivariate Imputation by Chained Equations in *R. J. Stat. Softw.* [Internet]. 2011 [cited 2024 Aug 14];45. Available from: http://www.jstatsoft.org/v45/i03/

47. Strawbridge RJ, Ward J, Bailey MES, Cullen B, Ferguson A, Graham N, Johnston KJA, Lyall LM, Pearsall R, Pell J, et al. Carotid Intima-Media Thickness: Novel Loci, Sex-Specific Effects, and Genetic Correlations With Obesity and Glucometabolic Traits in UK Biobank. *Arterioscler. Thromb. Vasc. Biol.* 2020;40:446–461.

48. Inoue E. NRI for Risk Prediction Models with Time to Event and Binary Response Data.

49. Uno H, Tian L, Cai T, Kohane IS, Wei LJ. A unified inference procedure for a class of measures to assess improvement in risk prediction systems with survival data. *Stat. Med.* 2013;32:2430–2442.

50. Harrell FE. Evaluating the Yield of Medical Tests. *JAMA J. Am. Med. Assoc.* 1982;247:2543.

51. Ganz P, Heidecker B, Hveem K, Jonasson C, Kato S, Segal MR, Sterling DG, Williams SA. Development and Validation of a Protein-Based Risk Score for Cardiovascular Outcomes Among Patients With Stable Coronary Heart Disease. *JAMA*. 2016;315:2532.

52. Austin PC, Steyerberg EW. The Integrated Calibration Index (ICI) and related metrics for quantifying the calibration of logistic regression models. *Stat. Med.* 2019;38:4051–4065.

53. Demler OV, Paynter NP, Cook NR. Tests of calibration and goodness-of-fit in the survival setting. *Stat. Med.* 2015;34:1659–1680.

54. Kooter AJ, Kostense PJ, Groenewold J, Thijs A, Sattar N, Smulders YM. Integrating Information From Novel Risk Factors With Calculated Risks: The Critical Impact of Risk Factor Prevalence. *Circulation*. 2011;124:741–745.

55. Byron Jaeger, Olga Demler. survival.calib: Tests of Calibration in the Survival Setting.

56. Kundu S, Aulchenko YS, Van Duijn CM, Janssens ACJW. PredictABEL: an R package for the assessment of risk prediction models. *Eur. J. Epidemiol.* 2011;26:261–264.

57. Turner S, Armstrong LL, Bradford Y, Carlson CS, Crawford DC, Crenshaw AT, de Andrade M, Doheny KF, Haines JL, Hayes G, et al. Quality control procedures for genome-wide association studies. *Curr. Protoc. Hum. Genet.* 2011;Chapter 1:Unit1.19.

58. Mbatchou J, Barnard L, Backman J, Marcketta A, Kosmicki JA, Ziyatdinov A, Benner C, O'Dushlaine C, Barber M, Boutkov B, et al. Computationally efficient whole-genome regression for quantitative and binary traits. *Nat. Genet.* 2021;53:1097–1103.

59. Willer CJ, Li Y, Abecasis GR. METAL: fast and efficient meta-analysis of genomewide association scans. *Bioinformatics*. 2010;26:2190–2191.

60. Hemani G, Zheng J, Elsworth B, Wade KH, Haberland V, Baird D, Laurin C, Burgess S, Bowden J, Langdon R, et al. The MR-Base platform supports systematic causal inference across the human phenome. *eLife*. 2018;7:e34408.

61. Hemani G, Tilling K, Davey Smith G. Orienting the causal relationship between imprecisely measured traits using GWAS summary data. *PLOS Genet.* 2017;13:e1007081.

62. The 1000 Genomes Project Consortium. A map of human genome variation from population-scale sequencing. *Nature*. 2010;467:1061–1073.

63. He Y, Koido M, Shimmori Y, Kamatani Y. GWASLab: a Python package for processing and visualizing GWAS summary statistics [Internet]. 2023 [cited 2024 Aug 14];Available from: https://doi.org/10.51094/jxiv.370

64. Georgakis MK, Malik R, Richardson TG, Howson JMM, Anderson CD, Burgess S, Hovingh GK, Dichgans M, Gill D. Associations of genetically predicted IL-6 signaling with cardiovascular disease risk across population subgroups. *BMC Med.* 2022;20:245.

65. Bowden J, Davey Smith G, Burgess S. Mendelian randomization with invalid instruments: effect estimation and bias detection through Egger regression. *Int. J. Epidemiol.* 2015;44:512–525.

66. Hartwig FP, Davey Smith G, Bowden J. Robust inference in summary data Mendelian randomization via the zero modal pleiotropy assumption. *Int. J. Epidemiol.* 2017;46:1985–1998.

67. Verbanck M, Chen C-Y, Neale B, Do R. Detection of widespread horizontal pleiotropy in causal relationships inferred from Mendelian randomization between complex traits and diseases. *Nat. Genet.* 2018;50:693–698.

68. Skrivankova VW, Richmond RC, Woolf BAR, Yarmolinsky J, Davies NM, Swanson SA, VanderWeele TJ, Higgins JPT, Timpson NJ, Dimou N. Strengthening the Reporting of Observational Studies in Epidemiology Using Mendelian Randomization: The STROBE-MR Statement. *Jama.* 2021;326:1614–1621.

69. Skrivankova VW, Richmond RC, Woolf BAR, Davies NM, Swanson SA, VanderWeele TJ, Timpson NJ, Higgins JPT, Dimou N, Langenberg C, et al. Strengthening the reporting of observational studies in epidemiology using mendelian randomisation (STROBE-MR): explanation and elaboration. *BMJ.* 2021;n2233.

70. Coffey S, Lewandowski AJ, Garratt S, Meijer R, Lynum S, Bedi R, Paterson J, Yaqub M, Noble JA, Neubauer S, et al. Protocol and quality assurance for carotid imaging in 100,000 participants of UK Biobank: development and assessment. *Eur. J. Prev. Cardiol.* 2017;24:1799–1806.

71. Cook NR. Quantifying the added value of new biomarkers: how and how not. *Diagn. Progn. Res.* 2018;2:14, s41512-018-0037–2.

72. Pencina MJ, D' Agostino RB, D' Agostino RB, Vasan RS. Evaluating the added predictive ability of a new marker: From area under the ROC curve to reclassification and beyond. *Stat. Med.* 2008;27:157–172.

73. Pencina MJ, D'Agostino RB, Pencina KM, Janssens ACJW, Greenland P. Interpreting incremental value of markers added to risk prediction models. *Am. J. Epidemiol.* 2012;176:473–481.

74. Franceschini N, Giambartolomei C, De Vries PS, Finan C, Bis JC, Huntley RP, Lovering RC, Tajuddin SM, Winkler TW, Graff M, et al. GWAS and colocalization analyses implicate carotid intima-media thickness and carotid plaque loci in cardiovascular outcomes. *Nat. Commun.* 2018;9:5141.

75. Malick WA, Goonewardena SN, Koenig W, Rosenson RS. Clinical Trial Design for Lipoprotein(a)-Lowering Therapies. *J. Am. Coll. Cardiol.* 2023;81:1633–1645.

76. Swerdlow DI, Rider DA, Yavari A, Wikström Lindholm M, Campion GV, Nissen SE. Treatment and prevention of lipoprotein(a)-mediated cardiovascular disease: the emerging potential of RNA interference therapeutics. *Cardiovasc. Res.* 2022;118:1218–1231.

77. Reyes-Soffer G, Ginsberg HN, Berglund L, Duell PB, Heffron SP, Kamstrup PR, Lloyd-
Jones DM, Marcovina SM, Yeang C, Koschinsky ML, et al. Lipoprotein(a): A Genetically
Determined, Causal, and Prevalent Risk Factor for Atherosclerotic Cardiovascular Disease:
A Scientific Statement From the American Heart Association. *Arterioscler. Thromb. Vasc.*
Biol. [Internet]. 2022 [cited 2024 Aug 14];42. Available from:
https://www.ahajournals.org/doi/10.1161/ATV.0000000000000147

78. Trinder M, Uddin MM, Finneran P, Aragam KG, Natarajan P. Clinical Utility of
Lipoprotein(a) and *LPA* Genetic Risk Score in Risk Prediction of Incident Atherosclerotic
Cardiovascular Disease. *JAMA Cardiol.* 2021;6:287.

79. Van Der Harst P, Verweij N. Identification of 64 Novel Genetic Loci Provides an Expanded
View on the Genetic Architecture of Coronary Artery Disease. *Circ. Res.* 2018;122:433–
443.

80. Sinnott-Armstrong N, Tanigawa Y, Amar D, Mars N, Benner C, Aguirre M, Venkataraman
GR, Wainberg M, Ollila HM, Kiiskinen T, et al. Genetics of 35 blood and urine biomarkers
in the UK Biobank. *Nat. Genet.* 2021;53:185–194.

81. Sakaue S, Kanai M, Tanigawa Y, Karjalainen J, Kurki M, Koshiba S, Narita A, Konuma T,
Yamamoto K, Akiyama M, et al. A cross-population atlas of genetic associations for 220
human phenotypes. *Nat. Genet.* 2021;53:1415–1424.

82. Song P, Fang Z, Wang H, Cai Y, Rahimi K, Zhu Y, Fowkes FGR, Fowkes FJI, Rudan I.
Global and regional prevalence, burden, and risk factors for carotid atherosclerosis: a
systematic review, meta-analysis, and modelling study. *Lancet Glob. Health.* 2020;8:e721–
e729.

83. McGill HC, McMahan CA, Herderick EE, Malcom GT, Tracy RE, Strong JP. Origin of
atherosclerosis in childhood and adolescence. *Am. J. Clin. Nutr.* 2000;72:1307s–1315s.

84. Peters SAE, Bakker M, Den Ruijter HM, Bots ML. Added value of CAC in risk
stratification for cardiovascular events: a systematic review. *Eur. J. Clin. Invest.*
2012;42:110–116.

85. Visseren FLJ, Mach F, Smulders YM, Carballo D, Koskinas KC, Bäck M, Benetos A, Biffi
A, Boavida J-M, Capodanno D, et al. 2021 ESC Guidelines on cardiovascular disease
prevention in clinical practice. *Eur. Heart J.* 2021;42:3227–3337.

86. Nielsen RV, Fuster V, Bundgaard H, Fuster JJ, Johri AM, Kofoed KF, Douglas PS,
Diederichsen A, Shapiro MD, Nicholls SJ, et al. Personalized Intervention Based on Early
Detection of Atherosclerosis. *J. Am. Coll. Cardiol.* 2024;83:2112–2127.

87. Ihle-Hansen H, Vigen T, Berge T, Walle-Hansen MM, Hagberg G, Ihle-Hansen H,
Thommessen B, Ariansen I, Røsjø H, Rønning OM, et al. Carotid Plaque Score for Stroke
and Cardiovascular Risk Prediction in a Middle-Aged Cohort From the General Population.
J. Am. Heart Assoc. 2023;12:e030739.

88. Saba L, Saam T, Jäger HR, Yuan C, Hatsukami TS, Saloner D, Wasserman BA, Bonati LH,
Wintermark M. Imaging biomarkers of vulnerable carotid plaques for stroke risk prediction
and their potential clinical implications. *Lancet Neurol.* 2019;18:559–572.

89. Sudlow C, Gallacher J, Allen N, Beral V, Burton P, Danesh J, Downey P, Elliott P, Green J, Landray M, et al. UK Biobank: An Open Access Resource for Identifying the Causes of a Wide Range of Complex Diseases of Middle and Old Age. *PLOS Med.* 2015;12:e1001779.

90. Ridker PM. From RESCUE to ZEUS: will interleukin-6 inhibition with ziltivekimab prove effective for cardiovascular event reduction? *Cardiovasc. Res.* 2021;117:e138–e140.

91. Ridker PM, Devalaraja M, Baeres FMM, Engelmann MDM, Hovingh GK, Ivkovic M, Lo L, Kling D, Pergola P, Raj D, et al. IL-6 inhibition with ziltivekimab in patients at high atherosclerotic risk (RESCUE): a double-blind, randomised, placebo-controlled, phase 2 trial. *The Lancet.* 2021;397:2060–2069.

92. Nordestgaard BG, Langsted A. Lipoprotein(a) and cardiovascular disease. *The Lancet.* 2024;404:1255–1264.

93. Lyall DM, Quinn T, Lyall LM, Ward J, Anderson JJ, Smith DJ, Stewart W, Strawbridge RJ, Bailey MES, Cullen B. Quantifying bias in psychological and physical health in the UK Biobank imaging sub-sample. *Brain Commun.* 2022;4:fcac119.

94. Cook NR, Demler OV, Paynter NP. Clinical risk reclassification at 10 years. *Stat. Med.* 2017;36:4498–4502.

95. Pencina MJ, D'Agostino RB, Steyerberg EW. Extensions of net reclassification improvement calculations to measure usefulness of new biomarkers. *Stat. Med.* 2011;30:11–21.

96. Burgess S, Davies NM, Thompson SG. Bias due to participant overlap in two-sample Mendelian randomization. *Genet. Epidemiol.* 2016;40:597–608.

97. Burgess S, Labrecque JA. Mendelian randomization with a binary exposure variable: interpretation and presentation of causal estimates. *Eur. J. Epidemiol.* 2018;33:947–952.

Table 1. Characteristics of the study cohort at the time of initial carotid ultrasound assessment.

	Total	Plaque absent	Plaque present	P-value for comparisons*
N	**19,499**	**10,718**	**8,781**	
Female, N (%)	9912 (50.8)	5688 (53.1)	4224 (48.1)	<0.001
Age, mean (SD)	64.6 (7.6)	63.4 (7.4)	66.1 (7.5)	<0.001
Smoking status, N (%)				<0.001
Never	12169 (62.4)	6951 (64.9)	5218 (59.4)	
Previous	6564 (33.7)	3384 (31.6)	3180 (36.2)	
Current	700 (3.6)	350 (3.3)	350 (4.0)	
Unknown	66 (0.3)	33 (0.3)	33 (0.4)	
SBP (mmHg), mean (SD)	138.6 (17.94)	137.18 (17.43)	140.42 (18.39)	<0.001
DBP (mmHg), mean (SD)	79.32 (9.78)	79.51 (9.75)	79.09 (9.81)	0.003
Total cholesterol (mmol/L), mean (SD)	5.72 (1.09)	5.70 (1.07)	5.74 (1.10)	0.009
LDL-C (mmol/L), mean (SD)	3.58 (0.83)	3.56 (0.82)	3.59 (0.84)	0.028
HDL-C (mmol/L), mean (SD)	1.48 (0.37)	1.47 (0.37)	1.48 (0.37)	0.708
HbA1c (mmol/mol), mean (SD)	35.13 (4.95)	34.91 (4.85)	35.40 (5.05)	<0.001
eGFR (mL/min/1.73m^2), mean (SD)	93.58 (12.63)	94.52 (12.54)	92.42 (12.65)	<0.001
BMI (kg/m^2), mean (SD)	26.60 (4.41)	26.80 (4.54)	26.37 (4.23)	<0.001
History of diabetes, N (%)	1073 (5.5)	511 (4.8)	562 (6.4)	<0.001
Statin usage, N (%)	4631 (23.7)	2131 (19.9)	2500 (28.5)	<0.001
Antihypertensive therapy, N (%)	4826 (24.7)	2312 (21.6)	2514 (28.6)	<0.001
Previous history of CVD, N (%)	1389 (7.1)	568 (5.3)	821 (9.3)	<0.001
Ethnicity, N (%)				<0.001
White	18932 (97.1)	10371 (96.8)	8561 (97.5)	
Asian	189.00 (1.0)	107 (1.0)	82 (0.9)	
Black	121.00 (0.6)	90 (0.8)	31 (0.4)	

Mixed	77.00 (0.3)	53 (0.5)	24 (0.3)	
Other/unknown	180.00 (0.9)	97 (0.9)	83 (0.9)	

SBP – Systolic Blood Pressure; DBP – Diastolic Blood Pressure; LDL-C – Low-Density Lipoprotein

Cholesterol; HDL-C – High-Density Lipoprotein Cholesterol; HbA1c – Glycated Hemoglobin; eGFR –

Estimated Glomerular Filtration Rate; BMI – Body Mass Index; CVD – Cardiovascular Disease.

* P-values were calculated using a two-sided t-test for continuous and a chi-square test for categorical

variables.

Table 2. Metrics of discrimination, reclassification and overall model fit between Cox regression models with and without plaque information.

Model	Control / Cases	Log-likelihood ratio test p-value	cfNRI [95% CI] (p-value)	IDI [95 % CI] (p-value)	C-index [95% CI]	C-index difference (p-value)
CVD factors		ref	ref	ref	0.745 [0.723; 0.767]	ref
CVD factors + Plaque presence	19069 / 430	4.8×10^{-4}	0.331 [0.217; 0.445] (1.26×10^{-8})	0.0022 [0.0002; 0.0050] (0.019)	0.747 [0.725; 0.769]	0.002 (0.222)
CVD factors + Plaque count		5.6×10^{-4}	0.369 [0.260; 0.478] (2.97×10^{-11})	0.0024 [2×10^{-5}; 0.0063] (0.048)	0.748 [0.726; 0.770]	0.003 (0.173)
CVD factors		ref	ref	ref	0.744 [0.719; 0.770]	ref
CVD factors + Plaque presence	17775 / 335	0.003	0.329 [0.206; 0.452] (1.65×10^{-7})	0.0019 [0.0003; 0.0052] (0.0079)	0.746 [0.721; 0.771]	0.002 (0.357)
CVD factors + Plaque count		0.006	0.330 [0.209; 0.450] (7.73×10^{-8})	0.0022 [0.0004; 0.0065] (0.0139)	0.746 [0.721; 0.772]	0.002 (0.349)
CVD factors		ref	ref	ref	0.761 [0.732; 0.790]	ref
CVD factors + Plaque presence	14283 / 234	0.014	0.308 [0.159; 0.457] (4.88×10^{-5})	0.0018 [0.0002; 0.0059] (0.014)	0.763 [0.734; 0.792]	0.002 (0.453)
CVD factors + Plaque count		0.039	0.308 [0.159; 0.458] (5.31×10^{-5})	0.0021 [0.0003; 0.0075] (0.0119)	0.763 [0.734; 0.792]	0.002 (0.471)

CVD – cardiovascular disease; cfNRI – category-free net reclassification improvement; C-index – concordance index; IDI – integrated discrimination improvement; ref – reference.

Table 3. Reclassification table showing the distribution of individuals who went on to develop or not major adverse cardiovascular events into two risk groups based on the recommended 7.5% threshold from PCE risk assessment, both with and without incorporating plaque information.

PCE	PCE + Plaque presence		% reclassified	PCE + Plaque Count		% reclassified
	<7.5%	≥7.5%		<7.5%	≥7.5%	
Controls						
<7.5%	16341	213	1	16300	254	2
≥7.5%	90	485	16	82	493	14
Cases						
<7.5%	250	17	6	247	20	7
≥7.5%	4	47	8	4	47	8

Upward movement for cases indicates correct reclassification, while downward movement indicates incorrect reclassification. For controls, the opposite applies.

PCE – Pooled Cohort Equations; blue background indicates correctly reclassified individuals; orange background indicates wrongly reclassified individuals.

2 **GRAPHICAL ABSTRACT.** ASCVD – Atherosclerotic Cardiovascular Disease, CVD – Cardiovascular

3 disease, PCE – Pooled Cohort Equations, TP– true positive, FN – False Negative, FP – False Positive, TN

4 – True Negative, GWAS – Genome-Wide Association Study.

Figure 1. Development and performance of the plaque detection model. A. Confusion matrix for model's classification performance for plaque presence at each image. The confusion matrix is based on comparing the presence of plaque in the annotations with the model's predictions. Specifically, if there is a plaque annotation for an image and the prediction contains a bounding box, then the prediction is annotated as true positive. B. Examples of model predictions as compared to manual annotations for three UK Biobank participants in the test set. Each image depicts the longitudinal view of the common carotid artery extending toward the bifurcation area (the left part of the images).

Figure 2. Prevalence and predictors of carotid plaque in the UK Biobank. A. Distribution of plaques in
the left and right carotid arteries. B. Percentage of plaque presence across age and sex groups. Error bars
represent 95% confidence intervals. C. Forest plot of the associations of demographic and vascular risk
factors with the odds of carotid plaque presence, as derived by a multivariable logistic regression model
that includes all variables in the figure. The results are presented as odds ratios (OR) and 95% confidence
intervals (CI). SBP – systolic blood pressure; CVD – cardiovascular disease; LDL – low-density lipoprotein
cholesterol; SD – standard deviation.

* History of hypertension is defined by the use of antihypertensive medication

Figure 3. Performance of the fine-tuned plaque detection model in the BiDirect sample. A. Confusion matrix for model's classification performance for plaque presence at each image. **B.** Examples of model predictions as compared to manual annotations for four BiDirect participants in the test set. **C.** Distribution of plaques in the left and right carotid arteries. **D.** Percentage of carotid plaque across age and sex groups. Error bars represent 95% confidence intervals.

2 **Figure 4. Survival curves of cumulative major adverse cardiovascular event (MACE) rates by total**

3 **count of carotid plaques predicted by the model.** The presented hazard ratios (HRs) were estimated

4 using Cox regression, adjusted for: sex, age, systolic blood pressure, use of statins, history of

5 antihypertensive therapy, current smoking, history of diabetes, HDL cholesterol and total cholesterol levels.

2 **Figure 5. Manhattan plot of the GWAS meta-analysis for carotid plaque presence (29,790 cases;**

3 **36,847 controls).** Loci highlighted in red point to novel significant associations for carotid plaque whereas

4 loci highlighted in green represent validation of previously described associations.

Exposure		# of SNPs	IVW OR [95% CI]	
Hb1Ac (SD-increment)		75	1.263 [0.993;1.607]	
Type 2 diabetes (2-fold increment in prevalence)		276	1.038 [1.010;1.067]	*
Smoking initiation (2-fold increment in prevalence)		248	1.204 [1.069;1.357]	*
BMI (SD-increment)		542	0.999 [0.921;1.085]	
Waist-to-hip ratio (SD-increment)		349	1.108 [0.992;1.238]	
LDL cholesterol (SD-increment)		303	1.401 [1.296;1.515]	*
HDL cholesterol (SD-increment)		411	0.851 [0.792;0.914]	*
SBP (10mmHg-increment)		429	1.327 [1.258;1.401]	*
DBP (10mmHg-increment)		426	1.279 [1.168;1.400]	*
IL-6 signaling (lnCRP-increment)		21	1.362 [1.151;1.611]	*

* denotes FDR-adjusted p-value
for IVW estimate < 0.05

Figure 6. Forest plot for Mendelian randomization results. The results are presented as odds ratios (OR) and 95% confidence intervals (CI) derived from random-effects inverse-variance weighted Mendelian randomization analyses. Two-fold increments in prevalence for binary exposures (type 2 diabetes and smoking initiation) were derived by multiplying the IVW betas and corresponding confidence intervals by 0.693, as described by Burgess and Labrecque[97]. HbA1c – Glycated hemoglobin; BMI – Body Mass Index; LDL – Low-Density Lipoprotein Cholesterol; HDL – High-Density Lipoprotein Cholesterol; SBP – Systolic Blood Pressure; DBP – Diastolic Blood Pressure; IL-6 – Interleukin-6.