Altered amyloid plasma profile in patients with disabling headaches after SARS-CoV-2 infection and vaccination

Anne Hege Aamodt,[1,2] Thor Ueland,[3,4,5] Marion Boldingh,[1] Burcu Ella Bezgal,[1,4] Maria Bengtson Argren,[1,4] Cecilia Adele Dunne,[4] Kari Otterdal,[3] Ida Gregersen,[3] Vigdis Bjerkeli,[3,4] Annika Elisabet Michelsen,[3,4] Andreas Husøy,[2,6] Åse Hagen Morsund,[7] Kristina Devik,[1,8] Anne Christine Poole[9], Kristine Bodding Gjendemsjø[10], Katrin Schlüter[11], Sara Maria Mathisen[11], Mari Aalstad-Johansen[12], Thor Håkon Skattør[1,4,13], Julie Sønnervik[14], Turid Birgitte Boye[15,16], Trine Haug Popperud[1], Einar August Høgestøl[1,4], Hanne Flinstad Harbo[1,4], Fridtjof Lund-Johansen[4,17], Pål Aukrust[3,4,18], Erling Tronvik[2,6], Tuva Børresdatter Dahl[3], Bente Evy Halvorsen[3,4,19]

[1]Department of Neurology, Oslo University Hospital, Oslo, Norway
[2]Department of Neuromedicine and Movement Science, the Norwegian University of Science and Technology, Trondheim, Norway
[3]Research Institute of Internal Medicine, Oslo University Hospital Rikshospitalet, Oslo, Norway
[4]Institute of Clinical Medicine, Faculty of Medicine, University of Oslo, Oslo, Norway
[5]Thrombosis Research Center (TREC), Division of internal medicine, University Hospital of North Norway, Tromsø, Norway
[6]Department of Neurology of Clinical Neurophysiology, St. Olav Hospital, Trondheim, Norway
[7]Department of Neurology, Molde Hospital, Molde, Norway
[8]Department of Neurology, Namsos Hospital, Molde, Norway
[9]Oslo Headache Centre, Oslo, Norway
[10]Sandvika Neuro Centre, Sandvika, Norway
[11]Department of Neurology, Stavanger University Hospital, Stavanger, Norway
[12]Department of Neurology, Innlandet Hospital Trust, Lillehammer, Norway
[13]Department of Radiology, Oslo University Hospital, Oslo, Norway
[14]Hodeverket, Sandnes, Norway
[15]Unit Psychosomatic medicine and CL psychiatry, Oslo University Hospital, Oslo, Norway
[16]Institute of Basic Medical Sciences, Faculty of Medicine, University of Oslo, Oslo, Norway
[17]Department of Immunology, Oslo University Hospital, Oslo, Norway.
[18]Section of Clinical Immunology and Infectious Diseases, Oslo University Hospital, Oslo, Norway
[19]Department of Medicine, Division of Cardiovascular Medicine, Brigham and Women's Hospital, Harvard Medical School, Boston, MA, USA

Abstract

Background and objectives: New onset persistent headache has been reported following

acute COVID-19 and to some degree also after SARS-CoV-2 vaccination. The mechanisms

for these headache types are unclear. The purpose of this study was to assess levels of

amyloid related biomarkers in patients with persistent headache after COVID-19 and SARS-

CoV-2 vaccine.

Methods: In this prospective observational cohort, patients with severe headache as the

dominating symptom after COVID-19 (n=29) and SARS-CoV-2 vaccination (n=31), had

neurological assessments with reassessments after 6 months. Plasma levels of amyloid

precursor protein (APP), pregnancy zone protein (PZP), cathepsin L1 (CTSL) and serum

Amyloid A (SAA1) were measured by ELISA in relation to levels in healthy controls (n=16).

Results: We found a strong and persistent upregulation of APP in patients with headache

after COVID-19 as compared to the two other groups. At both inclusion and after 6 months

APP levels were also increased in those with accompanying cognitive symptoms. In contrast,

plasma levels of PZP were elevated in patients with headache after SARS-CoV-2 vaccination

at both inclusion and after 6 months as compared to healthy controls. CTSL was only elevated

in those with COVID-19 associated as compared with those with vaccine associated headache

at baseline, whereas SAA1 showed levels comparable in all groups.

Conclusion: Altered plasma levels of soluble markers potentially reflecting changes in

amyloid processing was found in patients with persistent headache after SARS-CoV-2

vaccine and particular in those with persistent headache after COVID-19

where we also found some association with cognitive symptoms.

NCT04576351

NCT05235776

What is already known on this topic

New onset persistent headache occurs in a subset of individuals after COVID-19 and to some

extent after SARS-CoV-2 vaccine. However, the pathophysiological mechanisms are

unknown.

What this study adds

Altered plasma levels of soluble markers that potentially could reflect changes in amyloid

processing was found in patients with persistent headache after SARS-CoV-2 vaccine and

particular in those with persistent headache after COVID-19 with association to cognitive

symptoms.

How this study might affect research, practice or policy

Our data underscore the need for more long-time follow-up of patients with new onset

headache following COVID-19 or SARS-CoV-2 vaccination and this follow-up might also

include blood tests for amyloid processing and neuroinflammation.

Introduction

During the coronavirus disease (COVID)-19 pandemic, headache emerged as an important health issue, increasing the burden of headache disorders.[1, 2] Both severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2)) infection and, although to a lesser degree, SARS-CoV-2 vaccination, induced headache as a secondary headache disorder as well as negatively impact already existing primary headache disorder.[1, 3] Those with primary headache disorder experience headache in the acute phase of COVID-19 and after vaccination more often than other SARS-CoV-2-infected or vaccinated individuals.[1] Approximately one in five patients who present headache during the acute phase of COVID-19 get persistent headache, often with accompanying symptoms including cognitive symptoms and fatigue.[1] Also, a subset of individuals with headache attributed to SARS-CoV-2 vaccinations develop long lasting headaches.[1, 3] Furthermore, in patients with primary headache disorders, SARS-CoV-2 vaccination and in particular an acute SARS-CoV-2 infection worsened their chronic headache burden.[3] The mechanisms for persistent headaches after SARS-CoV-2 infection or vaccines are, however, unknown. One proposed mechanism is persistent immune activation and inflammation triggered by SARS-CoV-2 or vaccination[4, 5] with the trigeminovascular system as a potential target.[1] Additionally, the inflammatory responses during acute COVID-19, involving the release of prostaglandins, interleukin (IL)-6 and other inflammatory cytokines could increase blood-brain barrier permeability leading to neuroinflammation with accompanying headache.[6, 7]

Serum Amyloid A (SAA1), an acute phase protein and systemic marker of inflammation, was early reported as a biomarker of poor prognosis in COVID-19.[8] Recently, amyloid precursor protein (APP), a membrane protein considered to play a main role in Alzheimer disease pathology,[9] was described as a new receptor for SARS-CoV-2 infection and revealed a previously unknown mechanistic insight into COVID-19 related neuropathological sequelae.[10] Moreover, Pregnancy Zone Protein (PZP), originally known as a broad-spectrum immune-suppressive protein that suppresses T-cell function during pregnancy to prevent foetal rejection, has also been proposed as a biomarker for various inflammatory disorders,[11, 12] including the prognosis of COVID-19.[13] With relevance to CNS involvement, serum levels of PZP were higher in women with pre-symptomatic Alzheimer's disease.[14] Furthermore, Cathepsin L1 (CTSL), a lysosomal cysteine protease that plays a major role in intracellular protein catabolism, has been demonstrated to enhance SARS-CoV-2 virus entry, at least in pre-clinical models,[15] and interestingly, CTSL has also been suggested to be involved in the degradation of certain amyloid fibrils.[16] To this end, however,

118   circulating levels of these biomarkers (i.e., SAA1, APP, PZP, and CTSL) in patients with

119   persistent headaches after SARS-CoV-2 infection or vaccination, with or without headache

120   during the acute phase of COVID-19 or immediately after vaccination, have not been

121   reported.

122   In the present study, with the aim to give novel insight into the pathological

mechanisms leading to persistent headache following COVID-19 and SARS-CoV-2

vaccination, we hypothesize that plasma levels of soluble markers that potentially could

reflect changes in amyloid processing are related to persistent headache after SARS-CoV-2

vaccine and COVID-19 and particular in those with accompanying cognitive symptoms. This

hypothesis was tested by measurement of relevant markers in plasma in patients with new

onset persistent headache based on clinical assessments and validated questionnaires. The

levels were compared in healthy controls that had undergone COVID-19 or SARS-CoV-2

vaccination but without headaches. The levels of these molecules were also assessed in

relation to accompanying fatigue or cognitive symptoms, and in the COVID-19 group, the

need for hospitalization during the acute infection.

Methods

In this prospective observational cohort study performed at Norwegian neurological

departments and led by Oslo University Hospital, Oslo, Norway, clinical characteristics and

blood biomarker profiles were assessed in patients with severe headaches for a minimum of

six months after COVID-19 or SARS-CoV-2 vaccination and compared to healthy controls.

The inclusion period lasted from September 2020 to June 2023. We hypothesized that plasma

levels of soluble markers that potentially could reflect changes in amyloid processing such as

APP, PZP, CTSL, and SAA are linked to persistent headaches after SARS-CoV-2 vaccination

and COVID-19 in those with accompanying cognitive symptoms. This hypothesis was tested

by relevant plasma markers longitudinally patients with newly onset persistent headaches

using clinical assessments and validated questionnaires. Differences in plasma levels were

compared with healthy controls, and their relationship to cognitive symptoms was a major

outcome.

Study population and clinical assessments of the participants

The subgroup with **persistent headache after COVID-19 (n=29)** was recruited from the

cohort of 149 individuals aged $\geq$ 18 years with persistent neurological symptoms 6 months

after COVID-19 infection in the Norwegian observational multicentre study of nervous

system manifestations and sequelae after COVID infection (NNC), a sub study of the
ENERGY study led by European Academy of Neurology.[17] The subgroup with **persistent**
headache after SARS-CoV-2 vaccine (n=31) were participants in the Norwegian
observational multi-centre study CovaxHEAD: New-onset severe headache after COVID-19
vaccine (NCT05235776) - a parallel study led by the same team as the NNC study collecting
comparable variables.

Inclusion criteria for both subgroups were age $\geq$ 18 years and new-onset severe
headache, or severe worsening of pre-existing headache, within one week after SARS-CoV-2
infection or vaccination. The degree of headache severity had to be so pronounced that the
headaches were daily or almost daily and had resulted in significant disability or sick leave
and still were the dominant symptom at 6-month follow-up. Exclusion criteria were short-
lasting headaches, no follow-up assessments, or where other diagnoses were considered a
better explanation for their symptoms (e.g. infections including COVID-19, head trauma, or
somatoform disorders). Participants with COVID-19 who reported their headaches starting
after SARS-CoV-2 vaccine were not included. None in this COVID-19 subgroup had SARS-
CoV-2 vaccine induced headache although 24 participants (82.8%) had received SARS-CoV-
2 vaccine. Types of vaccines were registered (BNT162b2 from BioNTech/Pfizer, mRNA-
1273 from Moderna and ChAdOx1 nCoV-19 from AstraZeneca).

For both headache groups, an inclusion criterion was available plasma samples that
could be used in biomarker analyses. CovaxHEAD included a total of 62 participants from 7
Norwegian centres. However, as blood samples for amyloid profiling were available only at
Oslo University Hospital, only the participants included from Oslo University Hospital were
included. In the NeuroCOVID, a total of 149 patients were included, with 46 (30%)
categorised as having headache as the dominant symptom. However, in the present study,
only participants with available samples for biomarker analyses were included, i.e., only
patients from Oslo University Hospital were included in this biomarker substudy.

The cognitive symptoms were evaluated using questionnaires from the ENERGY
study, led from the European Academy of Neurology.[17] Additionally, all participants in the
group with persistent headache after COVID-19 (n=29) were assessed with the Montreal
Cognitive Assessment (MoCA), administered by a trained neurologist and defined as normal
when MoCA score was above 26 . The same questions were applied to the patients with
chronic headaches following SARS-CoV- psychologist, neuropsychologist or psychiatrist as
part of the multidisciplinary headache treatment.

185   The clinical data, including the headaches, were reviewed by a senior neurologist with

186 headache expertise (AHA). The headaches were classified according to their phenotype. The

187 clinical visits and first set of blood samples were collected at inclusion, mean 181.0 (SD

188 145.8) days after COVID-19 in the group with headache after COVID-19 and mean 290.3

189 (SD 145.8) days after SARS-CoV-2 vaccine in the group with headache after SARS-CoV-2

190 vaccine. Both groups had 6-month follow-up with clinical visits and blood samples thereafter.

191   Controls: individuals above 18 years of age, without any history migraine or other

192 disabling headache disorders and who had never experienced neurological symptoms after

193 SARS-CoV-2 vaccine or COVID-19. Furthermore, other exclusion criteria were

194 anyimmunological disorders or use of immunosuppressive drugs. A total of 16 plasma

195 samples were collected in the period April to July 2023 from 8 men and 8 women, mean age

196 40.6 years (SD 11.2). The healthy controls did not undergo detailed questionnaire or

197 neurologic, neuropsychiatric or neuropsychological examination. Serum antibody

198 assessments of the control group showed high levels in all participants (median 15831

199 BAU/ml, minimum 4309, maximum 64 676 BAU/ml) indicating strong or very strong

200 immune response indicating that all had undergone COVID-19 or received SARS-CoV-2

201 vaccine.

205 **Blood sampling protocol**

206 Peripheral venous blood was drawn into pyrogen-free blood collection tubes with

207 ethylenediamine tetra acetic acid (EDTA) as an anticoagulant, immediately immersed in

208 melting ice and centrifuged at 4°C 2500 $\times$ g for 20 minutes within 60 minutes to obtain

209 platelet-poor plasma. The plasma samples were stored in multiple aliquots and thawed only

210 once.

212 **Biomarker assessments**

213 Plasma markers were assessed in all three groups – patients with persistent headache after

214 COVID-19 infection (n=29), patients with persistent headache after the COVID-19 vaccine

215 (n=31), and healthy controls (n=16). The participants had blood samples drawn when they

216 were included in the study. Additionally, 6-month follow-up blood sampling was done in the

217 headache groups. Plasma levels of APP (Cat# DY850), PZP (Cat# DY8280-05), CTSL (Cat#

218 DY952), and SAA1 (Cat# DY3019-05) were measured by enzyme-linked immunosorbent

assay (ELISA) using commercially available antibodies (R&D Systems, Minneapolis, MN) in

a 384-format using a combination of a SELMA pipetting robot (Analytik Jena AG, Jena,

Germany) and a BioTek dispenser/washer (BioTek Instruments, Winooski, VT). Absorption

was read at 450 nm by using an EIA plate reader (BioTek Instruments) with wavelength

correction set to 540 nm. Intra- and inter-assay coefficients of variation for these analyses

were all <10%.

Ethics approval

This study was conducted in accordance with the Helsinki Declaration as revised in 2013.

Ethical approval for this study was obtained from The Norwegian South-Eastern Ethical

Committee (152727 NNC, 351097 CovaxHEAD). Written informed consent was obtained

from all subjects before inclusion into the study.

Statistics

Statistical Package for Social Science (IBM SPSS Inc., version 29 for Windows) was used for

statistical analysis. Differences in patient characteristics were tested using the Chi-square test

or Fisher's exact test for categorical variables, and the Mann-Whitney U test for continuous

variables. Prevalence of characteristics (frequency and percentages) were calculated for each

group. Plasma marker levels were compared between groups using a multivariate general

linear model using markers as dependent, group as fixed, and age as covariate. A two-sided p-

value of <0.05 was deemed statistically significant.

Patient and public involvement.

the user panel at the Department of Neurology, Oslo University Hospital and the user

organization "Hodepine Norge" has contributed to the project. Hodepine Norge has been

contributing to public meetings about the project.

Results

Characteristics and headache phenotypes in the study group

Demographics such as age, gender and co-morbidities are shown in Table 1. There was equal

age and sex distribution in both headache groups (i.e., persistent headache following COVID-

19 or SARS-CoV-2 vaccination, respectively) with 3 out of 4 participants being female.

Comorbidity of frequent diseases, such as pre-existing migraine, allergies, anxiety, and

depression were common in both groups. Few had cardiovascular or metabolic disorders. On

third of the patients with persistent headache after COVID-19 infection was hospitalised in

the acute phase of the infection. In the group with a persistent headache after SARS-CoV-2

vaccine 17 (54.8%) reported previous COVID-19 at baseline, but importantly, without any

persistent headache after the infection (Table 1).

Table 1. Demographic and clinical characteristics at inclusion.

Characteristics	Headaches after COVID-19 n = 29	Headaches SARS-CoV-2 vaccine n = 31	Control group n=16	P value*
Mean age at inclusion, (SD)	43.5 (11.2)	43.8 (12.4)	40.6 (11.2)	0.836
Female, n (%)	22 (75.9)	24 (77.4)	8 (50.0)	0.862
Education				
Secondary school n (%)	2 (6.8)	7 (22.6)		0.193
College/university ≤ 4 years, n (%)	14 (48.3)	12 (38.7)		
College/university > 4 years, n (%)	13 (44.8)	12 (38.7)		
Hospitalisation during COVID-19, n (%)	9 (31.0)	3 (9.7)	0	0.357
Smoking, n (%)	1 (4.4)	2 (6.5)		0.610
Pre-existing migraine, n (%)	11 (31.0)	12 (38.7)	0	0.871
Comorbidities				
Allergy, n (%)	16 (55.1)	13 (41.9)		0.240
Asthma, n (%)	2 (6.9)	6 (19.4)		0.150
Depression, n (%)	8 (27.6)	10 (32.3)		0.759
Anxiety, n (%)	7 (24.1)	13 (41.9)		0.170
Diabetes, n (%)	0	1 (3.3)	0	0.330
Cardiovascular disease, n (%)	0	1 (3.3)	0	0.330
Stroke, n (%)	0	0	0	
Associated symptoms				
Fatigue, n (%)	22 (75.9)	20 (64.5)		0.204

	22 (75.9)	17 (54.8)		0.119
Subjective cognitive symptoms, n (%)	22 (75.9)	17 (54.8)		0.119
Sick leave at first visit, n (%)	21 (72.4)	18 (58.1)	.	0.244
Sick leave at 6-month follow-up, n (%)	19 (65.5)	14 (45.2)		0.207

*Comparison of the two headache groups.

Chronic migrainous headache was the dominant headache phenotype in both headache

groups while tension-type headache was the second most common phenotype in both

headache groups followed by episodic migrainous headache (Figure 1). Cranial neuralgia

often accompanied by painful paraesthesia in the extremities was the dominant phenotype in 5

(17%) of the participants in the group with headaches after COVID-19.

More than two thirds of the participants with headache post COVID-19 and more than half the

participants with headache post SARS-CoV-2 vaccination were on sick-leave due to the

current symptoms at inclusion with only a modest reduction in the percentages on sick-leave

at 6-month follow-up (72.4 % to 65.5 % and 58.1 % to 45.2 %, post COVID-19 and SARS-

CoV-2 vaccination, respectively; Table 1). Subjective cognitive symptoms and fatigue were

common, but not dominating (see Methods), in the headache groups (Table 1). All the patients

with COVID-19 had normal MoCA score except for one patient with critical illness myopathy

in the acute phase of COVID-19 with a score of 24 at follow-up. The patients with chronic

headaches after COVID-19 vaccine were assessed by a psychologist, neuropsychologist, or

psychiatrist as part of the multidisciplinary headache treatment whereof three were diagnosed

with depression, one with functional neurological disorder accompanying deterioration of

chronic migraine, and two were not found to have any psychiatric disorders.

In the group with new onset headache after SARS-CoV-2 vaccine 34.5% reported the

headache starting after their first vaccine, 41.4% after the second vaccine and 24.1% reported

the headache to start after their third vaccine. Although the BNT162b2 vaccine was the most

used vaccine, 89.7% reported their headache starting after RNA-1273 vaccine whereas 10.3%

after BNT162b2 and none after ChAdOx1 nCoV-19.

Biomarkers in relation to post-COVID-19 headache and headache following SARS-

CoV-2 vaccination

At both inclusion and after 6 months, we found a strong and persistent upregulation of APP in

patients with headache after COVID-19 as compared to individuals with vaccine induced

headache and healthy controls (Figure 2A). In contrast, there was upregulation of CTSL in

patients with headaches after vaccine compared to patients with headaches after COVID-19,

but these differences were not seen at 6 months follow-up (Figure 2B). Furthermore, plasma

levels of PZP were elevated in patients with headache after SARS-CoV-2 vaccination at both

inclusion and after 6 months compared to controls, whereas a more modest increase was seen

in those with COVID-19 associated headache only at inclusion (Figure 2C). As for SAA1,

there were no differences between the two headache groups and healthy controls either at

baseline or during follow-up (Figure 2D). Finally, whereas post-COVID-19 symptoms have

been related to disease severity during the acute phase,[18] we found no differences in the actual

biomarkers between those who were hospitalized and those who were not hospitalized during

the acute phase of COVID-19 (Supplemental Table 1).

Biomarker levels in relation to associated fatigue and cognitive symptoms

There was a significant upregulation of APP among participants with accompanying cognitive

symptoms in the group with persistent headache after COVID-19, but this pattern was not

seen in those with headache after SARS-CoV-2 vaccination (Figure 3). Although those with

fatigue also had higher levels of APP compared with those without, the difference did not

reach statistical significance (Figure 3). As for PZP and CTSL, there were no differences

between those with or without associated cognitive symptoms or fatigue, neither in the

COVID-19 nor the SARS-CoV-2 group (Table 2).

Table 2. Median biomarker levels (ng/mL) with interquartile range (QR) of Cathepsin L,

Pregnancy Zone Protein (PZP) and Serum Amyloid A (SAA1) in relation to **A.** subjective

cognitive symptoms and **B.** fatigue in participants with persistent headache after SarsCoV-2

vaccine (COvax) (n=31) and participants with persistent headache after COVID-19 (C19)

(n=29).

A.

Biomarker levels	Headache group	No subjective cognitive symptoms (IQR)	Subjective cognitive symptoms (IQR)	P-value
Cathepsin L	COvax	1.6 (1.5, 2.1)	2.0 (1.5, 2.2)	0.451
	C19	2.2 (1.9, 2.7)	2.2 (1.9, 2.5)	0.760
PZP	COvax	36.5 (4.7, 52.9)	24.2 (9.1, 37.9)	0.889
	C19	19.3 (4.3, 79.1)	17.2 (4.2, 75.8)	0.308
SAA1	COvax	17.7 (10.8, 42.0)	17.3 (8.2, 29.1)	0.165
	C19	23.3 (9.2, 72.4)	12.0 (5.7, 19.7)	0.575

318
319 **B**.

Biomarker levels	Headache group	Fatigue (IQR)	No fatigue reported (IQR)	P-value
PZP	COvax	10.7 (9.2, 12.4)	10.8 (9.5, 11.7)	0.397
	C19	15.6 (15, 16.7)	23.5 (15.2, 27.9)	0.683
Cathepsin L	COvax	26.8 (4.7, 48.7)	27.7 (7.8, 49.3)	0.967
	C19	15.4 (4.2, 77.4)	24.3 (5.5, 75.8)	0.646
SAA1	COvax	2.9 (2.49, 3.46)	2.4 (1.97, 2.82)	0.088
	C19	2.4 (1.06, 3.25)	3.3 (1.96, 5.54v	0.248

Discussion

Persistent headache has been reported following acute COVID-19 and to some extent also

after SARS-CoV-2 vaccination.[1, 3] In the present study, we observed higher plasma levels of

APP in patients with persistent headache after COVID-19, but not in patients with persistent

headache following SARS-CoV-2 vaccination, compared to healthy controls. Remarkably,

elevated APP levels persisted at 6-months follow-up and was associated with accompanying

cognitive symptoms. Notably, plasma levels of CTSL were higher in patients with headaches

after SARS-CoV-2 vaccination compared to those with headaches after COVID-19.

Furthermore, PZP levels were persistently elevated in patients with headache after SARS-

CoV-2 vaccination, while a more modest increase was observed in those with COVID-19

associated headache as compared to healthy controls. To the best of our knowledge, this is the

first report linking proteins in amyloid formation to persistent headache following COVID-19,

and CTSL and PZP to persistent headache after SARS-CoV-2 vaccination.

As depicted in Figure 4, our data point to plausible mechanism of serum APP and PZP

proteins in relation to COVID-19 and SARS-CoV-2 vaccine induced headache. In brief, APP

is produced by neurons, and proteolysis by various secretases can generate a soluble (s)

fragment, sAPP, as well as release the neurotoxic Aβ peptide. SARS-CoV-2 infection may

increase APP expression via the ACE2 receptor[19, 20] and APP promotes cellular entry of the

virus enhancing Aβ-associated pathology.[10] The Aβ peptide may accumulate to form amyloid

plaques[21], leading to activation of astrocytes and microglia, and promote neuroinflammation

by different mechanisms[22] with potentially affecting headache symptoms.[23]

Increased levels of APP are reported as marker for preclinical Alzheimer disease;

however, the biological function of this protein extends beyond amyloid formation[24] and

elevated levels have been found in various metabolic and cardiovascular disorders.[25] Like in

some other viruses, amyloids are identified in the SARS-CoV-2 proteins, i.e., the structural

spike, the nucleocapsid, and the accessory ORF6/ORF10 proteins and this may also be of clinical relevance.[26] There are several reports on the worsening of preexisting dementia including Alzheimer`s Disease and the development of cognitive impairment following SARS-CoV-2 infection, but the pathogenesis of these changes is still incompletely understood.[26-28] However, in a recent study on the relationship between COVID-19 and Alzheimer's disease the changes in blood biomarkers were linked to brain structural imaging patterns associated with Alzheimer's disease, lower cognitive test scores, and poorer overall health evaluations.[26-29]

In the present study few or none had such clinically verified co-morbidities. Still, we found significant upregulation of APP among participants with accompanying cognitive symptoms in the group with persistent headache after COVID-19 compared to those without cognitive symptoms. Moreover, Camacho et al. reporting network analyses of SARS-CoV-2 infection in brain microvascular endothelial cells suggested increased expression of APP.[19] In a previous study of patients with COVID-19 associated neurological syndromes, signs of impaired amyloid processing were found with significantly reduced levels of soluble amyloid precursor protein in CSF from patients compared to controls.[30] However, the study primarily included patients with severe neurological diseases such as Guillain Barre syndrome and encephalopathy and only one patient was described with central pain syndrome.[30] Furthermore, cognitive symptoms were not reported. At present, however, the reason for as well as the consequence of the raised APP in patients with persistent headache following COVID-19 is elusive and follow-up studies are clearly needed.

Recent studies suggest that serum markers, known to be associated with cognitive impairment when measured in CSF, are also linked to cognitive impairment.[31] Thus, Urbano T et al. found a positive correlation between neurofilament light chain protein levels in serum and CSF in patients with neurocognitive disorders.[31] Moreover, a recent study revealed that the changes in blood biomarkers were linked to brain structural imaging patterns associated with Alzheimer's disease, as well as and lower cognitive test scores. Several of these proteins such as Tau, are known to be associated with Alzheimer's disease when detected in CSF.[29] To the best of our knowledge, whether a similar pattern will be seen for APP and PZP has not been studied.

Although a modest increase in the COVID-19 related headache group at inclusion, plasma levels of PZP were persistently elevated only those with prolonged headache following SARS-CoV-2 vaccination. The reason for this pattern remains unclear. PZP is known to promote a beneficial immunosuppression during pregnancy to prevent rejection of

the foetus. More recently, increased PZP levels have been observed during airway

infections,[12] and have been suggested as a marker of certain cancers.[32] Moreover, previous

studies with mouse models have indicated that APP may increase the entry of the SARS-Cov-

2 virus into cells, while PZP has been described to inhibiting amyloid aggregation.[33] Since

APP was lower in the vaccine group than the COVID-19 group, one intriguing hypothesis is

that PZP may have inhibited amyloid aggregation to a higher extent in the vaccine group than

in the COVID-19 group. However, PZP has also been reported to stabilize misfolded proteins

like amyloid β protein, and its role in amyloid formation remains unclear.[33] Based on this

relatively small study, which is at least hypothesis generating, it is hard to draw any firm

conclusion regarding primary and secondary effects. In our opinion data in the literature

regarding APP are more convincing in relation to its role in cognitive impairment compared

to PZP. Therefore, we currently consider elevated APP as the most important finding, with

potential some "additive" effects from PZP.

Since one of the three vaccines was disproportionately associated with headache, it

warrants a discussion of potential mechanisms. While the ChAdOx1 nCoV-19 vaccine has

been associated with severe cerebral events,[34] none of the patients in the present study

experienced new-onset persistent headaches after ChAdOx1 nCoV-19 vaccines. Moreover,

although the BNT162b2 vaccine was the most commonly vaccine, 89.7% of the participants

reported that their headache started after RNA-1273 vaccine. The reason for this pattern is not

clear, but interestingly, an online survey on self-reported adverse reactions in individuals who

had received two doses of either the BNT162b2 or mRNA-1273 vaccine reported more side

effects among those receiving the mRNA-1273 vaccine.[35] One could speculate that it was

related to the higher mRNA content in mRNA-1273 as compared with the BNT162b2

vaccine.[36] However, based on the low number of patients in the present study, the results

should be interpreted with caution.

The strengths of this study were the comprehensive examinations of the participants

done by headache specialists with precise phenotyping of the headache syndromes and the

strong multidisciplinary collaboration. The major limitations were the lack of CSF samples in

both headache groups and small sample size. Moreover, we lack parallel samples from CSF

and measurements of the actual protein in individuals that have undergone COVID-19 and

SARS-CoV-2 vaccination without the development of persistent headache. In particular, the

control group was small and could have been matched better in relation to age and undergone

longitudinal testing and been interviewed with more clinical information as in the patient

groups.

In summary, altered plasma levels of soluble markers that potentially could reflect

changes in amyloid processing was found in patients with persistent headache after SARS-

CoV-2 vaccine and particular in those with persistent headache after COVID-19 with

association to cognitive symptoms. Future and larger studies that also include CSF samples

are needed to further elaborate the clinical consequences of these findings.

Acknowledgement: We thank Marcus Boateng for registering data and Katrine Persgård

Lund and Azita Rashidi for the processing of blood biomarkers.

References

1. Caronna E, van den Hoek TC, Bolay H, et al. Headache attributed to SARS-CoV-2 infection, vaccination and the impact on primary headache disorders of the COVID-19 pandemic: A comprehensive review. Cephalalgia 2023;43:3331024221131337.

2. Shen Q, Joyce EE, Ebrahimi OV, et al. COVID-19 illness severity and 2-year prevalence of physical symptoms: an observational study in Iceland, Sweden, Norway and Denmark. Lancet Reg Health Eur 2023;35:100756.

3. Atalar AC, Acarli ANO, Baykan B, et al. COVID-19 vaccination-related headache showed two different clusters in the long-term course: a prospective multicenter follow-up study (COVA-Head Study). J Headache Pain 2023;24:132.

4. Files JK, Sarkar S, Fram TR, et al. Duration of post-COVID-19 symptoms is associated with sustained SARS-CoV-2-specific immune responses. JCI Insight 2021;6.

5. Gaebler C, Wang Z, Lorenzi JCC, et al. Evolution of antibody immunity to SARS-CoV-2. Nature 2021;591:639-644.

6. Garces KN, Cocores AN, Goadsby PJ, Monteith TS. Headache After Vaccination: An Update on Recent Clinical Trials and Real-World Reporting. Curr Pain Headache Rep 2022;26:895-918.

7. Hervé CL, B.; Del Giudice, G.; Didierlaurent, AM.; Tavares Da Silva, F. . The how's and what's of vaccine reactogenicity. NPJ Vaccines 2019;4.

8. Li HX, X.; Ren, H.; Xu, L.; Zhao, L.; Chen, X.; Long, H.; Wang, Q.; Wua, Q. Serum Amyloid A is a biomarker of severe Coronavirus Disease and poor prognosis. J Infect 2020;80:646–655.

9. Orobets KS, Karamyshev AL. Amyloid Precursor Protein and Alzheimer's Disease. Int J Mol Sci 2023;24.

10. Chen J, Chen J, Lei Z, et al. Amyloid precursor protein facilitates SARS-CoV-2 virus entry into cells and enhances amyloid-beta-associated pathology in APP/PS1 mouse model of Alzheimer's disease. Transl Psychiatry 2023;13:396.

11. Shao J, Jin Y, Shao C, Fan H, Wang X, Yang G. Serum exosomal pregnancy zone protein as a promising biomarker in inflammatory bowel disease. Cell Mol Biol Lett 2021;26:36.

12. Finch S, Shoemark A, Dicker AJ, et al. Pregnancy Zone Protein Is Associated with Airway Infection, Neutrophil Extracellular Trap Formation, and Disease Severity in Bronchiectasis. Am J Respir Crit Care Med 2019;200:992-1001.

13. Villar M, Urra JM, Rodriguez-Del-Rio FJ, et al. Characterization by Quantitative Serum Proteomics of Immune-Related Prognostic Biomarkers for COVID-19 Symptomatology. Front Immunol 2021;12:730710.

14. Ijsselstijn L, Dekker LJ, Stingl C, et al. Serum levels of pregnancy zone protein are elevated in presymptomatic Alzheimer's disease. J Proteome Res 2011;10:4902-4910.

15. Zhao MM, Yang WL, Yang FY, et al. Cathepsin L plays a key role in SARS-CoV-2 infection in humans and humanized mice and is a promising target for new drug development. Signal Transduct Target Ther 2021;6:134.

16. Wang H, Sang N, Zhang C, Raghupathi R, Tanzi RE, Saunders A. Cathepsin L Mediates the Degradation of Novel APP C-Terminal Fragments. Biochemistry 2015;54:2806-2816.

17. Beghi E, Moro E, Davidescu EI, et al. Comparative features and outcomes of major neurological complications of COVID-19. Eur J Neurol 2023;30:413-433.

18. Headache Classification Committee of the International Headache Society (IHS) The International Classification of Headache Disorders, 3rd edition. Cephalalgia 2018;38:1-211.

19. Camacho RC, Alabed S, Zhou H, Chang SL. Network Meta-analysis on the Changes
of Amyloid Precursor Protein Expression Following SARS-CoV-2 Infection. J Neuroimmune
Pharmacol 2021;16:756-769.
20. Caradonna A, Patel T, Toleska M, Alabed S, Chang SL. Meta-Analysis of APP
Expression Modulated by SARS-CoV-2 Infection via the ACE2 Receptor. Int J Mol Sci
2022;23.
21. O'Brien RJ, Wong PC. Amyloid precursor protein processing and Alzheimer's disease.
Annu Rev Neurosci 2011;34:185-204.
22. Di Benedetto G, Burgaletto C, Bellanca CM, Munafo A, Bernardini R, Cantarella G.
Role of Microglia and Astrocytes in Alzheimer's Disease: From Neuroinflammation to Ca(2+)
Homeostasis Dysregulation. Cells 2022;11.
23. Kursun O, Yemisci M, van den Maagdenberg A, Karatas H. Migraine and
neuroinflammation: the inflammasome perspective. J Headache Pain 2021;22:55.
24. Zheng H, Koo EH. The amyloid precursor protein: beyond amyloid. Mol
Neurodegener 2006;1:5.
25. Guo Y, Wang Q, Chen S, Xu C. Functions of amyloid precursor protein in metabolic
diseases. Metabolism 2021;115:154454.
26. Navolokin N, Adushkina V, Zlatogorskaya D, et al. Promising Strategies to Reduce the
SARS-CoV-2 Amyloid Deposition in the Brain and Prevent COVID-19-Exacerbated
Dementia and Alzheimer's Disease. Pharmaceuticals (Basel) 2024;17.
27. Danta CC. Calcium Channel Blockers: A Possible Potential Therapeutic Strategy for
the Treatment of Alzheimer's Dementia Patients with SARS-CoV-2 Infection. ACS Chem
Neurosci 2020;11:2145-2148.
28. Zhou Y, Xu J, Hou Y, et al. Network medicine links SARS-CoV-2/COVID-19
infection to brain microvascular injury and neuroinflammation in dementia-like cognitive
impairment. Alzheimers Res Ther 2021;13:110.
29. Duff EP, Zetterberg H, Heslegrave A, et al. Plasma proteomic evidence for increased
beta-amyloid pathology after SARS-CoV-2 infection. Nat Med 2025;31:797-806.
30. Ziff OJ, Ashton NJ, Mehta PR, et al. Amyloid processing in COVID-19-associated
neurological syndromes. J Neurochem 2022;161:146-157.
31. Urbano T, Maramotti R, Tondelli M, et al. Comparison of Serum and Cerebrospinal
Fluid Neurofilament Light Chain Concentrations Measured by Ella and Lumipulse in Patients
with Cognitive Impairment. Diagnostics (Basel) 2024;14.
32. Huang J, Xu Y, Chen Y, et al. Revisiting the role of pregnancy zone protein (PZP) as a
cancer biomarker in the immunotherapy era. J Transl Med 2024;22:500.
33. Cater JH, Kumita JR, Zeineddine Abdallah R, et al. Human pregnancy zone protein
stabilizes misfolded proteins including preeclampsia- and Alzheimer's-associated amyloid
beta peptide. Proc Natl Acad Sci U S A 2019;116:6101-6110.
34. Schultz NH, Sorvoll IH, Michelsen AE, et al. Thrombosis and Thrombocytopenia after
ChAdOx1 nCoV-19 Vaccination. N Engl J Med 2021;384:2124-2130.
35. Kitagawa H, Kaiki Y, Sugiyama A, et al. Adverse reactions to the BNT162b2 and
mRNA-1273 mRNA COVID-19 vaccines in Japan. J Infect Chemother 2022;28:576-581.
36. Dickerman BA, Gerlovin H, Madenci AL, et al. Comparative Effectiveness of
BNT162b2 and mRNA-1273 Vaccines in U.S. Veterans. N Engl J Med 2022;386:105-115.
515

517   A.

519   **A.**

520   **Figure 1.** Phenotype of persistent headache reported after COVID-19 (A) and SARS-CoV-2

521   vaccine (B).

Figure 2. Plasma levels of **A.** Amyloid Precursor Protein (APP), **B**. Cathepsin L., **C.**
Pregnancy Zone Protein (PZP) and **D**. Serum Amyloid A (SAA1) in healthy controls (HC)
(n=16), participants with persistent headache after SarsCoV-2 vaccine (COvax) (n=31) and
participants with persistent headache after COVID-19 (C19) (n=29).

Figure 3. Plasma levels of Amyloid Precursor Protein (AAP) in relation to cognitive
symptoms and fatigue in those with persistent headache after SARS-CoV-2 vaccine (COvax)
or COVID-19 (C19).

Figure 4. Possible mechanisms of soluble Amyloid Precursor Protein (sAPP) and Pregnancy

Zone Protein (PZP) in relation to COVID-19 and SARS-CoV-2 vaccination associated

headache. The authors wish to acknowledge SERVIER Medical Art (www.servier.fr) for use

of their medical art kits when making the illustration in the article.

Study funding: Supported by Oslo University Hospital and the South-Eastern Norway

Regional Health Authority (2021051).

Acknowledgement: We thank Marcus Boateng for registering data and Katrine Persgård

Lund and Azita Rashidi with the sampling of blood biomarkers.

Statistical Analysis: Thor Ueland and Anne Hege Aamodt

Disclosure

Conflicts of interest: The authors report no disclosures relevant to this study.

A.H. Aamodt has received honoraria for advice or lecturing from Pfizer Allergan, Teva,

Novartis, Roche, Lundbeck and Teva and research grant from Boehringer Ingelheim.

Anne Christine Poole: lecturing, advisory boards and/or clinical studies: Abbvie, Lundbeck,

Pfizer, Novartis, Teva, Eli Lilly

Supplemental Table 1. Plasma levels (ng/mL) of Amyloid Precursor Protein (APP),

Cathepsin L, Pregnancy Zone Protein (PZP) and Serum Amyloid A (SAA1) in participants

with persistent headache after COVID-19 (C19) in relation to severity of the acute infection

(n=29).

| Test statistics | Biomarkers | | | |
	APP	Cathepsin L	PZP	SAA1
Mann-Whitney U	61.00	49.00	60.00	69.00
Wilcoxon W	97.00	280.0	96.00	96.00
Z	-1,12	-1.71	-1.17	-0.73
Asymptotic Sig. (2-tailed)	0.26	0.09	0.24	0.46
Exact Sig	0.28	0.93	0.26	0.49

553
554

555
556   Supplemental Table 2.

Co-investigators	Affiliations
Øyvind Torkildsen	Neuro-SysMed, Department of Neurology, Haukeland University Hospital, Bergen, Norway; Department of Clinical Medicine, University of Bergen, Bergen, Norway
Vojetch Novotny	Department of Neurology, Haukeland University Hospital, Bergen, Norway
Heidi Øyen Flemmen	Department of Neurology, Telemark Hospital Trust, Skien, Norway
Hilde Karen Ofte	Department of Neurology, Nordlandssykehuset, Bodø, Norway
Christian Samsonsen	Department of Neurology of Clinical Neurophysiology, St. Olav Hospital, Trondheim, Norway
Barbara Ratajczak-Tretel	Department of Neurology, Østfold Hospital Trust, Sarpsborg, Norway
Anette Huuse Farmen	Department of Neurology, Innlandet Hospital Trust, Lillehammer, Norway
Per Selnes	Department of Neurology, Akershus University Hospital, Nordbyhagen, Norway and Institute of Clinical Medicine, Faculty of Medicine, University of Oslo, Oslo, Norway
Stein Andersson	Department of Psychology, University of Oslo, Oslo, Norway
Erlend Bøen	Unit for Psychosomatics and C-L psychiatry for adults, Oslo University Hospital, Oslo, Norway
Siv Elin Pignatiello	Unit for Psychosomatics and C-L psychiatry for adults, Oslo University Hospital, Oslo, Norway and Institute of Clinical Medicine, Faculty of Medicine, University of Oslo, Oslo, Norway