Robust uncertainty quantification in popular estimators of the instantaneous reproduction number

Nicholas Steyn[*1] and Dr Kris V Parag[2]

[1]Department of Statistics, University of Oxford, Oxford, United Kingdom
[2]MRC Centre for Global Infectious Disease Analysis, Imperial College London, London, United Kingdom

Abstract

The instantaneous reproduction number (R_t) is a key measure of the rate of spread of an infectious disease. Correctly quantifying uncertainty in R_t estimates is crucial for making well-informed decisions. Popular R_t estimators leverage smoothing techniques to distinguish signal from noise. Examples include EpiEstim and EpiFilter, which are both controlled by a "smoothing parameter" that is traditionally selected by users. We demonstrate that the values of these smoothing parameters are unknown, vary markedly with epidemic dynamics, and show that data-driven smoothing is crucial for accurate uncertainty quantification of R_t estimates. We derive model likelihoods for the smoothing parameters in both EpiEstim and EpiFilter and develop a Bayesian framework to automatically marginalise these parameters when fitting to epidemiological time-series data. This yields novel marginal posterior predictive distributions which prove integral to rigorous model evaluation. Applying our methods, we find that default parameterisations of these widely-used estimators can negatively impact R_t inference, delaying detection of epidemic growth, and misrepresenting uncertainty (typically producing overconfident estimates), with implications for public health decision-making. Our extensions mitigate these issues, provide a principled approach to uncertainty quantification, improve the robustness of real-time R_t inference, and facilitate model comparison using observable quantities.

The instantaneous reproduction number R_t, defined as the average number of infections generated per primary infection at time t, is a popular measure of epidemic spread [1]. A value of $R_t < 1$ indicates a declining epidemic while $R_t > 1$ indicates a growing epidemic. This quantity is useful for policymakers as it gives the change in transmission required to control the epidemic, informing decisions about public health interventions [2, 3, 4, 5, 6]. As a stark example, in June 2020, an R_t estimate of 1.01 was used to justify continued school closures in Greater Manchester, England [7, 8]. Additionally, estimates of R_t are also used for forecasting, scenario analysis, and understanding the impact of interventions [9].

Many models exist to estimate R_t, including compartmental models, agent-based models, and the renewal-model [10], the latter of which underlies most contemporary methods for real-time estimation of R_t from reported case time-series, including EpiEstim [11] and EpiFilter [12]. Reflecting a community drive to improve existing methods and make public health pipelines more reliable, we focus on quantifying uncertainty in these two models, although our approach generalises to any real-time model.

Correct quantification of the uncertainty of R_t is crucial for making well-informed decisions. It is expected that the true value of R_t should fall within a 95% credible interval 95% of the time. Undercoverage occurs when this happens less often than expected, leading to over-confident and biased decision-making. Overcoverage occurs when this happens more often than expected, leading to under-confident and highly uncertain decision-making. A model that produces correct coverage is termed well-calibrated.

Despite the importance of uncertainty quantification,

*Corresponding author: nicholas.steyn@univ.ox.ac.uk

calibration is often neglected in epidemiological models. For example, during the COVID-19 pandemic, SPI-M in England pooled estimates from multiple groups to produce consensus R_t estimates. However, pooling proved difficult due to models "*providing estimates with lower levels of uncertainty that are not fully accounting for inherent uncertainties*" [13]. Even when correct coverage is explicitly targeted, epidemic models frequently fail to achieve it. For instance, nearly all models submitted to the *open challenge to advance probabilistic forecasting for dengue epidemics* [14] produced over-confident predictions across various forecasting targets. A baseline model (included for comparison) demonstrated superior calibration compared to all 16 submitted models when predicting the peak of the epidemic.

Smoothing assumptions, which include penalized likelihoods [15], piecewise constant/trailing window models [11, 16, 17, 18], and latent-space models [12, 19, 20, 21], are a key source of model miscalibration [22]. Oversmoothed estimates result in delayed and overconfident estimates, while undersmoothed estimates are noisy and lack precision. Even with perfect case reporting (i.e. no observation noise), inherent stochasticity in the transmission of infectious diseases necessitates the use of smoothing to distinguish signal from noise. All popular estimators of R_t employ some type of smoothing.

Despite the importance and ubiquity of these assumptions, the philosophical and practical treatment of smoothing parameters varies by method. Some methods treat these parameters as unknown quantities to be estimated alongside R_t [19, 22, 23, 24], while others treat the choice of smoothing parameter(s) as a model selection problem and seek to find a fixed, optimal parameter value [22, 25, 26]. Some methods allow the user to choose their own values or provide heuristic default values [11, 15, 24].

We argue that, because the true dynamics of R_t are always unknown and depend on both the pathogen and the population in which the pathogen is spreading, uncertainty about the nature of these dynamics should not be ignored. This uncertainty factors in both the choice of the dynamic model itself (structural uncertainty), and the parameters associated with the chosen dynamic model (parametric uncertainty). We focus on parametric uncertainty in this paper, which on its own can cause substantial model miscalibration, and demonstrate that correct marginalisation of smoothing parameters generally improves model robustness. This result holds even if the structure of the dynamic model does not accurately reflect reality, as the increased flexibility of the model allows it to better adapt to the data.

In this paper, we derive novel likelihoods for smoothing parameters in two popular models: Cori et al. [11] and Parag [12], commonly known as EpiEstim and EpiFilter respectively[1]. We use these likelihoods to marginalise the smoothing parameters, presenting estimates of R_t that appropriately account for uncertainty in these parameters. We also derive predictive posterior distributions and demonstrate their use in model comparison via the continuous ranked probability score (CRPS) [27], emphasising the importance of model comparison using observable quantities. We validate our methods on both simulated data, where model estimates can be compared to ground truths, and real-world data, where we investigate the practical implications on decision-making during the COVID-19 pandemic in New Zealand. For each dataset considered, we fit four models: EpiEstim and EpiFilter with default parameters and with our marginalised algorithm.

Methods

We summarise our methods here but provide full mathematical derivations and technical details in supplementary section 1.

Background

Both EpiEstim and EpiFilter leverage the *Poisson renewal model* for R_t estimation. Letting C_s be the number of cases reported at time s and w_u be the probability that a secondary case is reported u days after the primary case (often approximated by the serial interval). The *total infectiousness* is defined as $\Lambda_t = \sum_{u=1}^{t-1} C_{t-u} w_u$. Given Λ_t and the current value of R_t, the Poisson renewal model considers the number of cases at time t to be Poisson distributed:

$$C_t \sim \text{Poisson}(R_t \Lambda_t) \qquad (1)$$

EpiEstim assumes that, on each day t, R_t has been fixed for a trailing window of k days. Larger values of k imply that R_t has been fixed for a longer period, resulting in smoother estimates. The likelihood of observing cases between time-steps $t - k + 1$ and t (denoted $C_{t-k+1:t}$) is the product of the daily Poisson likelihoods (equation 1) along the trailing win-

---

[1]Technically, the terms EpiEstim and EpiFilter refer to the associated software packages. We follow popular convention and use these terms to describe the method.

dow. A conjugate $\text{Gamma}(\alpha, \beta)$ prior distribution is assumed for R_t, resulting in a $\text{Gamma}(\alpha_{t,k}, \beta_{t,k})$ posterior distribution for R_t given $C_{1:t}$, where the shape-parameter $\alpha_{t,k}$ and rate-parameter $\beta_{t,k}$ are:

$$\alpha_{t,k} = \alpha + \sum_{s=t-k+1}^{t} C_s, \quad \beta_{t,k} = \beta + \sum_{s=t-k+1}^{t} \Lambda_s \quad (2)$$

Alternatively, EpiFilter assumes that R_t follows a Gaussian random walk with standard deviation equal to $\eta\sqrt{R_{t-1}}$. Larger values of η allow R_t to vary faster, resulting in less smooth estimates. A grid-approximation to the exact Bayesian filtering equations is used to derive the posterior distribution of R_t given $C_{1:t}$. While EpiFilter also allows for the estimation of the smoothing distribution (R_t given past and future data), we focus on real-time estimation (R_t given data up to time t). Further discussion is included in supplementary section 2.

Model likelihoods

We use the same framework to derive likelihoods for the smoothing parameters of both methods. Model-specific derivations are included in supplementary section 1. Letting θ denote an arbitrary smoothing parameter, we begin with the predictive decomposition of the likelihood:

$$\log P(C_{1:T}|\theta) = \sum_{t=1}^{T-1} \log P(C_{t+1}|C_{1:t}, \theta) \quad (3)$$

The one-step-ahead-likelihood can be written as:

$$P(C_{t+1}|C_{1:t}, \theta) = \int P(C_{t+1}|R_{t+1}, C_{1:t})P(R_{t+1}|C_{1:t}, \theta)\, dR_{t+1} \quad (4)$$

where $P(C_{t+1}|R_{t+1}, C_{1:t})$ is the renewal model (equation 1). Thus, deriving model likelihoods relies on the derivation of the one-step-ahead predictive distribution for R_{t+1}: $P(R_{t+1}|C_{1:t}, \theta)$.

For EpiEstim, R_t depends on reported cases only on days $t-k+1$ to t, however the predictive distribution explicitly ignores data on day t, so EpiEstim's predictive distribution for R_t is Gamma-distributed with shape $\alpha_{t-1,k-1}$ and rate $\beta_{t-1,k-1}$. In this case, equation 4 is a Gamma-Poisson mixture, hence $C_{t+1}|C_{1:t}$ follows a negative binomial distribution with parameters $r = \alpha_{t,k-1}$ and $p = \frac{\beta_{t,k-1}}{\beta_{t,k-1}+\Lambda_{t+1}}$. The likelihood of

EpiEstim's k is then the sum of log-negative binomial probability mass functions for each day t.

Figure 1: Schematic demonstrating the derivation of parameter likelihoods using the predictive decomposition (equation 3). R_t estimates are projected forward one time-step according to the relevant dynamic model (panel A), giving predictive distributions for R_{t+1} (panel B). These are combined with the renewal model to give the probability of observing C_{t+1} conditional on past data $C_{1:t}$ and the chosen smoothing parameter θ (panel C). In this time-step, the undersmoothed model is more likely than the oversmoothed model. Repeating this process and summing the log-predictive probabilities for all time-steps produces the model likelihood (Panel D). These log-likelihoods are later used to marginalise out uncertainty about θ, allowing for robust reporting of uncertainty about R_t. The form of the R_t estimates and predictive distributions for R_t depend on the specific model.

For EpiFilter, the predictive distribution of R_{t+1} is a by-product of the Bayesian filtering equations, found by propagating the estimated distribution of R_t given $C_{1:t}$ forward according to the assumed Gaussian random walk. The one-step-ahead likelihood of C_{t+1} is found by taking a weighted average of the Poisson

likelihood of C_{t+1} with respect to the predictive distribution of $R_{t+1}|C_{1:t}$.

Figure 1 provides a schematic of the model likelihood calculation. Implementations of these methods, alongside worked examples, are provided in the GitHub repository.

Posterior distributions

These methods admit log-likelihoods for k and η given $C_{1:t}$, denoted $\ell(k|C_{1:t})$ and $\ell(\eta|C_{1:t})$. We use these to derive posterior distributions (denoted $P(k|C_{1:t})$ and $P(\eta|C_{1:t})$), typically using a uniform prior distribution over $k \in \{1, 2, \ldots, 30\}$ (covering daily to monthly dependence) and $\eta \in [0, 1]$ (covering no noise to Poisson-type diffusion).

We are ultimately interested in estimates of R_t that account for uncertainty about k or η. To achieve this, we marginalise these parameters from the posterior distribution of R_t, a procedure that is rare in the literature. For EpiEstim, we leverage the discrete nature of k to write exactly:

$$P(R_t|C_{1:t}) = \sum_{k=1}^{30} P(R_t|C_{1:t}, k)P(k|C_{1:t}) \quad (5)$$

whereas for EpiFilter, we use a grid-approximation ($\eta \in \mathcal{E}$, supplementary section 1.1.2):

$$\begin{aligned} P(R_t|C_{1:t}) &= \int P(R_t|C_{1:t}, \eta)P(\eta|C_{1:t}) \, d\eta \\ &\approx \sum_{\eta \in \mathcal{E}} P(R_t|C_{1:t}, \eta)P(\eta|C_{1:t}) \end{aligned} \quad (6)$$

We can also marginalise the smoothing parameter from the predictive distributions for C_t (equation 4). This generates marginal one-step-ahead predictive distributions for C_t under both models, useful for model comparison and probabilistic forecasting.

Model evaluation

We argue that a "good" model is one that maximises precision (i.e. minimises the width of uncertainty intervals), subject to being well-calibrated [27]. Choosing the "best" model thus involves a trade-off: how much miscoverage are we willing to accept in exchange for more precise estimates? We use the Continuous Ranked Probability Score (CRPS), which measures the distance between the estimated predictive distribution and the empirical distribution of the data, to quantify this trade-off. Smaller distances signify closer alignment between the model's predictive uncertainty and observed data variability. Full details of the CRPS calculation are provided in supplementary section 1.4.

Data

We test our methods on three simulated datasets, each assuming a different dynamic model for R_t: a Gaussian random walk (matching the dynamic model assumed by EpiFilter), a sinusoidal curve, and a step-change model. These models cover a range of smooth to sharp changes in R_t. We also compare model outputs on real-world data from the COVID-19 pandemic in New Zealand [28], chosen as an example of high-quality data with limited reporting biases. We explicitly relate real-world decision-making to the inferences made by our models.

A common serial interval from the COVID-19 literature, a Gamma distribution with a mean of 6.5 days and standard deviation of 4.2 days [29, 30], is used for the simulation study, while a Weibull distribution with a mean of 5.0 days and standard deviation of 1.9 days is used when fitting to real-world data, matching the serial interval used in official modelling [31].

Further information on simulated data is provided in supplementary section 3. We also develop additional marginalisation routines to handle uncertainty in serial intervals, and test sensitivity to biases in the serial interval, in supplementary section 4.

Results

Simulation study

Fitting EpiEstim and EpiFilter to a single realisation from each of the three simulated epidemics (figure 2) demonstrates that default parameterisations of both models result in oversmoothed estimates of R_t, relative to the level of smoothing estimated from the data. The exception is EpiFilter in the random walk simulation, where the true value of η is deliberately chosen to match EpiFilter's default $\eta = 0.1$.

Using these default parameters results in credible intervals for R_t that typically undercover the true value. This is more noticeable for EpiEstim, where coverage of the 95% credible intervals for R_t in the default model ranges from just 8.9% in the sinusoidal simulation, to 74.4% in the step-change simulation. Marginalising out k dramatically improves coverage in these models to 81.1% and 94.4% respectively. Default EpiFilter is generally more robust, partially as

Figure 2: Simulated case data (A, B, C), posterior distributions for smoothing parameters at $t = 100$ (D, E, F), estimates of R_t (G, H, I), and estimates of predictive cases (J, K, L) for one realisation of each simulated epidemic. Estimates of R_t and predictive cases are shown for all four models (default EpiEstim, default EpiFilter, marginalised EpiEstim, marginalised EpiFilter). The first column (A, D, G, J) shows results for the Gaussian random walk simulation with $\eta = 0.1$, a dynamic model that precisely matches default EpiFilter. The second column (B, E, H, K) shows results for the sinusoidal simulation, and the third column (C, F, I, L) shows results for the step-change simulation. In panels (D, E, F), vertical dotted lines indicate default parameter values and the vertical solid line indicates both the default and true parameter value. In panels (G-L), solid coloured lines show central estimates (posterior means) and shading shows 95% credible intervals. Black lines (in R_t estimates) and black dots (in predictive C_t estimates) show the true values of R_t and C_t respectively. Predictive coverage of the 95% credible intervals (closer to 95% is better) and the CRPS (lower is better) are shown as text within each figure.

a result of the default $\eta = 0.1$ being less extreme with respect to the posterior distribution of η, although marginalising this parameter still improves coverage of R_t from 81.1% to 92.2% in the sinusoidal model.

The one-step-ahead predictive coverage of reported cases is also improved by marginalising the smoothing parameter. This is true for all models and simulations considered, but the effect is more pronounced in EpiEstim.

Finally, marginalising the smoothing parameter generally improves (decreases) the CRPS, suggesting that marginalised models produce more accurate predictive distributions of cases than default models. The sole exception is EpiFilter in the step-change simulation, where the CRPS worsens (increases) slightly, although we show in supplementary section 5 that, on average, marginalisation also improves EpiFilter when fit to step-change simulations. The higher CRPS in this specific simulation suggests that the narrower credible intervals produced by default EpiFilter may trade off favourably against its lower predictive coverage of cases. We observe a similar effect in the sinusoidal simulation when comparing EpiEstim and EpiFilter, where the CRPS is slightly lower in EpiEstim (indicating it as the better model), despite EpiFilter having better coverage of predictive cases.

These results depend on the simulated ground truth, and we consider different simulations in supplementary section 5, revealing how the appropriate level of smoothing depends on the underlying epidemic dynamics. Marginalisation becomes more important as the standard deviation of the simulated random walk increases, the frequency of the sinusoidal curve increases, or the step-change becomes more frequent. In all these scenarios, the true R_t is more dynamic and the default smoothing choices are progressively worse at adapting to these changes.

We also find, contrary to intuition, that fitting to greater numbers of daily cases does not necessarily improve inference quality. Supplementary figure S4 demonstrates that, while marginalised EpiFilter is largely robust to sample size, EpiEstim's coverage worsens as sample size increases in both the default and marginalised models. This occurs as guaranteed misspecification (R_t cannot be constant on $[t - k + 1, t]$ and then constant at a different value on $[t - k + 2, t + 2]$) results in the model becoming more confident in the incorrect estimate. For similar reasons, if observation noise is large, both models are also expected to degrade as sample size increases.

The COVID-19 pandemic in New Zealand, August-December 2021

After largely containing the spread of COVID-19, in August 2021 an outbreak of the delta-variant was detected in Auckland, New Zealand, triggering strict non-pharmaceutical interventions. The outbreak featured an initial peak in late August, followed by a subsequent period of decline, and then a second peak in mid-November (figure 3). R_t was repeatedly cited during decision-making, including by the Prime Minister and the Director-General of Health [3, 32, 33, 34, 4, 35, 36].

We fit all four models to reported cases (smoothed using a 5-day moving average to decrease reporting noise; supplementary section 6 tests sensitivity to this assumption) between 11 August and 1 December 2021 (figure 3). Both EpiEstim and EpiFilter exhibit improved CRPS after marginalisation, and EpiEstim exhibits improved predictive coverage after marginalisation. As R_t is unknown, we cannot evaluate its calibration.

As in the simulations, default models oversmooth relative to marginalised models. This is most notable for EpiEstim where almost all posterior mass is on $k = 2$. For EpiFilter, the MAP value of η was 0.12 with a 95% credible interval of $(0.102, 0.143)$, which nearly contains the default value. In this case, default EpiFilter is better calibrated than default EpiEstim. However, table 1 still highlights some delays in detection of epidemic growth or decline.

On 31 August, marginalised EpiFilter first signalled $R_t < 1$. Among the four models, default EpiEstim signalled this change last (3 days later). Notably, both marginalised models were confident that $R_t < 1$ by 2 September, before the possibility of decline had been detected by default EpiEstim. This was a period characterised by daily press conferences and strict stay-at-home orders, estimated to have cost the city NZ\$56m per day [37].

On 13 September, the Government announced that because restrictions had *"...reduced that value [R_t] down to consistently below one"*, they would be relaxed the following week on 21 September [34]. There was considerable interest in R_t over that week, as a resurgence could have triggered prolonged restrictions. This announcement occurred after the first detection of a potential resurgence by the marginalised models (11 September), while default EpiFilter first detected this resurgence on the day of the announcement, and default EpiEstim two days later. The marginalised models first produced central estimates

Figure 3: Reported case data and parameter posterior estimates (A), reproduction number estimates (B), predictive cases (C) and reproduction number estimates for selected periods (D, E, F) from fitting the four models to reported case data from the outbreak of the delta-variant of SARS-CoV-2 in Auckland, New Zealand between 11 August 2021 and 1 December 2021. In panel A, vertical dashed lines indicate default parameter values while coloured curves show parameter posterior distributions. In panels (B, C), solid lines show central estimates (posterior means) and shading shows posterior 95% credible intervals. Black points show observed reported cases. In panels (D, E, F), middle lines show central estimates (posterior means) while shading and outer lines show posterior 95% credible intervals. Results for default and marginalised EpiFilter in period 3 (Panel F) are nearly identical and overlap considerably. Our methods improve predictive coverage of EpiEstim and the CRPS of both models.

| | EpiEstim | | EpiFilter | |
	Default	Marginalised	Default	Marginalised
Period 1				
First detection	3 Sep (+3)	31 Aug (+0)	1 Sep (+1)	31 Aug (+0)
Expected $R_t < 1$	4 Sep (+3)	1 Sep (+0)	1 Sep (+0)	1 Sep (+0)
Confidence in $R_t < 1$	4 Sep (+2)	2 Sep (+0)	2 Sep (+0)	2 Sep (+0)
Period 2				
First detection	15 Sep (+4)	11 Sep (+0)	13 Sep (+2)	11 Sep (+0)
Expected $R_t > 1$	30 Sep (+17)	13 Sep (+0)	20 Sep (+7)	13 Sep (+0)
Confidence in $R_t > 1$	1 Oct (+1)	30 Sep (+0)	1 Oct (+1)	30 Sep (+0)
Period 3				
First detection	20 Nov (+5)	15 Nov (+0)	15 Nov (+0)	15 Nov (+0)
Expected $R_t < 1$	27 Nov (+9)	19 Nov (+1)	19 Nov (+1)	18 Nov (+0)
Confidence in $R_t < 1$	29 Nov (+1)	28 Nov (+0)	28 Nov (+0)	28 Nov (+0)

Table 1: Dates of the first detection of growth or decline (defined as the upper bound on the 95% credible interval of R_t crossing 1 for growth, or the lower bound crossing 1 for decline), the date that the central estimates of R_t first crossed 1, and the date that the models are first confident in growth or decline (defined as the lower bound on the 95% credible interval of R_t crossing 1 for growth, or the upper bound crossing 1 for decline) for the COVID-19 outbreak in Auckland, New Zealand that started in August 2021. The number of days after the first detection is shown in parentheses, highlighting the substantial delay often observed in default models.

of $R_t > 1$ on the same day as the announcement, while only 7 days and 17 days later that default Epi-Filter and default EpiEstim estimate $R_t > 1$, respectively.

Daily reported cases continued to increase until mid-November, before appearing to plateau. While the marginalised models (and default EpiFilter) became uncertain about continued epidemic growth around 15 November, default EpiEstim was still confident that $R_t > 1$ until 20 November (for 5 days longer), a clear example of oversmoothed models being over-confident.

We test our methods on additional real-world datasets in supplementary section 6. These examples highlight that real-world data feature many unknowns with no known ground truth. In every case but one, marginalisation decreased the CRPS, emphasising that the most robust models are those that are able to adapt to these unknowns, as well as the importance of validation on observable quantities.

Discussion

We have derived and analysed novel likelihoods for two popular R_t estimators. We used these likelihoods to develop posterior distributions for the corresponding smoothing parameters, and marginalised out this parametric uncertainty from estimates of R_t. Our algorithms generally improve uncertainty quantification for R_t, allowing increased confidence in the calibration of reported credible intervals.

Robust uncertainty quantification is crucial for real-world decision-making. Our methods provide one of the first principled and computationally efficient ways of ensuring the robustness of reported uncertainty in two popular models: EpiEstim [11] and Epi-Filter [12]. Furthermore, our methods can be applied to other sequential models (i.e., any model that estimates R_t using $C_{1:t}$ and then R_{t+1} using $C_{1:t+1}$ and so on), such as [20, 38, 39], all of which currently assume fixed values of smoothing parameters.

In addition to real-time decision-making, estimates of R_t are also used in models investigating the impact of non-pharmaceutical interventions [40, 41, 42, 43, 44, 45, 46, 47, 48], the effect of climate [49], or the relationship between mobility and transmission [50]. These models all use EpiEstim to estimate R_t, most using $k = 7$. While some correct for smoothing-induced delays by shifting estimates by $k/2$ days, this deterministic correction still ignores uncertainty about k. The Bayesian nature of our methods allows

for the propagation of uncertainty in k through to estimates of R_t and thus to downstream models, such as the consensus estimates of R_t produced by SPI-M in England during the COVID-19 pandemic.

While we focus on parametric uncertainty, by fitting our models to simulated data from a range of dynamic models, our work reveals that marginalising smoothing parameters improves model robustness, even when ignoring structural uncertainty. This is observed by improved coverage of predictive cases and decreased CRPS in simulations and on real data, by improved coverage of R_t in simulations, and is a key finding. As R_t is always unknown, it is not possible to compare coverage of this quantity on real data, and we rely on the CRPS on observables as a proxy for model performance (justified by our results on simulated data).

We used two metrics to evaluate model performance: coverage of the 95% credible intervals and CRPS. The former measures calibration only, while the latter also factors in precision (and simultaneously considers all credible intervals, not just the 95% level). Calibration and precision are often a trade-off: improved calibration can be obtained by decreasing precision. The CRPS is a principled way of balancing these two goals. Alternative scoring rules may be appropriate depending on context [51]. The documentation of [52] summarises scoring rules in an epidemiological context.

While our results suggest that improved CRPS implies improved estimates of R_t even when the model is misspecified, there is no guarantee of performance in such situations. Since misspecification is inevitable when modelling infectious diseases, any scoring rule should be interpreted with caution. One example faced by almost all R_t estimators is serial interval misspecification, addressed in supplementary section 4. Like existing literature [18, 26, 53], we find that misspecified serial interval distributions lead to biased estimates of R_t, although estimates near $R_t = 1$ are more robust. Another example of model misspecification comes from observation noise: EpiEstim and EpiFilter assume that reported cases follow the Poisson renewal model and that the appropriate level of smoothing is fixed over time (supplementary section 7). We demonstrate that our methods help with robustness to observation noise in supplementary section 5.3, but the lack of explicitly representing such noise remains a limitation of both models. We chose the New Zealand dataset as an example of high-quality data with limited reporting biases to reduce the impact of this limitation.

Alternative parameter selection procedures for Epi-Estim have previously been proposed. In the supplement of [11], the authors suggest selecting k such that the window contains sufficient cases to reduce the posterior coefficient-of-variation to a desired level. Depending on philosophy, this is either a subjective decision about the bias-variance trade-off, or a way of choosing parameters to obtain a desired confidence level. In either case, choosing a value of k with low likelihood will lead to poor model calibration. Alternatively, Parag and Donnelly [25] proposed an information-theoretic approach to selecting k called APEestim. While their approach is principled, it results in the selection of k shifted by one unit compared to the MAP value from our method, and does not allow for the marginalisation of uncertainty when there are multiple plausible values of k (supplementary section 8). EpiEstim and EpiFilter are Bayesian estimators, thus marginalising smoothing parameters is a more justified approach than selecting a single value. This is supported by supplementary section 2, where two implementations of a different R_t estimator (EpiLPS [22]) are compared: one implementation optimises the smoothing parameters while the other performs marginalisation. We find marginalisation continues to offer improved performance and robustness.

Other methods approach the smoothing problem differently. For example, EpiNow2 [19] models R_t using a Gaussian process. The smoothness of this model is determined by the covariance kernel, which is estimated alongside R_t. Comparisons with additional methods (EpiNow2 [19], EpiLPS [22], and rtestim [26]) are included in supplementary section 2. These methods include features such as explicitly representing observation noise, at the cost of increased mathematical complexity and decreased interpretability, when compared to EpiEstim and EpiFilter. We also provide novel methodology for finding posterior predictive distributions and calculating CRPS values for these models, providing model comparison techniques using observed data. While Epi-Estim is often outperformed by these methods, there is value in the simplicity and interpretability of a sliding window. Understanding the nuances of various approaches to smoothing and hence inference-based decision-making will form a future study.

The August 2021 outbreak of SARS-CoV-2 in New Zealand provides a pertinent example of the practical importance of smoothing assumptions. R_t was first reported by officials as being less than 1 on 29 August, two days before any of our models. While official models were also based on the renewal model, among other differences (e.g. accounting for asymptomatic infections), they approached the smoothing problem differently, assuming R_t was fixed prior to the lockdown on 18 August, and then step-changed to a different fixed value after the lockdown [31]. These piecewise-constant assumptions allowed more data to inform each estimate of R_t, reducing uncertainty. However, if these assumptions were incorrect then uncertainty about R_t will have been underestimated.

It is often argued that public health decision-making should be "data-driven", with R_t frequently featuring as an example of such "data" [54]. However, without an accurate representation of uncertainty, estimates of R_t risk being influenced more by assumptions than the underlying data. As demonstrated on both simulated and real-world data, public health decisions made using oversmoothed estimates of R_t will be delayed and overconfident relative to decisions made using estimates with more robust uncertainty quantification. Fortunately, our methods provide a simple and computationally efficient way to improve the robustness of these estimates, and to benchmark the uncertainty surrounding smoothing assumptions.

Additional material

In addition to the supplementary material, all code is available on the GitHub repository. This repository contains all code necessary to reproduce the results in this paper, as well as Julia implementations of both EpiEstim and EpiFilter. Tutorials for the use of these methods are provided on the corresponding website.

Acknowledgements

N.S. acknowledges support from the Oxford-Radcliffe Scholarship from University College, Oxford, the EPSRC CDT in Modern Statistics and Statistical Machine Learning (Imperial College London and University of Oxford), and A. Maslov for studentship support. K.V.P. acknowledges funding from the MRC Centre for Global Infectious Disease Analysis (Reference No. MR/X020258/1) funded by the UK Medical Research Council. This UK-funded grant is carried out in the frame of the Global Health EDCTP3 Joint Undertaking. The funders had no role in study design, data collection and analysis, decision to publish, or manuscript preparation.

Declaration of interests

The authors declare no competing interests.

Data availability statement

All data used in this study is publicly available from [28]. All data and code used in our analysis is available at https://github.com/nicsteyn2/RobustRtEstimators.

References

[1] Kris V. Parag, Robin N. Thompson, and Christl A. Donnelly. "Are Epidemic Growth Rates More Informative than Reproduction Numbers?" In: *Journal of the Royal Statistical Society Series A: Statistics in Society* 185.Supplement_1 (Nov. 2022), S5–S15. ISSN: 0964-1998. DOI: 10.1111/rssa.12867. (Visited on 06/27/2024).

[2] Ruth McCabe and Christl A. Donnelly. "Disease Transmission and Control Modelling at the Science–Policy Interface". In: *Interface Focus* 11.6 (Oct. 2021), p. 20210013. DOI: 10.1098/rsfs.2021.0013. (Visited on 09/12/2023).

[3] Ministry of Health NZ. *COVID-19 Update 30 August 4pm*. https://www.health.govt.nz/news/covid-19-update-30-august-4pm. Aug. 2021. (Visited on 09/04/2024).

[4] Ministry of Health NZ. *COVID-19 Update 11 October 2021 4pm*. https://www.health.govt.nz/news/covid-19-update-11-october-2021-4pm. Oct. 2021. (Visited on 09/04/2024).

[5] Center for Infectious Disease Research and Policy. *No 'reset' with Ebola Outbreak, WHO Official Says*. https://www.cidrap.umn.edu/ebola/no-reset-ebola-outbreak-who-official-says. June 2019. (Visited on 09/05/2024).

[6] 10 Downing Street Prime Minister's Office. *Prime Minister's Statement on Coronavirus (COVID-19): 10 May 2020*. https://www.gov.uk/government/speeches/pm-address-to-the-nation-on-coronavirus-10-may-2020. May 2020. (Visited on 09/05/2024).

[7] Lorenzo Pellis et al. "Estimation of Reproduction Numbers in Real Time: Conceptual and Statistical Challenges". In: *Journal of the Royal Statistical Society Series A: Statistics in Society* 185.Supplement_1 (Nov. 2022), S112–S130.

[8] ISSN: 0964-1998. DOI: 10.1111/rssa.12955. (Visited on 09/05/2024).

[8] P. A. Media. "Schools in North-West of England Postpone Reopening Plans after New Coronavirus Data". In: *The Guardian* (June 2020). ISSN: 0261-3077. (Visited on 10/09/2024).

[9] The Royal Society. *Reproduction Number (R) and Growth Rate (r) of the COVID-19 Epidemic in the UK*. Tech. rep. 2020.

[10] Anne Cori and Adam Kucharski. "Inference of Epidemic Dynamics in the COVID-19 Era and Beyond". In: *Epidemics* 48 (Sept. 2024), p. 100784. ISSN: 1755-4365. DOI: 10.1016/j.epidem.2024.100784. (Visited on 09/02/2024).

[11] Anne Cori et al. "A New Framework and Software to Estimate Time-Varying Reproduction Numbers During Epidemics". In: *American Journal of Epidemiology* 178.9 (Nov. 2013), pp. 1505–1512. ISSN: 0002-9262. DOI: 10.1093/aje/kwt133. (Visited on 09/09/2023).

[12] Kris V. Parag. "Improved Estimation of Time-Varying Reproduction Numbers at Low Case Incidence and between Epidemic Waves". In: *PLOS Computational Biology* 17.9 (Sept. 2021), e1009347. ISSN: 1553-7358. DOI: 10.1371/journal.pcbi.1009347. (Visited on 09/10/2023).

[13] Thomas Maishman et al. "Statistical Methods Used to Combine the Effective Reproduction Number, R(t), and Other Related Measures of COVID-19 in the UK". In: *Statistical Methods in Medical Research* 31.9 (Sept. 2022), pp. 1757–1777. ISSN: 0962-2802. DOI: 10.1177/09622802221109506. (Visited on 09/05/2024).

[14] Michael A. Johansson et al. "An Open Challenge to Advance Probabilistic Forecasting for Dengue Epidemics". In: *Proceedings of the National Academy of Sciences* 116.48 (Nov. 2019), pp. 24268–24274. DOI: 10.1073/pnas.1909865116. (Visited on 09/09/2023).

[15] Niel Hens et al. "Estimating the Effective Reproduction Number for Pandemic Influenza from Notification Data Made Publicly Available in Real Time: A Multi-Country Analysis for Influenza A/H1N1v 2009". In: *Vaccine* 29.5 (Jan. 2011), pp. 896–904. ISSN: 0264-410X. DOI: 10.1016/j.vaccine.2010.05.010. (Visited on 04/17/2024).

[16] Christophe Fraser et al. "Influenza Transmission in Households During the 1918 Pandemic". In: *American Journal of Epidemiology* 174.5

(Sept. 2011), pp. 505–514. ISSN: 0002-9262. DOI: 10.1093/aje/kwr122. (Visited on 04/17/2024).

[17] Seth Flaxman et al. "Estimating the Effects of Non-Pharmaceutical Interventions on COVID-19 in Europe". In: *Nature* 584.7820 (Aug. 2020), pp. 257–261. ISSN: 1476-4687. DOI: 10.1038/s41586-020-2405-7. (Visited on 04/17/2024).

[18] R. N. Thompson et al. "Improved Inference of Time-Varying Reproduction Numbers during Infectious Disease Outbreaks". In: *Epidemics* 29 (Dec. 2019), p. 100356. ISSN: 1755-4365. DOI: 10.1016/j.epidem.2019.100356. (Visited on 09/09/2023).

[19] Sam Abbott et al. "Estimating the Time-Varying Reproduction Number of SARS-CoV-2 Using National and Subnational Case Counts". In: *Wellcome Open Research* 5 (Dec. 2020), p. 112. ISSN: 2398-502X. DOI: 10.12688/wellcomeopenres.16006.2. (Visited on 09/09/2023).

[20] Shinsuke Koyama, Taiki Horie, and Shigeru Shinomoto. "Estimating the Time-Varying Reproduction Number of COVID-19 with a State-Space Method". In: *PLOS Computational Biology* 17.1 (Jan. 2021), e1008679. ISSN: 1553-7358. DOI: 10.1371/journal.pcbi.1008679. (Visited on 09/05/2024).

[21] Amin Azmon, Christel Faes, and Niel Hens. "On the Estimation of the Reproduction Number Based on Misreported Epidemic Data". In: *Statistics in Medicine* 33.7 (2014), pp. 1176–1192. ISSN: 1097-0258. DOI: 10.1002/sim.6015. (Visited on 09/05/2024).

[22] Oswaldo Gressani et al. "EpiLPS: A Fast and Flexible Bayesian Tool for Estimation of the Time-Varying Reproduction Number". In: *PLOS Computational Biology* 18.10 (Oct. 2022), e1010618. ISSN: 1553-7358. DOI: 10.1371/journal.pcbi.1010618.

[23] Richard Creswell et al. "A Bayesian Nonparametric Method for Detecting Rapid Changes in Disease Transmission". In: *Journal of Theoretical Biology* 558 (Feb. 2023), p. 111351. ISSN: 0022-5193. DOI: 10.1016/j.jtbi.2022.111351.

[24] Oliver Eales et al. "Appropriately Smoothing Prevalence Data to Inform Estimates of Growth Rate and Reproduction Number". In: *Epidemics* 40 (Sept. 2022), p. 100604. ISSN: 1755-4365. DOI: 10.1016/j.epidem.2022.100604. (Visited on 09/06/2023).

[25] Kris V. Parag and Christl A. Donnelly. "Using Information Theory to Optimise Epidemic Models for Real-Time Prediction and Estimation". In: *PLOS Computational Biology* 16.7 (July 2020), e1007990. ISSN: 1553-7358. DOI: 10.1371/journal.pcbi.1007990. (Visited on 04/18/2024).

[26] Jiaping Liu et al. "Rtestim: Time-varying Reproduction Number Estimation with Trend Filtering". In: *PLOS Computational Biology* 20.8 (Aug. 2024), e1012324. ISSN: 1553-7358. DOI: 10.1371/journal.pcbi.1012324.

[27] Tilmann Gneiting and Adrian E Raftery. "Strictly Proper Scoring Rules, Prediction, and Estimation". In: *Journal of the American Statistical Association* 102.477 (Mar. 2007), pp. 359–378. ISSN: 0162-1459. DOI: 10.1198/016214506000001437. (Visited on 09/13/2024).

[28] Ministry of Health NZ. *New Zealand COVID-19 Data.* (Visited on 09/13/2024).

[29] Kris V. Parag, Benjamin J. Cowling, and Christl A. Donnelly. "Deciphering Early-Warning Signals of SARS-CoV-2 Elimination and Resurgence from Limited Data at Multiple Scales". In: *Journal of The Royal Society Interface* 18.185 (Dec. 2021), p. 20210569. ISSN: 1742-5662. DOI: 10.1098/rsif.2021.0569. (Visited on 10/03/2024).

[30] N Ferguson et al. *Report 9: Impact of Non-Pharmaceutical Interventions (NPIs) to Reduce COVID19 Mortality and Healthcare Demand.* Tech. rep. Imperial College London, Mar. 2020. DOI: 10.25561/77482. (Visited on 09/09/2023).

[31] Nicholas Steyn et al. *Technical Report: Update to Modelling 7 September 2021.* Tech. rep. Sept. 2021. URL: https://www.covid19modelling.ac.nz/modelling-the-potential-spread-of-covid-19-during-the-august-2021-outbreak/.

[32] Ministry of Health NZ. *COVID-19 Update 1 September 2021.* https://www.health.govt.nz/news/covid-19-update-1-september-2021. Sept. 2021. (Visited on 09/15/2024).

[33] Ministry of Health NZ. *COVID-19 Update 2 September 2021.* https://www.health.govt.nz/news/covid-19-update-2-september-2021. Sept. 2021. (Visited on 09/15/2024).

[34] Ministry of Health NZ. *COVID-19 Update 13 September 2021 4pm.* https://www.health.govt.nz/news/covid-19-update-13-

september-2021-4pm. Sept. 2021. (Visited on 09/13/2024).

[35] Ministry of Health NZ. *COVID-19 Update 18 October 2021 4pm*. https://www.health.govt.nz/news/covid-19-update-18-october-2021-4pm. Oct. 2021. (Visited on 09/15/2024).

[36] Ministry of Health NZ. *COVID-19 Update 19 October 2021*. https://www.health.govt.nz/news/covid-19-update-19-october-2021. Oct. 2021. (Visited on 09/15/2024).

[37] *Covid Crunch: Auckland Lockdown Cost $8 Billion*. https://www.nzherald.co.nz/nz/covid-19-delta-outbreak-auckland-lockdown-cost-8-billion/AGURSE2ZSR475B4FZ2LBFRKPZI/. Jan. 2022. (Visited on 09/13/2024).

[38] Xian Yang et al. "Bayesian Data Assimilation for Estimating Instantaneous Reproduction Numbers during Epidemics: Applications to COVID-19". In: *PLOS Computational Biology* 18.2 (Feb. 2022), e1009807. ISSN: 1553-7358. DOI: 10.1371/journal.pcbi.1009807.

[39] M. J. Plank et al. *Robust Estimation of End-of-Outbreak Probabilities in the Presence of Delayed and Incomplete Case Reporting*. Sept. 2024. DOI: 10.48550/arXiv.2409.16531. eprint: 2409.16531.

[40] Louis Yat Hin Chan, Baoyin Yuan, and Matteo Convertino. "COVID-19 Non-Pharmaceutical Intervention Portfolio Effectiveness and Risk Communication Predominance". In: *Scientific Reports* 11.1 (May 2021), p. 10605. ISSN: 2045-2322. DOI: 10.1038/s41598-021-88309-1. (Visited on 09/04/2024).

[41] Chigozie A. Ogwara et al. "Impact of Public Health Policy and Mobility Change on Transmission Potential of Severe Acute Respiratory Syndrome Coronavirus 2 in Rhode Island, March 2020 – November 2021". In: *Pathogens and Global Health* 118.1 (Jan. 2024), pp. 65–79. ISSN: 2047-7724. DOI: 10.1080/20477724.2023.2201984. (Visited on 09/04/2024).

[42] Xiaoshuang Liu et al. "Differential Impact of Non-Pharmaceutical Public Health Interventions on COVID-19 Epidemics in the United States". In: *BMC Public Health* 21.1 (May 2021), p. 965. ISSN: 1471-2458. DOI: 10.1186/s12889-021-10950-2. (Visited on 09/04/2024).

[43] Vesna Barros et al. "A Causal Inference Approach for Estimating Effects of Non-Pharmaceutical Interventions during Covid-19 Pandemic". In: *PLOS ONE* 17.9 (Sept. 2022), e0265289. ISSN: 1932-6203. DOI: 10.1371/journal.pone.0265289. (Visited on 09/04/2024).

[44] Sylvia K. Ofori et al. "SARS-CoV-2 Transmission Potential and Rural-Urban Disease Burden Disparities across Alabama, Louisiana, and Mississippi, March 2020 — May 2021". In: *Annals of Epidemiology* 71 (July 2022), pp. 1–8. ISSN: 1047-2797. DOI: 10.1016/j.annepidem.2022.04.006. (Visited on 09/04/2024).

[45] Nina Haug et al. "Ranking the Effectiveness of Worldwide COVID-19 Government Interventions". In: *Nature Human Behaviour* 4.12 (Dec. 2020), pp. 1303–1312. ISSN: 2397-3374. DOI: 10.1038/s41562-020-01009-0. (Visited on 09/04/2024).

[46] Yacong Bo et al. "Effectiveness of Non-Pharmaceutical Interventions on COVID-19 Transmission in 190 Countries from 23 January to 13 April 2020". In: *International Journal of Infectious Diseases* 102 (Jan. 2021), pp. 247–253. ISSN: 1201-9712. DOI: 10.1016/j.ijid.2020.10.066. (Visited on 09/04/2024).

[47] David Rubin et al. "Association of Social Distancing, Population Density, and Temperature With the Instantaneous Reproduction Number of SARS-CoV-2 in Counties Across the United States". In: *JAMA Network Open* 3.7 (July 2020), e2016099. ISSN: 2574-3805. DOI: 10.1001/jamanetworkopen.2020.16099. (Visited on 09/04/2024).

[48] Michelle Kendall et al. "Epidemiological Changes on the Isle of Wight after the Launch of the NHS Test and Trace Programme: A Preliminary Analysis". In: *The Lancet Digital Health* 2.12 (Dec. 2020), e658–e666. ISSN: 2589-7500. DOI: 10.1016/S2589-7500(20)30241-7. (Visited on 09/04/2024).

[49] Rachel E. Baker et al. "Assessing the Influence of Climate on Wintertime SARS-CoV-2 Outbreaks". In: *Nature Communications* 12.1 (Feb. 2021), p. 846. ISSN: 2041-1723. DOI: 10.1038/s41467-021-20991-1. (Visited on 09/04/2024).

[50] Pierre Nouvellet et al. "Reduction in Mobility and COVID-19 Transmission". In: *Nature Communications* 12.1 (Feb. 2021), p. 1090. ISSN: 2041-1723. DOI: 10.1038/s41467-021-21358-2. (Visited on 09/06/2023).

[51] Nikos I. Bosse et al. "Scoring Epidemiological Forecasts on Transformed Scales". In: *PLOS Computational Biology* 19.8 (Aug.

2023), e1011393. ISSN: 1553-7358. DOI: 10 . 1371/journal.pcbi.1011393.

[52] Nikos Bosse et al. *Scoringutils: Utilities for Scoring and Assessing Predictions*. Nov. 2023. (Visited on 10/21/2024).

[53] Kris V. Parag, Benjamin J. Cowling, and Ben C. Lambert. "Angular Reproduction Numbers Improve Estimates of Transmissibility When Disease Generation Times Are Misspecified or Time-Varying". In: *Proceedings of the Royal Society B: Biological Sciences* 290.2007 (Sept. 2023), p. 20231664. DOI: 10.1098/rspb.2023. 1664.

[54] Gavin Freeguard. "The Story of the R Number: How an Obscure Epidemiological Figure Took over Our Lives. Part 4: The Politics of R". In: *Significance* 21.4 (Aug. 2024), pp. 19–22. ISSN: 1740-9705, 1740-9713. DOI: 10.1093/jrssig/ qmae058. (Visited on 09/11/2024).